Thierry, F., Bradley, K., & Warren-Smith, C. (2016). Prevalence of spinous process impingement in thoracic vertebrae on radiographs of clinically-unaffected dogs. *Journal of Small Animal Practice*, 57(12), 698-702. <https://doi.org/10.1111/jsap.12590>

Peer reviewed version

Link to published version (if available):
[10.1111/jsap.12590](https://doi.org/10.1111/jsap.12590)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Wiley at DOI [10.1111/jsap.12590](https://doi.org/10.1111/jsap.12590). Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**PREVALENCE OF SPINOUS PROCESS IMPINGEMENT IN THORACIC VERTEBRAE ON
RADIOGRAPHS OF CLINICALLY UNAFFECTED DOGS**

Running head: Spinous process impingement in dogs

FLORENCE THIERRY, KATE BRADLEY, CHRIS WARREN-SMITH

Corresponding Author

F. Thierry DVM MRCVS

The University of Edinburgh Easter Bush Campus, Small animal hospital, Edinburgh, Midlothian EH25 9RG

fthierry@exseed.ed.ac.uk

07553 199725

K. Bradley MA VetMB PhD DVR DipECVDI MRCVS

The University of Bristol, Radiology Department

Kate.Bradley@bristol.ac.uk

C. Warren-Smith BVetMed Dip ECVDI CertVDI MVetMed MRCVS

The University of Bristol, Radiology Department

chris.warren-smith@bristol.ac.uk

Previous presentation: EAVDI-BID meeting, April 2015, Birmingham, UK.

24 **Abstract**

25

26 **OBJECTIVES:** To assess the prevalence of impinged spinous processes in asymptomatic dogs. These lesions
27 are characterized by a narrowing of the interspinous space, associated with sclerosis and bone remodelling.
28 Such findings have only been reported in three dogs and one cat presenting with back pain. Impinged spinous
29 processes are also occasionally noted on radiographs of healthy dogs with unknown clinical significance.

30

31 **METHODS:** 190 lateral thoracic radiographs of asymptomatic dogs radiographed for reasons other than
32 spinal pain, were retrospectively reviewed by two boarded radiologists. Images were assessed for the presence
33 of impinged spinous processes and graded for the presence of narrowing, sclerosis, or remodelling of the
34 spinous processes.

35

36 **RESULTS:** The prevalence of impinged spinous processes in unaffected dogs was 33.2%. 95% (75/79) of
37 lesions were located between the spinous processes T8 and T11. Impingement of the spinous processes was
38 seen more frequently in older dogs. Size of dog was also related to the lesions, as larger breed dogs displayed
39 more frequent and more severe impingement of the spinous processes compared to smaller breeds.

40

41 **CLINICAL SIGNIFICANCE:** Spinous process impingement appears prevalent in animals with no history
42 of spinal pain, which indicates that this radiographic finding should be interpreted with caution.

43

44 **Keywords:** kissing spine, Baastrup's, spinous process, radiography, dog.

45 INTRODUCTION

46 Impingement of spinous processes is a common radiographic finding in horses and humans. It has been
47 reported in the literature by different names such as “kissing spine” syndrome or Baastrup’s disease and is a
48 common cause of back pain in horses (Jeffcott 1980), and people (Maes *et al.* 2008). Lesions are characterized
49 by a narrowing of the space between adjacent spinous processes associated with sclerosis, flattening or
50 remodelling of the cranial or caudal border of the spinous process. To the authors’ knowledge, radiologic
51 findings of spinous process impingement have only been described in one cat (Gutierrez-Quintana *et al.* 2011)
52 and three dogs (Beythien *et al.* 1994, Ragetly *et al.* 2009), all presenting with back pain. In our experience
53 impinged spinous processes are also occasionally noted on radiographs of dogs presented for reasons
54 unrelated to back pain. The clinical significance of such lesions has never been studied in dogs and no
55 quantitative data exists within the literature. In horses, the reported prevalence of radiographic impinged
56 spinous process lesions in clinically unaffected animals varies from 34% (Jeffcott 1979) to 91.5% (Holmer *et*
57 *al.* 2007) and lesions mainly occur between T13 and T18. The prevalence rises to 86% (Townsend *et al.* 1986)
58 or 92% (Haussler *et al.* 1999) when lesions are diagnosed post-mortem.

59 There have been two reports of canine spinous process impingement occurring either with a concomitant
60 chronic back pain or bilateral iliopsoas contracture. In the first report (Beythien *et al.* 1994) radiographs
61 showed a narrow interspinous space T10-T11 in two dogs, associated with sclerosis and radiolucencies within
62 the spinous process. Treatment consisted of the surgical resection of the spinous process. The second report
63 (Ragetly *et al.* 2009) described impinged spinous processes from T8 to L6. This unusual and wide localization
64 was thought to be secondary to a concurrent iliopsoas contracture and continuous flexion of the hips.

65 The aim of this study was to assess the prevalence of impinged spinous process lesions in a population of
66 asymptomatic dogs, which may help assessing their significance. We hypothesised that narrower interspinous
67 widths would be associated with more severe lesions of sclerosis and bone remodelling.

68

69 MATERIALS AND METHODS

70 Inclusion criteria

71 Canine lateral thoracic radiographs obtained between April 2012 and September 2013 were retrieved from the
72 database of Langford Veterinary Referral Hospital. Images were processed using either a Canon direct digital

radiography system (Xograph) or a Fuji computed radiography system (Fuji Capsula, Fuji Medical). Patient records for all animals were reviewed and animals with a clinical suspicion or a history of neck or back pain were excluded. Breed, age, sex, and reason for presentation were recorded. Dogs were excluded from the study if no history was available in the database. In addition radiographs were only included if all thoracic spinous processes were clearly visible. If several lateral projections were available, the one with the best exposure and least rotation was chosen. Any radiograph with a major overlap of the ribs onto the spinous processes or with major vertebral malformations, such as hemivertebrae or fused vertebrae, was excluded.

Scoring system

Two board-certified radiologists blindly and independently reviewed the radiographs. They were asked to grade from 0 to 3 each spinous process / interspinous space from T1 to T11 according to four criteria: sclerosis, radiolucency, remodelling, and interspinous width. A grade 0 was defined as an absence of sclerosis or radiolucency, no remodelling of the cranial or caudal aspect of the spinous process, and a normal interspinous width. Mild lesions were assigned a grade 1 (Fig.1A) and moderate modifications a grade 2 (Fig.1B, 1C). Severe lesions, such as sclerosis, bone remodelling and narrowed interspinous width, as described in the case report of a German Shepherd dog with bilateral iliopsoas muscle contracture (Ragety *et al.* 2009), were graded as 3. Post scoring, radiographs where there were any differences in scores were reviewed together by the reviewers and a consensus grade was determined.

In order to quantify the severity of the impingement of the spinous processes based on a single rating score, we defined for each pair of adjacent spinous processes a *severity index* computed as the sum of the three grades derived from the sclerosis, remodelling and radiolucency criteria. We chose not to include the interspinous width as part of the severity index, because this single criterion is not specific enough to define an impinged spinous process. Two spinous processes were considered impinged if the severity index was greater than 0.

Measurement method

We applied an additional method to obtain objective measurements of the interspinous widths from T8-T11, using reference lines perpendicular to each spinous process (Fig. 2). All measurements were done by the same

operator to an accuracy of one tenth of a millimetre using a DICOM viewer (Osirix, Geneva, Switzerland) and magnified images. A second set of measurements was performed by the same observer on a 10% sub-sample. These radiographs were chosen randomly, in order to assess the reproducibility of the method. We computed an intra-class correlation coefficient (two-way mixed, absolute agreement, single measures) to assess the reliability of measurements using SPSS 20 software for Macintosh (SPSS Inc, USA).

In order to take into account the size of the dog, the measurements were normalised by using the following method. Breeds were split into three categories according to body size (small, medium and large). To avoid a possible bias of the age, we tested only individuals having reached full body size by excluding from the analysis all individuals less than 1 year-old. We then established a ratio between each measured interspinous width and the mean interspinous width of dogs of same size with no radiographic sign of impinged spinous process.

RESULTS

190 canine thoracic radiographs were assessed; the mean age of dogs was 7 years (range 2 months to 15 years). The study involved 91 males and 99 females. The most common breeds were Labrador and golden retrievers ($N = 37$), cocker and springer spaniels ($N = 36$) and Jack Russell terriers ($N = 14$). 18 brachycephalic dogs such as bulldogs, cavalier King Charles spaniels and boxers met the inclusion criteria.

33.2% of dogs (63/190) displayed signs of spinous process *impingement*. Observers differed on 55/7600 grades (0.007%) on 43/190 radiographs, and each time by only one grade. A mean grade was established in these cases. 98% of the disagreed grades occurred at the T10-T11 space. Of the affected animals, 25.4% of dogs (16/63) had two interspinous spaces involved, which was the maximum found per animal in our study. Only 4 lesions (in 3 dogs) affected spinous processes T1 to T8, while 95% (75 lesions in 61 dogs) occurred at spinous processes T8 to T11 (Fig. 3). The interspinous width was scored as narrowed (i.e. interspinous width score >0.5) in 81 dogs between T8 to T11 and in only 1 dog between T5-T6. Thus, we focused subsequent measurements on the intervertebral spaces from T8 to T11: 6/75 lesions were found on the spinous processes of T8-T9, 39 lesions on T9-10, and 30 on T10-T11.

The width between spinous processes, from T8 to T11, was measured at the narrowest point on all radiographs (Fig. 4). Measurements of the interspinous width were performed twice for 19 radiographs to test

129 the repeatability of the method. The intra-observer agreements were good for the width of the three
130 intervertebral spaces T8-T9 ($ICC = 0.76$, 95% confidence interval 0.37–0.91), T9-T10 ($ICC = 0.93$, 95% *ci*
131 0.80–0.97), and T10-T11 ($ICC = 0.95$, 95% *ci* 0.87–0.98). Mean difference values between these two sets of
132 measurement were 0.6 mm for the interspinous width T8-T9 (range: 0-1.4 mm), 0.3 mm for T9-T10 (range: 0-
133 1 mm), and 0.4 mm for T10-T11 (range: 0-1.2 mm). As shown on Figure 4, the median interspinous width
134 consistently decreased from T8 to T11 in all breeds of dog, regardless of their body size.

135 To assess the relation between the severity of lesions and age and body size, we added up the severity
136 indices from T1 to T11 for each dog to obtain a *total severity index*. The mean age of dogs having
137 impingement of the spinous processes was 8.1 years (range: 10 months-15 years). Among the 63 dogs that had
138 a total severity index equal to or greater than 0.5, 71.4% (45/63) were 6 years old or older. Dogs younger than
139 6 years old with radiographic lesions had a mean total severity index of 2.1 ($N = 18$, range: 0.5-7). The older
140 population of dogs (6 years old or older) with impinged spinous processes had a mean total severity index of
141 2.0 ($N = 45$, range: 0.5-6). The percentage of dogs presenting a spinous process impingement were 22.2%
142 (12/54) for small breeds, 40.3% (25/62) for medium breeds, and 43.8% (25/57) for large breeds. (Note that we
143 excluded all individuals less than 1 year-old in these counts so that all individuals had reached their full body.)
144 There was a trend for mean total severity index to increase with breed size, being 0.3 for small-sized dogs
145 ($N = 54$, range: 0-4), 0.7 for medium-sized dogs ($N = 62$, range: 0-7), and 1.1 for large-sized dogs ($N = 57$,
146 range: 0-6).

147

148 **DISCUSSION**

149 This study showed a prevalence of 33% of impinged spinous process in dogs, between T1 and T11. This
150 prevalence may be underestimated due to the limitations of radiography and could actually be higher if
151 assessed by computed tomography or post-mortem. In horses without back pain, an even higher prevalence of
152 over 80% has been diagnosed at post-mortem (Townsend *et al.* 1986). Such a high prevalence in a population
153 of asymptomatic dogs questions the clinical significance of these findings. Medical histories of dogs were
154 thoroughly checked to exclude all dogs with any suspected neck or back pain. It should be noted, however,
155 that an exhaustive history was not available for all animals. Some symptomatic dogs may have been included
156 in the study population if their clinical signs were missed or not mentioned in the available clinical database.

157 Our results indicate that age could be related to the prevalence of spinous process impingement. Older
158 dogs presented with these lesions more frequently, whereas the severity of the impingement did not appear to
159 be linked to age. In comparison, the effect of age on the increase of radiological lesions remains controversial
160 in horses; a previous study did not find any correlation between the two (Jeffcott 1979), but recent studies
161 have suggested a link (Erichsen *et al.* 2004, Zimmerman *et al.* 2012). In humans the prevalence of Baastrup
162 lesions does increase with age (Kwong *et al.* 2011, Maes *et al.* 2008). In the present study, the severity of
163 radiographic lesions appears to be related to body size, with large breed dogs having more frequent and severe
164 impinged spinous processes compared to smaller breeds.

165 Implementing a reliable method to measure the interspinous width in dogs is problematic. Canine spinous
166 processes have a wide range of shapes, together with various inclinations, even within the same animal.
167 Widths appear more easily measured in horses, since interspinous spaces are quite regular with a narrowest
168 width often at the same level (Berner *et al.* 2012). An interspinous width less than 4 mm is considered
169 narrowed in horses (Erichsen *et al.* 2004, Sinding *et al.* 2010), whereas in dogs, normal interspinous widths at
170 the anticlinal vertebra were typically only 1-2 mm. Difficulties inherent in measuring at this order of
171 magnitude may explain why we did not manage to establish a minimum interspinous width for dogs
172 presenting lesions in this study. One limitation of this study is that measurements were only performed from
173 T8 to T11; however a subjective score evaluating the interspinous width was given for all thoracic spinous
174 processes.

175 Our results show that the interspinous width gets narrower between T8 and T11, which linked to
176 proximity to the anticlinal spinous process. Large breed dogs are more likely to have T11 identified as
177 anticlinal vertebra, and it has been reported to be at T10 in smaller breeds (Baines *et al.* 2009). Knowing that
178 most spinous process impingements were between T9 and T11, it seems logical that the distance between
179 spinous processes plays a major role in the occurrence of lesions. Grading the interspinous width T10-T11
180 was the most controversial for the observers. Indeed, this interspinous space was sometimes less clearly
181 delineated on radiographs than the majority of others, which was likely to be the cause of disagreement by one
182 point for 23% of radiographs.

183 An interesting finding is that no grade reached the maximum score of 3 for any of the criteria (sclerosis,
184 remodelling or radiolucency). All grades were between 0 and 2. We therefore assume that our study did not

185 include dogs with severe impingement of the spinous processes. Hence, we could hypothesise that
186 asymptomatic dogs may have less severe radiological changes than clinically affected animals. That feature
187 was generally observed by Jeffcott (1979) in horses. A further study should assess a group of dogs with spinal
188 pain, to see if these dogs have higher grade lesions.

189 Impinged spinous processes constitute a common finding in horses having back pain, especially among
190 jumping horses. It is believed that when the back is frequently required to extend and flex maximally,
191 particularly when jumping, it results in more severe lesions and possible back pain (Jeffcott 1980, Townsend
192 *et al.* 1986). A positive correlation has been reported between clinical signs and the severity of radiological
193 findings in horses (Zimmerman *et al.* 2012). In humans the pain induced by the impingement of the spinous
194 processes is thought to be mechanical, due to repetitive strain on the interspinous ligament. The degeneration
195 and collapse of the ligaments leads to neoarthrosis and bony erosions between adjacent spinous processes
196 (Mitra *et al.* 2007). The interspinous region forms a bursa with creation of a synovial joint, which will appear
197 on MRI as fluid-like signal between consecutive spinous processes (Maes *et al.* 2008). Similar
198 pseudoarticulation features on macroscopic examination were described in the only MRI report of such
199 lesions in a cat (Gutierrez-Quintana *et al.* 2011). The authors hypothesise that it is likely impingement of the
200 spinous processes results from the same process in dogs, with large breed dogs potentially putting more strain
201 on interspinous ligaments than small breed dogs. Impingement of the spinous processes represents a
202 progressive degenerative process in non-clinical animals, this is why it should not be considered as a disease
203 as such. In rare cases when impinged spinous processes cause pain, an active inflammatory process is likely to
204 trigger the clinical signs. True origin of the pain is still unclear in humans, since these lesions often have
205 concurrent degenerative spinal lesions, and do not always respond well to surgical treatment. Surgical
206 resection of spinous processes seems more successful in horses, with 72% of individuals in one study
207 (Walmsley *et al.* 2002) returning to work.

208 In conclusion, impingement of the spinous processes appears prevalent within this population of
209 asymptomatic dogs, with the changes concentrated in the T8-T11 region and being related to age and body
210 size. The clinical significance of these lesions is questionable and their presence should be interpreted with
211 caution.

References

- Baines, E., Grandage, J., Herrtage M. (2009) Radiographic definition of the anticlinal vertebra in the dog. *Veterinary Radiology & Ultrasound* 50, 69-73.
- Berner, D., Winter, K., Brehm, W., Gerlach, K. (2012) Influence of head and neck position on radiographic measurement of intervertebral distances between thoracic dorsal spinous processes in clinically sound horses. *Equine Veterinary Journal Supplement* 43, 21-26.
- Beythien, R. (1994) Baastrup's-syndrome in dogs – 2 cases – painful degeneration of touching spinous processes of the thoracal spine. *Praktische Tierarzt* 75, 651.
- Erichsen, C., Eksell, P., Holm, KR., *et al.* (2004) Relationship between scintigraphic and radiographic evaluations of spinous processes in the thoracolumbar spine in riding horses without clinical signs of back problems. *Equine Veterinary Journal* 36, 458-465.
- Gutierrez-Quintana, R., Lindley, S., Sullivan, M., *et al.* (2011) Dorsal spinous process impingement syndrome ('kissing spine') in a cat: imaging appearance and surgical management. *Journal of Feline Medicine and Surgery* 13, 618-621.
- Jeffcott, L. B. (1979) Radiographic features of the normal equine thoracolumbar spine. *Veterinary Radiology* 20, 140-147.
- Jeffcott, L. B. (1980) Disorders of the thoracolumbar spine of the horse – a survey of 443 cases. *Equine Veterinary Journal* 12, 197-210.
- Kwong, Y., Rao, N., Latief, K. (2011) MDCT findings in Baastrup disease: disease or normal feature of the aging spine? *American Journal of Roentgenology* 196, 1156-1159.
- Haussler, K.K., Stover, S.M., Willits, N.H. (1999) Pathologic changes in the lumbosacral vertebrae and pelvis in Thoroughbred racehorses. *American Journal of Veterinary Research* 60, 143-153.
- Holmer, M., Wollanke, B., Stadtbaumer, G. (2007) X-ray alterations on spinal processes of 295 warmblood horses without clinical findings. *Pferdeheilkunde* 23, 507-511.
- Maes, R., Morrison, W.B., Parker, L., *et al.* (2008) Lumbar interspinous bursitis (Baastrup disease) in a symptomatic population: prevalence on magnetic resonance imaging. *Spine* 33, E211-215.
- Mitra, R., Ghazi, U., Kirpalani, D., Cheng, I. (2007) Interspinous ligament steroid injections for the management of Baastrup's disease: a case report. *Archives of Physical Medicine and Rehabilitation* 88,

241 1353-1356.

242 Ragetly, G.R., Griffon, D.J., Johnson, A.L., *et al.* (2009) Bilateral iliopsoas muscle contracture and

243 spinous process impingement in a German shepherd dog. *Veterinary Surgery* 38, 946-953.

244 Sinding, M.F., Berg, L.C. (2010) Distances between thoracic spinous processes in Warmblood foals: a

245 radiographic study. *Equine Veterinary Journal* 42, 500-503.

246 Townsend, H.G., Leach, D.H., Doige, C.E., Kirkaldy-Willis, W.H. (1986) Relationship between spinal

247 biomechanics and pathological changes in the equine thoracolumbar spine. *Equine Veterinary Journal* 18,

248 107-112.

249 Walmsley, J.P., Pettersson, H., Winberg, F., McEvoy, F. (2002) Impingement of the dorsal spinous

250 processes in two hundred and fifteen horses: case selection, surgical technique and results. *Equine*

251 *Veterinary Journal* 34, 23-28.

252 Zimmerman, M., Dyson, S., Murray, R. (2012) Close, impinging and overriding spinous processes in the

253 thoracolumbar spine: the relationship between radiological and scintigraphic findings and clinical signs.

254 *Equine Veterinary Journal* 44, 178-184.

FIGURE LEGENDS

FIG. 1. (A) Impinged spinous process T9-T10 in a 6-year old female cocker spaniel (sclerosis: 1, radiolucency: 0, remodelling: 1, width: 1, severity index: 2); (B) Impinged spinous process T9-T10 in a 4-year-old female boxer (sclerosis: 2, radiolucency: 0, remodelling: 2, width: 1, severity index: 4); (C) Impinged spinous process T9-T10 in a 8-year-old male Labrador (sclerosis: 2, radiolucency: 0, remodelling: 2, width: 2, severity index: 4).

FIG. 2. Illustration of the method of measurement of interspinous width, for spaces T8-T9, T9-T10 and T10-T11. A line was drawn from the cranioventral border of the vertebral body to the caudal tip of its dorsal spinous process; a second line was drawn perpendicular (± 0.5 degree) to the first, at the tip of the dorsal spinous process. The interspinous width was determined from a third line (green) parallel to the second, drawn at the narrowest point between the two spinous processes. Measurements were recorded to the nearest one-tenth of a millimetre.

FIG. 3. Severity index for each pair of adjacent dorsal spinous processes from T1 to T11 in 190 dogs. Each dot represents one kissing spine lesion. Samples exceeding four lesions are represented by thick lines.

FIG. 4. Box plots of the interspinous width for spaces T8-T9, T9-T10, T10-T11 for small, medium and large body sizes ($N = 173$, medians, quartiles, range).

T9

T10

T11

T9

T10

T11

T9

T10

T11

T8

T9

T10

T11

2.3mm

2.1mm

1.1mm

RIGHT

Interspinous width (mm)