Paul, S. P., Kirkham, E. N., John, R., Staines, K., & Basude, D. (2016). Coeliac disease in children: an update for general dental practitioners. *British Dental Journal*, 220(9), 481-485.
<https://doi.org/10.1038/sj.bdj.2016.336>

Peer reviewed version

Link to published version (if available):
[10.1038/sj.bdj.2016.336](https://doi.org/10.1038/sj.bdj.2016.336)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Nature Publishing Group at DOI: 10.1038/sj.bdj.2016.336. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Coeliac disease in children – an update for general dental practitioners

Dr Siba Prosad Paul, Specialty Trainee Year 8 in Paediatric Gastroenterology, Bristol Royal Hospital for Children, Bristol

Miss Emily Natasha Kirkham, 5th Year Medical Student, University of Bristol

Dr Rebecca John, Consultant, Senior Lecturer in Paediatric Dentistry, School of Oral & Dental Sciences, University of Bristol.

Dr Konrad Staines, Consultant Senior Lecturer in Oral Medicine, School of Oral & Dental Sciences, University of Bristol.

Dr Dharamveer Basude, Consultant Paediatric Gastroenterologist, Bristol Royal Hospital for Children, Bristol

Word count: 3020

Number of tables: 2

Number of figures: 3

References: 39

Conflict of interest: None

Coeliac disease (CD) is an immune-mediated systemic disorder caused by ingestion of gluten found in wheat, rye and barley. It affects around 1% of children but 90% of cases are considered to remain undiagnosed. CD classically presents with gastrointestinal manifestations including diarrhoea, bloating, weight loss and abdominal pain, but extra-intestinal features (including oral and dental manifestations) are increasingly being reported. Dental and oral manifestations such as dental enamel defects, delayed eruption of teeth, recurrent aphthous ulcers are well recognized manifestations of CD. In patients with yet undiagnosed CD, these can sometimes be the only presenting features. Dentists have regular contact with well children, and therefore the visit to the dentist is an opportunity to suspect CD. When CD is suspected, Dental practitioners can liaise with the General Medical Practitioner to organise screening for coeliac disease. Positive serology will prompt onward referral to a paediatric gastroenterologist to confirm the diagnosis. The recent European Paediatric Gastroenterology Hepatology and Nutrition (ESPGHAN) guidelines have streamlined the diagnostic pathway for faster diagnosis of CD. Management involves strict adherence to a gluten free diet, which should lead to resolution of symptoms, recovery of intestinal mucosa and prevention of long-term complications associated with it. This article aims to describe CD, inform of recent changes to the diagnostic pathway and highlight the dental manifestations of the condition to equip Dental practitioners to aid early diagnosis and initiation of treatment for children with CD.

Introduction and historical perspectives

Coeliac disease (CD) is defined as an immune-mediated systemic disorder elicited by ingestion of gluten and related prolamines in genetically susceptible individuals. CD is characterised by the presence of a variable combination of gluten dependent clinical manifestations, specific antibodies, human leucocyte antigen HLA-DQ2 or HLA-DQ8 haplotypes and enteropathy^{1,2}. Children commonly present with gastrointestinal features such as abdominal pain, bloating, persistent diarrhoea, and weight loss although increasingly extra-intestinal features are being recognised².

The clinical features of CD were first accurately described in 1887 by Samuel Gee. The role of wheat and rye flour in causing CD was established by William Dicke, a Dutch paediatrician in the 1940's³. The concept of a gluten free diet (GFD) as a treatment for CD was devised after observing that an improvement in symptoms occurred in CD patients during the Second World War when there was reduced consumption of wheat flour³. The gluten protein in wheat, barley and rye was subsequently identified as the causative agent for CD³.

The revised European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) guidelines published in 2012 forms the current basis for diagnosing paediatric CD in children in Europe¹.

Oral and dental manifestations of CD are not uncommon in children with known and undiagnosed CD. The dental manifestations include dental enamel defects and delayed eruption of teeth. Children with CD have a greater prevalence of recurrent aphthous ulceration (RAU) which may cause significant discomfort and impede effectiveness of daily oral hygiene measures. Dentists have regular contact with well children, and therefore the visit to the dentist is an opportunity to raise suspicion of CD. Appropriate referral may allow interception of CD in its subclinical form⁴. It is important to remember that children with undiagnosed CD may have no other symptoms than the oral and dental abnormalities⁵. Early referral and diagnosis of CD is important to prevent associated complications.

This review article highlights the importance of considering CD as a multi-organ disorder and the importance of collaborative working between paediatricians, gastroenterologists and other health professionals, particularly dentists.

Epidemiology

Over the last two decades, understanding of CD has changed from an uncommon enteropathy (prevalence of 1 in 2500 to 3000) to a common multi-organ disease with a strong genetic predisposition associated with HLA-DQ2 and HLA-DQ8^{1,6}. The prevalence of CD in children based on a number of paediatric prevalence studies including the confidential Avon Longitudinal Study of Parents and Children (ALSPAC) is around 1%⁶. CD illustrates the 'tip of the iceberg' phenomenon as up to 90% of cases of CD still remain unidentified⁷. This was also evident from the ALSPAC study of 5470 children who underwent serological screening at 7.5 years of age; some of whom subsequently got diagnosed with CD⁸.

The overall prevalence of systemic dental enamel defects in CD patients with mixed or permanent dentition ranges from 9.5% to 95.9% (mean 51.1%); the prevalence in the deciduous dentition is 5.8% to 13.3% (mean 9.6%)⁹. An Italian study which assessed 166 patients between 2 and 17 years of age, reported the prevalence of RAU as 69% in CD patients versus 43% in the control group which was statistically significant. Interestingly, they reported the similar difference in prevalence of caries and dental abnormalities in both groups¹⁰.

Pathophysiology of CD

CD is caused by the ingestion of gluten and related prolamines (such as hordein in barley). The toxic component of gluten is contained in gliadin which traverses through the epithelial barrier of the small intestine during periods of increased intestinal permeability³. In genetically susceptible people, it may result in an inflammatory reaction mediated by the innate and adaptive immune system.

IgA-based anti-tissue transglutaminase antibodies (tTG) are CD specific antibodies and form part of the screening test for CD². Characteristic small intestinal mucosal changes are villous atrophy with crypt hyperplasia and increased intra-epithelial lymphocytes. These changes hamper absorption of nutrients, fat soluble vitamins and minerals².

Clinical presentation

CD is a multi-organ disorder with a variety of manifestations. Children diagnosed with CD may be classified as symptomatic and asymptomatic. Symptomatic children can present with a spectrum of both gastrointestinal and extra-intestinal symptoms from mild to severe.

CD is commonly diagnosed in children by the age of seven years, but can present later in life and in adults as well⁶. CD classically presents in young children with gastrointestinal symptoms including diarrhoea, abdominal pain and weight loss. Other gastrointestinal symptoms include bloating, flatulence, vomiting, steatorrhoea, and occasionally constipation¹¹. Extra-intestinal manifestations include unexplained iron-deficiency anaemia, idiopathic short stature, faltering growth, liver disease, arthropathy, muscle weakness, delayed menarche or delayed onset of puberty and dermatitis herpetiformis^{1,2,7}.

Oral and dental manifestations of CD

Specific dental and oral manifestations of CD are now increasingly being recognised and are highlighted below^{12,13}:

- Dental enamel defects
- Delayed eruption of teeth
- Dental caries
- Recurrent aphthous ulceration (RAU)
- Oral manifestations of dermatitis herpetiformis
- Angular cheilitis
- Atrophic glossitis
- Other oral manifestations

Dental enamel defects (DED) (Fig 2) are one of the most common dental manifestations seen in children¹⁴. The likely mechanism of development of DED is a combination of immune-mediated enamel damage and nutritional disturbance (e.g. hypocalcaemia)¹⁵. In a study of 53 children with confirmed CD (mean age 9.7 yrs) when compared with 28 children (mean age 10 yrs) where CD had been excluded, DED were identified in 29 in CD group against 5 in control group¹⁶. 'Localised' enamel defects (especially in incisors) were found in 20 children with CD and 1 in control group¹⁶. In a Dutch study with 4233 children (median age 6 years and not previously diagnosed with CD), serum samples for tTG-titres were collected along with clinical photographs of clean, moist teeth taken with an intra-oral camera. Children with positive tTG-titres tended to have more DED¹⁷. An Egyptian study with 140 children with DED and a control group of 720 children with no DED underwent serological screening for CD; both groups had no other features of CD. CD was diagnosed in 17.86% with DED in comparison to 0.97% in the control group¹⁸. An Italian study with 603 children with CD, systemic DED was detected in 32.4% (195/603) in comparison to 0.59% in healthy controls (52/6949)¹⁵. Another Italian study with 72 children with CD and 162 normal healthy subjects, DED was much higher in those with CD (20%) as compared to the healthy subjects (6.2%)¹⁹.

de Carvalho et al (2015) analysed the chemical composition of the dental enamel of the primary teeth of patients with CD²⁰. They found the Calcium/Phosphorous ratio to be significantly lower in the teeth of the children affected by CD²⁰. The decrease in Calcium/Phosphorous ratio could be explained by the incorporation of carbonate in the tissue structure, which would increase its solubility²¹.

Fig. 1 Clinical appearance of minor RAU

Fig. 2 - Clinical appearance of DED

The presence of CD when the permanent teeth are developing is a risk factor for DED's¹². These tend to appear symmetrically and chronologically in all 4 quadrants²². Both hypoplasia and hypomineralisation of the enamel can occur. DED may include pitting, grooving and complete loss of enamel. Table 1 highlights DED classification and has been adapted from Aine et al. (1990)²³:

Table 1 – Classification of DED²³	
Grade I	Defects in colour of enamel: single or multiple cream, yellow or brown opacities
Grade II	Slight structural defects: rough enamel surface, horizontal grooves, shallow pits
Grade III	Evident structural defects: deep horizontal grooves, large vertical pits
Grade IV	Severe structural defects: shape of tooth changed

Table 2 Few scenarios that may be encountered by a GDP	
Scenario 1	A child presenting with isolated DED or delayed dentition or RAU or dental caries without any other oral manifestations and no history of GI involvement – referral to GMP not indicated
Scenario 2	Child presenting with isolated DED or delayed dentition or RAU or dental caries with history of chronic GI symptoms – referral to GMP indicated.
Scenario 3	Child presenting with multiple oral manifestations with combinations of DED, delayed dentition, RAU, dental caries etc with no other obvious aetiology – referral to GMP indicated.
Scenario 4	Child presenting with DED, RAU with family history of CD in 1st degree relatives – referral to GMP indicated.
Scenario 5	Child presenting with DED, RAU in high risk group of CD (Fig. 3) – referral to GMP indicated.*
Scenario 6	A child with known CD with any combination of oral manifestations – referral to GMP to explore adherence to GFD.
Scenario 7	Child with any of the associated oral manifestations and presence of skin dermatitis herpetiformis – referral to GMP indicated. *GI involvement includes diarrhoea, abdominal pain, bloating, weight
*GI involvement includes diarrhoea, abdominal pain, bloating, weight loss and faltering growth	

Delayed dental development/eruption greater than 2 standard deviations occurs frequently in children with CD⁴. An Italian study²⁴ involving 120 female children where 60 children had CD and other 60 served as healthy controls, following were the percentages of children with skeletal and dental age delay - 20% in healthy subjects, 56.7% in CD (23% in children with CD with early diagnosis and in 90% with a late diagnosis [aged >8 years]). In children with a late diagnosis of CD, a difference of 11 months was noted between mean chronological age and mean dental age²⁴. In the absence of any local factors or genetic disorders, CD may be considered the commonest systemic disorder associated with this condition.

Dental caries occurs more often in children with CD. In a Turkish study of 64 children (mean age 8.2 years) with CD compared with another 64 age-sex matched control, incidence of caries was higher in CD group at 83% versus 62% in control group²². A similar result with higher incidence of caries was described in another Greek study with 27 children in each arm²⁵. In an Italian study, children (aged 4 – 13 years) with CD (n=300) had higher indexes of caries than healthy subjects (n=300), both in

deciduous teeth (dmft 2.31 ± 1.84 versus 1.42 ± 1.13 ; $p = 0.021$) and permanent teeth (DMFT 2.97 ± 1.74 versus 1.74 ± 1.64 ; $p = 0.0001$)²⁴. However, a different Italian study with 166 children (aged 2 – 17 years) in each arm reported similar prevalence of caries in the two groups (45% versus 45%)¹⁰. Another Italian study²⁶ compared the dental enamel fragment analysis of hypoplastic teeth obtained from 10 children with CD and 10 non-CD children who were treated for dental caries, dental extractions for extensive caries lesions or deciduous teeth exfoliation. The morphological analysis at scanning electron microscopy demonstrated that the enamel hypoplasia of deciduous and permanent teeth in CD is highly hypomineralised with shorter prisms, more irregularly distributed and less interprismatic substance than observed in the non-coeliac children²⁶.

Recurrent aphthous ulceration (RAU) may be associated with systemic predisposing conditions. A Canadian study with 168 children reported that 16% had mouth (aphthous) ulcers prior to diagnosis of CD²⁷. In a Turkish study with 81 children with CD and 20 healthy controls, RAU was present in 48.1% ($n=39$) cases with CD as compared to 5% ($n=1$) in healthy control²⁸. In another Italian study involving 72 children with CD and 162 normal healthy subjects, RAU was slightly higher in those with CD (24/72 [33.3%]) as compared to the healthy subjects (38/162 [23.4%])¹⁹.

Children with CD have a higher prevalence of RAU compared to unaffected children¹⁰. RAU in association with CD is predominantly of the minor variant with the average ulcer size being 5 mm with a non-keratinised mucosal distribution⁵. Clinically indistinguishable ulceration can be associated with conditions other than CD such as immunodeficiency states and Crohn's disease. However, children most commonly develop RAU in the absence of any disease or deficiency (idiopathic). While the exact cause of RAU in CD remains unknown; it may be potentially related to nutrient deficiencies associated with low serum iron, folic acid and vitamin B12.^{5, 12, 29} RAU often regresses with the implementation of a GFD²⁹.

Dermatitis Herpetiformis (DH) is a rare skin condition with clear association with CD. It has an annual reported incidence of 0.56 per 100,000³⁰. The skin lesions typically present as rashes affecting elbows, knees and buttocks³⁰. It is important to note that DH can present with oral mucosal erythema and vesiculobullous lesions which rupture immediately to form shallow painful ulcers. Gingival erythema may manifest as a patchy or diffuse erythematous or desquamative gingivitis. Oral lesions have been reported 6 months prior to onset of skin lesions in DH. Histopathological diagnosis and screening for CD may be useful when suspicion arises³¹⁻³³. DH resolves when the child with CD commences strict GFD.

Other oral/ perioral manifestations such as atrophic glossitis and angular cheilitis, may arise secondary to the development of nutritional deficiencies or anaemia in association with CD. Atrophic glossitis arises secondary to selective depapillation with atrophy of filiform papillae resulting in the clinical appearance of a smooth shiny tongue and has been reported as the only manifestation of CD in published case studies^{34,35}.

Diagnosis

Diagnosis of CD in children is usually made by a paediatric gastroenterologist or paediatrician based on symptoms, signs, serological screening with tTG-titres and typical histological features on small bowel biopsies⁷. Whilst the medical history within this context at primary care may be presented by the parents to their general medical practitioner, it is not unreasonable to expect a general dental practitioner (GDP) to probe into the symptoms during a dental consultation where appropriate. Table 2 highlights a few situations which a GDP may come across and where a referral to the general medical practitioner (GMP) is indicated.

Certain groups of children are at higher risk of developing CD and therefore the dentists needs to be aware of the high risk groups which are listed in Figure 3. Any suspicion of CD in such groups should also be followed up by an appropriate referral.

Fig. 3 High risk group where serological screening for CD is indicated ^{2,7,11}
First degree relatives of CD patient
Type 1 diabetes mellitus
Selective IgA deficiency
Down's, Williams and Turner Syndromes
Autoimmune thyroiditis
Autoimmune liver disease
Unexplained raised transaminases without known liver disease
Dermatitis herpetiformis

The referral to the GMP should prompt serological screening with tTG-titres. A raised tTG-titre warrants onward referral to the paediatric services (paediatrician or paediatric gastroenterologist if direct access available) for confirmation of a diagnosis of CD. The paediatrician will review the history again, plot the height and weight of the child on an age and sex appropriate growth chart and elicit signs of anaemia, jaundice, mouth ulcers, skin rashes, distended abdomen etc.². This will be followed by blood investigations (including a full blood count, liver function tests, urea and electrolytes, blood glucose, vitamin D level, thyroid function), particularly due to the association of CD with other

autoimmune conditions (such as diabetes and hypothyroidism)¹¹. The paediatric gastroenterologist confirms the diagnosis of CD based on serology and endoscopic small bowel mucosal biopsies.

The ESPGHAN guidelines for diagnosing CD in children recommend that in symptomatic children with tTG-titres of >10 times of the upper limit of normal (>10xULN), CD can be diagnosed without need for small bowel biopsies provided they have a positive HLA-DQ2/DQ8 haplotype and a positive anti-endomysial antibody (EMA) result^{1,2}. Small bowel biopsies are still required in all other cases as highlighted below while the child remains on a normal non-restrictive gluten containing diet [initiation of GFD leads to recovery of intestinal mucosa resulting in negative small bowel biopsies]^{1,2,7}:

- Symptomatic children with a positive tTG--titre of <10xULN.
- Symptomatic children with a positive tTG-titre of >10xULN but with a negative HLA DQ2/8 status.
- Asymptomatic children (see Fig 3) with a positive tTG-titre (irrespective of the actual tTG value)

Management

CD is a lifelong condition and time initially spent on detailed counselling and educating the child and family goes a long way. The only treatment currently available is strict adherence to a lifelong GFD. It is important that any nutrient deficiencies (e.g. vitamin-D deficiency, iron-deficiency anaemia) detected at initial diagnosis are corrected^{36,37}. Intestinal mucosa and tTG-titres will return to normal once GFD is initiated, this can however take up to a year depending on the initial level of tTG-titre¹.

Children should continue to be seen by the paediatrician or paediatric gastroenterologist initially at 3 – 6 monthly intervals post diagnosis of CD and then annually². Children are assessed for their growth and development, adherence to GFD and for early detection of other autoimmune associations such as diabetes and hypothyroidism¹¹. Dietitians are invaluable particularly in explaining the role of GFD in managing CD, risk of non-adherence to GFD and avoiding cross contamination. Dietitians ensure that the child is provided with a nutritionally balanced diet and make the family aware of how to cater for a culturally appropriate GFD, suggesting nutritional supplements where necessary and monitor growth when commenced on a GFD.

Families should be encouraged to join the charity Coeliac UK (www.coeliac.org.uk) as they can provide authentic information on CD along with comprehensive food and drink information and also has an expert helpline. GMPs play an important role in supporting the child with CD in the community and prescribe appropriate gluten-free food to maximise adherence to the GFD. GMPs also facilitate serological testing of first degree relatives (even if asymptomatic), after adequate counselling¹¹. The pneumococcal vaccine is now recommended for patients with CD; many children will have received this as part of their routine immunisation program².

The diagnosis of CD should be communicated to the child's dentist as they can play an important role in managing and monitoring children with CD¹². Children who have persistence of DED or RAU post diagnosis of CD, suspicion regarding poor adherence to GFD should be raised and communicated to the child's paediatrician. Dentists may also detect DED, RAU, etc. in first degree relatives or in children from high risk groups (table 2) and a suspicion of CD should be raised and highlighted to the GMP for serological screening¹². Furthermore, patients detected with DED at diagnosis of CD will require regular monitoring by dentists and adherence to GFD is likely to improve the symptoms¹⁸. Dentists also need to be aware of the different compounds that they use when carrying out various dental procedures, such as equipment's, dental fillings or retainers to ensure that they do not contain gluten. A case study highlighted a 9 year old girl with CD who continued to be symptomatic despite strict adherence to GFD and this was later found to be exposure to gluten from her orthodontic retainer that contained a plasticised methacrylate polymer³⁸. Although anecdotal, we have managed children who were asymptomatic at diagnosis of CD and subsequent poor adherence to GFD has led to development of DED and other oro-dental symptoms³⁹

Prognosis

Patients who adhere to a strict GFD, the prognosis of CD are generally good and will not develop further complications. Strict adherence to a GFD allows for healing of the intestinal mucosa. Symptomatic children often feel much better within a few weeks of excluding gluten and are likely to appreciate the advantages of the GFD. In children with ongoing symptoms including new onset dental manifestations³⁹, cross-contamination should be considered. Adherence to a GFD in children diagnosed when asymptomatic often needs more support for long-term adherence.

Missing a diagnosis of CD or non-adherence to GFD can lead to persistence of gastrointestinal symptoms, impaired nutrition, impaired growth and delayed onset or progress of puberty. Long-term risks include osteoporosis and low bone mineral density, increased risk of pathological fractures, bowel cancer, unfavourable pregnancy outcomes, low birth weight in offspring, spontaneous abortion and development of other autoimmune conditions^{2,7}.

Conclusion

CD is a lifelong condition caused by an immune-mediated reaction to the ingestion of gluten. Improved sensitivity and specificity of serological screening and increased awareness of the condition has led to improved identification of the disease. The recent ESPGHAN guideline should streamline the diagnostic pathway and allow faster diagnosis of CD in children. Dentists should be aware of the oral presentation of CD including DED and RAS and make an early referral where

suspicion of CD is raised. GFD is the only management available and strict adherence such a diet will lead to resolution of symptoms and likely complications.

References

1. Husby S, Koletzko S, Korponay-Szabó IR, et al. European Society for Pediatric Gastroenterology, Hepatology, and Nutrition guidelines for the diagnosis of coeliac disease. *J Pediatr Gastroenterol Nutr.* 2012;**54**: 136-60.
2. Murch S, Jenkins H, Auth M, et al. Joint BSPGHAN and Coeliac disease UK guidelines for the diagnosis and management of coeliac disease in children. *Arch Dis Child.* 2013;**98**: 806-11
3. Sandhu B, Onyeador N. Coeliac disease. Textbook of Pediatric Gastroenterology, Hepatology and Nutrition. Jaypee Brothers, 2015, 1st edition, 125-36
4. Condo R, Costacurta M, Maturo P, et al. The dental age in the child with coeliac disease. *Eur J Paediatr Dent.* 2011;**12**: 184-8
5. Baccaglini L, Lalla RV, Bruce AJ, et al. Urban Legends Series: Recurrent Aphthous Stomatitis. *Oral Dis.* 2011; **17**: 755–770.
6. Bingley PJ, Williams AJ, Norcross AJ, et al. Undiagnosed coeliac disease at age seven: population based prospective birth cohort study. *BMJ.* 2004;**328**: 322-3.
7. NICE Guidelines [NG20] (2015). Coeliac disease: recognition, assessment and management National Institute of Health and Care Excellence. Available at <https://www.nice.org.uk/guidance/ng20> (accessed 04/10/15)
8. Ravikumara M, Nootigattu VK, Sandhu BK. Ninety percent of celiac disease is being missed. *J Pediatr Gastroenterol Nutr.* 2007;**45**: 497-9.
9. Pastore L, Carroccio A, Compilato D, et al. Oral manifestations of celiac disease. *J Clin Gastroenterol.* 2008;**42**:224-32
10. Saraceno R, Perugia C, Ventura A, et al. Aphthous, coeliac disease and other dental disorders in childhood. *G Ital Dermatol Venereol.* 2015 Apr 9. [Epub ahead of print]
11. Paul SP, Spray C. Diagnosing coeliac disease in children. *Br J Hosp Med.* 2014;**75**: 268-70.
12. Rashid M, Zarkadas M, Anca A, Limeback H. Oral Manifestations of Celiac Disease: A Clinical Guide for Dentists. *J Can Dent Assoc.* 2011;**77**: 1-6
13. Muggiano F, Quaranta A, Giannantoni I. Major dental clinical pathological manifestation of celiac disease. *WebmedCentral ORAL MEDICINE.* 2013;4:WMC004448
14. Shteyer E, Berson T, Lachmanovitz O, et al. Oral health status and salivary properties in relation to gluten-free diet in children with celiac disease. *J Pediatr Gastroenterol Nutr.* 2013;**57**:49-52.
15. Ventura A, Martelossi S. Dental enamel defects and coeliac disease. *Arch Dis Child.* 1997;**77**:91.
16. Wierink CD, van Diermen DE, Aartman IH, et al. Dental enamel defects in children with coeliac disease. *Int J Paediatr Dent.* 2007;**17**:163-8.
17. Beth S. Specific developmental defects of enamel in children with elevated antitissue transglutaminase concentrations. *J Pediatr Gastroenterol Nutr.* 2015;**60**: 251.
18. El-Hodhod MA, El-Agouza IA, Abdel-Al H, et al. Screening for celiac disease in children with dental enamel defects. *ISRN Pediatr.* 2012;2012:763783. doi: 10.5402/2012/763783.
19. Bucci P, Carile F, Sangianantoni A, et al. Oral aphthous ulcers and dental enamel defects in children with coeliac disease. *Acta Paediatr.* 2006;**95**:203-7.
20. de Carvalho FK, de Queiroz AM, Bezerra da Silva RA, et al. Oral aspects in celiac disease children: clinical and dental enamel chemical evaluation. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;119(6):636-43.
21. De Menezes Oliveira MA, Torres CP, Gomes-Silva JM, et al. Microstructure and

- mineral composition of dental enamel of permanent and deciduous teeth. *Microsc Res Tech*. 2010;**73**:572-7
22. Avsar A, Kalayci AG. The presence and distribution of dental enamel defects and caries in children with celiac disease. *Turk J Pediatr*. 2008;**50**: 45-50
 23. Aine L, Maki M, Collin P, et al. Dental enamel defects in celiac disease. *J Oral Pathol Med*. 1990;**19**: 241-5
 24. Costacurta M, Condò R, Sicuro L, et al. Cervical vertebral maturation and dental age in coeliac patients. *Oral Implantol (Rome)*. 2011;**4**:11-7.
 25. Priovolou CH, Vanderas AP, Papagiannoulis L. A comparative study on the prevalence of enamel defects and dental caries in children and adolescents with and without coeliac disease. *Eur J Paediatr Dent*. 2004;**5**:102-6.
 26. Bossù M, Bartoli A, Orsini G, et al. Enamel hypoplasia in coeliac children: a potential clinical marker of early diagnosis. *Eur J Paediatr Dent*. 2007;**8**:31-7
 27. Rashid M, Cranney A, Zarkadas M, et al. Celiac disease: evaluation of the diagnosis and dietary compliance in Canadian children. *Pediatrics*. 2005;**116**:e754-9.
 28. Ertekin V, Sümbüllü MA, Tosun MS, et al. Oral findings in children with celiac disease. *Turk J Med Sci*. 2012;**42**:613-7
 29. Campisi G, Di Liberto C, Iacono G, et al. Oral pathology in untreated coeliac [corrected] disease. *Aliment Pharmacol Ther* 2007;**26**: 1529–1536.
 30. Salmi TT, Hervonen K, Kautiainen H, Collin P, Reunala T. Prevalence and incidence of dermatitis herpetiformis: a 40-year prospective study from Finland. *Br J Dermatol*. 2011 ;**165**:354-9
 31. Economopoulou P, Laskaris G. Dermatitis herpetiformis: Oral lesions as an early manifestation. *Oral Surg Oral Med Oral Pathol*. 1986;**62**:77-80.
 32. Lahteenoja H, Irjala K, Viander M, et al. Oral mucosa is frequently affected in patients with dermatitis herpetiformis. *Arch Dermatol*. 1998;**134**:756- 8.
 33. Nicolas ME, Krause PK, Gibson LE, et al. Dermatitis herpetiformis. *Int J Dermatol* . 2003;**42**:588-600.
 34. Erriu M, Canargiu F, Orrù G, et al. Idiopathic atrophic glossitis as the only clinical sign for celiac disease diagnosis: a case report. *J Med Case Rep*. 2012;**6**:185. doi: 10.1186/1752-1947-6-185.
 35. De D, Kanwar AJ. Eruptive lichen planus in a child with celiac disease. *Indian J Dermatol Venereol Leprol*. 2008;**74**:164-5
 36. Imam MH, Ghazzawi Y, Murray JA, et al. Is it necessary to assess for fat-soluble vitamin deficiencies in pediatric patients with newly diagnosed celiac disease? *J Pediatr Gastroenterol Nutr*. 2014;**59**:225-8.
 37. Winter S, Halsey C. Should children presenting with iron deficiency anaemia be screened for coeliac disease? *Arch Dis Child*. 2014;**99**:180-2.
 38. Memon Z, Baker SS, Khan A, et al. An orthodontic retainer preventing remission in celiac disease. *Clin Pediatr (Phila)*. 2013;**52**:1034-7.
 39. Paul SP, Harries SL, Basude D. Role of Dentists in Managing Pediatric Celiac Disease. *Indian J Pediatr*. 2016 Feb 18. [Epub ahead of print]