Dettmann, C. P., & Fain, V. (2017). Linear and nonlinear stability of periodic orbits in annular billiards. *Chaos*, 27, [043106].
<https://doi.org/10.1063/1.4979795>

Publisher's PDF, also known as Version of record

Link to published version (if available):
[10.1063/1.4979795](https://doi.org/10.1063/1.4979795)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via AIP at <http://aip.scitation.org/doi/10.1063/1.4979795#> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Linear and nonlinear stability of periodic orbits in annular billiards

Carl P. Dettmann and Vitaly Fain

Citation: *Chaos* **27**, 043106 (2017); doi: 10.1063/1.4979795

View online: <http://dx.doi.org/10.1063/1.4979795>

View Table of Contents: <http://aip.scitation.org/toc/cha/27/4>

Published by the American Institute of Physics

Welcome to a

Smarter Search

with the redesigned
Physics Today Buyer's Guide

Find the tools you're looking for today!

PHYSICS
TODAY

Linear and nonlinear stability of periodic orbits in annular billiards

Carl P. Dettmann^{a)} and Vitaly Fain^{b)}

School of Mathematics, University of Bristol, University Walk, Bristol BS8 1TW, United Kingdom

(Received 29 November 2016; accepted 23 March 2017; published online 11 April 2017)

An annular billiard is a dynamical system in which a particle moves freely in a disk except for elastic collisions with the boundary and also a circular scatterer in the interior of the disk. We investigate the stability properties of some periodic orbits in annular billiards in which the scatterer is touching or close to the boundary. We analytically show that there exist linearly stable periodic orbits of an arbitrary period for scatterers with decreasing radii that are located near the boundary of the disk. As the position of the scatterer moves away from a symmetry line of a periodic orbit, the stability of periodic orbits changes from elliptic to hyperbolic, corresponding to a saddle-center bifurcation. When the scatterer is tangent to the boundary, the periodic orbit is parabolic. We prove that slightly changing the reflection angle of the orbit in the tangential situation leads to the existence of Kolmogorov-Arnold-Moser islands. Thus, we show that there exists a decreasing to zero sequence of open intervals of scatterer radii, along which the billiard table is not ergodic.

Published by AIP Publishing. [<http://dx.doi.org/10.1063/1.4979795>]

A billiard is a dynamical system where a point particle moves with constant velocity inside a domain and experiences elastic collisions with the boundary of the domain. The shape of the boundary determines the dynamics of the billiard. Billiards in a disk on a plane are completely integrable, while annular billiard tables consisting of a particle confined between two nonconcentric disks generically display mixed phase space due to a family of regular orbits that never touch the scatterer. Billiard models find applications in a variety of problems in statistical,¹ classical, and quantum² physics. In this paper, we consider the annular billiard tables formed of a small circular scatterer placed in the interior of a unit circle; this is a popular geometry for microwave billiard experiments.³ Circular boundaries allow us to analytically examine the linear and nonlinear stability of some periodic orbits. Depending on the parameters of the problem, we find that there exist linearly stable orbits of an arbitrarily large period. We show the existence of a saddle-center bifurcation as the parameters vary, corresponding to a change in stability from the linearly elliptic type to the saddle type. Placing the scatterer tangentially to the external circle creates a cusp that is a source of singularities in the billiard. We use the Kolmogorov-Arnold-Moser (KAM) theory to establish that in the cusp case, the periodic orbits are nonlinearly stable.

The mathematical study of billiards was initiated by Birkhoff⁴ and later significantly extended by Sinai⁵ and his followers. Billiards arise in models for various physical phenomena, for example, in statistical mechanics models of hard balls due to Boltzmann.⁶ A billiard in a plane consists of a classical point particle moving with constant velocity in a bounded domain $Q \subset \mathbb{R}^2$, called the billiard table, and obeying the optical reflection law upon collisions with the boundary of the billiard table ∂Q . The shape of the boundary determines the dynamics of the billiard.

It was proved by Birkhoff⁴ that elliptic billiard tables are integrable. A long standing Birkhoff's conjecture, in fact, states that elliptic billiards are the only types of completely integrable strictly convex tables. Recently, it was shown by Avila *et al.*⁷ that this conjecture is true for small perturbations of elliptic billiards with small eccentricity. By using the KAM theory, Lazutkin⁸ proved the existence of a continuum set of caustics near the boundary of strictly convex $C^{5.53}$ boundaries, thus preventing ergodicity. Douady⁹ refined this result to C^6 boundaries. On the other hand, it was shown by Sinai⁵ that concave billiard tables are ergodic and hyperbolic, while later Bunimovich,¹⁰ by using the defocusing mechanism, showed that certain piecewise smooth convex tables (i.e., the stadium) are also hyperbolic and ergodic. It has also been recently conjectured by Bunimovich and Grigo^{11,12} that the presence of absolute focusing components is a requirement for ergodicity.

As noted in Foltin¹³ the method of defocusing requires the circular arcs of the boundary to be disjoint and thus does not apply to strictly convex billiard tables with inner scatterers. It was shown by Foltin¹³ and independently by Chen¹⁴ that the generically, strictly convex C^2 tables with small inner scatterers admit positive topological entropy.

In the class of convex billiards with scatterers, perhaps the simplest geometry is that of an annular billiard, that is, a circle billiard with a smaller inner scatterer. There appears to be a lack of published mathematically rigorous studies of

I. INTRODUCTION AND MAIN RESULTS

Billiards are dynamical systems modelling the motion of a classical particle moving with constant speed inside a bounded domain and performing elastic collisions with the boundary of the domain. They display a whole spectrum of dynamical behaviour ranging from completely integrable to

^{a)}Electronic mail: Carl.Dettmann@bris.ac.uk

^{b)}Electronic mail: vf13950@bristol.ac.uk

billiards of this type. Analytical and numerical methods were used to catalogue some symmetric periodic orbits up to order 6 in annular billiards in the work by Gouesbet *et al.*,¹⁵ while the coexistence of KAM islands and chaotic motions in annular billiards were studied numerically in Saitô *et al.*¹⁶ Recently, the related work by Correia and Zhang¹⁷ demonstrated the existence of stability of some periodic orbits in the so-called *moon* billiards and ergodicity of certain other tables in that class. The linear stability and bifurcations of some periodic orbits in oval and elliptic billiards with an inner scatterer were investigated by da Costa *et al.*¹⁸ Marginally unstable periodic orbits and relation to quantum chaos have been investigated by Altmann *et al.*¹⁹

In this work, we show that there exist certain linearly stable periodic orbits of an arbitrarily large period in a circle billiard with a small interior scatterer. Furthermore, we prove that in the case of the scatterer being tangential to the outer circle, the periodic orbits can be made to be nonlinearly (KAM) stable.

Take a unit disk D in the plane with boundary ∂D . The billiard in D is completely integrable.²⁰ For every positive integer $n \geq 3$, the billiard trajectory with the angle of reflection $\theta = \frac{\pi}{n}$ made with the positively oriented tangent to ∂D is n -periodic, tracing an n -sided regular polygon inscribed in D . The billiard trajectory with reflection angle $\theta = \frac{k\pi}{n}$ where k is an integer, $1 \leq k \leq \frac{n}{2}$, is also n -periodic but traces an n -pointed star polygon inscribed in D if (k, n) are coprime. Let us fix n and k . We obtain the annular billiard table by placing an inner circular scatterer D_R , of small radius $R \ll 1$ in the interior of D , centered on the middle of one of the sides of the polygon. Thus, D_R is normal to the billiard path. Let the boundary of D_R be ∂D_R . Since the circle and the n -polygon are rotationally symmetric, it makes no difference on which side of the polygon D_R is located. It is possible to perturb this configuration in two ways. One may vary R up to some maximum admissible values (to be specified in Section III) to ensure that D_R is in the interior of D , and in the case of star polygonal orbits, to avoid other sides of the same polygon. Another perturbation would be to make small displacements $\delta \geq 0$ of D_R along the side of the polygon away from the centre of the initial position of D_R , as long as D_R stays in the interior of D . Therefore, the maximum value of R depends on n , k , and δ , and we suppress this dependence for clarity of presentation. We will call the corresponding annular billiard table $Q(R)$.

With the scatterer located as described above, we obtain a $(2n+2)$ -period orbit, and we call it a type (a) orbit (see Fig. 1) in the following way. The billiard will undergo n consecutive collisions with ∂D , with the initial angle of reflection made with ∂D chosen to be $\theta_i = \frac{k\pi}{n}$ for $i \in \{0, \dots, n-1\}$. Suppose that D_R is located on the straight line billiard trajectory segment joining the n -th and $(n+1)$ -th collision points. Then, the $(n+1)$ -th collision is perpendicularly on ∂D_R . After collision with ∂D_R , the particle reverses its path, and the $(n+2)$ -th collision is again with ∂D . The particle now performs $(n-1)$ collisions on ∂D again, before colliding with ∂D_R perpendicularly and bouncing back to form a closed orbit of length $(2n+2)$. For the reversed direction of the trajectory, we have $\theta_i = \pi - \frac{k\pi}{n}$ for $i \in \{n+1, \dots, 2n\}$.

FIG. 1. Type (a) orbit with $\theta = \frac{k\pi}{n}$ with $k=1$, $n=4$, and $\delta > 0$, and a type (b) orbit with $\theta = \frac{\pi}{n} + \epsilon$; R_b is as in Eq. (18) below.

Let $\gamma_{a,k}$ denotes the type (a) orbit corresponding to fixed $k \geq 1$ for a given n . We suppress the dependence of the orbit $\gamma_{a,k}$ on the parameters n , R , and δ .

For case $k=1$ and $\delta=0$, one may take a certain maximum R such that ∂D_R is tangent to ∂D (thus forming a cusp). It is known that cusps can be a source of singularities in billiards.²¹ At the present time, cusps created by one focusing boundary and one dispersing boundary have not received much attention in the literature except in the recent work.¹⁸ Prior to that publication, studies were limited to the situation with two dispersing or one dispersing and one flat wall.^{21–23} In the cusp case, R depends on n only, and we obtain a one-parameter family of $(2n+2)$ -periodic orbits for an annular cusp billiard.

We have the following theorem concerning the linear stability of periodic orbits for type (a).

Theorem I. 1. *For any given $n \geq 3$, there exists a billiard table $Q(R)$ such that the orbit $\gamma_{a,k}$ is linearly stable for certain choices of R , $k < 7$, and small $\delta \neq 0$. Specifically, $\gamma_{a,1}$ is linearly stable when $\delta \neq 0$ for R as in Proposition III.2. When (k, n) are coprime, $\gamma_{a,k}$ is linearly stable for $\delta \neq 0$ and $n \geq n_k$ with n_k and R as in Proposition III.3. When $\delta = 0$, $\gamma_{a,k}$ is neutrally stable for all n and k at any admissible R and also when $\delta \neq 0$ for $R = \frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n}$.*

The proof of the Theorem I.1 is in Section III of the paper. Propositions III.2 and III.3 make up Theorem I.1.

We also introduce another type (b) of $(2n+2)$ -periodic orbits, with $n \geq 3$ (also see Fig. 1), by slightly changing the initial reflection angle of the billiard trajectory away from $\frac{\pi}{n}$ by some small $\epsilon > 0$ such that $\theta = \frac{\pi}{n} + \epsilon$ is not π -rational. In this case, the orbit in the D (without D_R) is not periodic, and the polygon traced by the billiard path does not quite close. We position ∂D_R tangentially to ∂D and perpendicularly to the billiard path as before, creating a closed $(2n+2)$ orbit. Thus, type (b) orbits may be created from $\gamma_{a,1}$ by changing the angle of reflection $\frac{\pi}{n}$ slightly. Since the scatterer is tangent to D , the radius is of the scatterer as defined by the choice of n and ϵ and will be specified in Section IV. Denote the radius by R_b and the scatterer by D_{R_b} for this situation. For every $n \geq 3$, the corresponding billiard table is denoted by $Q(R_b)_{n,\epsilon}$. The periodic orbit corresponding to type (b) will be denoted by γ_b . We suppress the dependence of γ_b on n .

For the type (b) configuration, we study the linear and nonlinear (KAM) stability of γ_b . We have the following theorem.

Theorem I.2. *For every fixed $n \geq 3$, there exists an open interval in ϵ such that the γ_b orbit in the billiard table $Q(R_b)_{n,\epsilon}$ is KAM stable, with R depending on ϵ . Therefore, each billiard table in the sequence $Q(R_b)_{n,\epsilon}$ is not ergodic, with R_b decreasing to zero as $n \rightarrow \infty$.*

The proof of Theorem I.2 is given in Section IV. While some heuristic and numerical papers on billiards treat the existence of linearly stable (elliptic) periodic orbits as a sufficient criterion to deduce the existence of elliptic islands (a set of invariant curves of positive measure surrounding the elliptic orbit) and hence the non-ergodicity of the billiard, for a rigorous mathematical investigation of the stability of elliptic orbits, one needs a more delicate analysis. Indeed, “linear ellipticity” is a fragile dynamical property: for example, it is known that in certain two dimensional maps, elliptic fixed points are not surrounded by invariant curves after a small perturbation.²⁴ Thus, one needs to consider the effect of higher order terms to ensure the (local) stability of periodic orbits.

To prove Theorem I.2, we apply the Birkhoff Normal Form with Moser’s Twist Theorem,²⁵ which is a commonly used approach to study the KAM stability in area-preserving maps. This technique has been used for establishing the stability of some periodic orbits in certain billiard systems before. The papers by Kamphorst *et al.*^{26,27} established the stability of 2-periodic orbits in billiards with strictly convex C^5 boundaries, while Donnay²⁸ showed the existence of elliptic islands in generalised Sinai billiards. Rom-Kedar and Turaev²⁹ proved the existence of islands for certain near-ergodic Hamiltonian flows limiting to a billiard flow and also for billiards with steep repelling potentials.³⁰ However, explicit computations with the Birkhoff normal form are not feasible for an arbitrary billiard boundary since a priori one needs to know its details (the form of the billiard map and the location of the periodic orbit). Because we are dealing with circular boundaries, our task is tractable in this regard.

We show that the Birkhoff coefficient³¹ of γ_b periodic orbits is nonzero, which implies the KAM stability, hence showing the nonergodicity of the billiard dynamics.

The paper is organised as follows: In Section II, we review the basic theory of billiards necessary for the study of the linear stability properties of our billiard tables. In Section III, we study the billiard geometry for type (a) periodic orbits and analytically prove Theorem I.1. Section IV is devoted to the study of type (b) orbits. First, we show their linear stability by the same methods as in Section III. Then, by using the KAM theory and Birkhoff normal form, we prove Theorem I.2. The Appendixes provide the derivation of the billiard map required for the computation of the Birkhoff coefficient. The Appendixes also include an auxiliary Lemma C.1 used in the proof of Proposition III.3.

II. PRELIMINARIES

We state some standard facts from the theory of billiards and area-preserving maps. The following information may be found in Chernov²⁰ or in Berry.³²

Let $Q \in \mathbb{R}^2$ be a bounded domain, with C^l -smooth, $l \geq 3$, and boundary ∂Q . We call Q the billiard table. An

orientation of ∂Q is such that Q is to the left on ∂Q . The billiard phase space M consists of the boundary ∂Q and unit velocity vectors $\vec{v}$ pointing inwards of ∂Q . A standard coordinate system on M is (s, θ) , where s is the arc length parameter on ∂Q and $\theta \in (0, \pi)$ is the angle between the positively oriented tangent to ∂Q at the point s and the vector $\vec{v}$. Then, M is the Poincaré section for the billiard flow, and we define the billiard map $B : M \rightarrow M$, $B(s, \theta) = (s_1, \theta_1)$. The billiard map preserves the measure $\sin \theta ds d\theta$ on M , and it is well known that B is area-preserving in the coordinates $(s, \cos \theta)$. Define the signed curvature of ∂Q by $\kappa = \kappa(s)$, such that $\kappa < 0$ for convex boundaries, $\kappa > 0$ for concave boundaries, and $\kappa = 0$ for flat boundaries. Let τ denotes the *flight distance* between two consecutive collision points on the boundary, s and s_1 ; κ is the curvature at s and κ_1 is the curvature at s_1 . Then, the derivative of B at $z = (s, \theta)$ is given by

$$DB(z) = - \begin{pmatrix} \frac{\tau \kappa + \sin \theta}{\sin \theta_1} & \frac{\tau}{\sin \theta_1} \\ \frac{\tau \kappa \kappa_1 + \kappa_1 \sin \theta}{\sin \theta_1} + \kappa & \frac{\tau \kappa_1}{\sin \theta_1} + 1 \end{pmatrix}. \quad (1)$$

To study the linear stability of an n -periodic point $B^n(z_0) = z_0$, where $z_0 = (s_0, \theta_0)$, we need to examine the product of n -matrices of the above type

$$DB^n(z_0) = DB(z_{n-1})DB(z_{n-2}) \dots DB(z_0). \quad (2)$$

The characteristic polynomial of $DB^n(z_0)$ is of the form $\lambda^2 - \lambda \text{tr}(DB^n(z_0)) + 1$, where $\{\lambda, \lambda^{-1}\}$ are eigenvalues of $DB^n(z_0)$. The corresponding periodic point is said to be elliptic and linearly stable if $|\text{tr}(DB^n(z_0))| < 2$, hyperbolic and unstable if $|\text{tr}(DB^n(z_0))| > 2$, and parabolic (neutrally stable) if $|\text{tr}(DB^n(z_0))| = 2$.

III. STABILITY ANALYSIS OF TYPE (A) ORBITS

In this section, we prove Theorem I.1. Consider the billiard table $Q(R)$ as defined in the introduction with the orbit $\gamma_{a,k}$. Define $\delta \geq 0$ to be the parallel displacement of D_R from the midpoint of the billiard trajectory segment and along it. R and δ have to be chosen such that to ensure that D_R stays in the interior of D . Thus, we obtain a (R, δ) -parameter family of $(2n+2)$ periodic orbits for fixed $n \geq 3$ and k (see Fig. 2).

The maximum value of R depends on the choice of δ , k , and n . From geometry, for a fixed $n \geq 3$, $\delta > 0$, and $k = 1$, we have the maximum possible $R = R_\delta$ such that D_R avoids collision with the other parts of the same billiard trajectory

$$R_\delta = 1 - \sqrt{\delta^2 + \cos^2 \frac{\pi}{n}}. \quad (3)$$

For $\delta = 0$, this expression yields R_0

$$R_0 = 1 - \cos \frac{\pi}{n}, \quad (4)$$

which corresponds to ∂D_{R_0} being *tangent* to ∂D , thus forming a cusp.

When $k \neq 1$, with the (k, n) coprime, we have $n \geq 5$, and it is well known²⁰ that the caustics for the orbit with the

FIG. 2. Some periodic orbits $\gamma_{a,k}$; $k=1$, $n=4$, and $\delta=0$ (left); $k=1$, $n=4$, and $\delta \neq 0$ (middle); $k=2$, $n=5$, and $R_{k,0}$ as in Eq. (6) (right).

angle of reflection $\theta = \frac{k\pi}{n}$ in the disk D (with D_R removed) are just inner circles given by the equation

$$x^2 + y^2 = \cos^2 \frac{k\pi}{n}.$$

Thus, the billiard orbit produces a regular star n -gon with the inscribed tangent circle given by

$$x^2 + y^2 = \cos^2 \frac{k\pi}{n}.$$

It is simple to calculate the length of the side of the n -gon as $2 \cos \frac{k\pi}{n} \tan \frac{\pi}{n}$.

For given $n \geq 5$, $\delta \geq 0$, and $k > 1$, we have the maximum possible radius $R = R_{k,\delta}$ for star orbits

$$R_{k,\delta} = \sin \frac{2\pi}{n} \left(\cos \frac{k\pi}{n} \tan \frac{\pi}{n} - \delta \right), \quad k > 1. \quad (5)$$

This yields, for $\delta = 0$,

$$R_{k,0} = 2 \cos \frac{k\pi}{n} \sin^2 \frac{\pi}{n}, \quad k > 1. \quad (6)$$

We note that $R_{k,\delta}$ is such that the other segments of the billiard orbit do not hit D_R .

Define the map B_D to be the composition of the $(n-1)$ iterate of the well-known²⁰ billiard map in a unit disk D

$$B_D : (s, \theta) \mapsto (s + 2(n-1)\theta, \theta). \quad (7)$$

Define B_{in} to be the billiard map that takes the phase point (s, θ) with $s \in \partial D$ to $(\bar{s}, \bar{\theta})$, where $\bar{s} \in \partial D_R$, and define B_{out} to be the map from $(\bar{s}, \bar{\theta})$ to a point $(\bar{s}, \bar{\theta})$ on ∂D again. Thus, we may write the $(2n+2)$ -periodic orbit as a square of the composition of B_D , B_{out} , and B_{in} :

$$(B_{out} \circ B_{in} \circ B_D)^2 = B^{2n+2}.$$

Remark III.1. Note that for linear stability computations, we do not require explicit formulae for B_{in} and B_{out} since we will be using formula (1). However, the explicit forms of B_{in} and B_{out} will be required for the study of nonlinear stability and thus will be provided in Appendix A.

The following Propositions III.2 and III.3 constitute Theorem I.1.

Proposition III.2. Fix $k=1$ and $n \geq 3$. For $\delta=0$ and $R \leq R_0$, $\gamma_{a,1}$ is neutrally stable. For $\delta \neq 0$ and $R < R_\delta$, the stability of $\gamma_{a,1}$ depends on the size of δ and R . In particular, for a small given $\delta > 0$, $\gamma_{a,1}$ is linearly stable for $\frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$

$< R \leq R_\delta$. There is a saddle-center bifurcation at $R = \frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$.

Proof. Consider the billiard geometry type (a), with $k=1$. Fix $n \geq 3$. We have a periodic orbit $\{z_0, \dots, z_{2n+1}\}$, where $z_i = (s_i, \theta_i)$. For $i \in \{0, \dots, n-1, n+1, \dots, 2n\}$, we have $s_i \in \partial D$, and for $i \in \{n, 2n+1\}$, we have $s_i \in \partial D_R$. The initial condition is $z_0 = (s_0, \frac{\pi}{n})$. We will calculate and establish the conditions on the trace of the derivative of the map $B^{2n+2}(z_0)$ that ensures the linear stability.

Assume that D_R is displaced by $\delta \neq 0$ parallel to the orbit's segment in the direction of z_{n-1} . As is well known,²⁰ the flight distance between the two consecutive impact points z_i and z_{i+1} for $i \in \{0, 1, \dots, (n-2), (n+1), \dots, (2n-1)\}$ is $\tau = 2 \sin \frac{\pi}{n}$ since the collisions are on D_R . The flight distance between z_i and z_{i+1} for $i \in \{n-1, n\}$ is $\tau_{R,\delta} = \sin \frac{\pi}{n} - R - \delta$, which corresponds to the length of the billiard trajectory segment between ∂D and ∂D_R . Accordingly, the flight distance for the reverse trajectory between z_i and z_{i+1} , $i \in \{2n, 2n+1\}$ is $\bar{\tau}_{R,\delta} = \sin \frac{\pi}{n} - R + \delta$. The signed curvature of ∂D is -1 , and the curvature of ∂D_R is $\kappa = \frac{1}{R}$.

For $(n-1)$ consecutive bounces along the outer circle, we have the stability matrix

$$DB_D(z_0) = \begin{pmatrix} 1 & -2(n-1) \\ 0 & 1 \end{pmatrix}. \quad (8)$$

For the n -th bounce from ∂D to ∂D_R , the stability matrix is

$$DB_{in}(z_{n-1}) = - \begin{pmatrix} -\tau_{R,\delta} + \sin \frac{\pi}{n} & \tau_{R,\delta} \\ -\tau_{R,\delta}\kappa - 1 + \kappa \sin \frac{\pi}{n} & \tau_{R,\delta}\kappa + 1 \end{pmatrix}. \quad (9)$$

The stability matrix of the billiard map back from ∂D_R to ∂D is

$$DB_{out}(z_n) = - \begin{pmatrix} \frac{\tau_{R,\delta}\kappa + 1}{\sin \frac{\pi}{n}} & \frac{\tau_{R,\delta}}{\sin \frac{\pi}{n}} \\ \frac{-\tau_{R,\delta}\kappa - 1}{\sin \frac{\pi}{n}} + \kappa & \frac{-\tau_{R,\delta}}{\sin \frac{\pi}{n}} + 1 \end{pmatrix}. \quad (10)$$

Similar formulae follow for $DB_D(z_{n+1})$, $DB_{in}(z_{2n-1})$, and $DB_{out}(z_{2n})$. Using expressions (1) and (2), we need to compute $\text{tr}(DB^{2n+2}(z_0))$, which turns out to be

$$\text{tr}(DB^{2n+2}(z_0)) = 2 - \frac{16n\delta^2 \left(nR^2 - R \sin \frac{\pi}{n} - n\delta^2 \right)}{R^2 \sin^2 \frac{\pi}{n}}. \quad (11)$$

Setting $\delta = 0$ shows the parabolic stability of the corresponding orbit for all $R \leq R_0$. Note that when ∂D_R is tangent to ∂D , $\delta = 0$ is necessary since otherwise D_R is no longer in the interior of D . This completes the proof of the first part of the proposition.

Let us consider the case $\delta \neq 0$. Fix n and small enough $\delta > 0$. For linear stability, we need to ensure $|\text{tr}(DB^{2n+2}(z_0))| < 2$. From (11), a necessary condition for the possibility of linearly stable orbits is

$$nR^2 - R \sin \frac{\pi}{n} - n\delta^2 > 0. \quad (12)$$

This yields, upon rejecting the unphysical negative value, $R > \frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$. Thus, R is determined from (3) and (12) by the inequalities (see Fig. 3)

$$\frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n} < R \leq R_\delta. \quad (13)$$

Let us also show that (12) is sufficient. Indeed, for sufficiency, (11) implies that we need $-2 < 2 - \frac{16n\delta^2(nR^2 - R \sin \frac{\pi}{n} - n\delta^2)}{R^2 \sin^2 \frac{\pi}{n}}$, which upon rearranging yields

$$\frac{R^2 \left(\sin^2 \frac{\pi}{n} - 4n^2 \delta^2 \right)}{4n\delta^2} + R \sin \frac{\pi}{n} + n\delta^2 > 0. \quad (14)$$

FIG. 3. Admissible region for the linear stability of $\gamma_{a,1}$. The shaded region shows the range of R defined by (13) with $n = 5, 20$ for which $\gamma_{a,1}$ is linearly stable. Here, $R_{\min} = \frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$ is the bifurcation value, and the stability of $\gamma_{a,1}$ is parabolic on this curve. Below $R_{\min}$, γ_a is hyperbolic. For $n = 5$, $R_{\min} = R_\delta$ at $\delta = 0.11004$ and for $n = 20$, $R_{\min} = R_\delta$ at $\delta = 0.00740$.

Denote $f = \frac{\sin^2 \frac{\pi}{n} - 4n^2 \delta^2}{4n\delta^2}$, the coefficient of R^2 in (14). The formal solutions of (14) are $R \in (-\infty, R^-) \cup (R^+, \infty)$ if $f > 0$ and $R \in (R^+, R^-)$ if $f < 0$, with $R > \frac{-n\delta^2}{\sin \frac{\pi}{n}}$ if $f = 0$. Here, $R^\pm = \frac{-2\delta^2 n}{\sin \frac{\pi}{n} \pm 2\delta n}$. In addition, we require that R satisfies (13).

Consider the case $f > 0$ corresponding to $\delta < \frac{\sin \frac{\pi}{n}}{2n}$. Then, it is clear that $R^\pm < 0$. Since in the billiard context we have $R > 0$, the physically allowed solutions of (14) correspond to R , satisfying (13). Consider now $f < 0$, corresponding to $\delta > \frac{\sin \frac{\pi}{n}}{2n}$. Then, $R^+ < 0 < R^-$, and so physically, the possible domain for R is $0 < R < R^-$. Tedious computations that we suppress show that $R^- > R_\delta$; hence, the admissible values for R lie in the set defined by (13) indeed. The last case $f = 0$ that implies $\delta = \frac{\sin \frac{\pi}{n}}{2n}$ also leads to (13).

Hence, $\gamma_{a,1}$ is linearly stable. Setting $R = \frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$, we see that $\text{tr}(DB^{2n+2}(z_0)) = 2$, and so at this value of R , there is a saddle-center bifurcation, where the stability of $\gamma_{a,1}$ changes from hyperbolic to elliptic. This completes the proof of the proposition. $\square$

Now consider the star polygonal orbits: this is when $k \neq 1$ and (k, n) are coprime. We have the following

Proposition III.3. Let $n \geq 5$ and $k \leq \frac{n}{2}$, (k, n) coprime. For $\delta = 0$ and $R < R_{k,0}$, $\gamma_{a,k}$ is neutrally stable. For $\delta \neq 0$ and $R < R_{k,0}$, the stability of $\gamma_{a,k}$ depends on the relative size of k , R , and δ . In particular, $\gamma_{a,k}$ is hyperbolic for $k \geq 7$. For $k < 7$ and small $\delta \neq 0$, $\gamma_{a,k}$ is linearly stable for $n \geq n_k$. Specifically, $n_2 = 5$, $n_3 = 9$, $n_4 = 13$, $n_5 = 21$, and $n_6 = 53$.

There is a saddle-center bifurcation at $R = \frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n}$.

Proof. We again need to examine $\text{tr}(DB^{2n+2}(z_0))$, where we now have $z_0 = (s_0, \frac{k\pi}{n})$, and the values of τ and κ are modified appropriately to account for $k \neq 1$ orbit configuration. The computations are identical to the above, and so we suppress them and proceed to give the result

$$\text{tr}(DB^{2n+2}(z_0)) = 2 - \frac{16n\delta^2 \left(nR^2 - R \sin \frac{k\pi}{n} - n\delta^2 \right)}{R^2 \sin^2 \frac{k\pi}{n}}. \quad (15)$$

Setting $\delta = 0$, we again see that the corresponding periodic orbit is parabolic. For $\delta \neq 0$, the same analysis as in the paragraph after (14) shows that the condition

$$nR^2 - R \sin \frac{k\pi}{n} - n\delta^2 > 0,$$

is necessary and sufficient for existence of linearly stable orbits. This yields the inequality $\frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n} < R$. From (5), we have $\delta < \cos \frac{k\pi}{n} \tan \frac{\pi}{n}$. The same arguments as the ones following (14) imply that for given k and n with $0 < \delta \leq \frac{\sin \frac{k\pi}{n}}{2n}$, the allowed radius range is

$$\frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n} < R \leq R_{k,\delta}, \quad (16)$$

while further laborious computations which we omit for the case $\frac{\sin \frac{k\pi}{n}}{2n} < \delta < \cos \frac{k\pi}{n} \tan \frac{\pi}{n}$ lead to $\frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n} < R \leq \tilde{R}$, where $\tilde{R}$ is at most $R_{k,\delta}$ (depending on the relative sizes of n and k). Setting $R = \frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n}$, we obtain $\text{tr}(DB^{2n+2}(z_0)) = 2$, and thus, there is a saddle-center bifurcation at this value of R for a given δ .

Let us find the range of k for given n such that (16) is satisfied. Since we may take $\delta > 0$ arbitrarily small, let us examine the limit $\delta \rightarrow 0$ in (16) to facilitate the computation of admissible range of values of k for a given n . Setting $\delta = 0$ in (16) yields

$$\frac{\sin \frac{k\pi}{n}}{n} < R < R_{k,0}.$$

From which we obtain the inequality

$$2n \sin^2 \frac{\pi}{n} > \tan \frac{k\pi}{n}. \quad (17)$$

One needs to choose k and n such that this inequality is satisfied to obtain stable periodic orbits. Let us first examine (17) for large n . Expanding (17) in the Taylor series for $n \rightarrow \infty$ gives the condition $2\pi > k$, i.e., $k \leq 6$ since $k \in \mathbb{Z}^+$.

Let us now determine admissible k values more rigorously by examining (17) for any $n \geq 5$. Using the function f in the Lemma C.1 of Appendix C, we put $k = \lfloor f(n) \rfloor$, and we obtain that (17) is only satisfied for $k \leq 6$ for any n . Numerically we find that $k = 2, n \geq 5$; $k = 3, n \geq 9$; $k = 4, n \geq 13$; $k = 5, n \geq 21$; $k = 6, n \geq 53$. Now $R_{k,0}$ is a decreasing function of k , while $\sin \frac{k\pi}{n}$ is an increasing function of k , and so if the inequality $nR_{k^*,0} - \sin \frac{k^*\pi}{n} > 0$ holds for some k^* , then it holds for all $k \leq k^*$. $\square$

Remark III.4. Note that setting $\delta = 0$ makes $\gamma_{a,1}$ parabolic for all $R < R_0$, and $\gamma_{a,k}$ becomes parabolic for all $R < R_{k,0}$. Geometrically, $\delta = 0$ corresponds to a completely symmetric orbit. When $\delta \neq 0$, the symmetry is lost, and the orbit $\gamma_{a,1}$ is only parabolic for $R = \frac{\sin \frac{\pi}{n} + \sqrt{\sin^2 \frac{\pi}{n} + 4n^2 \delta^2}}{2n}$, while $\gamma_{a,k}$ is only parabolic for $R = \frac{\sin \frac{k\pi}{n} + \sqrt{\sin^2 \frac{k\pi}{n} + 4n^2 \delta^2}}{2n}$; these values of R are precisely the saddle-center bifurcation values given in Propositions III.2 and III.3, respectively.

IV. STABILITY ANALYSIS OF TYPE (B) ORBITS

We have established for the $Q(R)$ table cusp case that $\gamma_{a,1}$ orbits are parabolic. Now for the cusp geometry, it is possible to construct a type (b) $(2n+2)$ -periodic orbit that corresponds to the case when the initial reflection angle on ∂D is not π -rational: $\theta_0 \neq \frac{\pi}{n}$ (see Fig. 1). We denote these orbits γ_b . Again, we create a closed orbit by positioning D_R in the orbit's path perpendicularly, such that the R value for the tangency condition now reads

$$R_b = 1 + \frac{\cos \theta_0}{\cos n\theta_0}. \quad (18)$$

We note that in this case, R_b depends on ϵ , and the billiard table is $Q(R_b)_{n,\epsilon}$. In the Secs. IV A and IV B that follow, we will prove the linear and nonlinear stability of γ_b , thus proving Theorem I.2.

A. Linear stability of γ_b

First, we investigate the linear stability of γ_b . In the general case for $\theta_0 \neq \frac{\pi}{n}$, the expression for DB^{2n+2} is complicated, and so it is difficult to draw any conclusions for the stability of the periodic orbit. Instead, let us pick $\epsilon > 0$ and investigate the limit $\theta_0 = \frac{\pi}{n} + \epsilon$ as $\epsilon \rightarrow 0$.

Proposition IV.1. *There exists $\epsilon^*(n) > 0$ such that for all $\epsilon < \epsilon^*(n)$, γ_b orbits are linearly stable for the initial reflection angle $\theta_0 = \frac{\pi}{n} + \epsilon$.*

Proof. We consider the trace of $(DB_{out}DB_{in}DB_D(z_0))^2$ as before, modifying the values of τ , κ , R , and θ_0 as appropriate. Expanding the trace in the Taylor series in ϵ with the aid of Mathematica, we find

$$\begin{aligned} \text{tr}(DB^{2n+2}(z_0)) &= 2 - \frac{16n\epsilon \left(\cos \frac{\pi}{n} - n \cot \frac{\pi}{n} + n \cos \frac{\pi}{n} \cot \frac{\pi}{n} \right)}{\cos \frac{\pi}{n} - 1} + O(\epsilon^2). \end{aligned} \quad (19)$$

Since $(\cos \frac{\pi}{n} - n \cot \frac{\pi}{n} + n \cos \frac{\pi}{n} \cot \frac{\pi}{n})$ is negative, we may ensure that γ_b is elliptic if we take a small enough positive ϵ . $\square$

Remark IV.2. *The formula (19) implies that for the linear stability of γ_b , one has to take $\epsilon \lesssim \frac{\cos \frac{\pi}{n} - 1}{4n(\cos \frac{\pi}{n} - n \cot \frac{\pi}{n} + n \cos \frac{\pi}{n} \cot \frac{\pi}{n})}$, which implies $\epsilon \simeq \frac{\pi^2}{4n^3(\pi-2)}$ for very large n .*

B. KAM stability of γ_b

To show the KAM stability of γ_b , we make use of the following well-known result regarding the Birkhoff normal form:

Proposition IV.3.^{26,27,33} *Suppose that the map $B^n(s, p)$ is area-preserving and has an n -periodic point at $(0, 0)$. Assume that B^n is C^k with $k \geq 4$. Writing its Taylor expansion up to order 3 in the neighbourhood of $(0, 0)$*

$$\begin{aligned} B^n(s, p) &= \begin{pmatrix} a_{10}s + a_{01}p + a_{20}s^2 + a_{11}sp + \dots + a_{03}p^3 \\ b_{10}s + b_{01}p + b_{20}s^2 + b_{11}sp + \dots + b_{03}p^3 \end{pmatrix} \\ &+ O(|(s, p)|^4). \end{aligned} \quad (20)$$

If the point $(0, 0)$ is elliptic with eigenvalues $\lambda = \exp(\pm i\mu)$ satisfying the nonresonant condition $\lambda^3, \lambda^4 \neq 1$, there is a real-analytic coordinate change that takes the map to its Birkhoff normal form $z \rightarrow \lambda z e^{iA|z|^2} + O(|z|^4)$. The first Birkhoff coefficient A is

$$A = \Im c_{21} + \frac{\sin \mu}{\cos \mu - 1} \left(3|c_{20}|^2 + \frac{2 \cos \mu - 1}{2 \cos \mu + 1} |c_{02}|^2 \right), \quad (21)$$

where

$$\begin{aligned}\Im c_{21} &= \frac{1}{8}a_{10} \left[-a_{12} + 3 \frac{b_{10}a_{03}}{a_{01}} - 3 \frac{a_{01}b_{30}}{b_{10}} + b_{12} \right] \\ &\quad - \frac{1}{8}b_{10} \left[a_{12} - 3 \frac{a_{01}a_{30}}{b_{10}} - \frac{a_{01}b_{21}}{b_{10}} + 3b_{03} \right] \\ |c_{20}|^2 &= \frac{1}{16} \sqrt{\frac{-a_{01}}{b_{10}}} \left[\frac{b_{10}}{a_{01}} a_{02} + a_{20} + b_{11} \right]^2 \\ &\quad + \frac{1}{16} \sqrt{\frac{-b_{10}}{a_{01}}} \left[\frac{a_{01}}{b_{10}} b_{20} + b_{02} + a_{11} \right]^2 \\ |c_{02}|^2 &= \frac{1}{16} \sqrt{\frac{-a_{01}}{b_{10}}} \left[\frac{b_{10}}{a_{01}} a_{02} + a_{20} - b_{11} \right]^2 \\ &\quad + \frac{1}{16} \sqrt{\frac{-b_{10}}{a_{01}}} \left[\frac{a_{01}}{b_{10}} b_{20} + b_{02} - a_{11} \right]^2.\end{aligned}$$

If A is non-zero, the fixed point is nonlinearly stable.³¹

Let us compute A for γ_b . The boundaries ∂D and ∂D_R are analytic except at the *cusp*—which corresponds to the point tangency of ∂D_R to ∂D . However, our periodic orbits are bounded away from the cusp, and so we do not have to deal with this issue. Observe that the billiard geometry is symmetrical about the x -axis, and γ_b is also symmetric with respect to D_R . Define $z_0 = (\pi + \pi/n + \epsilon(1-n), \pi/n + \epsilon)$ to be the fixed point of B^{2n+2} , corresponding to γ_b . We have $B_{out} \circ B_{in} \circ B_D(z_0) = B^{n+1}(z_0)$ and $B^{2n+2}(z_0) = (B^{n+1}(z_0))^2$, with the explicit expressions for B_{out} , B_{in} , and B_D given in [Appendix A](#).

We have the following:

Proposition IV.4. *For every fixed $n \geq 3$ and sufficiently small $\epsilon > 0$, the point $z_0 = (\pi + \pi/n + \epsilon(1-n), \pi/n + \epsilon)$ is KAM stable for B^{2n+2} .*

Proof. We see that z_0 can be moved to the origin via the linear change of coordinates $Z = z - z_0$. Let us define the Reflection map $\mathcal{R}$

$$\mathcal{R}(s, \theta) = (-s, \pi - \theta). \quad (22)$$

It is obvious that $\mathcal{R}$ is a diffeomorphism and $\mathcal{R}^2 = \text{Id}$. Furthermore, it is a reversing involution for $B^{n+1}(z)$ since $\mathcal{R} \circ B = B \circ \mathcal{R}$. Observe that by composing $B_{out} \circ B_{in} \circ B_D(z_0)$ with $\mathcal{R}(s, \theta)$, we obtain z_0 . Thus, z_0 is a fixed point of the map $\mathcal{R} \circ B^{n+1}$. Hence, we have $B^{2n+2} = (\mathcal{R} \circ B^{n+1})^2$,

and the stability of the fixed point of $\mathcal{R} \circ B^{n+1}$ corresponds to the stability of the fixed point of B^{2n+2} .

Let us check the properties of the linearized map $\mathcal{R} \circ B^{n+1}$. In particular, we are interested in the linear stability of z_0 . Let us change coordinates $(s, \theta) \mapsto (s, r) \equiv (s, \cos \theta)$. We remark that $\mathcal{R}$ in terms of (s, r) is $\mathcal{R}(s, r) = (-s, -r)$, and thus, indeed $\mathcal{R} \circ B^{n+1}$ is area-preserving. The tangent map is

$$D(\mathcal{R} \circ B^{n+1}) = \begin{pmatrix} \frac{\partial s}{\partial s_0} & \frac{\partial s}{\partial r_0} \\ \frac{\partial r}{\partial s_0} & \frac{\partial r}{\partial r_0} \end{pmatrix}. \quad (23)$$

The determinant of which is equal to 1. For small enough ϵ , the modulus of the trace of (23) evaluated at z_0 is $|-2 - 2\epsilon \left(\frac{n \cos \frac{\pi}{n}}{\sin \frac{\pi}{n} \sin^2 \frac{\pi}{n}} (n(\cos \frac{\pi}{n} - 1) + \sin \frac{\pi}{n}) \right) + O(\epsilon^2)| < 2$, and hence, z_0 is linearly stable. Let the eigenvalues of (23) be $\lambda_{\pm} = \exp(\pm i\mu) = u \pm iv$. Then,

$$\lambda_{\pm} = \frac{\frac{\partial s}{\partial s_0} + \frac{\partial r}{\partial r_0} \pm i \sqrt{4 - \left(\frac{\partial s}{\partial s_0} + \frac{\partial r}{\partial r_0} \right)^2}}{2},$$

which gives, since $\mu = \arctan \frac{v}{u}$,

$$\mu = -\frac{\sqrt{2}\epsilon}{\sin \frac{\pi}{n}} \sqrt{n \left(n \sin \frac{2\pi}{n} - \left(1 + 2 \cos \frac{\pi}{n} + \cos \frac{2\pi}{n} \right) \right)} + O(\epsilon^{3/2}), \quad (24)$$

and μ tends to 0 as ϵ tends to 0, thus signifying in the limit $\epsilon = 0$, and we have the parabolic stability corresponding to γ_a orbit, as expected.

Using the explicit form of B_{in} , B_{out} , and B_D given in [Appendix A](#), Eqs. (A2), (A3), (A4), and [Appendix B](#) for partial derivatives, we are able to find higher terms in Taylor expansion $\mathcal{R} \circ B^{n+1}$. By plugging the resulting expressions in Mathematica and expanding for small $\epsilon > 0$ and fixed $n \geq 3$, we find that the Birkhoff coefficient A (21), is, to leading order

$$A(n, \epsilon) = \frac{5 \csc^5 \left(\frac{\pi}{2n} \right) \sec^3 \left(\frac{\pi}{2n} \right) \sin^{5/2} \left(\frac{\pi}{n} \right) \left(\cos \left(\frac{\pi}{2n} \right) - n \sin \left(\frac{\pi}{2n} \right) \right)^2}{192 n^2 \epsilon^2 \sqrt{n \sin \left(\frac{2\pi}{n} \right) - \left(1 + 2 \cos \left(\frac{\pi}{n} \right) + \cos \left(\frac{2\pi}{n} \right) \right)}} \sqrt{\frac{\sin \left(\frac{2\pi}{n} \right)}{n \sin \left(\frac{\pi}{n} \right) - \left(1 + \cos \left(\frac{\pi}{n} \right) \right)}} + O(\epsilon^{-1}). \quad (25)$$

It is obvious that $A(1, \epsilon)$ and $A(2, \epsilon)$ are undefined. Numerically, we see that $A(n, \epsilon) \neq 0$ for all $n \geq 3$. Hence, z_0 is KAM stable. $\square$

Proof of Theorem 1.2. The KAM stability of γ_b immediately follows from Proposition IV.4. The existence of an open interval in ϵ for which the table $Q(R_b)_{n,\epsilon}$ has an

elliptic island follows from the form of Birkhoff coefficient (25), as it is clear that by changing ϵ slightly, $A(n, \epsilon)$ stays nonzero. It is obvious that with increasing n , R_b tends to 0. $\square$

Let us examine the coefficient in the limit $n \rightarrow \infty$. Expanding (25) in the Taylor series for $n \rightarrow \infty$ gives

$$A = \frac{5}{24\epsilon^2} \left(\frac{\pi-2}{\pi^2} + \frac{\pi-1}{6n^2} \right) + O\left(\frac{1}{n^4\epsilon^2}\right), \quad (26)$$

i.e., a non-vanishing function of n and ϵ that grows unboundedly as ϵ tends to 0.

Let us fix ϵ and plot $\tilde{A} = \lim_{\epsilon \rightarrow 0} \epsilon^2 A$, ignoring terms of $O(\epsilon^{-1})$ as a function of n for $3 \leq n \leq 1000$ with logarithmic scale for n (see Fig. 4). For $n=3$, we have $\tilde{A} = -\frac{5(-2+\sqrt{3})}{54(-1+\sqrt{3})} \simeq 0.0338912$. As $n \rightarrow 1000$, $\tilde{A} \rightarrow 0.024 \simeq \frac{5(\pi-2)}{24\pi^2}$, as expected from (26). These calculations imply that A is, to leading order, $O(\epsilon^{-2})$ and thus tends to infinity as ϵ decreases and the period n increases.

Remark IV.5. It is known from Grigo's thesis¹¹ that for certain small local perturbations of the scatterer boundary in the normal direction, the elliptic periodic orbit will survive and remain nonlinearly stable.

Remark IV.6. Constraining the radius $R = R_b$ given by (18) enabled us to make explicit computations with the Birkhoff normal form in terms of n and ϵ only. Setting $R < R_b$ while n and ϵ are fixed will correspond to a nontangential position of D_R to D . Similar computations to the ones in Section IV A show that the corresponding orbit will be linearly stable for $\frac{\sin \frac{\pi}{n}}{n} \leq R < R_b$. We expect that Proposition IV.4 will also hold for such R ; however, the Birkhoff coefficient will depend on R and as such, relevant computations would be more laborious. Therefore, we believe that there exists a sequence of non ergodic billiard tables with the scatterers of radius $\frac{\sin \frac{\pi}{n}}{n} \leq R \leq R_b$ for every n .

Remark IV.7. We believe that analogous computations could be used to verify the KAM stability of the $\gamma_{a,k}$ orbits for $\delta \neq 0$ corresponding to billiard tables where the scatterer is not tangent to the boundary. However, we expect that the computations would be much more lengthy since $\delta \neq 0$ implies a loss of symmetry for the billiard orbit and as such the reduced billiard map $\mathcal{R} \circ B^{n+1}$ may not be utilised.

V. SUMMARY AND CONCLUSIONS

We have studied the stability properties of some periodic orbits in a certain case of annular billiards, where the

FIG. 4. A plot of the Birkhoff coefficient for γ_b . $\tilde{A} = \lim_{\epsilon \rightarrow 0} \epsilon^2 A$ with A as in Eq. (26).

radius of the scatterer is very small compared to the external boundary. We have also considered a special limit when the scatterer is just tangent to the outer boundary, forming a cusp. This situation has thus far received relatively little attention, with no published rigorous results concerning the billiard dynamics in the regions formed by such cusps. The advantage of circular boundaries is that they allow one to obtain the explicit formulae for the billiard map and perform direct computations to study the linear and nonlinear stability of periodic orbits. We have established that given any arbitrary $n \geq 3$, the resulting $(2n+2)$ -periodic orbit may be made linearly stable for an appropriate choice of the scatterer radius and small displacement. Further, we have shown that for the cusp geometry, orbits with π -rational reflection angles are neutrally stable. We have found via the application of the KAM theory that for the cusp geometry, orbits with non π -rational angles can be nonlinearly stable. We have also found a neutrally stable configuration of γ_a for a specific value of R for the non-symmetric scatterer position for a given $\delta \neq 0$, which corresponds to a direct parabolic bifurcation.

We note that the circular boundaries significantly simplified our investigation, and the straight forward application of the KAM theory is unlikely to be feasible for other convex billiards with small tangential scatterers. Lazutkin's theorem⁸ implies the non-ergodicity of strictly convex billiards. One might wonder whether this property of a strictly convex billiard other than a circle will be maintained when a small tangential scatterer is introduced. This general problem seems much more difficult due to the curvature of the boundary no longer being constant.

We hope that this work will serve as a motivation for future investigation into billiards with cusps formed by a dispersing and a focusing arc. There is a brief numerical investigation in Ref. 18 into the scaling of the number of collisions for excursions into such a cusp but as yet no published rigorous results.

ACKNOWLEDGMENTS

The authors would like to thank Peter Balint, George Fullman, Kyle Guan, and Dmitry Turaev for useful discussions. Carl Dettmann's research is supported by EPSRC Grant No. EP/N002458/1. Vitaly Fain's research is supported by the University of Bristol Science Faculty Studentship grant.

APPENDIX A: DERIVATION OF B_{in} AND B_{out}

We give some details leading up to the expressions for B_{out} , B_{in} , and B_D that were used in the computation of (21). We remark that similar formulae for the annular billiard map have been derived previously.^{15,16} However, we choose to derive the formulae used in our work since we are focusing on a very specific type of a billiard table with the scatterer tangential to the unit disk. The relevant sketch of the billiard geometry is shown in Fig. 5. We align D_R and D such that their centres fall on the horizontal axis $y=0$, and we position D_R such that ∂D_R is tangent to ∂D since we study the type (b) orbits. Let us parameterize ∂D by φ as

FIG. 5. Sketch of the billiard geometry for derivation of (A3) and (A4). D_R is symmetric on the x-axis.

$$\partial D = \partial D(\varphi) = \{(\cos \varphi, \sin \varphi) : \varphi \in (-\pi, \pi)\},$$

with ∂D traversed anticlockwise. Let us parameterize ∂D_R by γ where γ is as in Fig. 5

$$\partial D_R = \partial D_R(\gamma) = \{(R \cos \gamma - (1 - R), R \sin \gamma) : \gamma \in (0, 2\pi)\}.$$

Let us measure the arc length parameter on ∂D_R clockwise, from the point of tangency of ∂D_R to ∂D . The arc length s in terms of γ for ∂D_R is thus

$$s = \pi + R(\pi - \gamma). \quad (\text{A1})$$

Let us define $\phi = \pi/2 - \theta$, where $\phi \in [-\pi/2, \pi/2]$ is the angle made between the velocity vector of the particle at ∂D_R and the normal $\mathbf{n}$, chosen to be +ve in a clockwise direction, and $\theta \in [0, \pi]$ is the usual angle of reflection made with the +ve tangent vector $\mathbf{t}$ to ∂D_R . Also, define $\alpha = \phi + \pi/2$ as in Fig. 5.

Thus, we have the billiard map as follows: For $(n - 1)$ bounces along the circle, B_D is obtained from (7), i.e.

$$\begin{aligned} s_{n-1} &= s_0 + 2(n - 1)\theta_0 \\ \theta_{n-1} &= \theta_0. \end{aligned} \quad (\text{A2})$$

Now from Fig. 5, we have $\gamma_n = -\pi + \alpha_{n-1} + \theta_{n-1} - \phi_{n+1}$, and using this and (A1), we obtain $B_{in} : (s_{n-1}, \theta_{n-1}) \mapsto (s_n, \theta_n)$

$$\begin{aligned} s_n &= R(2\pi - (\theta_n + \theta_{n-1} + s_{n-1})) + \pi, \\ \theta_n &= \arccos\left(\frac{-\cos \theta_{n-1} - (1 - R)\cos(\theta_{n-1} + s_{n-1})}{R}\right), \end{aligned} \quad (\text{A3})$$

where subscript n denotes the n -th impact which is on ∂D_R , and $(n - 1)$ -th impact is on ∂D , as above. Likewise, the map B_{out} from the scatterer to circle is obtained by reversing time, $B_{out} : (s_n, \theta_n) \mapsto (s_{n+1}, \theta_{n+1})$

$$\begin{aligned} s_{n+1} &= \theta_n + \theta_{n+1} - \frac{s_n - \pi}{R} \\ \theta_{n+1} &= \arccos\left(-R \cos \theta_n - (1 - R)\cos\left(\theta_n - \frac{s_n - \pi}{R}\right)\right). \end{aligned} \quad (\text{A4})$$

Now for γ_b orbits (with $\theta_0 = \pi/n + \epsilon$), we need the particle to collide with ∂D_R perpendicularly, which can be achieved by choosing the initial position on the boundary ∂D to be $s_0 = -\pi + \pi/n + \epsilon(1 - n)$, which corresponds to $\theta_{n-1} = \theta_0 = \pi/n + \epsilon$, giving z_0 . It can be checked (by implicit differentiation and calculating the Jacobian) that the above maps satisfy the symplecticity condition, if we convert to the coordinates $(s, r) \equiv (s, \cos \theta)$.

APPENDIX B: DERIVATIVES OF $\mathcal{R} \circ B^{n+1}$

Let us compute, by the chain rule, the partial derivatives of $\mathcal{R} \circ B^{n+1}$ in coordinates (s, r) , that we use in Section IV B for computation of (21). We have $\mathcal{R} \circ B^{n+1}(s_0, r_0) = (s_{n+2}, r_{n+2})$.

To reduce the typographical clutter, we write $s = s_{n+2}$, $r = r_{n+2}$, $\theta = \theta_{n+2}$. We do not require chain rule to compute $\frac{\partial s}{\partial s_0}$, $\frac{\partial^2 s}{\partial s_0^2}$ and $\frac{\partial^3 s}{\partial s_0^3}$. The other derivatives are

$$\frac{\partial s}{\partial r_0} = -\frac{1}{\sin \theta_0} \frac{\partial s}{\partial \theta_0}; \quad \frac{\partial^2 s}{\partial s_0 \partial r_0} = -\frac{1}{\sin \theta_0} \frac{\partial^2 s}{\partial s_0 \partial \theta_0}, \quad (\text{B1})$$

$$\begin{aligned} \frac{\partial^2 s}{\partial r_0^2} &= \frac{1}{\sin^2 \theta_0} \left(\frac{\partial^2 s}{\partial \theta_0^2} - \frac{\cos \theta_0}{\sin \theta_0} \frac{\partial s}{\partial \theta_0} \right); \\ \frac{\partial^3 s}{\partial s_0^2 \partial r_0} &= -\frac{1}{\sin \theta_0} \frac{\partial^3 s}{\partial s_0^2 \partial \theta_0}, \end{aligned} \quad (\text{B2})$$

$$\begin{aligned} \frac{\partial^3 s}{\partial r_0^3} &= -\left(\frac{1}{\sin^3 \theta_0} + \frac{3 \cos^2 \theta_0}{\sin^5 \theta_0} \right) \frac{\partial s}{\partial \theta_0} + \frac{3 \cos \theta_0}{\sin^4 \theta_0} \frac{\partial^2 s}{\partial \theta_0^2} \\ &\quad - \frac{1}{\sin^3 \theta_0} \frac{\partial^3 s}{\partial \theta_0^3}; \end{aligned}$$

$$\frac{\partial^3 s}{\partial r_0^2 \partial s_0} = \frac{1}{\sin^2 \theta_0} \left(\frac{\partial^3 s}{\partial \theta_0^2 \partial s_0} - \frac{\cos \theta_0}{\sin \theta_0} \frac{\partial^2 s_0}{\partial s_0 \partial \theta_0} \right), \quad (\text{B3})$$

$$\frac{\partial r}{\partial s_0} = -\sin \theta_0 \frac{\partial \theta}{\partial s_0}; \quad \frac{\partial r}{\partial r_0} = \frac{\sin \theta}{\sin \theta_0} \frac{\partial \theta}{\partial \theta_0}, \quad (\text{B4})$$

$$\begin{aligned} \frac{\partial^2 r}{\partial s_0^2} &= -\cos \theta \left(\frac{\partial \theta}{\partial s_0} \right)^2 - \sin \theta \frac{\partial^2 \theta}{\partial s_0^2}; \\ \frac{\partial^2 r}{\partial r_0^2} &= -\frac{\cos \theta}{\sin^2 \theta_0} \left(\frac{\partial \theta}{\partial \theta_0} \right)^2 + \frac{\cos \theta_0 \sin \theta}{\sin^3 \theta_0} \frac{\partial \theta}{\partial \theta_0} - \frac{\sin \theta}{\sin^2 \theta_0} \frac{\partial^2 \theta}{\partial \theta_0^2}, \end{aligned} \quad (\text{B5})$$

$$\begin{aligned} \frac{\partial^2 r}{\partial s_0 \partial r_0} &= \frac{\cos \theta}{\sin \theta_0} \frac{\partial \theta}{\partial s_0} \frac{\partial \theta}{\partial \theta_0} + \frac{\sin \theta}{\sin \theta_0} \frac{\partial^2 \theta}{\partial s_0 \partial \theta_0} \\ \frac{\partial^3 r}{\partial s_0^3} &= \sin \theta \left(\frac{\partial \theta}{\partial s_0} \right)^3 - 3 \cos \theta \frac{\partial \theta}{\partial s_0} \frac{\partial^2 \theta}{\partial s_0^2} - \sin \theta \frac{\partial^3 \theta}{\partial s_0^3}, \end{aligned} \quad (\text{B6})$$

$$\begin{aligned} \frac{\partial^3 r}{\partial s_0^2 \partial r_0} &= \frac{\sin \theta}{\sin \theta_0} \left(\frac{\partial^3 \theta}{\partial s_0^2 \partial \theta_0} - \left(\frac{\partial \theta}{\partial s_0} \right)^2 \frac{\partial \theta}{\partial \theta_0} \right) \\ &\quad + \frac{\cos \theta}{\sin \theta_0} \left(\frac{\partial^2 \theta}{\partial s_0^2} \frac{\partial \theta}{\partial \theta_0} + 2 \frac{\partial \theta}{\partial s_0} \frac{\partial^2 \theta}{\partial s_0 \partial \theta_0} \right), \end{aligned} \quad (\text{B7})$$

$$\begin{aligned} \frac{\partial^3 r}{\partial r_0^2 \partial s_0} &= \frac{-\sin \theta}{\sin^2 \theta_0} \left(\frac{\partial^3 \theta}{\partial \theta_0^2 \partial s_0} - \left(\frac{\partial \theta}{\partial \theta_0} \right)^2 \frac{\partial \theta}{\partial s_0} \right) \\ &\quad + \frac{\cos \theta_0}{\sin^3 \theta_0} \left(\sin \theta \frac{\partial^2 \theta}{\partial s_0 \partial \theta_0} + \cos \theta \frac{\partial \theta}{\partial s_0} \frac{\partial \theta}{\partial \theta_0} \right) \\ &\quad - \frac{\cos \theta}{\sin^2 \theta_0} \left(2 \frac{\partial^2 \theta}{\partial s_0 \partial \theta_0} \frac{\partial \theta}{\partial \theta_0} + \frac{\partial \theta}{\partial s_0} \frac{\partial^2 \theta}{\partial \theta_0^2} \right), \end{aligned} \quad (\text{B8})$$

$$\begin{aligned} \frac{\partial^3 r}{\partial r_0^3} &= \frac{\sin \theta}{\sin^3 \theta_0} \left(- \left(\frac{\partial \theta}{\partial \theta_0} \right)^3 + \frac{\partial^3 \theta}{\partial \theta_0^3} \right) - \frac{3 \cos \theta_0}{\sin^4 \theta_0} \\ &\quad \times \left(\cos \theta \left(\frac{\partial \theta}{\partial \theta_0} \right)^2 + \sin \theta \frac{\partial^2 \theta}{\partial \theta_0^2} \right) + \frac{3 \cos \theta}{\sin^3 \theta_0} \frac{\partial^2 \theta}{\partial \theta_0^2} \frac{\partial \theta}{\partial \theta_0} \\ &\quad + \frac{\sin \theta}{\sin^4 \theta_0} \frac{\partial \theta}{\partial \theta_0} \left(\sin \theta_0 + \frac{3 \cos^2 \theta_0}{\sin \theta_0} \right). \end{aligned} \quad (\text{B9})$$

APPENDIX C: AUXILIARY LEMMA FOR PROPOSITION III.2

Lemma C.1. *The function $f(x) = \frac{x}{\pi} \arctan\left(2x \sin^2 \frac{\pi}{x}\right)$ is strictly increasing on $(1, \infty)$ and $\lim_{x \rightarrow \infty} f(x) = 2\pi$.*

Proof. Let us show that $f(x)$ is strictly increasing on $(1, \infty)$. Let us define the auxiliary function

$$g(y) = (1 + y^2) \arctan y - y.$$

Since $g(0) = 0$ and $g'(y) = 2y \arctan y > 0$ for $y > 0$, $g(y)$ is strictly increasing and positive on $(0, \infty)$. Let

$$y = 2x \sin^2 \frac{\pi}{x}, \quad x > 1.$$

Then, rewriting g , we have

$$\left(1 + 4x^2 \sin^4 \frac{\pi}{x}\right) \arctan\left(2x \sin^2 \frac{\pi}{x}\right) - 2x \sin^2 \frac{\pi}{x} > 0,$$

thus

$$\begin{aligned} &\left(1 + 4x^2 \sin^4 \frac{\pi}{x}\right) \arctan\left(2x \sin^2 \frac{\pi}{x}\right) \\ &\quad + 2x \sin^2 \frac{\pi}{x} > 4x \sin^2 \frac{\pi}{x} > 4\pi \sin \frac{\pi}{x} \cos \frac{\pi}{x}. \end{aligned} \quad (\text{C1})$$

Now observe that

$$f'(x) = \frac{\arctan\left(2x \sin^2 \frac{\pi}{x}\right)}{\pi} + \frac{2x \sin^2 \frac{\pi}{x} - 4\pi \sin \frac{\pi}{x} \cos \frac{\pi}{x}}{\pi \left(1 + 4x^2 \sin^4 \frac{\pi}{x}\right)} > 0,$$

by the above. Hence, $f(x)$ is strictly increasing on $(1, \infty)$.

Since $\arctan x < x$ and $\sin x < x$ for $x > 0$, we have that $f(x)$ is bounded above by 2π and $\lim_{x \rightarrow \infty} f(x) = 2\pi$. $\square$

exceptional point under broken time-reversal invariance," *Phys. Rev. E* **89**, 032909 (2014).

⁴G. Birkhoff, *Dynamical Systems* (American Mathematical Society Colloquium Publications, 1927), Vol. 9.

⁵Ya. G. Sinai, "Dynamical systems with elastic reflections. Ergodic properties of dispersing billiards," *Uspekhi Mat. Nauk* **25:2**(152), 141–192 (1970); [Russian Math. Surveys, **25:2**, 137–189 (1970)].

⁶Y. Sinai, "Development of Krylov's ideas," *Afterword to NS Krylov's "Works on the Foundations of Statistical Physics,"* see reference [K (1979)] (Princeton University Press, 1979).

⁷A. Avila, J. De Simoi, and V. Kaloshin, "An integrable deformation of an ellipse of small eccentricity is an ellipse," preprint [arXiv:1412.2853](https://arxiv.org/abs/1412.2853) (2014).

⁸V. Lazutkin, "The existence of caustics for a billiard problem in a convex domain," *Math. USSR - Izvestiya* **7**, 185–214 (1973).

⁹R. Douady, "Application du théorème des tores invariants," *These 3eme Cycle* (Université Paris VII, 1982).

¹⁰L. Bunimovich, "On ergodic properties of certain billiards," *Funct. Anal. Appl.* **8**, 254–255 (1974).

¹¹A. Grigo, "Billiards and statistical mechanics," Ph.D. thesis (Georgia Institute of Technology, 2009).

¹²L. Bunimovich and A. Grigo, "Focusing components in typical chaotic billiards should be absolutely focusing," *Commun. Math. Phys.* **293**, 127–143 (2010).

¹³C. Foltin, "Billiards with positive topological entropy," *Nonlinearity* **15**, 2053 (2002).

¹⁴Y. Chen, "On topological entropy of billiard tables with small inner scatterers," *Adv. Math.* **224**, 432–460 (2010).

¹⁵G. Gouesbet, S. Meunier-Guttin-Cluzel, and G. Grehan, "Periodic orbits in Hamiltonian chaos of the annular billiard," *Phys. Rev. E* **65**, 016212 (2001).

¹⁶N. Saitô, H. Hirooka, J. Ford, F. Vivaldi, and G. Walker, "Numerical study of billiard motion in an annulus bounded by non-concentric circles," *Physica D* **5**, 273–286 (1982).

¹⁷M. Correia and H. Zhang, "Stability and ergodicity of moon billiards," *Chaos* **25**, 083110 (2015).

¹⁸D. da Costa, C. Dettmann, J. de Oliveira, and E. Leonel, "Dynamics of classical particles in oval or elliptic billiards with a dispersing mechanism," *Chaos* **25**, 033109 (2015).

¹⁹E. Altmann, T. Friedrich, A. Motter, H. Kantz, and A. Richter, "Prevalence of marginally unstable periodic orbits in chaotic billiards," *Phys. Rev. E* **77**, 016205 (2008).

²⁰N. Chernov and R. Markarian, *Chaotic Billiards* (American Mathematical Society, 2006), Vol. 127.

²¹N. Chernov and R. Markarian, "Dispersing billiards with cusps: slow decay of correlations," *Commun. Math. Phys.* **270**, 727–758 (2007).

²²P. Bálint, N. Chernov, and D. Dolgopyat, "Limit theorems for dispersing billiards with cusps," *Commun. Math. Phys.* **308**, 479–510 (2011).

²³P. Bálint and I. Melbourne, "Decay of correlations and invariance principles for dispersing billiards with cusps, and related planar billiard flows," *J. Stat. Phys.* **133**, 435–447 (2008).

²⁴D. V. Anosov and A. B. Katok, "New examples in smooth ergodic theory, Ergodic Diffeomorphisms," *Trudy Mosk. Math. Obs.* **23**, 3–36 (1970), English version in *Trans. Mosc. Math. Soc.*, *AMS* **23**, 1–35 (1972).

²⁵C. Siegel and J. Moser, *Lectures on Celestial Mechanics*, Reprint of the 1971 Edition (Springer Science & Business Media, 2012).

²⁶S. Kamphorst and S. Pinto-de Carvalho, "The first Birkhoff coefficient and the stability of 2-periodic orbits on billiards," *Exp. Math.* **14**, 299–306 (2005).

²⁷M. Carneiro, S. Kamphorst, and S. De Carvalho, "Elliptic islands in strictly convex billiards," *Ergodic Theory Dyn. Syst.* **23**, 799–812 (2003).

²⁸V. Donnay, "Elliptic islands in generalized Sinai billiards," *Ergodic Theory Dyn. Syst.* **16**, 975–1010 (1996).

²⁹D. Turaev and V. Rom-Kedar, "Elliptic islands appearing in near-ergodic flows," *Nonlinearity* **11**, 575 (1998).

³⁰V. Rom-Kedar and D. Turaev, "Big islands in dispersing billiard-like potentials," *Physica D* **130**, 187–210 (1999).

³¹J. Moser, *Stable and Random Motions in Dynamical Systems: With Special Emphasis on Celestial Mechanics* (Princeton University Press, 2001), Vol. 1.

³²M. Berry, "Regularity and chaos in classical mechanics, illustrated by three deformations of a circular 'billiard,'" *Eur. J. Phys.* **2**, 91 (1981).

³³A. Grigo, "Billiards and statistical mechanics," Ph.D. thesis (Citeseer, 2009).

¹C. P. Dettmann, "Diffusion in the Lorentz gas," *Commun. Theor. Phys.* **62**, 521 (2014).

²F. Haake, *Quantum Signatures of Chaos* (Springer Science & Business Media, 2013), Vol. 54.

³S. Bittner, B. Dietz, H. Harney, M. Miski-Oglu, A. Richter, and F. Schäfer, "Scattering experiments with microwave billiards at an