Matlova, V., Roffet-Salque, M., Pavlu, I., Kyselka, J., Sedlarova, I., Filip, V., & Evershed, R. (2017). Defining pottery use and animal management at the Neolithic site of Bylany (Czech Republic). *Journal of Archaeological Science: Reports*, 14, 262-274.
<https://doi.org/10.1016/j.jasrep.2017.05.028>

Peer reviewed version

Link to published version (if available):
[10.1016/j.jasrep.2017.05.028](https://doi.org/10.1016/j.jasrep.2017.05.028)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Elsevier at DOI: 10.1016/j.jasrep.2017.05.028. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Defining pottery use and animal management at the Neolithic site of Bylany (Czech Republic)

Author names: Matlova Veronika ^{a,*}, Roffet-Salque Mélanie ^b, Pavlu Ivan ^c, Kyselka Jan ^a, Sedlarova Ivona ^d, Filip Vladimir ^a, Evershed Richard P. ^b

^a Department of Dairy, Fat and Cosmetics, University of Chemistry and Technology Prague, Technická 3, 166 28 Prague 6, Czech Republic

^b Organic Geochemistry Unit, School of Chemistry, University of Bristol, Cantock's Close Bristol BS8 1TS, United Kingdom

^c Institute of Archaeology of the Academy of Sciences of the Czech Republic, Prague, V.V.I. Letenská 4, 118 01 Prague 1, Czech Republic

^d Department of Inorganic Technology, University of Chemistry and Technology Prague, Technická 5, 166 28 Prague 6, Czech Republic

Corresponding author

Brychova Veronika, MSc.

Department of Dairy, Fat and Cosmetics

University of Chemistry and Technology Prague

Technická 3

166 28 Prague 6

Czech Republic

Phone number: +420 220 443 266

E-mail: matlovav@vscht.cz

Abstract

Archaeological potsherds have become a valuable source of information about diet and the wider economies of ancient communities, especially through the analysis of lipids preserved in the microporous matrix of the ceramic vessels. This study investigated >160 potsherds recovered from settlement phase 19 dated to 5160-5100 cal. BC from the Neolithic site of Bylany, one of the largest *Linearbandkeramik* (LBK) settlement in Central Europe. The aim was to investigate vessel use and animal management at the site and explore variations in organic residue composition and thus human activity at the household level. Pottery technology was also studied revealing a predominance of micro- and mesopores, indicating an advanced level of pottery production technology. More than 70% of the analysed potsherds yielded appreciable amounts of lipids dominated by C_{16:0} and C_{18:0} fatty acids, with compound-specific

carbon isotope compositions indicating origins predominantly from ruminant and non-ruminant animal fats. Detection of very long fatty acids, fatty alcohols and traces of terpene compounds originating from plants suggested a combination of meat- and plant-based diet components and specialised use of some vessels. However, evidence of the use of vessels for milk collection or processing was not detectable at Bylany, at least during the settlement phase investigated herein.

Highlights

- Lipids were extracted from late LBK sherds from Bylany (Czech Republic).
- Findings were interpreted in relation to pottery typology and households.
- No difference in food processing practices between house types could be identified.
- The processing of ruminant and non-ruminant carcass products was confirmed.
- The lack of dairy fats pointed at the absence of milk exploitation.

Key words: LBK pottery, organic residue analysis, gas chromatography, fatty acids, stable carbon isotope analyses, vessel use, porosity.

1. Introduction

The Neolithic period saw major changes in the way food and natural resources were used. It is well-known that the early farmers cultivated crops and bred livestock, although many of the details of plant agriculture and animal management are yet to be elucidated (Pavlu and Zapotocka 2007). Neolithic settlements of the central European Linear Pottery culture (*Linearbandkeramik*, LBK) consist of small, middle or long houses with thatched roofs, supported by rows of poles (Coudart, 1998). It was thought that the size of the houses corresponded to the status of their inhabitants (e.g. Modderman, 1986; van de Velde, 1990), although recent research proposed that the three basic types of houses correspond to different household activities and roles within the settlement (Hachem, 2000; Gomart et al., 2015). The different house sizes may reflect the size of animal herds, the proportion of hunted animal species and/or the type and volume of cultivated or gathered crops. Notwithstanding, variable local environmental conditions which not have been particularly suitable for stable subsistence strategies (Pavlu, 2014b), some large LBK settlements persisted more than 400 years. Social groups with different economies would have coexisted responding to the fluctuations and pressures associated with the beginnings of Neolithic agriculture (Pavlu, 1987, 2014b).

The well-described site of Bylany (Kutna Hora, Czech Republic; see Fig. 1) is one of the largest central European Neolithic settlements of the LBK and following STK (*Stichbandkeramik*) cultures, comprising more than 100 house-plans. The settlement area was discovered in 1950s and over 7 ha of settlement remains were uncovered during excavations. The LBK period alone included 25 settlement phases (resp. ceramic phases), chronologically classified between 5350 and 4900 cal. BC (Podborsky, 1997; Pavlu and Zapotocka, 2007). Phase 19 (examined here) is dated to 5160-5100 cal. BC and falls within the 5th interval of the LBK settlement (late LBK, phases 18-20). Phase 19 of the settlement exhibited a complicated house development

79 consisting of small ($n = 3$), middle ($n = 4$) and long ($n = 3$) houses, with many associated clay
80 pits (i.e. large pits used as a source of clay) and grain pits, covering areas A and B of the site
81 (Fig. 2). A pit containing ceramic artefacts was found alongside each house. The pottery
82 assemblage of Bylany is very large, comprising >76,000 classified fragments of vessels
83 (Kvetina and Koncelova, 2012), allowing functional classification of the vessels based on
84 ethnoarchaeological markers (Varien and Mills, 1997), primarily the shape and rim diameter of
85 the vessels (Fig. 5; Rice, 2006). The archaeozoological assemblage recovered at the site is
86 particularly scarce, with only 1.6 kg of poorly preserved animal remains having been discovered
87 during the 40-year long excavations (Pavlu, 2014a). Archaeobotanical remains at Bylany are
88 lacking for environmental and historical reasons. Furthermore, no evidence of burial sites have
89 been detected. The type and position of the houses, pits and trenches, macrolithic tools and
90 pottery are thus the only source of information that have been examined so far regarding
91 household economies and other human activities at the Neolithic site of Bylany.

92
93 The increasing range of biomolecular methods used in archaeology are proving particularly
94 effective for investigating the diet and subsistence economies of Neolithic communities
95 (Evershed et al., 1992). In particular, organic residue analysis of pottery vessels (e. g. Evershed,
96 2008b) has been successfully applied widely, allowing various levels of information to be
97 revealed, ranging from vessel use and technological innovation (Roffet-Salque et al., 2013) to
98 specialised animal management strategies (Copley et al., 2003; Evershed et al., 2008) and the
99 exploitation of wild resources (Cramp et al., 2014a, b; Craig et al., 2015; Roffet-Salque et al.,
100 2016). Organic residues accumulated during vessel use, mainly of lipophilic origin persist for
101 millennia absorbed into the ceramic fabric of ancient pottery vessels (e. g. Evershed et al., 2002;
102 Regert et al., 2003). The presence of characteristic lipidic biomarkers, including saturated fatty
103 acids, triacylglycerols and a range of fatty acyl derivatives, waxes, long-chain ketones and

triterpenoid components, allow the type and origin of the residue to be assessed (for reviews see Evershed, 1993, 2008b; Regert et al., 2003; Mukherjee et al., 2005). Traces of animal fats, plant oils, beeswax, resins, tars, pitches, etc. have been identified in archaeological pottery vessels. A combination of chromatographic techniques (GC/FID) and mass spectrometric techniques (GC/MS) is used to separate and identify the compounds. Animal fat residues occur widely, however, the molecular composition of degraded animal fats alone does not allow fat type to be identified and thus compound-specific stable carbon isotopic analyses with GC-C-IRMS (gas chromatography-combustion-isotope ratio mass spectrometry) are carried out to enable ruminant adipose, ruminant dairy or non-ruminant adipose fats to be distinguished (e. g. Dudd and Evershed, 1998; Dudd et al., 1999; Copley et al., 2003). The possibility also exists of identifying milk residues in pottery vessels, which has opened up new avenues of investigation on the beginnings of dairying across Europe and the Near East (e.g. Evershed et al., 2008). Evidence for dairying has been revealed dating as early as the 7th millennium BC in the Near East and lipid residue analyses of potsherds from the 6th LBK have shown that some Central European LBK communities were processing milk into cheese using cheese-strainers (Salque et al., 2013).

Notwithstanding the latter findings, extensive work is required to identify the temporal and spatial patterning of milk use in Europe order to understand the lactase persistence allele amongst the first farmers of Central Europe (Itan et al., 2009). As a contribution to this endeavour, herein, we focus on the organic residue analysis of pottery vessels from the LBK site of Bylany (phase 19) in order to assess: (i) pottery vessel use, and (ii) animal management. The investigation also aims to explore variations in pottery vessel use at household and site levels. Further, given the differences observed in lipid preservation, porosimetry was used in an attempt to explore the mechanisms of preservation of lipids. The focus of this study was

phase 19 from the Bylany excavation, chosen as sherds of all functional categories (Pavlu, 2000) were recovered and, overall, the sherds were less-fragmented than in other phases, suggesting a simpler taphonomic history for this assemblage, which legitimises the comparisons presented (Pavlu, 2010).

2. Material and Methods

2.1. Selection of pottery sherds

A total of 1,842 rim potsherds were excavated from phase 19, from which 1,539 sherds were classified into 14 categories according to their presumed function (Fig. 5; Pavlu, 2010). Only 842 potsherds (46%) could be simultaneously categorized using one of 14 classes and associated to a house type. From this set, a subset of 163 upper rim sherds (20%) were sampled and submitted to organic residue analysis. The potsherds originated from presumed water storage/processing vessels ($n = 60$; categories F6, F9 - F11, F13), from food processing and serving vessels ($n = 88$; categories F1-F5, F7, F8, F12) and from vessels for storage of dry commodities ($n = 15$, category F14). The potsherds were sampled from pits alongside 9 different households, with 74 potsherds originating from long houses (houses 96, 162, 1246), 49 from middle houses (houses 361, 619, 702) and 40 from small houses (houses 959, 1161, 1240; see Table I).

Porosimetry studies were carried out on 7 sherds from Bylany phase 19, of which 3 (categories F9, F10, F13) were presumed to be water storage/processing vessels and 4 (categories F4, F7, F12) were presumed to be food processing and serving vessels. A portion (0.1-0.2 g) of each potsherd was sampled and subjected to mercury porosimetry using an AutoPore IV 9500 V1.06 instrument. Each sherd was placed in the porosimeter, evacuated, and porosity determined using

pressure ranges 0. 0003-0. 01 MPa for macropores, 0. 13-200 MPa for mesopores. The pressure was gradually increased simultaneously while the volume of mercury entering the pore of the sherds was recorded.

2.2. Lipid extraction of potsherds

All solvents used for lipid analyses were HPLC grade and all the glassware was furnace at 450 °C for a minimum of 4 h. The surface of a sub-sample of the archaeological potsherd was cleaned with a manual modelling drill to remove exogenous lipids (from the soil and post-excavation handling). The portion of 2-3 g of potsherd was then crushed and ground in a glass mortar using a pestle to obtain a fine powder, which was accurately weighed and 20 µg of internal standard (*n*-tetratriacontane, Supelco Analytical, Bellefonte, USA) added. Lipids were extracted via the direct methanolysis method described in Correa-Ascencio and Evershed (2014). The method combines hydrolysis and transesterification reactions to obtain fatty acid methyl esters (FAMES) from triacylglycerols and their derivatives simultaneously during the extraction of potsherds. Aliquots of total lipid extracts (TLEs) were taken and free hydroxyl groups trimethylsilylated by treatment with *N*, *O*-bis(trimethylsilyl)trifluoroacetamide (BSTFA, 20 µL, 70 °C, 1 h) to obtain trimethylsilylated (TMS) derivatives prior to GC analyses. The excess BSTFA was evaporated under a gentle stream of nitrogen and methylated/trimethylsilylated extracts dissolved in hexane for analyses by GC/FID, GC/MS and GC-C-IRMS.

2.3. GC analyses

GC/FID analyses were performed on an Agilent Technologies 6890N gas chromatograph equipped with a (5%-phenyl)methylpolysiloxane coated fused silica capillary column (Agilent 19091S-433 HP-5MS; 30 m x 0.32 mm i.d., 0.25 µm film thickness). One microlitre of

177 methylated/trimethylsilylated extract dissolved in hexane was introduced using a split/splitless
178 injector at 220 °C. The temperature of GC oven was programmed from 120 °C for 5 min, to
179 175 °C at 5 °C min⁻¹, followed by an isothermal hold for 25 min, then to 300 °C at 7 °C min⁻¹,
180 followed by a third isothermal hold for 25 min. Helium was used as carrier gas with a constant
181 flow of 1 mL min⁻¹. The GC/MS analyses were performed with the same temperature program
182 on a GC system Agilent Technologies 7890A with a 5975C VL MSD detector. The GC/MS
183 system was equipped with the same column as the GC-FID system.

185 Compound-specific stable carbon isotope analyses were performed using an Agilent Industries
186 7890A gas chromatograph coupled to an IsoPrime 100 isotope ratio mass spectrometer. One
187 microlitre of methylated/trimethylsilylated portions of extract dissolved in hexane were
188 introduced via a split/splitless injector operated in the splitless mode onto a 50 m x 0.32 i.d.
189 fused silica capillary column coated with a 100% dimethylpolysiloxane stationary phase
190 (Agilent HP 1; 0.17 µm film thickness,). The GC oven temperature programme was held at 40
191 °C for 2 min, followed by a gradient increase to 300 °C at 10 °C min⁻¹, after which the oven
192 was held isothermally for 10 min. Helium was used as carrier gas at a constant flow of
193 2 mL min⁻¹. The combustion reactor consisted of a quartz tube filled with copper oxide pellets
194 maintained at a temperature of 850 °C. Data processing was carried out using the Ion Vantage
195 software (version 1.5.6.0, Isoprime).

196 **3. Results**

197 *3.1. Lipid recovery*

198 More than 70% of potsherds from the 163 analysed yielded >5 µg g⁻¹ of (µg of lipid per gram
199 of potsherd) TLE, while 18% yielded >100 µg g⁻¹. This high recovery rate is comparable to that
200 observed at the LBK site of Kopydłowo (Poland; Roffet-Salque and Evershed, 2015).

201

202 3.2. *Lipid compositions*

203 Most of the TLEs were dominated by palmitic (C_{16:0}) and stearic (C_{18:0}) acids. Odd-numbered
204 and branched-chain fatty acids (C_{17:0} and C_{17:0br}), biomarkers of bacterial population from the
205 rumen and characteristic of ruminant fats (Keeney et al., 1962), were detected in 23% of the
206 extracts. The relatively high abundance of the C_{18:0} fatty acid compared to the C_{16:0} fatty acid
207 suggests that these lipids derive from animal fats (Copley et al., 2001). Low concentrations of
208 oleic acid (C_{18:1}) and its degradation products (9,10 - dihydroxyoctadecanoic acid and azelaic
209 acid) were detected in most of extracts. The presence of unsaturated fatty acids at high
210 concentration is often considered as arising from modern contamination due to the lability of
211 the double bond in oxidative conditions. However, considering that oleic acid can be found in
212 animal fat triacylglycerols at high concentration (in the case of modern reference porcine or
213 ruminant fats it ranges from 31 to 44%; Gunstone, 2007; Velisek, 2013), its occurrence at low
214 concentration in well-preserved archaeological pottery is thus possible. The presence of its
215 degradation products (Table 2) also points towards altered (archaeological) animal fats.

216

217 Myristic (C_{14:0}) and arachidic (C_{20:0}) acids were also detected. Longer branched fatty acids (C₁₅
218 to C₁₈) detected in 23% of the sherds could originate from microbial flora of the rumen and
219 originate from domestic ruminant adipose or venison fats (e.g. Duncan and Garton, 1978;
220 Velisek, 2013). The presence of mid-chain ketones (C₃₁ and C₃₅) in a single sherd from a pot
221 of type F13, indicated that the pot was heated at high temperatures, leading to the pyrolysis of
222 acyl lipids and their ketonic decarboxylation (Evershed et al., 1995; Raven et al., 1997).

223

224 Nearly 23% of the samples with the appreciable amount of lipids had significant concentrations
225 of long-chain fatty acids and fatty alcohols, such as behenic (C_{22:0}), lignoceric (C_{24:0}) or cerotic

(C_{26:0}) fatty acids (Fig. 3) and hexa- and octacosanol. These compounds derived mainly from plant tissues and plant waxes and suggest a combination of meat- and plant-based foodstuffs in some of the vessels (20 of 88 samples). Plant lipid residues were present in every type of vessels, in bowls, dishes and jars, and could have been used as flavouring (Filipović and Tasić, 2012) or waterproofing agents (Heron et al., 1994; Roffet-Salque et al., 2016). In 13 extracts *n*-alkanes (C₁₆ to C₂₉), resinous compounds and dicarboxylic acids were also detected (see Fig. 4b). Shorter *n*-alkanes (C₁₆-C₁₉) might have arisen in the potsherds through pyrolysis (Eckmeier and Wiesenberg, 2009; Schellekens et al., 2013), longer *n*-alkanes (C₂₈, C₂₉) could originate from waxes of higher plants (Gunstone, 2007). Resinous diterpenic (abietic acid derivatives) and triterpenic (betulin and friedelin) compounds were detected in small jars (type F13) from the long house 162 and small house 959 (Fig. 4b) providing evidence for the presence of tar adhesives possibly originating from a birch bark tar in case of betulin (e.g. Urem-Kotsou et al., 2002; Grünberg, 2002; Regert et al., 2003) or from beach and oak barks in case of friedelin (Chandler and Hooper, 1979; Urem-Kotsou et al., 2002; Prost et al., 2011) or from altered pine resin in case of abietic acid derivatives (Regert, 2004).

3.3. Stable carbon isotope compositions of fatty acids

A total of 38 total lipid extracts identified as pure animal fats and with an appreciable concentration of lipids (>10 µg g⁻¹) were analysed to determine the carbon isotopic composition of the C_{16:0} and C_{18:0} fatty acids and identify the source of the animal fats. The δ¹³C_{16:0} values of archaeological animal fats ranged between -27.9 and -23.8‰, while δ¹³C_{18:0} values range between -30.6 and -22.9‰ (Fig. 8a). These δ¹³C values are in agreement with pure fats and mixtures of carcass fats from non-ruminant and ruminant animals raised on C₃ diets (Copley et al., 2003). The Δ¹³C (= δ¹³C_{18:0} - δ¹³C_{16:0}) proxy was used in order to identify the fat types by emphasising the influence of animal metabolism (Evershed et al., 1999; Copley et al., 2003).

The archaeological animal fats extracted from the pots from Bylany exhibit $\Delta^{13}\text{C}$ values ranging between -3.1 and 1.4‰ (Fig. 8b), consistent with pure non-ruminant adipose fats ($n = 14$) or mixtures of ruminant adipose (carcass) fats and non-ruminant adipose fats ($n = 24$; Table 3). No dairy residues were detected in the extracts.

3.4. Porosity

The mercury porosimetry analyses revealed the presence of mostly mesopores ($10^{-7}\text{m} > \text{diameter} > 2 \cdot 10^{-9}\text{ m}$) and micropores (diameter $< 2 \cdot 10^{-9}\text{ m}$). The total pore surface of potsherds ranged between 4 and 11 $\text{m}^2\text{ g}^{-1}$, the average radius of pores ranged between 0.02 and 0.06 μm and the mean porosity was 23%.

4. Discussion

4.1. Analysis of pottery and lipid preservation

The presence of mesopores ($10^{-7}\text{ m} > \text{diameter} > 2 \cdot 10^{-9}\text{ m}$) and micropores (diameter $< 2 \cdot 10^{-9}\text{ m}$) in pots from the site of Bylany (phase 19) detected by porosimetry analyses agrees with the hypothesis that pots were fired in open kilns with firing temperatures ranging between 700-800 °C (Pavlu and Zapotocka, 2007). Indeed, these temperatures are sufficient for creating hard microporous fabric and vessels with ‘suitable’ porosity. The well-developed technology of pottery manufacturing at Bylany would have had an influence on the everyday activities at the settlement, e.g. by decreasing the amount of fuel needed for heating cooking vessels. High porosity allows liquids to permeate easily through the vessel walls, extending the time for liquid contents to boil by cooling the outer surface of the vessel wall. On the contrary, low vessel wall porosity and reduction of wall thickness (mean thickness value of Bylany potsherds without counting of storage vessels of type F14 was 0.8 cm) increases thermal

shock resistance during repeated heating (Gosselain, 1992; Tite et al., 2001; Nelson, 2010) and heat conduction (Braun, 1983) while it decreases heat loss (Schiffer, 1990). Those properties had to be controlled also by a type of tempers and clays (Tite et al., 2001). Postfiring treatments can be applied to decrease permeability (Rice, 2006) and repetitive cooking of plant and animal tissues has been shown to seal vessel walls, improving the heat transfer during cooking (Charters and Evershed, 1997; Evershed, 2008b).

The basic assumption is that lipophilic compounds are capable of binding into submicron pores of ceramic fabric due, the hydrophobicity, hence, insoluble properties of lipids, the presence of carboxylic and hydroxy moieties enhancing their propensity to bind to the polar ceramic fabric. However, the question of enhanced physico-chemisorption of lipids was recognised nearly twenty years ago, with the use of a caustic methanol extractant follow $\text{CHCl}_3/\text{MeOH}$ extraction was effective in revealing highly functionalised lipids produced via oxidation of unsaturated fatty acids (Regert et al., 1998). Building on this approach the new extraction method recently proposed by Correa-Ascencio and Evershed (2014) uses an acidified solution of methanol to extract chemisorbed compounds, which could not be extracted using the commonly used organic solvents (chloroform, dichloromethane and methanol). As with the caustic methanol extraction method, the new protocol results in higher extraction yields than a solvent extraction with a mixture of chloroform/methanol (Evershed et al., 1990), suggesting that some lipids are strongly bound into ceramic pores or on surfaces, e.g. to metal ions such as Ca^{2+} , Fe^{3+} , Al^{3+} (as salts) or to SiOH (via hydrogen bonding), which create an inner ceramic lattice. In the sherds from Bylany, it appears that well-shaped micro- or mesopores protect adsorbed lipids from microbial utilization, as microbial flora cannot utilize unreachable substrates (mean size of bacteria $>10^{-6}$ m). Degradation of lipids in the clay walls of potsherds is thus only driven by

outer environmental conditions (humidity, air access, temperature, redox condition etc.; Evershed, 2008b).

Some of the sherds from Bylany contained >1.5 mg of lipids per gram. While, such concentrations of lipid are not uncommon, the highest lipid concentrations are observed in potsherds excavated from arid (Dunne et al., 2012) or acidic (Smyth and Evershed, 2015) burial environments. Within a site higher concentration of lipids in some vessels could reflect a frequency of use of certain vessels (Smyth and Evershed, 2015). The burial conditions at Bylany are neutral or slightly acidic, and the loess soil is considered reasonable water-permeable (Pye, 1995), but clearly these conditions lead to favourable preservation of lipids in potsherds. Not surprisingly, however, these same conditions cause extensive decalcification of bones at Bylany, dissolving the vast majority archaeozoological and human skeletal remains.

4.2. Concentrations of lipid residues in different vessel types

The function of the pottery vessels comparing the Bylany assemblage has been assessed in detail using typological analysis, correlating the shape of vessels to their potential use, with reference to ethnoarchaeological studies (Varien and Mills, 1997). Three basic shapes have been identified in the ceramic assemblage, namely: bowls, dishes and jars, and their volumes reconstructed based on orifice diameters and rim angles (Pavlu, 2000). This ethnoarchaeological approach has allowed the vessels to be classified into several categories of presumed use, e.g. cooking vessels, storage vessels or vessels for storing water, and a range of respective subcategories (Pavlu, 2000; Fig. 5). All the functional categories exhibited high recovery rate of lipids (above 60%), except sherds from the type F2 (31%, 4 residues extracted from 13 potsherds). The highest recoveries rates of lipids are observed for small dishes of types F1 and F3 (90% and 88%, respectively) but the median concentration of total lipid extract was <50 $\mu\text{g g}^{-1}$.

324

325 The typological set 1 (F6, F9, F10, F11 and F13) was interpreted as having been used for water
326 storage and handling. Lipids were recovered from >75% of the sherds from this vessel category
327 (45 residues extracted of 60 potsherds, one-sample χ^2 test, $p < 0.01$), which is the highest
328 recovery rate of lipids detected in functional sets from Bylany. Lipids detected in those sherds
329 were identified as being animal fats ($n = 38$; 23% of examined potsherds), with traces of plant
330 waxes ($n = 25$; 15%; Fig. 8-9). Some of the vessel could have been used for water storage, with
331 the animal fats present resulting from post-firing waterproofing (e.g. Evershed et al., 1997;
332 Skibo, 2013). However, more than 11% of sherds contained >500 μg of lipids per gram,
333 suggesting that animal products were processed in these vessels. The presence of mid-chain
334 ketones in jar F13 (Fig. 4a), provides compelling evidence for the heating of animal fats at high
335 temperatures (Evershed et al., 1995; Raven et al., 1997) that could have occurred when animal
336 fats were spread on the inner surface of the pots for waterproofing just after the firing or when
337 animal products were processed in those large pots.

338

339 The typological vessel set 5 (F14) are large vessels proposed to be storage pots, as their
340 substantial size would have prevented them being easily manoeuvred. Moreover, substantial
341 thickness of their vessel wall (mean 1.6 cm) and toughness of the fired clay would have been
342 important properties of storage vessels (Tite et al., 2001). Although lipids were recovered from
343 15 sherds of this category (73%; one-sample χ^2 test, $p < 0.01$), the concentrations were very
344 low (average 11 $\mu\text{g g}^{-1}$), which is entirely consistent with the hypothesis that the F14 pots were
345 used for storage of dry goods or water.

346

347 Based on typological assessments liquid and solid food would have been served in pots from
348 sets 3 and 4, respectively. However, cold or hot contacts of foodstuffs lead in both cases to lipid

adsorption, although concentrations may differ. Odd-carbon number mid-chain ketones, usually interpreted as demonstrating that pots were heated at high temperature (Raven et al., 1997) were not detected in these potsherds. Although it is not possible to determine whether the foodstuffs contained in these pots were hot or not, the hypothesis remains that such vessels were used as tableware suitable for serving (e.g. Urem-Kotsou and Kotsakis, 2007). Significantly, the potsherds from set 3 exhibited relatively high concentration of lipids (average $135 \mu\text{g g}^{-1}$), which were skewed by high lipid concentrations in the potsherds from vessel types F3, F5 and F7 (Fig. 6a). This contrasted with the lower mean lipid concentrations in potsherds of set 4 (average TLE concentration: $39 \mu\text{g g}^{-1}$) implying these pots were used either less frequently or for processing different foodstuffs.

Finally, the vessels from the set 2 (F4 and F12) were hypothesised to be the most commonly used ware for cooking, food processing and serving. Repetitive use of pots for cooking foodstuffs leads to the accumulation of lipids in the clay walls (Evershed, 2008a). Lipids were recovered from 71% of potsherds from this set (17 residues were extracted from 24 potsherds, one-sample χ^2 test, $p < 0.01$). Moreover, the mean lipid concentrations determined for the sherds from pots F4 and F12 were 121 and $152 \mu\text{g g}^{-1}$, respectively, which are relatively high compared to sherds from other sets except those from sets 1 or 2 (Fig. 6b). Significantly, the lipids detected in 8 potsherds from this set were identified as being animal fats (Table II) indicating that the original pots were used for food processing, likely cooking.

No evidence was obtained from the analyses the specialisation in the use of pottery for processing specific types of foodstuff (Fig. 8), except for small bowls ($n = 4$, analysed categories F7, F9 and F11) where only pure non-ruminant fats were detected. In contrast to the inferences based on typological assessments, the lipid concentrations (Fig. 6) and compositions

suggests no detectable hierarchy existed in vessel use at Bylany (Pavlu, 2014a; Fig. 5). The inference is that either: (i) lipid analyses lack the resolution to reveal specialisation, or (ii) that vessels were highly utilitarian and used for a wide range of purposes. The high concentrations of lipids in some vessels possibly reflect the repetitive use of vessels for cooking, as well as the use of vessels for longer time periods. However, in any discussion of concentrations of lipid observed we must be mindful of the taphonomic history of the sherds. Notwithstanding the latter, both repeated use and extended use-life will result in lipids being accumulated in appreciable concentrations in vessel walls. In contrast, vessels with lower recovery rates and low concentrations of lipids were either: (i) not used for cooking or processing of fat-rich foodstuffs, or (ii) had shortened use-lives due to early breakage of the vessels.

Consideration of the lipid recovery rates and concentrations in different vessel types in the context of households (Fig. 7, Tab. 1), shows highest lipid concentrations were exhibited by the vessels from middle house 702 and small house 959 (recovery rates 67% and 100%, respectively). Relatively constant recovery rates were observed for the different house types, with rates being 70%, 76 % and 73% for small, middle and long houses, respectively (one-sample χ^2 test, $p < 0.04$; Table 1). Sherds from long houses thus present slightly lower recovery rates than sherds from middle and small houses. However, this observation is biased as large and small bowls, which are presumed to be cooking pots, represented just over 24% (18/74 sherds) of the potsherds sampled from long houses.

4.3. Stable carbon isotope analysis of lipid residues and animal management

The faunal assemblage at Bylany is relatively scarce, with only 1.6 kg of poorly preserved bone and fragments recovered during the entire excavation. Archaeozoological remains of cattle (*Bos taurus*), aurochs (*Bos primigenius*), pigs (*Sus domesticus*), goats (*Capra aegargus*), sheep (*Ovis*

399 *orientalis*), wild boar (*Sus scrofa*), wild horses (*Equus ferus*) and roe deer (*Capreolus*
400 *capreolus*) were identified (Pavlu, 2014a). The remains were irregularly spread over the
401 settlement mainly concentrated in pits situated alongside and between houses. Statistical
402 correspondence analysis of the bone fragments found at the whole site of Bylany and at specific
403 households showed that cattle and pigs were mainly associated with long and middle houses,
404 while goats and sheep were more common in small houses (Pavlu, 2014a).

405
406 All potsherds examined herein were excavated from pits alongside long, middle or small houses
407 or from the immediate surroundings of houses, and not from the inner space of houses
408 (Soudsky, 1966; Pavlu, 2010). Animal fats were detected in sherds from across all types of
409 houses (Fig. 8). Animal fats extracted from sherds from long, middle and short houses exhibited
410 $\Delta^{13}\text{C}$ values ranging between -3.1 and 1.1‰ ($n = 13$), 1.1 and 2.3 ‰ ($n = 16$) and 1.4 and 2.8‰
411 ($n = 9$), respectively. The $\Delta^{13}\text{C}$ values of animal fats extracted from sherds from long and
412 middle houses are similar to those from short houses. The median of $\Delta^{13}\text{C}$ values for small
413 houses (-0.6‰) and middle and long houses (-0.9‰) were statistically comparable (Mann-
414 Whitney U test, $U = 115$, $P < 0.05$). Animal fats detected in sherds from long and middle houses
415 exhibited $\delta^{13}\text{C}$ and $\Delta^{13}\text{C}$ values consistent with pure non-ruminant carcass ($n = 10$; 34%) fats
416 and mixtures of non-ruminant and ruminant carcass fats ($n = 18$; 44%). However, non-ruminant
417 carcass fats and mixtures of ruminant and non-ruminant carcass fats were also detected in sherds
418 from pits alongside small houses ($n = 9$), providing evidence that both non-ruminant and
419 ruminant products were processed in the small houses. Products from ruminant
420 (cattle/sheep/goats) and non-ruminants (pigs) could have thus been processed at the small
421 houses. Mixtures of ruminant and non-ruminant products occur across all types of houses
422 without any preference, although the exploitation of pigs seems to have been restricted to long
423 and middle houses (Pavlu, 2014a). The inhabitants likely share their economies to support also

their neighbours (probably birth related), e.g. at the level of two or three houses, to maintain effective subsistence. For instance, if a cow, a pig or a large wild animal was slaughtered before winter or during some ceremonial rites, meat could have been divided between a bigger groups of people and processed at the level of each household, independently of house size, as evidenced in other Central European LBK settlements (Halstead, 2011; Marciniak 2011, 2008b).

Interestingly, no evidence for dairy fat residues was detected in potsherds from long, middle and small houses at the later settlement phase of Bylany (phase 19). The absence of dairy fats in sherds from Bylany can relate to: (i) the absence of milk use and processing at the site, (ii) the mixing of dairy and non-ruminant carcass products in pots, masking the milk signal, or (iii) the use of dairy products in perishable containers, which did not survive, e.g. in wooden vessels (Maigrot, 2003) or in leather containers (Morris, 2013). It should be also considered, dairy fat triacylglycerols with lower carbon number are more susceptible to degradation, especially when present in fresh milk and not concentrated e.g. in butter or cheese (see Dudd and Evershed 1998; Copley et al., 2005). However, per compound-specific stable carbon isotope analysis results, the high number of animal fats detected in the pots (38 potsherds of 94 with concentration of lipids $> 10 \mu\text{g g}^{-1}$) provide a relatively secure evidence for the absence of dairy fat residues in pots from phase 19 at Bylany. Unfortunately, the absence of sufficient archaeological remains prevents herd structures to be reconstructed and herding management to be assessed (Vigne and Helmer, 2007). Thus, milk exploitation at the site seems of rather low intensity or even non-existent. This agrees with lipid residue analysis studies carried out at other central European LBK sites (Salque et al., 2012; Roffet-Salque and Evershed, 2015) and with aDNA analyses performed on human skeletal remains from LBK sites in Germany, Hungary or Poland (Burger et al., 2007; Oelze et al., 2011).

449

450 Fresh milk drinking by early farming economies has been suggested to have offered an
451 evolutionary advantage to Neolithic societies with a long tradition of cattle herding (Gerbault
452 et al., 2007; Itan et al., 2009). Although Bylany inhabitants clearly exploited cattle and other
453 ruminants, thus, raw milk would have been available to them. However, based upon lipid
454 residues they appeared not to be processing dairy products in pottery. Explanations for this
455 include absence of the genetic disposition to digest milk and/or the lack of knowledge of how
456 to process it into a digestible form (Spangenberg et al., 2006; Evershed et al., 2008; Salque et
457 al., 2013; Budja et al., 2013).

458

459 **Conclusions**

460 In order to extend our knowledge about diet and household economies at the later LBK phases
461 of the Neolithic site of Bylany (phase 19) >160 potsherds were submitted to lipid residue
462 analyses. The sherds studied originated from bowls, dishes and jars excavated from long,
463 middle and small houses. More than 70% of potsherds exhibited detectable lipids, confirming
464 that many of the vessels were used for cooking or serving of food, or the processing/storage of
465 animal products.

466

467 The favourable preservation of lipids is enhanced by porous nature of the pottery microstructure
468 and chemistry (Heron et al., 1991) as evidenced further by the porosimetry analyses reported
469 herein. Mean porosity of the studied potsherds was 23% and they contained a predominance of
470 micro- and mesopores, which would aid the protection of adsorbed lipids during burial. The
471 porosity measurements also indicated that pottery production technology at Bylany was well-
472 developed, which would have influenced the ways vessels were used and the extent to which

they survived in the archaeological record (Tite et al., 2001). The slightly acid pH burial conditions at the site appear favourable for lipid preservation (e.g. Smyth and Evershed, 2015).

Comparing lipid residue compositions to the pot shapes, indicate some differences in the foodstuffs processed in different types of pots, although some hypotheses regarding vessel specialisation appear not to hold.

Stable carbon isotopic analyses of the animal fats detected in potsherds from Bylany demonstrated that carcass products were obtained from both ruminant and non-ruminant sources. Mixtures of carcass fats from both ruminant and non-ruminant animals were detected across all types of vessels and houses without any significant context specificity. It appears that the inhabitants of Bylany shared their economies between each other, independent of the size of the houses, herds or fields they managed, thus, suggesting maintenance of extensive cultural relationships needed to maintain essential sustainable community-level subsistence (Richards, 2002; Halstead, 1999). Notwithstanding this, individual households may still have played a specific roles in the overall settlement economy.

Evidence for dairy fats was not detected in any of the sherds analysed in this study neither by lipid composition nor by compound-specific stable carbon isotope analysis, suggesting that cattle or other ruminants were likely not milked at Bylany, at least during the later LBK phase 19. The importance of dairying to the economy at Bylany, thus, remains an open question. Further work in progress will shed light on the exploitation of animal resources at earlier phases of the site.

496

497 **Acknowledgements**

498 We would like to thank Institute of Archaeology in Kutna Hora for providing the potsherds
499 from Bylany. We also wish to thank Doc. Ing. Jan Vidensky, CSc at the Department of
500 Inorganic Technology of UCT Prague for the mercury porosimetry analyses. We thank the
501 European Research Council (FP7-IDEAS-ERC/324202 attributed to R.P.E.) for funding and
502 the Natural Environment Research Council (R8/H10/63) for partial funding of the mass
503 spectrometry facility at Bristol.

504 **References**

- 505 Braun, D. P., 1983. Pots as tools. In: Keene, A., Moore, J. (Eds.), *Archaeological Hammers and*
506 *Theories*. Academic Press, New York, 107-134.
- 507
- 508 Budja, M., Ogrinc, N., Gasparic, A. Z., Potocnik, D., Zigon, D., Mlekuz, D., 2013. Transition
509 to farming—transition to milk culture: a case study from Mala Triglavca, Slovenia. *Documenta*
510 *Praeistorica* 40, 97-117.
- 511
- 512 Burger, J., Krichner, M., Bramanti, B., Haak, W., Thomas, M. G., 2007. Absence of the lactase-
513 persistence-associated allele in early Neolithic Europeans. *Proceedings of National Academy*
514 *of Sciences of the United States of America* 104 (10), 3736-3741.
- 515
- 516 Correa-Ascencio, M., Evershed, R. P., 2014. High throughput screening of organic residues in
517 archaeological potsherds using direct acidified methanol extraction, *Analytical Methods* 6 (5),
518 1330-1340.
- 519
- 520 Copley, M. S., Rose, P. J., Clapham, A., Edwards, D. N., Horton, M. C., Evershed, R. P., 2001.
521 Detection of palm fruit lipids in archaeological pottery from Qasr Ibrim, Egyptian Nubia.
522 *Proceedings of the Royal Society of London B*. 268 (1467), 593-597.
- 523
- 524 Copley, M. S., Berstan, R., Dudd, S. N., Docherty, G., Mukherjee, A. J., Straker, V., Evershed,
525 R. P., 2003. Direct chemical evidence for widespread dairying in prehistoric Britain.
526 *Proceedings of the National Academy of Sciences of the United States of America* 100 (4),
527 1524-1529.
- 528
- 529 Copley, M. S., Berstan, R., Dudd, S. N., Aillaud, S., Mukherjee, A. J., Straker, V., Payne, S.,
530 Evershed, R. P., 2005. Processing of milk products in pottery vessels through British prehistory.
531 *antiquity* 79 (306), 895-908.
- 532
- 533 Coudart, A., 1998. *Architecture et société néolithique. L'unité et la variance de la maison*
534 *danubienne*. Documents d'Archéologie Française 67. Paris.
- 535
- 536 Dudd, S. N., Evershed, R. P., 1998. Direct demonstration of milk as an element of
537 archaeological economies. *Science* 282 (5393), 1478-1481.
- 538
- 539 Dudd, S. N., Evershed, R. P., Gibson, A. M., 1999. Evidence for varying patterns of exploitation
540 of animal products in different Prehistoric pottery traditions based on lipids preserved in surface
541 and absorbed residues. *Journal of Archaeological Science* 26 (12), 1473-1482.
- 542
- 543 Duncan, W. R. H., Garton, G. A., 1978. Differences in the proportions of branched-chain fatty
544 acids in subcutaneous triacylglycerols of barley-fed ruminants. *British Journal of Nutrition* 40
545 (01), 29-33.
- 546
- 547 Dunne, J., Evershed, R. P., Salque, M., Cramp, L. J. E., Bruni, S., Ryan, K., Biagetti, S., di
548 Lernia, S., 2012. First dairying in green Saharan Africa in the fifth millennium BC. *Nature* 486
549 (7403), 390-394.

550

551 Eckmeier, E., Wiesenberg, G. L.B., 2009. Short-chain n-alkanes (C 16–20) in ancient soil are
 552 useful molecular markers for prehistoric biomass burning. *Journal of Archaeological*
 553 *Science* 36 (7), 1590-1596.

554

555 Evershed, R. P., Heron, C., Goad, L. J., 1990. Analysis of organic residues of archaeological
 556 origin by high-temperature gas chromatography and gas chromatography-mass spectrometry.
 557 *The Analyst* 115 (10), 1339-1342.

558

559 Evershed, R. P., Heron, C., Charters, S., Goad, L. J., 1992. The survival of food residues, new
 560 methods of analysis, interpretation and application. In *Proceedings of the British Academy*.
 561 Vol. 77, No. 187, 2).

562

563 Evershed, R. P., Stott, A. W., Raven, A., Dudd, S. N., Charters, S., Leyden, A., 1995. Formation
 564 of long-chain ketones in ancient pottery vessels by pyrolysis of acyl lipids. *Tetrahedron Letters*
 565 36 (48), 8875-8878.

566

567 Evershed, R. P., Mottram, H. R., Dudd, S. N., Charters, S., Stott, A. W., Lawrence, G. J.,
 568 Reeves, V., 1997. New criteria for the identification of animal fats preserved in archaeological
 569 pottery. *Naturwissenschaften* 84 (9), 402-406.

570

571 Evershed, R. P., Dudd, S. N., Copley, M. S., Berstan, R., Stott, A. W., Mottram, H., Crossman,
 572 Z., 2002. Chemistry of archaeological animal fats. *Accounts of Chemical Research* 35 (8), 660-
 573 668.

574

575 Evershed, R. P., Copley, M. S., Dickson L., Hansel F. A., 2008. Experimental evidence for the
 576 processing of marine animal products and other commodities containing polyunsaturated fatty
 577 acids in pottery vessels. *Archaeometry* 50 (1), 101-113.

578

579 Evershed, R. P., 2008a. Experimental approaches to the interpretation of absorbed organic
 580 residues in archaeological ceramics, *World Archaeology* 40 (1), 20-47.

581

582 Evershed, R. P., 2008b. Organic residue analysis in archaeology: The archaeological biomarker
 583 revolution. *Archaeometry* 50 (6), 895-924.

584

585 Filipović, D., Tasić, N. N., 2012. Vinča-Belo Brdo, a Late Neolithic Site in Serbia:
 586 Consideration of the Macro-Botanical Remains as Indicators of Dietary Habits. *Balkanica*
 587 (XLIII), 7-28.

588

589 Gerbault, P., Liebert, A., Itan, Y., Powell, A., Currat, M., Burger, J., Swallow, M. D., Thomas,
 590 G. M., 2011. Evolution of lactase persistence: an example of human niche construction.
 591 *Philosophical Transactions of the Royal Society B: Biological Sciences* 366 (1566), 863-877.

592

593 Gomart, L., Hachem, L., Hamon, C., Giligny, F., Ilett, M., 2015. Household integration in
 594 Neolithic villages: A new model for the Linear Pottery Culture in west-central Europe. *Journal*
 595 *of Anthropological Archaeology* 40, 230-249.

596

- Gosselain, O.P., 1992. Technology and Style: potters and pottery among Bafia of Cameroon. *Man*, New Series 27, 559-586.
- Gregg, M. W., Banning, E. B., Gibbs, K., Slater, G. F., 2009. Subsistence practices and pottery use in Neolithic Jordan: molecular and isotopic evidence. *Journal of Archaeological Science* 36 (4), 937-946.
- Grünberg, J. M., 2002. Middle Palaeolithic birch-bark pitch. *Antiquity* 76 (291), 15-16.
- Gunstone, F. D., Hardwood, J. O., Dijkstra, A., D., 2007. The lipid handbook, 3rd edition, CRC Press, Taylor & Francis group, USA.
- Halstead, P., 1999. Neighbours from hell? The household in Neolithic Greece. In Halstead, P. (ed.) *Neolithic society in Greece. Sheffield Studies in Aegean Archaeology* 2, 77-95. Sheffield, Sheffield Academic Press.
- Halstead, P., 2012. Feast, food and fodder in Neolithic-Bronze Age Greece: commensality and the construction of value, in *Between Feasts and Daily Meals. Towards an Archaeology of Commensal Spaces* (ed. S. Pollock). *eTopoi. Journal for Ancient Studies*, special volume 2, 21-51.
- Hachem, L., 2000. New observations on the Bandkeramik house and social organization, *Antiquity* 74 (284), 308-312.
- Heron, C., Evershed, R. P., Goad, L. J., 1991. Effects of migration of soil lipids on organic residues associated with buried potsherds. *Journal of. Archaeological Science* 18 (6), 641-659.
- Heron, C., Nemcek, N., Bonfield, M. K., 1994. Chemistry of Neolithic beeswax. *Naturwissenschaften* 81 (6), 266-269.
- Chandler, R. F., Hooper, S. N., 1979. Friedelin and associated triterpenoids. *Phytochemistry* 18 (5), 711-724.
- Charters, S., Evershed, R. P., Quye, A., Blinkhorn, P. W., Reeves, V., 1997. Simulation experiments for determining the use of ancient pottery vessels: the behaviour of epicuticular leaf wax during boiling of a leafy vegetable. *Journal of Archaeological Science* 24 (1), 1-7.
- Itan, Y., Powell, A., Beaumont, M. A., Burger, J., Thomas, M. G., 2009. The origins of lactase persistence in Europe. *PloS computational biology* 5 (8), e1000491.
- Keeney, M., Katz, I., Allison, M. J., 1962. On the probable origin of some milk fat acids in rumen microbial lipids. *Journal of the American Oil Chemists Society* 39 (4), 198-201.
- Kvetina, P., Pavlu, I., 2007. Neolitické sídlisko v Bylanech - základní databáze. Neolithic settlement at Bylany – essential database. Institute of Archaeology, Prague.

Kvetina, P., Koncelova, M., 2012. Neolithic LBK Intrasite Settlement Patterns. A Case Study from Bylany (Czech Republic). *Journal of Archaeology*, 1-7.

Kolattukudy, P. E., 1976. *Chemistry and Biochemistry of Natural Waxes*. Amsterdam: Elsevier.

Maigrot, Y., 2003. Ivory, bone and antler tools production systems at Chalain 4 (Jura, France): late Neolithic site, 3rd millennium. From Hooves to Horns, from Mollusc to Mammoth—Manufacture and Use of Bone Artefacts from Prehistoric Times to the Present. *Proceedings of the 4th Meeting of the ICAZ WBRG at Tallinn 2003*, 113-126.

Marciniak, A., 2008b. Miejsca ceremonialne w kulturze ucharow lejkowatych na Kujawach: Krusza Podlotowa, stan. 2 I Podgaj, stan. 7A w swietle wynikov nowych badan archeozoologicznych i tafonomicznych. In J. Bednarczyk, J. Czebreszuk, P. Makarowicz, & M. Szmyt (Eds.), *Na pograniczu swiatow: Studia z pradziejow miedzymorza baltycko-pontynjskiego* (pp. 382-394). Poznan: Wydawnictwo Poznanskie.

Marciniak, A., 2011. The Secondary products revolution: Empirical evidence and its current zooarchaeological critique. *Journal of World Prehistory* 24 (2-3), 117-130.

Mitkidou, S., Dimitrakoudi, E., Urem-Kotsou, D., Papadopoulou, D., Kotsakis, K., Stratis, J. A., Stephanidou-Stephanatou, I., 2008. Organic residue analysis of Neolithic pottery from North Greece. *Microchimica Acta* 160 (4), 493-498.

Modderman, P. J. R., 1986. On the typology of the house plans and their European setting. *Pamatky Archeologicke* 77, 3853-394.

Morris, S., 2013. From Clay to Milk in Mediterranean Prehistory: Tracking a Special Vessel. *Backdirt. Annual Review of the Cotsen Institute of Archaeology* 2013, 70-79.

Mukherjee, A. J., Copley, M. S., Berstan, R., Evershed, R. P., 2005. Interpretation of $\delta^{13}\text{C}$ values of fatty acids in relation to animal husbandry, food processing and consumption in Prehistory, in *The zooarchaeology of fats, oils, milk and dairying* (ed. J. Mulville and A. K. Outram), 77-92, Oxbow Books, Oxford

Nelson, K., 2010. Environment, cooking strategies and containers. *Journal of Anthropological Archaeology* 29 (2), 238-247.

Oelze, V. M., Siebert, A., Nicklisch, N., Meller, H., Dresely, V., Alt, K. W., 2011. Early Neolithic diet and animal husbandry: stable isotope evidence from three Linearbandkeramik (LBK) sites in Central Germany. *Journal of Archaeological Science* 38 (2), 270-279.

Outram, A. K., Stear, N. A., Bendrey, R., Olsen, S., Kasparov, A., Zaibert, V., Evershed, R. P. 2009. The earliest horse harnessing and milking. *Science* 323 (5919), 1332-1335.

Pavlu, I., 1987. Das Modell der neolithischen Siedlung in Bylany. Bylany. *International Seminar Liblice*.

- Pavlu, I., 2000. Life on a Neolithic site: Bylany-situational analysis of artefacts. Institute of Archaeology, Czech Academy of Sciences. Praha.
- Pavlu, I., Zapotocka, M., 2007. Archeologie pravekych Cech 3 Neolit. Institute of Archaeology, Czech Academy of Sciences, v.v. i. Praha.
- Pavlu, I., 2010. Cinnosti na neolitickem sidlisti Bylany. Prostorova analyza keramiky. Activities on a Neolithic Site of Bylany - An Intrasite Spatial Analysis of Pottery. Institute of Archaeology, Czech Academy of Sciences. Praha.
- Pavlu, I., 2014a. Spolecnost a lide na neolitickem sidlisti Bylany – Society and People on a neolithic Site of Bylany. Institute of Archaeology, Czech Academy of Sciences. Praha.
- Pavlu, I., 2014b. The origins of the Neolithic of Bohemia in space, time and probability. In: M. Popelka – R. Smidtova (eds.), Neolitizace aneb setkani generaci, 165 -174. Praha
- Podborsky, V., 1997. Dejiny praveku a rane doby dejinne. Masarykova univerzita. Brno.
- Prost, D., Langevin, O., Lucquin, A., March, R. J., Verdin, P., Lepinay, D., 2011. Le site Neolithique recent-final de. Revue archeologique de Picardie. Numero special 28 (1), 221-248.
- Pye, K., 1995. The nature, origin and accumulation of loess. Quaternary Science Reviews 14 (7), 653-667.
- Raven, A. M., van Bergen, P. F., Stott, A. W., Dudd, S. N., Evershed, R. P., 1997. Formation of long-chain ketones in archaeological pottery vessels by pyrolysis of acyl lipids. Journal of Analytical and Applied Pyrolysis 40, 267-285.
- Regert, M., Garnier, N., Decavallas, O., Cren-Olivé, C., Rolando, Ch., 2003. Structural characterization of lipid constituents from natural substances preserved in archaeological environments. Measurement Science and Technology 14 (9), 1620-1630.
- Regert, M., 2004. Investigating the history of prehistoric glues by gas chromatography–mass spectrometry. Journal of separation science 27 (3), 244-254.
- Rice, P. M., 2006. Pottery analysis: a sourcebook. University of Chicago Press.
- Richards, M. P., 2002. A brief review of the archaeological evidence for Palaeolithic and Neolithic subsistence. European Journal of Clinical Nutrition 56 (12), 1270-1278.
- Roffet-Salque, M., Evershed, R. P., 2015. Shifting pottery use and animal management at Kopydłowo (Poland) traced through lipid residue analyses of pottery vessels. Kopydłowo, stanowisko 6. Osady neolityczne z pogranicza Kujaw i Wielkopolski. A. Marciniak, I. Sobkowiak-Tabaka, M. Bartkowiak and M. Lisowski. Pękowice-Poznań, Wydawnictwo Profil-Archeo: 133-142.

- Roffet-Salque, M., Dunne, J., Altoft, T. D., Casanova, E., Cramp, J. E. L., Smyth, J., Whelton, H., Evershed, R. P., 2016. From the inside out: Upscaling organic residue analyses of archaeological ceramics. *Journal of Archaeological Science: Reports*. <http://dx.doi.org/10.1016/j.jasrep.2016.04.005>
- Salque, M., Radi, G., Tagliacozzo, A., Pino Uria, B., Wolfram, S., Stäuble, H., Hohle, I., Whittle, A., Hofmann, D., Pechtl, J., Schade-Lindig, S., Eisenhauer, U. and Evershed, R. P., 2012. New insights into the early Neolithic economy and management of animals in Southern and Central Europe revealed using lipid residue analyses of pottery vessels. *Anthropozoologica* 47 (2), 45-61.
- Salque, M., Bogucki, P. I., Pyzel, J., Sobkowiak-Tabaka, I., Grygiel, R., Szmyt, M., Evershed, R. P., 2013. Earliest evidence for cheese making in the sixth millennium BC in northern Europe. *Nature* 493 (7433), 522-525.
- Schellekens, J., Barbera, G. G., Buurman, P., 2013. Potential vegetation markers—analytical pyrolysis of modern plant species representative of Neolithic SE Spain. *Journal of Archaeological Science* 40 (1), 365-379.
- Schiffer, M. B., 1990. Technological change in water-storage and cooking pots: some predictions from experiment. In: King, W. D. (Ed.). *The Changing Roles of Ceramics in Society: 26 000 B.P. to the Present. Ceramic and Civilization 5*. American Ceramic Society Inc., Westerville, Ohio, 119-136.
- Skibo, J. M., 2013. Understanding pottery function. In: *Understanding Pottery Function*. Springer New York, 1-25.
- Smyth, J., Evershed, R. P., 2015. Milking the Megafauna: Using organic residue analysis to understand early farming practice. *Environmental Archaeology* 0, 1-16.
- Soudsky, B., 1966. Bylany, osada nejstarsich zemedelcu z mladsi doby kamenne. Czechoslovak Academy of Sciences. Academia Praha.
- Spangenberg, J. E., Jacomet, S., Schibler, J., 2006. Chemical analyses of organic residues in archaeological pottery from Arbon Bleiche 3, Switzerland—evidence for dairying in the late Neolithic. *Journal of Archaeological Science* 33 (1), 1-13.
- Stacey, R. J., 2013. Organic residues: origins, analysis and scope—an overview for the archaeological ceramicist. *The Old Potter's Almanack* 14 (1), 1-8.
- Tite, M. S., Kilikoglou, V., Vekinis, G., 2001. Strength, toughness and thermal shock resistance of ancient ceramics, and their influence on technological choice. *Archaeometry* 43 (3), 301-324.
- Urem-Kotsou, D., Stern, B., Heron, C., Kotsakis, K., 2002. Birch-bark tar at Neolithic Makriyalos, Greece. *Antiquity* 76 (294), 962-967.

783 Urem-Kotsou, D., Kotsakis, K., 2007. Pottery, cuisine and community in the Neolithic of north
784 Greece. *Cooking up the Past: food and culinary practices in the Neolithic and Bronze Age*
785 *Aegean*, Oxbow, Oxford, 225-246.
786
787 Van de Velde, P., 1990. Bandkeramik social inequality – a case study. *Germania* 68 (1), 19-38.
788
789 Varien, M. D., Mills, B. J., 1997. Accumulations research: Problems and prospects for
790 estimating site occupation span. *Journal of Archaeological Method and Theory* 4 (2), 141-191.
791
792 Vigne, J.-D., Helmer, D., 2007. Was milk a “secondary product” in the Old World
793 Neolithisation process? Its role in the domestication of cattle, sheep and goats.
794 *Anthropozoologica* 42 (2), 9-40.
795
796 Velisek, J., 2013. *The Chemistry of Food*. John Wiley & Sons, New York.
797

798 **Fig.1.** Map showing the location of Bylany within the Early Neolithic (LBK) settlement area
 799 in Bohemia, Czech Republic (after Kvetina, Koncelova, 2012).

Fig. 2. Map of households from which potsherds were selected for analysis from the settlement of Bylany (Area A and B - settlement phases 9-25; area F - settlement phases 1-8).

Fig. 3. Partial gas chromatogram of total lipid extracts from sherds from Bylany phase 19 typical of animal fat. Peak identities: FAME = fatty acid methyl esters, br-branched, C16-MAG - monopalmitoylglycerol, C18-MAG-monostearoylglycerol, IS-internal standard (*n*-tetratriacontane).

Fig. 4. Partial gas chromatograms of (a) animal fats with mid-chain ketone and (b) traces of fatty acids and resinous components (magnified). Peak identities: FAME = fatty acid methyl esters, TMS-trimethylsilylated, br-branched, C16-MAG - monopalmitoylglycerol, C18-MAG-monostearoylglycerol, 9 DCA – azelaic acid, 9,10- C18 FAOH – 9,10-dihydroxystearic acid, 31(35)-K – mid-chain ketones with 31 and 35 carbons, IS-internal standard (*n*-tetratriacontane), PL – plasticizer (contamination).

FUNCTIONAL SETS POTTERY	
WATER PROCESSING	<p>Volume (N)</p> <p>water manipulation</p> <p>3.22(343) F6 SET1</p> <p>4.93(1582) F10</p> <p>2.92(204) F13</p> <p>earlier F11 1.27(1264)</p> <p>later F9 0.94(1669)</p>
	<p>6.94(1226) F12</p> <p>2.66(592) F4 SET2</p>
SERVING AND FOOD PROCESSING	<p>serving liquid food</p> <p>0.62(484) F3</p> <p>0.76(318) F5</p> <p>3.55(841) F8</p> <p>0.75(784) F7 SET3</p>
	<p>serving solid food</p> <p>0.49(258) F1</p> <p>2.02(289) F2 SET4</p>
STORAGE	<p>storage pot</p> <p>15.10(150) F14 SET5</p>

Fig. 5 Functional classification of archaeological pottery from Bylany according to their typologies (after Pavlu, 2000).

Fig. 6. Box plot of total lipid concentrations [$\mu\text{g g}^{-1}$] in different categories of vessels from Bylany (A) and functional sets (B); $\circ$ remote values; * extremes. Set 1 - Water processing (cat. F6, F9, F10, F11, F13), Set 2 - Serving and food processing (cat. F4, F5), Set 3 - Serving liquid food (cat. F3, F5, F7, F8), Set 4 - Serving solid food (cat. F1, F2), Set 5 - Storage (cat. F14).

Fig. 7. Box plot of total lipid concentrations [$\mu\text{g g}^{-1}$] in pottery recovered from specific houses examined in this study. Recovery rates of lipids in brackets; ○ remote values; * extremes.

Fig. 8. (A) $\delta^{13}\text{C}$ values for the $\text{C}_{16:0}$ and $\text{C}_{18:0}$ fatty acids from the TLEs extracted from potsherds from Bylany phase 19 according to the type of houses (long ▲, middle ■ and small ○). The three fields correspond to the $P = 0.684$ confidence ellipses for animals raised on a strict C_3 diet in Britain (Copley et al., 2003). (B) Difference in the $\delta^{13}\text{C}$ values of the $\text{C}_{18:0}$ and $\text{C}_{16:0}$ fatty acids ($\Delta^{13}\text{C} = \delta^{13}\text{C}_{18:0} - \delta^{13}\text{C}_{16:0}$) for the same archaeological fats. The ranges represent the mean $\pm$ s.d. for a global database comprising modern animal fats from across the globe (Copley et al., 2003; Outram et al., 2009; Spangenberg et al., 2006; Gregg et al., 2009; Dunne et al., 2012). Each data point represents an individual potsherd. Analytical precision ± 0.3

Fig. 9. Scatter plots of stable carbon isotope compositions of animal fat residues extracted from the potsherds of Bylany phase 19 in a context of presumed **functional sets** (A) where ● represent water processing vessels, ■ processing and food serving vessels and ▲ represent storage vessel, and **vessel shapes** (B) where ▲ represent big jar, ■ big dish, □ small dish, ● big bowl and ● small bowl.

Table 1 Summary of the results of the lipid residue analyses of potsherds from Bylany (phase 19).

			Potsherds analysed	Residues detected	Recovery rate
			<i>n</i>	<i>n</i>	[%]
Houses	Size				
96	Long		40	29	73
162	Long		16	12	75
1246	Long		18	11	61
361	Middle		17	10	59
619	Middle		23	21	91
702	Middle		9	6	67
959	Small		21	21	100
1161	Small		13	6	46
1240	Small		6	2	33
Total			163	118	72
Category	Shape	Functional set			
F1	Small dish	4	10	9	90
F2	Big dish		13	4	31
F3	Small dish		8	7	88
F4	Big dish		12	8	67
F5	Small dish	3	11	7	64
F6	Big dish	1	9	7	78
F7	Small bowl	3	11	9	82
F8	Big bowl		11	9	82
F9	Small bowl		12	10	83
F10	Big bowl	1	14	10	71
F11	Small bowl		12	8	67
F12	Big bowl	2	12	9	75
F13	Jar	1	13	10	77
F14	Big jar	5	15	11	73
Total			163	118	72

Table 2 Summary of extracted lipids in the context of houses and vessel types

House type	House number	Vessel type	Lipids detected ^a
Small house	959	Big dish	FA(l,vl,br), MAG	Small dish (F1, F3, F5)
		Big bowl	FA(m,l,vl,br), MAG	
		Jar	FA(m,l,vl,br), MAG, OH, TOH, ALK, DCA	Big dish (F4,F6)
	1161	Small bowl	FA(l,br,vl), MAG
Middle house	361	Small dish	FA(l,br,vl), MAG, OH(tr)	Small bowl (F7, F9, F11)
		Big dish	FA(l,br,vl), MAG, OH(tr)	
		Small bowl	FA(l,br,vl), MAG, DCA (tr), OH(tr)	
		Big bowl	FA(m,l,br), MAG, DCA, OH	
	619	Jar	FA(l,br), MAG	Big bowl (F8, F10, F12)
		Small dish	FA(l,br), MAG, OH(tr)	
		Big dish	FA(l,br,vl), MAG	
		Small bowl	FA(l,br), MAG	
		Big bowl	FA(m,l,br), MAG	
	702	Big dish	FA(l,br), MAG	Jar (F13)
Long house	96	Small dish	FA(m,l,br,vl), MAG, FAOH, DCA	
		Big dish	FA(m,l,br,vl), MAG, DCA	
		Small bowl	FA(tr)	
		Big bowl	FA(tr), MAG(tr)	
		Big jar	FA(tr)	
	162	Small dish	FA(l,br,vl), MAG, DCA	Big jar (F14)
		Jar	FA(m,l,br,vl), MAG, DCA, ALK, K, FAOH, TOH	
	1246	Small dish	FA(l,br,vl), MAG	
		Big bowl	FA(l,br,vl), MAG	

^aFA, *n*-alkanoic fatty acids (m-middle, l-long, br-branched, vl-very long); MAG, monoacylglycerols; OH, *n*-alcohols; DCA, dicarboxylic acids; ALK, *n*-alkanes; K, mid-chain ketones; FAOH, hydroxy fatty acids; TOH, triterpene alcohols; tr, traces.

Table 3 Details of potsherds selected for GC-C-IRMS, results and interpretation of the isotopic analyses.

<i>i</i>	Lab number	House number	Category	Rim diameter [cm]	TLE [$\mu\text{g g}^{-1}$]	$\delta^{13}\text{C}_{16:0}$ [‰]	$\delta^{13}\text{C}_{18:0}$ [‰]	$\Delta^{13}\text{C}$ [‰]	Classification by $\Delta^{13}\text{C}$ values
1	BYL-C-1321	702	F4	unknown	193	-25.9	-27.7	-1.8	Mixture N/R ^a
2	BYL-C-1322	702	F12	unknown	895	-26.1	-28.4	-2.3	Ruminant adipose
3	BYL-C-1323	702	F12	unknown	350	-24.8	-25.8	-0.9	Mixture N/R
4	BYL-C-1325	702	F4	unknown	498	-24.8	-25.8	-0.9	Mixture N/R
5	BYL-C-1329	1161	F5	unknown	822	-25.8	-27.9	-2.2	Ruminant adipose
6	BYL-C-1334	1246	F4	unknown	43	-26.4	-28.2	-1.7	Mixture N/R
7	BYL-C-1345	1161	F7	unknown	1975	-25.7	-25.4	0.2	Non-ruminant
8	BYL-C-1353	1246	F3	unknown	1245	-25.7	-25.1	0.6	Non-ruminant
9	BYL-C-1354	1246	F10	unknown	77	-26.4	-27.2	-0.8	Mixture N/R
10	BYL-C-1375	361	F10	18	11	-26.4	-28.1	-1.7	Mixture N/R
11	BYL-C-1376	361	F2	20	322	-26.2	-26.1	0.1	Non-ruminant
12	BYL-C-1383	619	F6	unknown	99	-26.1	-27.1	-0.9	Mixture N/R
13	BYL-C-1389	959	F10	25	61	-23.8	-22.9	0.9	Non-ruminant
14	BYL-C-1395	959	F12	16	298	-26.9	-29.3	-2.4	Ruminant adipose
15	BYL-C-1400	619	F11	12	99	-25.6	-24.7	0.9	Non-ruminant
16	BYL-C-1402	619	F10	28	33	-25.7	-26.6	-0.8	Mixture N/R
17	BYL-C-1408	619	F9	11	499	-24.9	-24.6	0.3	Non-ruminant
18	BYL-C-1411	619	F3	17	148	-24.9	-24.3	0.6	Non-ruminant
19	BYL-C-1416	959	F10	16	1634	-27.8	-30.6	-2.8	Ruminant adipose
20	BYL-C-1417	959	F10	19	600	-27.9	-28.5	-0.6	Mixture N/R
21	BYL-C-1419	959	F8	16	955	-25.3	-26.5	-1.2	Mixture N/R
22	BYL-C-1423	959	F6	20	77	-25.5	-24.1	1.4	Non-ruminant
23	BYL-C-1424	959	F6	20	53	-25.6	-24.7	0.8	Non-ruminant
24	BYL-C-1425	361	F1	9	558	-26.2	-28.2	-1.9	Ruminant adipose
25	BYL-C-1426	361	F2	26	40	-25.9	-26.1	-0.2	Mixture N/R
26	BYL-C-1427	361	F4	27	109	-25.7	-26.9	-1.2	Mixture N/R
27	BYL-C-1429	361	F10	24	232	-25.3	-24.3	1.1	Non-ruminant
28	BYL-C-1433	361	F12	16.1	1867	-25.6	-26.9	-1.3	Mixture N/R
29	BYL-C-1436	162	F1	unknown	202	-26.4	-28.4	-2.0	Ruminant adipose
30	BYL-C-1437	162	F1	unknown	49	-25.4	-24.9	0.5	Non-ruminant
31	BYL-C-1440	162	F5	unknown	43	-27.3	-30.1	-2.8	Ruminant adipose
32	BYL-C-1449	162	F14	unknown	33	-26.9	-28.8	-1.9	Ruminant adipose
33	BYL-C-1461	96	F3	13	17	-25.9	-25.7	0.2	Non-ruminant
34	BYL-C-1464	96	F5	10	824	-25.3	-26.3	-1.0	Mixture N/R
35	BYL-C-1470	96	F5	14	35	-27.6	-29.0	-1.4	Mixture N/R
36	BYL-C-1472	96	F6	20	126	-26.1	-25.9	0.2	Non-ruminant
37	BYL-C-1474	96	F6	20	166	-25.3	-28.4	-3.1	Ruminant adipose
38	BYL-C-1476	96	F7	unknown	78	-25.0	-23.9	1.1	Non-ruminant

^a Mixture N/R - mixture of non-ruminant adipose and ruminant adipose fats