Grassi, G., House, J., Dentener, F., Federici, S., den Elzen, M., & Penman, J. (2017). The key role of forests in meeting climate targets requires science for credible mitigation. *Nature Climate Change*, 7(3), 220-228. <https://doi.org/10.1038/nclimate3227>

Peer reviewed version

Link to published version (if available):
[10.1038/nclimate3227](https://doi.org/10.1038/nclimate3227)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Springer Nature at <https://www.nature.com/nclimate/journal/v7/n3/full/nclimate3227.html>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Key role of forests in meeting climate targets but science needed for credible mitigation

Authors

Giacomo Grassi*, European Commission, Joint Research Centre, Ispra (VA), Italy

Jo House, Cabot Institute, Bristol University, UK

Frank Dentener, European Commission, Joint Research Centre, Ispra (VA), Italy

Sandro Federici, FAO consultant, Italy

Michel den Elzen, PBL Netherlands Environmental Assessment Agency, The Hague, The Netherlands

Jim Penman, University College London, UK

*Corresponding author: giacomo.grassi@ec.europa.eu

ABSTRACT

Forest-based climate mitigation may occur through conserving and enhancing the carbon sink and through reducing greenhouse gas emissions from deforestation. Yet the inclusion of forests in international climate agreements has been complex, often treated separately or considered a secondary mitigation option. In the lead up to the Paris Climate Agreement, countries submitted their (Intended) Nationally Determined Contributions ((I)NDCs), including climate mitigation targets. Assuming full implementation of (I)NDCs, we show that land use, and forests in particular, emerge as a key component of the Paris Agreement: turning globally from a net anthropogenic source during 1990-2010 (1.3 ± 1.1 GtCO₂e/y) to a net sink of carbon by 2030 (up to -1.1 ± 0.5 GtCO₂e/y), and providing a quarter of emission reductions planned by countries. Realizing and tracking this mitigation potential requires more confidence in numbers, including reconciling estimates between country reports and scientific studies. This represents a challenge and an opportunity for the scientific community.

MAIN TEXT

In December 2015, 195 countries adopted the Paris Climate Agreement¹ at the 21st Conference of Parties (COP-21) of the United Nations Framework Convention on Climate Change (UNFCCC). As part of the process, 187 countries, representing more than 96% of global net emissions in 2012², submitted their Intended National Determined Contributions³ (INDCs, which become NDCs with the ratification of the Paris Agreement⁴). The NDCs are the basis for implementing actions under the Agreement, and the vast majority include commitments in the land-use sector.

Land use, including agriculture and forests, accounts for about 10% of global greenhouse gas (GHG) emissions as CO₂, and nearly quarter including CH₄ and N₂O⁵⁻⁹. Also, about one third of the current anthropogenic CO₂ emissions are removed by terrestrial ecosystems, mainly forests. While deforestation is estimated to be the main GHG source in many tropical countries, forest sinks are important globally with net sinks dominating in temperate and boreal countries¹⁰.

Including land use in the UNFCCC process has been long and complex. For forests, uncertainties of GHG estimates and methodological issues such as additionality (i.e. showing that proposed mitigation efforts go beyond Business-as-Usual (BAU) and separation of non-anthropogenic effects) and leakage (displacement of land-use activities to other areas) have often led to controversies and compromises¹¹.

The UNFCCC requires that all countries report GHG inventories of anthropogenic emissions and removals using methodologies developed by the Intergovernmental Panel on Climate Change (IPCC) and adopted by UNFCCC¹². Developed countries report annual GHG inventories¹³, using mandatory and voluntary land-use activities towards meeting their emission reduction targets where applicable under the Kyoto Protocol¹⁴. Developing countries' GHG inventories have historically been reported less frequently¹⁵, though biennial updates are now required¹⁶, and may undertake voluntary mitigation activities, notably through the REDD+ process (Reducing Emissions from Deforestation, forest Degradation, and other forest activities).

The Paris agreement is a potential game changer for land use mitigation. It calls explicitly for all countries to make use of a full-range of land-based mitigation options, and to take action on REDD+.

Based on country information, this analysis quantifies the expected GHG mitigation role of the land-use sector in the (I)NDCs to the year 2030, including activities conditional on finance, technology and capacity-building support. It does not assess specific country policies. It focuses on CO₂ emissions and removals and non-CO₂ emissions from Land Use, Land-Use Change and Forestry (LULUCF, primarily deforestation and forest management), encompassing most of the land-use sector identified in (I)NDCs. Harvested wood products are included for most developed countries. Non-CO₂ emissions from agriculture are not included.

Country mitigation targets are expressed in different ways

Countries express their (I)NDC targets with different combinations of the following elements¹⁷⁻¹⁹ (Supplementary Tables 1-2), reflecting different national circumstances, i.e.:

- Quantifier - targets are expressed as either an *absolute quantity* e.g. amount of GHG reduction in tonnes of CO₂ equivalent (tCO₂e), or as a change in the *emission intensity*, e.g. China and India express a reduction of emission intensity per unit of GDP.
- Reference point – Emissions in the target year (e.g. 2025 or 2030) are compared to either a historic base year (e.g. 1990, 2005) or to the target year in a BAU scenario. The BAU scenario assumes either no mitigation activity, or some existing mitigation activity.
- Conditionality - While developed country (I)NDC targets are unconditional, most developing countries expressed at least part of their targets as conditional on finance, technology or capacity-building support.

The (I)NDCs vary in the way they include LULUCF. It may be fully included as part of the overall target like other sectors, or partially included (e.g., only deforestation), or considered separately with

special mitigation actions or accounting rules. Consequently, evaluating the expected effect of LULUCF on the (I)NDC mitigation targets is complex.

Our analysis is based on information provided on LULUCF in the (I)NDCs³, and also (in order of priority) other country reports to UNFCCC^{13,15,16,20,21}, other official country documents, and FAO-based datasets for forest^{8,22} and for other land uses²³ (Supplementary Tables 4-5). Given the Paris Agreement context of our analysis, we prioritized (I)NDCs and those country reports which are formally reviewed or technically assessed by UNFCCC, with FAO-based datasets used for gap filling, allowing global estimates covering 195 countries (see Methods). We found sufficient information to analyse the LULUCF mitigation contribution for 68 countries (or 41 (I)NDCs, with the EU's NDC representing 28 countries), representing around 78% of global net emissions in 2012² and 83% of the global forest area²². For the remaining countries, where LULUCF is not expected to offer a large mitigation potential (Supplementary Section 1), the future LULUCF mitigation contribution was assumed to be zero.

Historical and projected forest emissions and removals

Fig. 1 shows, for all 195 UNFCCC countries, historical and future anthropogenic LULUCF emissions and removals from this analysis, based on official country data. The Supplementary Sections 2 and 3 provide, respectively, additional country-specific assessments and an analysis of uncertainties for the absolute level of net emissions and their trend^{24,25}, based on information from countries' reports.

While country information on uncertainty up to 2030 is not available, we conservatively assumed that the uncertainty estimated for historical net emissions would also hold for the future.

Historically, global LULUCF net emissions decreased from 1.54 ± 1.06 GtCO₂e/y (95% CI) in 1990 to 0.01 ± 0.86 GtCO₂e/y in 2010 (slope of linear trend: -0.08 GtCO₂e/y). The trend and the inter-annual variability over this period are influenced by: (i) deforestation in Brazil, with peak years in 1995 and 2002-2004 followed by a steep reduction of emissions by about -1.3 GtCO₂e/y till 2010; (ii) high deforestation rates (1997-1999) and peak years in peat fire emissions (e.g., 1997) in Indonesia;

(iii) an increasing sink in managed temperate and boreal forests, of about $-0.8 \text{ GtCO}_2\text{e/y}$ from 1990 to 2010. By splitting the 1990-2010 period (average emissions: $1.28 \pm 1.15 \text{ GtCO}_2\text{e/y}$) into four sub-periods, we conclude that the historical trend is statistically significant after 2000 (Supplementary Section 3).

The wide range of future LULUCF net emissions depends on policy scenarios (Fig. 1). The ‘country-BAU’ scenario foresees a marked increase in global net emissions (Supplementary Table 6), reaching $1.94 \pm 1.53 \text{ GtCO}_2\text{e/y}$ in 2030. This is because several developing countries assumed BAU to be a no-measures scenario, e.g. ignoring the existing policies to reduce deforestation. Under the ‘pre-(I)NDC scenario’, i.e. considering policies in place prior to COP-21 (including the earlier Copenhagen pledges²¹), global net emissions increase moderately, up to $0.36 \pm 0.94 \text{ GtCO}_2\text{e/y}$ in 2030. For the ‘unconditional (I)NDC scenario’ the global net emissions slightly decrease, reaching a sink of $-0.41 \pm 0.68 \text{ GtCO}_2\text{e/y}$ in 2030. An additional reduction of net emissions is estimated for the ‘conditional (I)NDC’ scenario, leading to a sink of $-1.14 \pm 0.48 \text{ GtCO}_2\text{e/y}$ in 2030.

The analysis of the emission trend over the entire period shows that the difference between the 1990-2010 average and the net emissions in 2030 is not significant for the pre-(I)NDC scenario, but is significant (95% CI) for both the unconditional and the conditional (I)NDC scenarios (Supplementary Figure 3b). This indicates that the reduction of net emissions assumed by the (I)NDCs relative to the historical period, if achieved, is statistically robust.

Comparison with global datasets

Fig. 2 compares the historical LULUCF trend from our analysis to other three well-known global LULUCF datasets: (i) latest country reports to UNFCCC (ref^{13,15,16,20}); (ii) FAOSTAT for all land uses²³; and (iii) IPCC Fifth Assessment Report (AR5) Working Groups (WG) I⁵ and III⁶ data used for the global carbon budget.

The difference between this analysis and the UNFCCC country reports is because several (I)NDCs updated past datasets, and because we used FAO-based data for gap-filling, instead of pre-2010 National Communications.

Differences between this analysis and FAOSTAT include the definition of forest (UNFCCC vs. FAO); coverage of areas and of carbon pools; and differing estimation methods by reporting agencies⁸ (see Methods).

There is a large difference of about 3 GtCO₂e/y between this analysis, based on country reports following the IPCC Guidelines for national GHG inventories^{25,26} (IPCC GL), and the scientific studies summarized by the IPCC AR5^{5,6}. For the period 2000-2009, the level of net emissions is on average 0.90 ± 1.11 GtCO₂e/y (95 % CI) in our analysis and 4.03 ± 2.93 GtCO₂e/y (90 % CI, reflecting both methodological and terminological choices²⁷⁻²⁹) in IPCC AR5 (Fig. 2). The above differences are linked to different scopes of the two IPCC work streams³⁰: the GL focus on internationally agreed methodologies for national anthropogenic GHG estimation, recognizing different countries' definitions and technical capabilities, whilst the AR5 focuses on assessing the state of the science on the global carbon budget using globally applied data, definitions and modeling methods.

Specifically, LULUCF in the IPCC GL includes estimates of GHG emissions and removals from all land uses, reported under either a stable or changed land-use status (typically in the last 20 years), e.g. “forest remaining forest” or “forest converted to cropland” (or vice versa). There is a large scientific challenge of providing a practicable methodology to factor out direct human-induced mitigation action from indirect human-induced and natural effects^{31,32}, such as the natural aging of forests, natural disturbances and environmental change (e.g. climate change, extended growing seasons, fertilizing effects of increased [CO₂] and nitrogen deposition). Therefore, the IPCC GL^{25,26} use the category of “managed land” as a default first order approximation of “anthropogenic” emissions and removals, based on the rationale that the preponderance of anthropogenic effects occurs on managed land³². The GHG inventories should report all emissions and removals for managed land, while GHG

fluxes from unmanaged land are excluded. What is included in “managed land” varies from country to country, although the countries’ definition must be applied consistently over time.

In contrast, global models such as those used in IPCC AR5 and the Global Carbon Project take a different approach to separate anthropogenic from natural effects. Anthropogenic fluxes (referred as “net land-use change”^{5,9}, or “Forestry and Other Land Uses”⁶), are estimated by a bookkeeping model²⁷ or by dynamic global vegetation models⁹ based on changes in land cover (i.e. between forest and agriculture), forest regrowth and, depending on the modeling capability, some forms of management (wood harvest and shifting cultivation). The difference between this modeled “anthropogenic” flux and the estimated total net flux of CO₂ between the land and atmosphere⁹ is the “residual terrestrial sink”^{5,6,9}, which is generally assumed to be a natural response of primary or mature regrowth forests to environmental change^{9,27}.

The above methodological differences are reflected in the net emissions from developed countries, where most of the ≈ 3 GtCO₂e/y difference between our analysis and IPCC AR5 occurs for the period 2000-2009: while these countries report a substantial “anthropogenic” sink (-1.9 GtCO₂e/y in “UNFCCC Annex 1” countries), the bookkeeping model (IPCC AR5) finds a small net source (0.1 GtCO₂e/y, “OECD” in Fig. 11.7 of ref.⁶). This difference lies essentially in whether the large sinks in areas designated by countries as “managed forest” (following IPCC GL), well documented in forest inventories¹⁰, are attributed to “anthropogenic” (in the GHG inventories) or to “natural” fluxes (in IPCC AR5).

To explore, at least in part, the impact of these different attribution methods, Fig. 3a compares what is considered undisputedly “anthropogenic” by both IPCC AR5 (land-use change) and the country reports (land converted to other land uses). These estimates, both predominated by tropical deforestation, are of similar magnitude, especially after 2000. The other fluxes, where the attribution differs more between IPCC AR5 and the countries, are shown in Fig 3b. Thus much of the sink that countries report under ‘forest remaining forest’, the global models consider part of the natural flux. This disaggregation suggests that the residual sink is at least partly influenced by management

practices not captured by global carbon models³³, but also that countries consider anthropogenic what is partly influenced by environmental change and by recovery from past disturbances.

There are many reasons for the lower sink reported by countries in Fig 3b compared to the residual sink from IPCC AR5³⁰, including the fact that countries do not report sinks for unmanaged lands (e.g., a large sink in tropical and boreal unmanaged forests¹⁰) and their reporting for managed land may be incomplete, i.e. ignoring fluxes (e.g. sink in grasslands, wetlands or forest regrowth) or carbon pools. There would be other factors to consider, including treatment of legacy fluxes from past land-use and other definitional and methodological differences. These would require a more detailed analysis, which is outside the scope of this paper.

Finally, the projections from this analysis can be compared to RCP scenarios used in IPCC AR5 up to 2030 (Fig. 3, dashed lines). For the undoubtedly “anthropogenic” fluxes (Fig. 3a), our country data analysis falls broadly within the IPCC AR5 scenarios, supporting previous qualitative findings³⁴.

Overall, our analysis shows 1) that various global LULUCF datasets may be more consistent than apparent at first glance, 2) unless the scientific and GHG inventory community appreciate these definitional and methodological issues, conflicting numbers and messages are likely to appear in the coming years, and 3) that several reasons for the differences among datasets can be further reconciled in collaboration between the two communities, which would be a very useful contribution to science and policy.

Different perspectives on mitigation contribution by forests

To reflect the complexity of approaches to (I)NDCs, this analysis assesses three different perspectives on LULUCF mitigation:

(A) 2030 (I)NDC vs. 2005, i.e., the expected impact of full (I)NDC implementation. The year 2005 is chosen as historically reliable in terms of data. Fig. 1 shows that the global LULUCF net emissions to the atmosphere would transition from an estimated net anthropogenic source of +0.8 GtCO₂e/y in

2005 to a net sink of -0.4 GtCO₂e/y (unconditional (I)NDCs) or -1.1 GtCO₂e/y (conditional (I)NDCs) in 2030.

(B) 2030 (I)NDC vs. 2030 alternative scenarios: country-BAU or pre-(I)NDC, i.e., the additional LULUCF contribution relative to alternative scenarios (Fig. 1). The magnitude of the difference between country-BAU and pre-(I)NDC (1.6 GtCO₂/y) may raise concerns about the expected results-based payments under REDD+, which should be based on credible baselines and not on a no-measures scenario. Clarification of the role of REDD+ in (I)NDCs should therefore be seen as a priority by countries. Compared to the estimated pre-(I)NDC scenario, net emissions in 2030 are lower by 0.8 GtCO₂e/y or 1.5 GtCO₂e/y for unconditional and conditional (I)NDCs, respectively. For the ‘conditional (I)NDC vs. 2030 pre-(I)NDC’ scenario (Fig. 4a), this LULUCF contribution of 1.5 GtCO₂e/y (Fig. 4a, last column) represents 26% of the total mitigation expected from all GHG sectors (5.9 GtCO₂e/y³⁵, Fig. 4a, third column). The countries contributing most to LULUCF mitigation under this perspective are Brazil and Indonesia, followed by other countries focusing either on avoiding carbon emissions (e.g. Ethiopia, Gabon, Mexico, DRC, Guyana and Madagascar) or on promoting the sink through large afforestation programs (e.g. China, India).

(C) Country perspective on emissions reduction in the (I)NDC, i.e. what each country might consider its “LULUCF contribution to the overall (I)NDC”, as part of its mitigation package, e.g. if a country commits to reduce its all-sectors emissions by $x\%$ relative to y (reference point: base year or BAU-scenario), what fraction of x is attributable to LULUCF? This approach looks at the way countries define their (I)NDCs (e.g. reference point) and the way LULUCF is included within the (I)NDC (as any other sector or with special accounting rules). Globally, under this perspective the LULUCF contribution is 3.1 GtCO₂e/y (unconditional) or 3.8 GtCO₂e/y (conditional). The latter case (Fig. 4b, last column) corresponds to 24% of total all-sectors emission reduction relative to the reference point (i.e. 15.8 GtCO₂e/y, Fig. 4b, third column).

The emission reductions from a country perspective (Fig. 4b) are greater than the deviation from the pre-(I)NDC scenario (Fig. 4a), because countries’ choices of reference point in their (I)NDCs tend to

maximize the accounted mitigation, i.e. countries that already reduced emissions used a historical base year, whereas countries expecting a future increase of emissions used a future BAU-scenario. This is evident under perspective C, where nearly one third of the contribution comes from Brazil, followed by Indonesia and Russia (Fig 4b, last column). In Brazil, where total emissions have declined after 2004 due to successful implementation of policies to reduce deforestation³⁶, the NDC target (-43%) is relative to 2005. Our analysis suggests that in Brazil the LULUCF contribution to NDC is greater than the all-sectors NDC target for 2030, i.e. the NDC allows emissions from other sectors to increase. In Indonesia the conditional NDC target (-41%) is relative to the BAU-scenario in 2030. LULUCF represents about 65% of current (2010) total emissions and is expected to contribute nearly two-thirds of the NDC emission reduction (relative to BAU) foreseen in for 2030. Brazil and Indonesia are representative examples of GHG emission trends in developing countries: with an expanding and industrializing economy, the currently high LULUCF emissions are expected to decrease, and be superseded by growing emissions from the energy sector. The (I)NDC target of Russia (-30%) is relative to 1990, with LULUCF contributing by about two-fifths to this emission reduction. Russia is more important in perspective C than in B because its specific accounting method for LULUCF gives prominence to the contribution of the current forest sink to climate mitigation.

The (I)NDCs of the three countries above may be assessed also in terms of clarity and trust of information provided (see Supplementary Section 2). Overall, Brazil's NDC is transparent on the land-use sector and the underling GHG estimates are based on a well-developed monitoring system. The recent relevant upward revision of historical deforestation emissions in Brazil opens new questions on the implementation of the NDC target and on how and when data consistency between NDC, REDD+ and National Communications will be ensured. The relative ambiguity of Indonesia's NDC on how it would address land use emissions is improved by the information in more recent documents. Furthermore, recent monitoring efforts have improved the GHG emission estimates, especially from peatland drainage and from forest degradation, whereas emissions from peat fires remain very uncertain. These improvements are mainly due to the REDD+ process, which in many developing countries is triggering unprecedented monitoring efforts. The challenge is increasingly to

transfer these improvements into the NDC process, and to clarify the often uncertain relationship between the financially-supported REDD+ activities and the NDCs. For Russia, transparency of mitigation efforts will crucially depend on clarifying the accounting method chosen for LULUCF. In addition, credible GHG estimates will require reconciling or explaining the currently large difference in the forest sink between the reports submitted by Russia to UNFCCC and to FAO.

In summary, the full implementation of (I)NDCs would turn LULUCF globally from a net source during 1990-2010 (1.3 ± 1.1 GtCO₂e/y) to a net sink by 2030 (up to -1.1 ± 0.5 GtCO₂e/y). The absolute LULUCF mitigation contribution in 2030 is very different depending on the way that mitigation is calculated, ranging from 0.8 to 3.1 GtCO₂e/y for unconditional (I)NDCs and from 1.5 to 3.8 GtCO₂e/y for conditional (I)NDCs (Supplementary Table 3). However, in relative terms, LULUCF would provide about a quarter of total emission reductions planned in countries' (I)NDCs irrespective of the approach to calculating mitigation.

Whereas a similar trend of decreasing LULUCF net emissions with full (I)NDCs implementation has been suggested also by other analyses (ref^{34,37}), the absolute level of net emissions differs significantly: e.g., ref³⁷ reports net emissions about 3 GtCO₂e/y higher than ours, due to the 'harmonization' of different datasets (country projections and (I)NDCs were aligned to historical FAOSTAT data). By contrast, our study is the first so far showing a global picture of country-based LULUCF net emissions that is consistent between historical and projected periods, including discussing the differences with other global datasets and different mitigation perspectives.

Science can help countries to keep the forest mitigation promise

Several studies suggest a theoretical mitigation potential from land use^{6,35,38} higher than in this analysis, others suggest limits posed by ecological and socio-economic constraints (including land availability)^{39,40}. Irrespective of the potential, in the past UNFCCC negotiations the LULUCF sector has often been treated separately and considered as a secondary mitigation option, largely due to its complexity and limited trust in data.

284 Our analysis shows a wide range of future LULUCF net emissions, depending on policy scenarios.
285 Through the implementation of (I)NDCs countries (especially developing ones) expect a key
286 contribution from LULUCF in meeting their (I)NDC targets, with a clear focus on forests. Achieving
287 this will require increasing the credibility of LULUCF mitigation, through more transparency in
288 commitments and more confidence in estimates. To this regard, the Paris Agreement includes a
289 “Framework for transparency of actions”, key for its credibility⁴¹, aimed at providing clarity on GHG
290 estimates and tracking of progress toward achieving countries’ individual targets.

291 More transparent commitments means that future updates of the NDCs should provide more details
292 on how LULUCF mitigation is calculated towards meeting the target and how the financially-
293 supported REDD+ activities contribute to the pledges. More confidence in LULUCF estimates will
294 require improving the country GHG inventories in terms of transparency, accuracy (including
295 information on uncertainties), consistency, completeness and comparability⁴², especially in
296 developing countries.

297 This is a challenge and an opportunity for the scientific community. Supporting country GHG
298 estimation includes regular reviews of the latest science (e.g. ref⁴³), expanding the scope of the
299 operational methods in the IPCC guidance, as has been done for REDD+⁴⁴, and incorporating
300 opportunities offered by emerging satellite data^{45,46} available through highly accessible products⁴⁷.

301 More confidence also requires independent checks of the transparency and reliability of data, e.g. by
302 reproducing and verifying countries’ GHG estimates. According to IPCC guidance²⁵, verification of
303 GHG inventories is key to improve scientific understanding and to build confidence on GHG
304 estimates and their trends. This can be achieved by comparing GHG inventories with scientific
305 studies using partially or totally independent datasets and/or different methods (e.g. ref⁴⁸), including
306 greater integration of modeling and measurement systems of land use-related net emissions⁹.

307 Meaningful verification requires improving mutual understanding and cooperation between the
308 scientific community and the developers of national GHG inventories.

Finally, increasing trust in proposed LULUCF mitigation options will require reconciling the current differences in global LULUCF net emissions between country reports and scientific studies (as reflected in IPCC reports). Among the many possible reasons for these differences^{30,49}, we suggest that what is considered “anthropogenic sink” is key and deserves further analyses. While recognizing differences in scopes among these communities, reconciling differences in estimates is a necessity, as the “Global stocktake”, i.e. the foreseen five-yearly assessment of the collective progress toward achieving the long-term goals of the Paris Agreement, will be based on both country reports and IPCC reports. Without speaking the same language, the “balance between anthropogenic GHG emissions by sources and removals by sinks in the second half of this century”¹, needed to reach the ambitious “well-below 2°C” target, cannot be properly assessed.

Correspondence and requests for materials: giacomo.grassi@ec.europa.eu

Disclaimer: The views expressed are purely those of the writers and may not in any circumstances be regarded as stating an official position of the European Commission.

Author Contributions G.G. conceived the analysis on (I)NDCs, executed the calculations and drafted the paper. J.H., F.D., M.d.E. and J.P. contributed to the analysis and to the writing of the paper. S.F. provided data from FAO FRA-2015 and contributed to the analysis. J.H. was supported by Leverhulme Foundation and EU FP7 through project LUC4C (GA603542).

Competing financial interests. The authors declare no competing financial interests.

References

- 1 UNFCCC. Adoption of the Paris Agreement. Report No. FCCC/CP/2015/L.9/Rev.1, <http://unfccc.int/resource/docs/2015/cop21/eng/l09r01.pdf>. (UNFCCC, 2015).
- 2 European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). Emission Database for Global Atmospheric Research (EDGAR), release version 4.3.1 <http://edgar.jrc.ec.europa.eu/overview.php?v=431>, 2016.
- 3 UNFCCC. INDCs as communicated by Parties, <http://www4.unfccc.int/submissions/indc/Submission%20Pages/submissions.aspx>. (UNFCCC, 2015).
- 4 UNFCCC. Paris Agreement - Status of Ratification. http://unfccc.int/paris_agreement/items/9444.php (UNFCCC 2016)
- 5 Ciais, P. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds T.F. Stocker *et al.*) Ch. 6, 465-522 (Cambridge University Press, 2013).
- 6 Smith, P. *et al.* in *Climate change 2014: mitigation of climate change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds O. Edenhofer *et al.*) Ch. 11, 811-886 (Cambridge University Press, 2014).
- 7 Tubiello, F. N. *et al.* The Contribution of Agriculture, Forestry and other Land Use activities to Global Warming, 1990-2012. *Global Change Biology* **21**, 2655-2660 (2015).
- 8 Federici, S., Tubiello, F. N., Salvatore, M., Jacobs, H. & Schmidhuber, J. New estimates of CO₂ forest emissions and removals: 1990-2015. *Forest Ecology and Management* **352**, 89-98 (2015).
- 9 Le Quéré, C. *et al.* Global Carbon Budget 2015. *Earth System Science Data* **7**, 349-396 (2015).
- 10 Pan, Y. *et al.* A large and persistent carbon sink in the world's forests. *Science* **333**, 988-993 (2011).

- 357 11 Schlamadinger, B. *et al.* Options for including land use in a climate agreement post-2012:
358 improving the Kyoto Protocol approach. *Environmental Science and Policy* **10**, 295-305
359 (2007).
- 360 12 UNFCCC. Decision 2/CP.17. Outcome of the work of the Ad Hoc Working Group on Long-
361 term Cooperative Action under the Convention, FCCC/CP/2011/9/Add.1,
362 <http://unfccc.int/resource/docs/2011/cop17/eng/09a01.pdf> (UNFCCC, 2011).
- 363 13 UNFCCC. Greenhouse Gas Inventories,
364 [http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8812.php)
365 [items/8812.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8812.php). (UNFCCC, 2015).
- 366 14 Grassi, G., den Elzen, M. G. J., Hof, A. F., Pilli, R. & Federici, S. The role of the land use,
367 land use change and forestry sector in achieving Annex I reduction pledges. *Climatic Change*,
368 1-9 (2012).
- 369 15 UNFCCC. National Communications Non-Annex 1, [http://unfccc.int/nationalreports/non-](http://unfccc.int/nationalreports/non-annexinatcom/submittednatcom/items/653.php)
370 [annexinatcom/submittednatcom/items/653.php](http://unfccc.int/nationalreports/non-annexinatcom/submittednatcom/items/653.php) (UNFCCC, 2015).
- 371 16 UNFCCC. Biennial Update Reports, [http://unfccc.int/national_reports/non-](http://unfccc.int/national_reports/non-annex_i_natcom/reporting_on_climate_change/items/8722.php)
372 [annex_i_natcom/reporting_on_climate_change/items/8722.php](http://unfccc.int/national_reports/non-annex_i_natcom/reporting_on_climate_change/items/8722.php) (UNFCCC, 2015).
- 373 17 WRI. CAIT Climate Data Explorer. Paris Contributions Map, <http://cait.wri.org/indc/>. (World
374 Resources Institute, 2015).
- 375 18 Admiraal, A. *et al.* Assessing Intended Nationally Determined Contributions to the Paris
376 climate agreement – what are the projected global and national emission levels for 2025–
377 2030? Report No. PBL 1879, [http://www.pbl.nl/en/publications/assessing-intended-](http://www.pbl.nl/en/publications/assessing-intended-nationally-determined-contributions-to-the-paris-climate-agreement)
378 [nationally-determined-contributions-to-the-paris-climate-agreement](http://www.pbl.nl/en/publications/assessing-intended-nationally-determined-contributions-to-the-paris-climate-agreement). (PBL, 2015).
- 379 19 Rogelj, J. *et al.* Paris Agreement climate proposals need a boost to keep warming well below
380 2 °C. *Nature* **534**, 631-639, doi:10.1038/nature18307 (2016).
- 381 20 UNFCCC. National Communications Annex 1,
382 <http://unfccc.int/nationalreports/annexinatcom/submittednatcom/items/7742.php>; (UNFCCC,
383 2015).

- 384 21 UNFCCC. Compilation of information on nationally appropriate mitigation actions to be
 385 implemented by Parties not included in Annex I to the Convention,
 386 FCCC/AWGLCA/2011/INF.1, <http://unfccc.int/resource/docs/2011/awglca14/eng/inf01.pdf>.
 387 (UNFCCC, 2011).
- 388 22 FAO-FRA. Country reports (2015), [http://www.fao.org/forest-resources-assessment/current-](http://www.fao.org/forest-resources-assessment/current-assessment/country-reports/en/)
 389 [assessment/country-reports/en/](http://www.fao.org/forest-resources-assessment/current-assessment/country-reports/en/). (Food and Agricultural Organization of the United Nations,
 390 2015).
- 391 23 FAOSTAT. Land use emissions. (Food and Agricultural Organization of the United Nations
 392 (FAO), Rome, Italy, http://faostat3.fao.org/download/G2/*/E, 2015).
- 393 24 Grassi, G., Monni, S., Federici, S., Achard, F., and Mollicone, D. Applying the
 394 conservativeness principle to REDD to deal with the uncertainties of the estimates.
 395 Environmental Research Letters, 3(3) (2008)
- 396 25 IPCC. in *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (eds H.S.
 397 Eggleston *et al.*) (National Greenhouse Gas Inventories Programme, Institute for Global
 398 Environmental Strategies, Hayama, Japan, 2006).
- 399 26 IPCC. in *Good Practice Guidance for Land Use, Land-Use Change and Forestry* (eds J.
 400 Penman *et al.*) (Institute for Global Environmental Strategies, 2003).
- 401 27 Houghton, R. A. *et al.* Carbon emissions from land use and land-cover change.
 402 *Biogeosciences* **9**, 5125-5142 (2012).
- 403 28 Hansis, E., Davis, S. J. & Pongratz, J. Relevance of methodological choices for accounting of
 404 land use change carbon fluxes. *Global Biogeochemical Cycles* **29**, 1230-1246 (2015).
- 405 29 Pongratz, J., Reick, C. H., Houghton, R. A. & House, J. I. Terminology as a key uncertainty
 406 in net land use and land cover change carbon flux estimates. *Earth System Dynamics* **5**, 177-
 407 195 (2014).
- 408 30 Federici, S. *et al.* GHG fluxes from forests: An assessment of national reporting and
 409 independent science in the context of the Paris Agreement,

410 <http://www.climateandlandusealliance.org/reports/ghg-fluxes-from-forests/> (Climate and
 411 Land Use Alliance, 2016).

412 31 IPCC. in *IPCC Meeting on Current Scientific Understanding of the Processes Affecting*
 413 *Terrestrial Carbon Stocks and Human Influences on them* (eds D. Schimel & M. Manning)
 414 (National Oceanic & Atmospheric Administration, 2003).

415 32 IPCC. in *Revisiting the use of managed land as a proxy for estimating national*
 416 *anthropogenic emissions and removals* (eds S Eggleston, N Srivastava, K Tanabe, & J
 417 Baasansuren) (IGES, 2010).

418 33 Erb, K. H. *et al.* Bias in the attribution of forest carbon sinks. *Nature Climate Change* **3**, 854-
 419 856 (2013).

420 34 UNFCCC. Synthesis report on the aggregate effect of the intended nationally determined
 421 contributions, FCCC/CP/2015/7, <http://unfccc.int/resource/docs/2015/cop21/eng/07.pdf>
 422 (UNFCCC, 2015).

423 35 UNEP. The Emissions Gap Report 2015,
 424 http://uneplive.unep.org/media/docs/theme/13/EGR_2015_301115_lores.pdf. 1-98 (United
 425 Nations Environment Programme, 2015).

426 36 Lapola, D. M. *et al.* Pervasive transition of the Brazilian land-use system. *Nature Clim.*
 427 *Change* **4**, 27-35 (2014).

428 37 Forsell, N. *et al.* Assessing the INDCs' land use, land use change, and forest emission
 429 projections. *Carbon Balance Manage* 11:26, doi: 10.1186/s13021-016-0068-3 (2016).

430 38 Houghton, R. A., Byers, B. & Nassikas, A. A. A role for tropical forests in stabilizing
 431 atmospheric CO₂. *Nature Climate Change* **5**, 1022-1023 (2015).

432 39 Mackey, B. *et al.* Untangling the confusion around land carbon science and climate change
 433 mitigation policy. *Nature Climate Change* **3**, 552-557 (2013).

434 40 Clarke, L. *et al.* in *Climate Change 2014: Mitigation of Climate Change. Contribution of*
 435 *Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
 436 *Change* (eds O. Edenhofer *et al.*) Ch. 6, 413-510 (Cambridge University Press, 2014).

437 41 Tollefson, J. Paris climate deal hinges on better carbon accountancy. *Nature* **529**, 450 (2016).

438 42 UNFCCC. Guidelines for the preparation of national communications from Parties not
439 included in Annex I to the Convention, Bonn, Decision 17/CP.8,
440 <http://unfccc.int/resource/docs/cop8/07a02.pdf> (UNFCCC, 2003).

441 43 GOFC-GOLD. A sourcebook of methods and procedures for monitoring and reporting
442 anthropogenic greenhouse gas emissions and removals associated with deforestation, gains
443 and losses of carbon stocks in forests remaining forests, and forestation, GOFC-GOLD
444 Report version COP21-1, University of Wageningen, the Netherlands,
445 <http://www.gofcgold.wur.nl/redd/>. (GOFC-GOLD, 2015).

446 44 GFOI. Integrating remote-sensing and ground-based observations for estimation of emissions
447 and removals of greenhouse gases in forests: methods and guidance from the Global Forest
448 Observations Initiative. (Global Forest Observations Initiative, 2014).

449 45 Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change.
450 *Science* **342**, 850-853 (2013).

451 46 Achard, F. & House, J. I. Reporting carbon losses from tropical deforestation with Pan-
452 tropical biomass maps. *Environmental Research Letters* **10**, doi:10.1088/1748-
453 9326/10/10/101002 (2015).

454 47 Global Forest Watch <http://www.globalforestwatch.org/>. (2016).

455 48 Pilli, R., Grassi, G., Kurz, W.A., Abad Viñas, R., Guerrero, N. Modelling forest carbon stock
456 changes as affected by harvest and natural disturbances. I. Comparison with countries'
457 estimates for forest management. *Carbon Balance Manage*, 11:5, doi: 10.1186/s13021-016-
458 0047-8 (2016)

459 49 Roman-Cuesta, R.M. *et al.* Multi-gas and multi-source comparisons of six land use emission
460 datasets and AFOLU estimates in the Fifth Assessment Report, for the tropics for 2000–2005.
461 *Biogeosciences*, **13**, 5799–5819 (2016).

462 50 Friedlingstein, P. *et al.* Update on CO₂ emissions. *Nature Geoscience* **3**, 811-812 (2010).

463 51 IPCC. in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group*
464 *I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds
465 T.F. Stocker *et al.*) Ch. SPM, 1-29 (Cambridge University Press, 2013).
466

METHODS

This analysis quantifies the mitigation role of Land Use, Land Use Change and Forestry (LULUCF, mainly forests), based on the (I)NDCs^{3,4} submitted by Parties in the context of the Paris Climate Agreement¹, complemented with information from other countries' official documents. This analysis does not aim to assess specific country policies or the quality of country data in comparison with independent sources.

Our analysis of LULUCF net emissions over time covered all 195 UNFCCC countries, with assumptions necessary in some cases (i.e. using the latest historical data where no (I)NDC projection was available, see below). However, due to constraints, our estimation of the LULUCF mitigation contribution was possible only for 68 countries (41 (I)NDCs), covering 83% of global forest area (based on FAO-FRA 2015²²). Other countries were not included either because LULUCF was not clearly included in the target or because the LULUCF contribution was not entirely clear or directly quantifiable (see Supplementary Section 1, Supplementary Information).

Our analysis is based on countries' documents submitted up to February 2016. However, the most relevant recalculations made by countries after that date and before December 2016 (e.g. Brazil, Indonesia and USA) are briefly discussed in the Supplementary Section 2.

Information used in this analysis

The methodological approach applied in this analysis required collecting information on:

- (i) Countries' historical data and projections up to 2030 (for all 195 UNFCCC countries), using countries' documents submitted up to end of February 2016, with the following priority: (I)NDCs³; other country data submitted to UNFCCC (2015 GHG Inventories¹³

(GHGI) for developed countries, and GHGIs included in recent National Communications^{15,20} (NC) and in Biennial Update Reports¹⁶ (BUR) for developing countries); other official countries' documents (e.g. ref.²¹); FAO-based datasets (for forests^{8,22} and non-forest emissions²³). Despite gaps in country reports (especially for non-forest land uses in developing countries), this priority is justified by the fact that country reports to UNFCCC are formally reviewed or technically assessed by UNFCCC (GHGIs of developed countries are formally reviewed annually, with biennial technical assessment for developing country inventories), and are the means by which countries assess their progress towards targets. Furthermore, FAO-FRA reports²² are not primarily for reporting CO₂ emissions and removals, while UNFCCC country reports specifically address emissions and removals. The range of historical country datasets (dotted line in Fig. 1) reflects alternative selections of country sources, i.e. GHGIs for developed countries (selected for both the lower and the upper range), plus FAO-based datasets (upper range) or NCs (lower range) for developing countries. This alternative selection assumes a high reliability of GHGIs for developed countries, while providing an idea of the impact of choosing only NCs (including old NCs) vs. FAO-based datasets for developing countries. See Supplementary Table 4 for an overview of historical datasets used.

For **historical data**, GHGIs with a time series from 1990 to 2013 were available for all developed countries, in most cases including Harvested Wood Products. For developing countries, data are from BURs when available or from latest NCs, typically not including Harvested Wood Products. When only few years were available (typically at least two between 1990 and 2010), 5 or 10 years averages were used. Sometimes, especially for older NCs, data from NCs contain ambiguities, or are not fully comparable across countries (e.g. while most countries implicitly report only emissions and removals from “managed forests”, in accordance with the recent IPCC guidance, a few countries include the sink from apparently unmanaged forests). To reduce the risk of using old or

inappropriate data, the more recent FAO datasets were used instead of NCs prior to 2010. Net emissions from forests (e.g., sink from forest management and emissions from deforestation) usually dominate the LULUCF fluxes in country reports, although in some case emissions from croplands and grasslands (rarely reported by developing countries) are also relevant, especially from organic soils.

Based on available information from countries' reports to UNFCCC complemented by expert judgment, we performed an analysis of the uncertainties for LULUCF absolute GHG net emissions (level) and for the associated trends (see Supplementary Section 3, Supplementary Table 7 and Supplementary Figures 2 and 3).

The FAO-based datasets include country data on forest carbon stock change from the Forest Resource Assessment (FAO-FRA 2015²², as elaborated by ref⁸) and FAOSTAT²³ data on country-level non-forest land use emissions (CO₂, CH₄ and N₂O from biomass fires, including peatlands fires, and from drainage of organic soils). The overall small difference between the FAO-FRA 2015 forest carbon stock data used in our analysis (based on ref.⁸) and the FRA-2015 forest carbon stock data included in FAOSTAT²³ is that the gap-filling methods differ (although for the biomass pools such difference does not impact the total net CO₂ emissions/removals across the time series), and that we include both living biomass (above and below-ground) and dead organic matter if reported by countries, while FAOSTAT only considers living biomass. Overall, for the historical period we only used FAO-based datasets to fill the gaps for a relative large number (60), but typically rather small developing countries (covering 11% of global forest area). The significant difference between this analysis and FAOSTAT (Fig. 2 of the paper) is due to several factors, including higher non-forest land use emissions in FAOSTAT for developing countries (especially in Indonesia, Sudan, Zambia) and higher forest land use emissions in FAOSTAT for both developing countries (e.g. Colombia, Liberia, Madagascar, Myanmar, Nigeria, Philippines, Zimbabwe) and developed ones (USA and Russia).

For **projections**, data from (I)NDCs (with some expert-judgment interpretation when needed), or NCs²⁰ were available for almost all developed countries. For developing countries, if no projection was available in the (I)NDCs, BURs or NCs, FAO-FRA 2015 country projections^{8,22} were used in few cases. Where no projection was available, the latest historical country data available were used (i.e. continuing the recent estimates).

While almost no country provided formal information on uncertainties in their projections, in the analysis of uncertainties (see Supplementary Section 3) we conservatively assumed that the uncertainties estimated for the past will hold for the future. In addition, the different scenarios that our analysis identified up to 2030 (Fig. 1) may also give an order of magnitude of the uncertainties. The range “LULUCF projections min-max” shown in Fig. 1 is slightly broader than the various scenarios (by about 500 MtCO₂e/y, or 0.5 GtCO₂e/y, in 2030) because in few cases countries provide a range of projections and not all the various projections can be associated with the four scenarios analyzed. The overall difference of about 500 MtCO₂e/y is essentially due to the range of projections from the US (the difference between the “high” and a “low” sequestration scenario in their latest National Communication amounts to 370 MtCO₂e/y in 2030), and due to Russia (the difference between the various sequestration scenario in their latest National Communication amounts to about 150 MtCO₂e/y in 2030).

With regards to the GHGs considered, this paper focuses on CO₂ emissions and removals and on available data on non-CO₂ emissions (CH₄ and N₂O), based on the information included in countries’ documents. National GHGIs are required to report on all GHGs, but in some developing countries the information on non-CO₂ gases is incomplete. Based on available information, and excluding agricultural emissions, the importance of non-CO₂ gases is typically small relative to the total GHG fluxes (see ref³⁰ for details), representing about 2-3% of total CO₂-equivalent forest flux, with slightly higher values found where forest fires are important in the overall GHG budget. This suggests that, when comparing different datasets (Fig. 2), the possible different coverage in the

(I)NDCs and other documents of non-CO₂ gases does not represent a major reason for discrepancy.

(ii) Type of mitigation target elaborated in each countries' (I)NDC (Supplementary Table 1), i.e. change in absolute emissions or intensity, either relative to a base year or to a BAU scenario (i.e. 2025 or 2030 scenario year); target 'unconditional' or 'conditional' (i.e. related to the provision of finance, technology or capacity-building support). (I)NDCs expressing only 'policies and measures' (without quantitative targets) were not taken into account.

(iii) Modality of inclusion of LULUCF within each countries' (I)NDC (Supplementary Table 1), i.e. it may be treated in the same way as other sectors (fully included as part of the overall target), or partially included (only forest activities), or considered separately with special mitigation actions and/or accounting rules.

Some additional expert evaluation was included where necessary.

(I)NDC cases

The (I)NDCs were classified into four '(I)NDC cases' (Supplementary Table 2). Based on the availability of country LULUCF information, enough information was found to assign 68 countries to these different "(I)NDC cases", and to quantify directly the expected LULUCF mitigation. These 68 countries include all countries with a major forest coverage and correspond to 78% of global emissions in 2012 (including LULUCF emissions and international aviation and marine emissions)².

Different mitigation perspectives

The quantification of the mitigation role of LULUCF has been undertaken using different approaches, reflecting different perspectives, according to the questions addressed (Supplementary Table 3).

596

597 **Estimation of LULUCF mitigation**

598 Whereas estimates for perspective ‘A’ (LULUCF net emissions over time) could be made for all 195
599 UNFCCC countries, the information needed for the LULUCF mitigation contribution under
600 perspectives ‘B’ ((I)NDC compared to alternative future scenarios) and ‘C’ (country perspective on
601 calculating emissions reduction (I)NDC) was available only for the 68 countries (41 (I)NDCs)
602 included in Supplementary Table 1. For the remaining countries, the additional mitigation in
603 perspectives ‘B’ and ‘C’ were assumed to be zero relative to other sectors. This assumption is
604 probably conservative (see Supplementary Section 1).

605 Based on the four (I)NDC cases (Supplementary Table 2), and using the available country
606 information (generally with limited expert judgment), this analysis quantified the LULUCF mitigation
607 perspectives (Supplementary Table 3) following the method illustrated in Supplementary Fig. 1. In
608 the very few cases where the target is expressed for 2025, we assumed that the same target applies to
609 2030, allowing us to sum up all the countries’ contribution to 2030.

610

611 **Contribution of the land sector to mitigation activity across all sectors**

612 The LULUCF mitigation perspectives ‘B’ and ‘C’ were compared to the expected (I)NDC mitigation
613 efforts across all sectors, for each country and at a global level. The global-level all-sectors ‘pre-
614 (I)NDC’ and ‘(I)NDC unconditional + conditional’ are taken from UNEP³⁵. All-sector emissions at
615 the ‘reference point’ (i.e. base year or BAU scenario for target year 2025 or 2030) are from: (i)
616 countries or (ii) from ref¹⁸ (for the BAU estimates for China and India). These two sources of
617 information were sufficient for countries representing 87% of global GHG emission in 2012.
618 Emissions for the remaining countries were approximated by assuming the same ratio of emissions at
619 reference point (i.e. estimates from available sources were multiplied by 100/87).

620

Comparison of this analysis with IPCC AR5

In order to make a meaningful comparison of country data (this analysis) with IPCC AR5^{5,6}, we disaggregated country data between “land converted to another land use” and “land remaining under the same land use”. While this disaggregation was directly available in all developed country reports, and was largely available for the most important developing countries (e.g. Brazil, Indonesia, India, China, Mexico), for the remaining developing countries information was generally available only for deforestation. In these cases, unless specified otherwise, the other emissions and removals were assigned to “land remaining under the same land use”.

Data availability

This study is primarily based on countries’ (I)NDCs^{3,4} and other GHG reports submitted to UNFCCC^{13,15,16,20,21}, complemented by FAO-based datasets^{8,22,23}. A large part of elaborated data used to support our findings are available in the Supplementary Information, including:

- (i) *Country-specific information* for 68 countries (41 (I)NDCs), in terms of general features of the (I)NDCs (Supplementary Tables 1 and 2) and of data and sources of information of LULUCF net emissions for the historical period 1990-2010 and for 2030, as expected for unconditional and conditional (I)NDC targets (Supplementary Table 5).
- (ii) *Aggregated information* on uncertainties (Supplementary Figures 2 and 3), on LULUCF mitigation perspectives (Supplementary Table 3) and on LULUCF net emissions (Supplementary Table 6).

Any other raw or elaborated data used in this study are available from the corresponding author upon request.