Mittal, T., Atack, N., Naish, N. E., Williams, J., Puryer, J., Sandy, J., & Ireland, A. (2018). The aberrant canine part 2: treatment. *Orthodontic Update*, 11(1), 6-12. <http://www.orthodontic-update.co.uk/issueArticle.asp?aKey=208>

Peer reviewed version

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via George Warman Publications at <http://www.orthodontic-update.co.uk/issueArticle.asp?ec=av+1&aKey=208>. Please refer to any applicable terms of use of the publisher.
Educational use only

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Generic Heading: Orthodontics

Title: The Aberrant Canine Part 2: Treatment

Authors:

TK Mittal

BDS, Specialty Registrar, Derriford Hospital, Plymouth

NE Atack

BDS, MSc, MOrth RCS, FDS RCS, Consultant Orthodontist, Musgrove Park Hospital, Taunton and School of Oral and Dental Sciences, University of Bristol

H Naish

BSc, BDS, MFDS RCS(Ed), MOrth RCS(Ed) Specialist Practitioner, Cathedral Orthodontics, Cardiff

JC Williams

BDS, MSc, DDS, MOrth RCS, FGDP, Academic Clinical Lecturer in Orthodontics, School of Oral and Dental Sciences, University of Bristol

J Puryer

BDS, DPDS, FDS RCS(Eng), PGDip, FHEA, Clinical Lecturer in Restorative Dentistry, School of Oral and Dental Sciences, University of Bristol

JR Sandy

BDS, MSc, PhD, MOrth RCS, FDS RCS, FDS RCS(Ed), FFD RCS, Professor of Orthodontics and Dean of Health Sciences, University of Bristol

AJ Ireland

BDS, MSc, PhD, MOrth RCS, FDS RCS, FHEA Professor of Orthodontics, School of Oral and Dental Sciences, University of Bristol,

The Aberrant Canine Part2: Treatment

Abstract: The eruption of the permanent canine, particularly the maxillary tooth is a milestone in dental development. Although often uneventful, occasionally there are disturbances in eruption, the management of which can be one of the more challenging aspects of orthodontics. This article gives an overview of the possible treatments of the aberrant canine tooth.

Clinical Relevance: Awareness of the possible sequelae of unerupted canines is important in diagnosis and treatment planning.

Objectives statement: To inform the reader of possible treatment of the aberrant canine.

1. Introduction

The permanent canine usually erupts uneventfully, but occasionally it may fail to do so.

When this occurs there is a potential for the adjacent teeth to be damaged. Even when it does not cause any damage, treatment of the ectopically positioned canine can present a substantial challenge to the orthodontist. Part 1 addressed the aetiology and diagnosis of the aberrant canine. Part 2 will discuss the various treatment options for the aberrant canine tooth.

2. Treatment of the aberrant canine

Following clinical and radiographic investigations, treatment decisions can be made based not only on the type of malocclusion, but more particularly on the presenting features associated with the aberrant canine. It is the management of this canine that will now be described.

2.1 Treatment for the developmentally absent canine

The treatment options for the developmentally absent canine are either to accept the resulting malocclusion, to reopen the space prior to prosthetic replacement, or to close the space.

Space closure may be prudent, especially if a good contact can be made between the lateral incisor and first premolar, which is not always possible in the lower arch. The first premolar can provide an aesthetic camouflage solution for the missing canine, but in the upper arch may require buccal root torque, mesial rotation and selective grinding of the palatal cusp to achieve the desired aesthetic outcome (Figure 1)

Figure 1 - A DPT of a patient where upper right canine is developmentally absent. The upper right deciduous canine was extracted and the space was closed with orthodontic fixed appliances. Notice how the gingival heights of the upper left canine and upper right first premolar are not the same

2.2 Treatment of the impacted unerupted canine

When the canine is present but unerupted, there are again a number of different treatment options available, ranging from accept and monitor, through to simple interceptive procedures and finally more complex orthodontic treatments.

2.2.1 Accept and monitor

If the canine is extremely ectopically positioned it may not be wise to attempt alignment as this is likely to be a lengthy and challenging treatment for both the patient and the

orthodontist. The risks of treatment may also outweigh the potential benefits. If the unerupted canine is left *in situ*, it should be monitored with a radiograph every couple of years, as there is a risk of resorption of the roots of the adjacent teeth (up until 14 years of age), and less commonly cystic change within the follicle of the canine.

2.2.1.1 Cystic change within the follicle

A rare complication (0.4%) of leaving an unerupted tooth *in situ* is the risk that the follicle undergoes cystic change¹. In a study by Mourshed¹ cystic change was defined as a visible pericoronal space within the follicle, larger than 2.5mm in diameter on radiographic imaging. If there are signs of cystic change the canine should either be removed or aligned within the arch. If left untreated it can lead to the canine becoming more ectopically positioned or result in movement of adjacent erupted teeth.

2.2.1.2 Resorption of adjacent teeth and subsequent treatment

Resorption of the roots of the adjacent teeth is a more common consequence of aberrantly positioned unerupted teeth. Root resorption will only occur if the crown of the unerupted tooth is in close proximity to the adjacent tooth roots (*i.e.* in the line of the arch or close to it) and results from direct tooth contact rather than under pressure from an enlarging follicle². It occurs more commonly when the canine is closer to the midline and has advanced root formation³. The incidence of resorption of lateral incisors, in the presence of palatally ectopic maxillary canines, is reported to be 12% on plain film⁴ (Figure 2). However, the incidence of root resorption being detected on CBCT ranges from 38% to 66%^{5, 2, 6}, as CBCT will permit a full 3D assessment of the lateral incisor roots (Figure 4). Resorption commonly occurs in the middle and apical thirds of the lateral incisor and is often classified

as slight ($< \frac{1}{2}$ dentine thickness), moderate ($> \frac{1}{2}$ dentine thickness) or severe (pulp exposure).

If no resorption has occurred by the age of 14, it is unlikely to occur⁵. Resorption may lead to increased mobility of the adjacent teeth and hence be a clinical indicator of canine position.

If an aberrant canine tooth is causing resorption to either the lateral or central incisor, the long-term prognosis of the damaged tooth should be assessed, along with the chances of canine alignment. A tooth with extensive resorption may not have a good long term prognosis and it may be prudent to sacrifice it in order to allow the canine to be aligned and camouflaged. If the canine is so far mesial that it has caused resorption of the central incisor, it is unlikely that alignment to the correct position will be successful. In this instance the aberrant canine should either be extracted, or if resorption is extensive, aligned in the position of the central incisor and camouflaged (Figure 3). Alignment may possibly occur spontaneously if the resorbed tooth is extracted, or it may require active alignment with fixed appliances. It is not just aberrant palatal canines that can cause extensive root resorption of adjacent teeth, it can also occur with buccally positioned maxillary canines⁷.

Figure 2 – Anterior Occlusal radiograph showing extensive resorption of the lateral incisor root. There has been so much resorption of the lateral incisor that the canine tip can just be seen erupting at the palatal gingival margin

Figure 3 - The upper left canine has caused significant resorption of the upper left central incisor. Following alignment of the canine into the central incisor position it can be seen that the gingival margin is at a good level. The canine crown can now be camouflaged as a central incisor.

2.2.2 Extraction of the ectopic canine

In instances where there is an unerupted upper or lower ectopic canine and extraction is indicated (Figure 4), it is important to be aware of potential surgical complications. Risks include swelling, bruising, post-operative paraesthesia, damage to adjacent teeth and bone loss during the extraction^{8,9}. There is little published literature reporting the incidence of damage to the roots of adjacent teeth during the removal of impacted canines. If the aberrant canine has caused resorption of an adjacent tooth, the crown will be in direct contact with the roots². Common sense dictates that the removal of such teeth will have increased risk of damage to the already resorbed adjacent tooth roots, due to their close proximity.

When the aberrant canine is extracted, the occlusion might be accepted, the space can be opened for prosthetic replacement or it can be closed as in the case of the developmentally absent canine (Section 2.1)

Figure 4 – The lower left canine has transmigrated to the symphyseal region and is not amenable to alignment. Extraction and space closure is the only option for this malocclusion

2.2.3 Alignment of the palatally placed canine

The vast majority of unerupted maxillary canines are palatally ectopic⁴ and The Royal College of Surgeons of England have developed guidelines dealing with the management of these teeth¹⁰. Treatment will depend not only on the position of the ectopic canine, but also the age of the patient and their stage of dental development. Suggested treatment can

range from simple interceptive extractions to surgical exposure of the canine and orthodontic alignment.

2.2.3.1 Interceptive extraction of the deciduous canine

Simple interceptive extraction of the deciduous canine, provided the ectopic permanent canine is favourably positioned, will hopefully result in successful eruption of the permanent tooth. This is reported to be successful in up to 80% of cases in the maxillary arch, provided sufficient space already exists¹¹. Power and Short¹² found that, in the presence of crowding, only 62% of canines erupted but a further 19% still improved position following interceptive deciduous extraction. The difference between these two studies highlights the importance of space being available for the unerupted canine (Figure 5). Although a Cochrane review concluded that there was no high quality evidence supporting the interceptive extraction of deciduous canines in such cases¹³, a more recently published RCT has provided evidence supporting the practice of interceptive deciduous canine extractions¹⁴.

2.2.3.2 Interceptive extraction of the deciduous canine and space provision

Power and Short¹² highlighted the effect crowding has on the eruption of ectopic canines following deciduous extractions. There have since been several investigations into the effect of space provision on ectopic canine eruption. Olive¹⁵ used fixed appliances to provide space for the canine following interceptive extraction and found that 75% erupted into a midalveolar position, and nearly all improved in position. Baccetti and his co-workers^{16, 17} used headgear to create space, also finding that a high proportion of ectopic canines successfully erupted into the arch. However, space gained through expansion did not significantly increase successful eruption rates when compared to the use of headgear¹⁸.

or even just a space maintainer¹⁹. Figure 6 illustrates a case where space was created for an aberrant maxillary canine, which subsequently erupted into the line of the arch

Figure 5 - Interceptive extraction of the maxillary deciduous canines has allowed the maxillary permanent canines to improve in position.

Figure 6 - The upper left permanent canine is unerupted and palatally placed. The deciduous canine was extracted and sufficient space opened with a fixed appliance to permit the canine to erupt.

2.2.3.3 Exposure and alignment of the ectopic canine

Surgical exposure and alignment of the ectopic canine may be indicated when interceptive measures fail, or the tooth is in such an unfavourable position to begin with that it is bound to fail to erupt. There are two main surgical methods of exposure of ectopic canine, namely the Open and the Closed Techniques and these will be described first.

Open vs. Closed exposure

Open exposure technique²⁰ (Figure 7) requires the surgeon to expose the crown of the unerupted canine, allowing the orthodontist to directly place an attachment on the tooth at a later date. Closed exposure²¹ (Figure 8) requires the surgeon to expose the crown of the

canine, bond an attachment and gold chain and then replace the soft tissue flap. In this case the orthodontist can then place traction to the chain at a later date.

Open exposure and direct visualisation of the crown can aid the orthodontist in applying traction with appropriate force vector to move it into the line of the arch, but this may be impractical if the tooth is very ectopic. Some clinicians favour early open exposure of the tooth in order to promote autonomous eruption and disimpaction before the placement of fixed appliances^{22, 23}. However, if the exposure is insufficient, the mucosa may heal over the canine crown such that by the time the patient returns to the orthodontist, there is no sign of the unerupted canine in the mouth. In the UK there is a roughly even split between open and closed exposures²⁴.

Figure 7 – Open canine exposure

Figure 8 - The upper right canine has been exposed with a closed technique. The gold chain is attached to the fixed appliance with elastic thread.

There are advantages and disadvantages of both techniques as discussed by Mathews and Kokich²³ and by Becker and Chaushu²⁵ and these are summarised in Table 1. A Cochrane review in 2008²⁶ concluded that there were no studies comparing open and closed exposure that could be included in the review and therefore there was an insufficiency of evidence. As a result, the same author conducted a trial comparing open and closed exposure and reported that there were no differences in surgical outcome, operating time or patient related outcome measures (PROMs)²⁷ between the two techniques. Nor were there differences between the techniques in terms of the final periodontal condition of the exposed canines²⁸ and whether orthodontists could identify which technique was used, although they could identify an exposed tooth²⁹⁻³¹.

	Open Exposure	Closed Exposure
Advantages	May reduce total time with fixed appliances Autonomous eruption and disimpaction of the canine without active force Allows the periodontal ligament to establish before orthodontic tooth movement	Follicle can remain and influence movement Immediate traction May not be as sore postoperatively
Disadvantages	May be sore or require coverplate Patient must keep clean Surgeon may not remove enough bone around the canine crown. Follicle removed	Force vector applied to tooth 'blind' May drag tooth along root surfaces Chain rarely bonded buccally – tooth erupts rotated Tooth movement only through pressure necrosis (crown-bone contact)

Table 1: Advantages and disadvantages of open and closed canine exposure^{25, 23}

Alignment of the canine following exposure

Whatever method of exposure is carried out, the task of alignment of the canine still remains. The root surface area of the canine is substantial³² and therefore movement of this tooth is anchorage demanding, both vertically and horizontally.

Occasionally the ectopic canine may be the only substantial feature of the patient's malocclusion. In these instances traction can be applied to the exposed canine using a sectional TMA archwire attached to a transpalatal arch³³ (Figure 9) in the case of the upper, or a lingual arch in the case of a lower. This will allow majority of tooth movement required to align the canine without the need to place an appliance on the rest of the teeth. Once brought to the dental arch, finishing can be achieved with a full fixed appliance.

Alternatively in the upper arch an upper removable appliance can be used to apply vertical

traction to the canine and once it is in the mouth a fixed appliance can be fitted to fully align the tooth and treat the rest of the occlusion (Figure 10).

If a fixed appliance is required for correction of other features of the malocclusion, or to create space, alignment of the aberrant canine can be performed either with a ballista spring³⁴, elastic chain or ligature (Figure 11) or a nickel titanium (NiTi) sectional archwire “piggy back” (Figure 12). The use of a NiTi wire or ballista spring will have the advantage of providing relatively constant force to the canine compared to elastic, which will experience significant force decay³⁵. Once aligned, the canine may require labial root torque to restore the canine eminence and create a good gingival margin.

Figure 9 - Occlusal and buccal views of a TMA sectional being used to align a maxillary canine, supported by a transpalatal arch

Figure 10 – An upper removable appliance being used to apply traction to the aberrant maxillary canine

Figure 11 - Traction applied to the upper left canine with a 'slingshot' of elastic chain. A button is bonded to the palatal surface of the left canine. Note that the upper right canine has also been exposed with a closed technique and the chain attached to the appliance with elastomeric thread.

Figure 12 - Traction is being applied to the upper left canine using a sectional NiTi piggy back with a steel base archwire.

2.2.4 Alignment of the buccally placed canine

If the canine tooth is unerupted and palpable buccally, its failure to erupt is most commonly due to insufficient space however, these crowded teeth often erupt high in the buccal sulcus (Figure 13). In such cases when the canine cups tip is mesially positioned and the tooth is also mesially angulated, extraction of the first premolar will often allow the canine to drop down spontaneously into the line of the arch. As part of this treatment it may be necessary to fit a fixed or removable space maintainer (Figure 14). If the canine is distally angulated it is unlikely to improve a great deal following the premolar extraction, and correction with fixed appliances will be required. If significant root translation is necessary this can be extremely anchorage demanding. If there is already good contact between the lateral incisor and the first premolar, extraction of the canine may well be the treatment of choice.

Figure 13 - The upper left canine has managed to erupt but remains high and buccally excluded from the arch. The canine is favourably placed for alignment following removal of the left first premolar.

Figure 14 - There is insufficient space for the upper right (and left) canine. A removable space maintainer was been fitted prior to extraction of the first premolars. Space maintenance allowed eruption of both upper canines before commencing upper and lower fixed appliances.

2.2.5 Treatment of the transposed canine

A transposition can either be a true transposition, where the whole tooth is transposed with the adjacent tooth (Figure 15), or a pseudo transposition where only the crown is in the wrong place, with the root being in the correct position within the line of the arch (Figure 16). Whereas a pseudo transposition is often amenable to orthodontic treatment, a true

transposition can usually only be treated following the extraction of the transposed canine, or the adjacent tooth. This is because it would otherwise require movement of the transposed canine out of the alveolus and around the adjacent tooth. Extraction of the transposed tooth is often the treatment of choice in the crowded dentition and in the uncrowded dentition it may be wise to accept the transposition.

Figure 15 - A true transposition of the canine with the lateral incisor

Figure 16 – A pseudotransposition of the canine with the lateral incisor

3. Treatment duration

On average patients with bilateral impacted canines take 10 months longer to treat than orthodontic patients who do not have impacted canines³⁶. In the case of impacted canine patients the average length of treatment is slightly greater than 2 years^{36, 37}. Other factors that might increase treatment duration in such cases include the severity of the impaction and the age of the patient³⁸. This is perhaps supported by the finding that only 70% of adults with impacted canines are successfully treated when compared to a matched group of children³⁹. This may be due to ankylosis of the canine teeth and can occur in one third of patients referred for failed eruption of aberrant canines⁴⁰. Ankylosis of canine teeth following exposure is more likely with increased patient age, depth of impaction and a closed exposure technique⁴¹, but is not completely predictable.

4. Conclusion

Eruption of the canine is a significant milestone in dental development. Often this occurs without incidence, but the aberrant canine can be one of the most challenging treatment scenarios commonly faced by the orthodontist. Early detection and interceptive treatment for aberrant canines can reduce the risk of adverse events, such as resorption, and the complexity of definitive treatment. Clinicians should be aware of the possible sequelae of unerupted canines and the management strategies.

Acknowledgements

The Authors would like to thank Mrs Karen Drage (Consultant Orthodontist, Derriford Hospital, Plymouth) and Mr Christian Day (Consultant Orthodontist, Bristol Dental Hospital, Bristol) for the use of some of their clinical cases

References

1. Mourshed F. A roentgenographic study of dentigerous cysts: I. Incidence in a population sample. *Oral Surgery, Oral Medicine, Oral Pathology*. 1964;18(1):47-53.
2. Ericson S, Bjerklin K, Falahat B. Does the canine dental follicle cause resorption of permanent incisor roots? A computed tomographic study of erupting maxillary canines. *Angle Orthod*. 2002;72(2):95-104.
3. Ericson S, Kurol J. Resorption of maxillary lateral incisors caused by ectopic eruption of the canines: A clinical and radiographic analysis of predisposing factors. *American Journal of Orthodontics and Dentofacial Orthopedics*. 1988;94(6):503-13.
4. Ericson S, Kurol J. Radiographic examination of ectopically erupting maxillary canines. *American Journal of Orthodontics and Dentofacial Orthopedics*. 1987;91(6):483-92.
5. Ericson S, Kurol J. Resorption of Incisors After Ectopic Eruption of Maxillary Canines: A CT Study. *The Angle Orthodontist*. 2000;70(6):415-23.
6. Walker L, Enciso R, Mah J. Three-dimensional localization of maxillary canines with cone-beam computed tomography. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*. 2005;128(4):418-23.
7. Abela S, Day C, Attack N. The Unusual Loss of Upper Lateral Incisor Teeth. *Orthodontic Update*. 2009;2:87-9.
8. Alling CC, Catone GA. Management of impacted teeth. *Journal of Oral and Maxillofacial Surgery*. 1993;51(1):3-6.
9. Frank CA. TREATMENT OPTIONS FOR IMPACTED TEETH. *The Journal of the American Dental Association*. 2000;131(5):623-32.
10. Husain J, Burden D, McSherry P. The management of the palatally ectopic maxillary canine. *National Clinical Guidelines, Faculty of Dental Surgery, Royal College of Surgeons of England*. 2010.
11. Ericson S, Kurol J. Early treatment of palatally erupting maxillary canines by extraction of the primary canines. *The European Journal of Orthodontics*. 1988;10(4):283-95.
12. Power SM, Short MB. An investigation into the response of palatally displaced canines to the removal of deciduous canines and an assessment of factors contributing to favourable eruption. *British journal of orthodontics*. 1993;20(3):215-23.
13. Parkin N, Furness S, Shah A, Thind B, Marshman Z, Glenroy G, et al. Extraction of primary (baby) teeth for unerupted palatally displaced permanent canine teeth in children. *The Cochrane database of systematic reviews*. 2012;12:Cd004621.
14. Naoumova J, Kurol J, Kjellberg H. Extraction of the deciduous canine as an interceptive treatment in children with palatal displaced canines - part I: shall we extract the deciduous canine or not? *Eur J Orthod*. 2015;37(2):209-18.

15. Olive RJ. Orthodontic treatment of palatally impacted maxillary canines. *Australian orthodontic journal*. 2002;18(2):64-70.
16. Leonardi M, Armi P, Franchi L, Baccetti T. Two Interceptive Approaches to Palatally Displaced Canines: A Prospective Longitudinal Study. *The Angle Orthodontist*. 2004;74(5):581-6.
17. Baccetti T, Leonardi M, Armi P. A randomized clinical study of two interceptive approaches to palatally displaced canines. *The European Journal of Orthodontics*. 2008;30(4):381-5.
18. Armi P, Cozza P, Baccetti T. Effect of RME and headgear treatment on the eruption of palatally displaced canines: A randomized clinical study. *The Angle Orthodontist*. 2011;81(3):370-4.
19. Baccetti T, Sigler LM, McNamara JA. An RCT on treatment of palatally displaced canines with RME and/or a transpalatal arch. *The European Journal of Orthodontics*. 2011;33(6):601-7.
20. Lewis PD. Preorthodontic surgery in the treatment of impacted canines. *American Journal of Orthodontics*. 1971;60(4):382-97.
21. Kokich VG, Mathews DP. Surgical and orthodontic management of impacted teeth. *Dental Clinics of North America*. 1993;37(2):181-204.
22. Kokich VG. Surgical and orthodontic management of impacted maxillary canines. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*. 2004;126(3):278-83.
23. Mathews DP, Kokich VG. Palatally impacted canines: the case for preorthodontic uncovering and autonomous eruption. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*. 2013;143(4):450-8.
24. Spencer HR, Ramsey R, Ponduri S, Brennan PA. Exposure of unerupted palatal canines: A survey of current practice in the United Kingdom, and experience of a gingival-sparing procedure. *British Journal of Oral and Maxillofacial Surgery*. 2010;48(8):641-4.
25. Becker A, Chaushu S. Palatally impacted canines: The case for closed surgical exposure and immediate orthodontic traction. *American Journal of Orthodontics and Dentofacial Orthopedics*. 2013;143(4):451-9.
26. Parkin N, Benson PE, Thind B, Shah A. Open versus closed surgical exposure of canine teeth that are displaced in the roof of the mouth. *The Cochrane database of systematic reviews*. 2008;4.
27. Parkin NA, Deery C, Smith AM, Tinsley D, Sandler J, Benson PE. No difference in surgical outcomes between open and closed exposure of palatally displaced maxillary canines. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2012;70(9):2026-34.
28. Parkin NA, Milner RS, Deery C, Tinsley D, Smith A-M, Germain P, et al. Periodontal health of palatally displaced canines treated with open or closed surgical technique: A multicenter, randomized controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics*. 2013;144(2):176-84.

29. Woloshyn H, Artun J, Kennedy DB, Joondeph DR. Pulpal and periodontal reactions to orthodontic alignment of palatally impacted canines. *Angle Orthod.* 1994;64(4):257-64.
30. Schmidt AD, Kokich VG. Periodontal response to early uncovering, autonomous eruption, and orthodontic alignment of palatally impacted maxillary canines. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics.* 2007;131(4):449-55.
31. Parkin NA, Freeman JV, Deery C, Benson PE. Esthetic judgments of palatally displaced canines 3 months postdebond after surgical exposure with either a closed or an open technique. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2015;147(2):173-81.
32. Jepsen A. Root surface measurement and a method for x-ray determination of root surface area. *Acta Odontol Scand.* 1963;21:35-46.
33. Roberts-Harry DP, Harradine NW. A sectional approach to the alignment of ectopic maxillary canines. *British journal of orthodontics.* 1995;22(1):67-70.
34. Jacoby H. The 'ballista spring' system for impacted teeth. *Am J Orthod.* 1979;75(2):143-51.
35. Taloumis LJ, Smith TM, Hondrum SO, Lorton L. Force decay and deformation of orthodontic elastomeric ligatures. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics.* 1997;111(1):1-11.
36. Stewart JA, Heo G, Glover KE, Williamson PC, Lam EWN, Major PW. Factors that relate to treatment duration for patients with palatally impacted maxillary canines. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2001;119(3):216-25.
37. Fleming PS, Scott P, Heidari N, DiBiase AT. Influence of Radiographic Position of Ectopic Canines on the Duration of Orthodontic Treatment. *The Angle Orthodontist.* 2009;79(3):442-6.
38. Zuccati G, Ghobadlu J, Nieri M, Clauser C. Factors associated with the duration of forced eruption of impacted maxillary canines: A retrospective study. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2006;130(3):349-56.
39. Becker A, Chaushu S. Success rate and duration of orthodontic treatment for adult patients with palatally impacted maxillary canines. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2003;124(5):509-14.
40. Becker A, Chaushu G, Chaushu S. Analysis of failure in the treatment of impacted maxillary canines. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2010;137(6):743-54.
41. Koutzoglou SI, Kostaki A. Effect of surgical exposure technique, age, and grade of impaction on ankylosis of an impacted canine, and the effect of rapid palatal expansion on eruption: A prospective clinical study. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics.* 2013;143(3):342-52.