Watson, J. (2017). All tests can sometimes cause more harm than good. *BMJ*, 358(j4070). <https://doi.org/10.1136/bmj.j4070>

Publisher's PDF, also known as Version of record

Link to published version (if available):
[10.1136/bmj.j4070](https://doi.org/10.1136/bmj.j4070)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via BMJ at <http://www.bmj.com/content/358/bmj.j4070.full>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

LETTERS

NEW DIAGNOSTIC TESTS: MORE HARM THAN GOOD

All tests can sometimes cause more harm than good

Jessica C Watson *GP and NIHR doctoral research fellow*

Canyngge Hall, University of Bristol, Bristol BS8 2PS, UK

Hofmann and Welch warn of the downsides of diagnostic tests.¹ “All screening programmes do harm . . . [but] some do more good than harm at a reasonable cost.”² This provocative statement is arguably true of any test, not just screening or new diagnostic tests. The challenge for clinicians is to share decisions with patients—choosing the right test for the right patient at the right time.

Inflammatory marker tests, such as C reactive protein, are simple, inexpensive, and commonly used in both primary and secondary care.³ When we interviewed GPs about their use of inflammatory marker blood tests in primary care, all doctors mentioned potential harms as well as benefits of testing.⁴ These could be divided into several categories; harms of the test itself, workload and financial costs, medicalisation of a patient's problem, patient anxiety, downstream costs, and harms of overdiagnosis.

Harms from the test itself are often overlooked and include cost of patients' time off work, needle phobia, bruising, and vasovagal syncope. One GP described the psychological cost: “There is always this pass or fail thing . . . because it's looking inside your body, and there might be something wrong there. It's not something one should do lightly.”

In UK primary care, rising workload is a major concern,⁵ and many GPs in our study thought that time spent dealing with pathology results was often inefficient or wasteful. Medicalisation can occur when tests cause “somatic fixation,”⁶ with patients or clinicians focusing only on the biomedical aspects of complex illness. Patient anxiety is perhaps the hardest to capture yet most important harm of unnecessary testing. Anecdotal stories, such as Fran's story,⁷ can be powerful, but the overall cost is hard to estimate. We need further research

into the benefits and harms of diagnostic tests, not just for new diagnostic tests, but also for many commonly used “routine” tests.

Overuse of diagnostic tests might reflect a societal culture of increasing risk aversion and rising medical litigation, as well as a culture in medicine that “evinces a deep rooted unwillingness to acknowledge and embrace uncertainty.”⁸ Technological advances and new tests provide ever more information, yet without wisdom⁹ we risk making well people sick, rather than sick people well.

Competing interests: None declared.

- 1 Hofmann B, Welch HG. New diagnostic tests: more harm than good. *BMJ* 2017;358:j3314. doi:10.1136/bmj.j3314 pmid:28720607.
- 2 Gray JAM, Patnick J, Blanks RG. Maximising benefit and minimising harm of screening. *BMJ* 2008;358:480-3. doi:10.1136/bmj.39470.643218.94. pmid:18310003.
- 3 Watson J, Round A, Hamilton W. Raised inflammatory markers[published Online First: 2012/02/07]. *BMJ* 2012;358:e454. doi:10.1136/bmj.e454. pmid:22306478.
- 4 Watson J, de Salis I, Hamilton W, Salisbury C. 'I'm fishing really'—inflammatory marker testing in primary care: a qualitative study. *Br J Gen Pract* 2016;358:e200-6. doi:10.3399/bjgp16X683857. pmid:26852797.
- 5 Hobbs FDR, Bankhead C, Mukhtar T, et al. National Institute for Health Research School for Primary Care Research. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007-14. *Lancet* 2016;358:2323-30. doi:10.1016/S0140-6736(16)00620-6. pmid:27059888.
- 6 Biderman A, Yeheskel A, Herman J. Somatic fixation: the harm of healing. *Soc Sci Med* 2003;358:1135-8. doi:10.1016/S0277-9536(02)00108-9. pmid:12593884.
- 7 Fran's story. An experience of unnecessary medical tests. <https://fabnhsstuff.net/2016/02/10/pathology-optimisation/>
- 8 Simpkin AL, Schwartzstein RM. Tolerating uncertainty - the next medical revolution?*N Engl J Med* 2016;358:1713-5. doi:10.1056/NEJMp1606402. pmid:27806221.
- 9 Heath I. Information without wisdom. *BMJ* 2017;358:j3203. doi:10.1136/bmj.j3203. pmid:28679507.

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>