Taylor, D., Sumner, S., & Bentley, M. (2018). Social wasps as models to study the major evolutionary transition to superorganismality. *Current Opinion in Insect Science*, 28, 26-32.
<https://doi.org/10.1016/j.cois.2018.04.003>,
<https://doi.org/10.1016/j.cois.2018.04.003>

Peer reviewed version

Link to published version (if available):

[10.1016/j.cois.2018.04.003](https://doi.org/10.1016/j.cois.2018.04.003)
[10.1016/j.cois.2018.04.003](https://doi.org/10.1016/j.cois.2018.04.003)

[Link to publication record in Explore Bristol Research](#)

PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via [insert publisher name] at [insert hyperlink] . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Social wasps as models to study the major evolutionary transition to superorganismality

Daisy Taylor¹, Michael A. Bentley² and Seirian Sumner^{2*}

¹ Life Sciences Building, University of Bristol, 24 Tyndall Avenue, Bristol, BS8 1TQ, United Kingdom

² Centre for Biodiversity & Environment Research, Department of Genetics, Evolution & Environment, University College London, Gower Street, London, WC1E 6BT, United Kingdom

*corresponding author: s.sumner@ucl.ac.uk

Abstract

The major evolutionary transition to superorganismality has taken place several times in the insects. Although there has been much consideration of the ultimate evolutionary explanations for superorganismality, we know relatively little about what proximate mechanisms constrain or promote this major transition. Here we propose that Vespid wasps represent an understudied, but potentially very useful, model system for studying the mechanisms underpinning superorganismality. We highlight how there is an abundance of behavioural data for many wasp species, confirming their utility in studies of social evolution; however, there is a sparsity of genomic data from which we can test proximate and ultimate hypotheses on this major evolutionary transition.

Box 1: Glossary of terms

Superorganism: A concept proposed by Wheeler [1] to suggest that some social insect colonies represent an entirely new type of higher-level organism. The idea behind this is that irreversible caste differentiation in social insects is akin to the germline and soma split among cells of multicellular organisms. See [2] for full review.

Vespidae: The family Vespidae (order: Hymenoptera, suborder: Aculeata, superfamily: Vespoidea) contains the subfamilies Vespinae, Polistinae, Stenogastrinae, Eumeninae and Masarinae [3,4]. Of these subfamilies, only the Vespinae, Polistinae and Stenogastrinae exhibit sociality.

Inclusive fitness theory: A framework focussing on how the reproductive interests of individuals depend both upon the impact of their behaviour on their own reproductive success, (direct fitness effects) and on that of individuals to whom they are related (indirect fitness effects) [5,6]. Often used to explain adaptations to social environments [7,8].

23

24 Introduction

25 Life on earth has been shaped by a series of rare but important events termed major evolutionary
26 transitions [9]. The hallmark of a major transition is a change in the way that biological information
27 is stored and transmitted, leading to a new level of biological organization in life's hierarchy. Major
28 transitions are predicted to develop in a series of stages under specific types of ecological
29 conditions [10,11] but we know little about the mechanisms by which they occur. In recent years
30 sociogenomics (the study of sociality in molecular terms [12]) has transformed our understanding
31 of the molecular basis of sociality, raising the likelihood that we will soon be able to understand the
32 kinds of molecular mechanisms that promote or constrain major transitions.

33

34 The evolution of superorganismality (Box 1) among some social hymenopteran (bees, wasps and
35 ants) insect lineages is one of the most striking examples of a major transition [1]. Whilst there has
36 been much empirical research on insect species that have already undergone the major transition
37 to superorganismality (e.g. the honeybee *Apis mellifera*, the fire ant *Solenopsis invicta*), we still
38 know relatively little about what happens at the molecular level during the earliest and intermediate
39 stages of a major transition [13,14]. Here, we make the case that Vespid wasps (Box 1) provide
40 unique opportunities for testing hypotheses about the molecular mechanisms underlying the
41 transitional stages in a major evolutionary transition.

42

The importance of studying the early and intermediate stages of the major transition to superorganismality

A major evolutionary transition is predicted to occur in several stages [10,11]. Firstly, natural selection must favour lower-level entities to come together to form a cooperative group. Secondly, a cooperative group must be irreversibly transformed into a cohesive whole that can be considered a new higher-level entity or 'individual'. The process of a major transition therefore marks important changes in the level at which natural selection predominantly acts, and the types of adaptations it is predicted to give rise to.

51

Traditionally, most research on major transitions has focused on identifying the ultimate evolutionary explanations for their origin [10,11]; in recent years there have been a growing number of empirical studies attempting to understand their underlying proximate mechanisms [13,15]. To date, these empirical studies in insects have, however, tended to focus on species which have already undergone a major transition to superorganismality, e.g. [16–18]. A potential limitation of these studies is that the kinds of adaptations that occur before the transition are unlikely to be the same as those that happen after the transition [2]. Ancestral traits that were present before the major transition to superorganismality may have been lost, altered, or masked by the emergence of novel traits once a major transition has occurred. It may be the case, therefore, that we need to look to species exhibiting characteristics of being at early or intermediate stages in the evolution of superorganismality, rather than solely superorganisms themselves, if our goal is to understand the mechanistic details underlying this major transition. Figure 1 highlights how proximate mechanisms and the stages of the transition to eusociality may evolve. It is important, however, to stress that our suggestion here is not that species exhibiting these characteristics are necessarily on an evolutionary trajectory towards greater complexity, but rather that these species might provide important clues about the mechanisms that were present in the ancestors of modern day superorganisms.

69

Vespid wasps as a model system

Here we discuss how the Vespidae provide an excellent phylogenetic context for testing hypotheses about the major transition to superorganismality. Wasps are a relatively understudied group, but they play important ecological roles and exhibit a remarkable diversity in social complexity, from species with the simplest of social groups (where all individuals can reproduce,

but some act as helpers; e.g. *Polistes* paper wasps) to species with the most complex societies in which the colony can be considered an individual (super)organism in its own right (where division of labour is fixed during development, irreversibly; e.g. *Vespula* yellow-jacket wasps) [4].

Crucially, new genetic data has confirmed that sociality has evolved twice in the Vespidae; once in the Stenogastrinae and once in the sister group Vespinae + Polistinae [3,19]. This recent revelation brings the Vespidae into sharp focus as a model group for understanding the evolution of sociality as they provide two independent evolutionary events of the same set of innovations [4,20]. Moreover, there are many species from both lineages, representing different stages, whose ecology and behaviour have been well studied. These species and the innovations they display provide exciting opportunities for future sociogenomic research, and the potential to help reconcile ultimate and proximate explanations for the major evolutionary transition to superorganismality.

In Table 1 we provide examples of social behaviours in Vespidae that may represent important adaptations in the early and intermediate stages of the major transition to superorganismality. We highlight how little is known about the underlying molecular mechanisms of these behaviours.

Social innovations in Vespid wasps

Evolution of group living from a solitary, ancestral state

At the very early stages in social evolution is the shift of individuals coming together to form groups [10,11]. From the highly plastic, solitary phenotype, in which individuals display the full repertoire of behavioural traits, the first conversion to sociality involves a decoupling of behaviours between the reproductive ('queens') and those that delay or abstain from reproduction altogether ('workers') [21]. Facultatively social species, such as members of the Stenogastrinae [22,23], are excellent models for studying this innovation as all individuals have the ability to initiate their own nest and become the reproductive. There is, however, usually a single dominant egg-layer in these societies [24,25]; most female offspring choose to remain on their natal nest as adults, as non-egg-laying helpers [22] although they retain the capacity to develop their ovaries throughout their life, and thus the option to disperse and found a nest alone. This reproductive strategy is likely to have arisen due to ecological constraints on independent founding [24], and the high fitness payoffs from indirect reproduction [24,25]. An outstanding question (which has been broached in halictid bees [26]), is what molecular processes are required for the shift from solitary living to facultative group living?

107 Most simply, differences in the timing of expression of the molecular processes regulating the
108 reproductive and provisioning subsets of the solitary phenotype could explain this innovation;
109 alternatively modification in the use of genes or gene networks may be required [12–15] (Figure 1).

110

111 Evolution of altruistic behaviour from ancestral, selfish behaviour

112 From an ultimate perspective, the evolution into group living involves a shift from direct to indirect
113 fitness for non-reproductive group members (Box 1). This shift is best studied in the very simple
114 social societies, as exhibited by the Stenogastrinae [24,27] and some Polistinae [28–30], where
115 helpers can inherit the nest when the dominant females dies. When this happens, the inclusive
116 fitness interests of an individual changes from indirect (when they refrain from reproducing and
117 engage in the cooperative brood care of siblings) to direct (when they produce their own offspring).
118 Who becomes queen during this process is determined by a reproductive hierarchy which can
119 either be age- [29,31], or dominance-based [29]. Individuals who never get the opportunity to
120 inherit the nest are said to behave ‘altruistically’ [8], since their inclusive fitness interests can only
121 be met through indirect fitness benefits. Hamilton [6] explained how natural selection can lead to
122 the evolution of such behaviours (reviewed in [8]), but we continue to know very little about their
123 molecular basis [12]. The search for genes encoding altruistic behaviours has been attempted in
124 several species, including the honeybee, ants and termites (reviewed in [32]). In the context of
125 understanding the processes by which the major transition to superorganismality arises, however,
126 we need to look for evidence of a molecular basis of altruism in the first stages of group living,
127 where altruism first evolves.

128

129 Loss of reproductive plasticity

130 One of the hallmarks of the major transition to superorganismality is the loss of plasticity with
131 respect to reproduction [2]; specifically, this is the evolution of a commitment to reproductive
132 potency (as a queen) or reproductive sterility (as a worker). In the simplest social groups,
133 individuals retain both behavioural totipotency, and reproductive potential. The interesting question,
134 therefore, is when, why and how a loss of reproductive plasticity arises. During the major
135 transition, the point at which plasticity is lost (or ‘commitment’ [2]) is pushed back in the
136 developmental time of individuals, from adulthood (e.g. in species that show a loss of reproductive
137 potential *during* their adult life, as in some Polistinae [33,34]) to brood development (e.g. in species
138 with pre-imaginal caste determination, effective sterility is committed during larval development in

139 the Vespinae wasps). To understand the process by which plasticity is lost at the outset of this
140 innovation, species which exhibit a loss in reproductive potential within adult life are potentially
141 important models. Documentation of such losses are rare as they require behavioural experiments
142 to test of reproductive capacity; however, there is evidence of this in *Metapolybia* sp. and also
143 *Polistes canadensis*, where replacement queens are drawn from the youngest females on the nest,
144 and where experiments show that females are incapable of taking advantage of egg-laying
145 opportunities as they age [28,34]. This loss of plasticity, and of reproductive potential, marks the
146 mechanistic basis for the initial shift from direct to indirect fitness (Box 1), and transcriptomic data
147 shows that differences between queens and workers significantly increase at this stage (Taylor et
148 al. unpublished). Analyses of the molecular processes that regulate this loss of plasticity during
149 adulthood would provide insights into the first stages of this critical innovation. These changes may
150 involve epigenetic processes that impose limitations on the reversibility in the expression of
151 molecular pathways [35].

152

153 Pre-imaginal caste differences

154 The most recognisable, and most studied, shift in sociality is the so-called 'point of no return' at
155 which true superorganismality emerges [2]; distinct morphological castes are exhibited in species
156 living in complex societies of wasps, such as some Polistinae [36–38] and Vespinae [39]. This
157 innovation marks the commitment of individuals during development to a fixed reproductive or non-
158 reproductive role, where ultimate investments in direct or indirect fitness are sealed by mechanistic
159 processes (Figure 1). The nutritional basis of caste-determination has been studied in species
160 across all levels of social complexity (reviewed in [40]), with the differential feeding of queens and
161 workers causing significant molecular changes in gene expression [41], and the epigenetic
162 mechanisms that regulate gene expression [17,42]. While the evolution of pre-imaginal caste
163 differentiation is well studied in the bees and ants, a long-standing question is whether molecular
164 processes underpinning fixed caste commitment are the same across social lineages and
165 ecological life-histories [13,43]. Remarkably, we lack any large-scale studies on the molecular
166 basis of pre-imaginal caste differentiation in the wasps. The Vespines are the obvious model clade
167 for this analysis, where worker reproductive potential is limited by the inability to mate and strong
168 differences in morphology, physiology and behaviour. The Polistinae are less well studied, but
169 some species show evidence of allometric caste differentiation (e.g. *Agelaia* and *Brachygastra*
170 [37,38]), suggesting pre-imaginal commitment of individuals to specific social roles. The gradient of

171 pre-imaginal caste differentiation in the Polistines therefore may provide valuable insights into the
172 *process* of this key innovation.

173

174 *Mutual dependence in the life cycle - Evolution of swarm founding behaviour*

175 Independent nest founding is the ancestral state of social insects. This requires significant levels of
176 behavioural plasticity from the founding queen, as she is required to found, build and maintain a
177 nest, and provision her brood. A key alteration once superorganismality has evolved, therefore, is
178 the reduced ability by reproductive females to independently found a nest, and instead evolve
179 mutual dependency on group founding (with a cohort of workers or other potential reproductives),
180 as a swarm. Swarm-founding is a key trait in many bee species (e.g. honeybees; stingless bees)
181 where a single queen leaves the natal nest to found a new nest, accompanied by a group of
182 workers. In ants another form of reduction in behavioural plasticity is found in the claustrally-
183 founding queens, who must build a nest, but then use their own body's resources to raise the first
184 brood, thus precluding the need to provision. In the wasps, swarm-founding is a key behavioural
185 trait of the Epiponini in the Polistinae [4,44,45]. In multi-queen colonies, swarm founding has the
186 benefit of maintaining high relatedness through cyclical oligyny, whereby only a single queen is
187 present at the time of sexual reproduction [46]. Swarm founding is obligatory in these species: the
188 queens have lost the ancestral behavioural repertoire required to successfully initiate their own
189 nest. This behaviour may therefore have evolved as: a) a novel trait to maintain high relatedness
190 within the colony, and b) a loss of ancestral plasticity and complete co-dependence of queen and
191 worker castes. This trait may therefore be underlain by both novel genes for swarm-founding
192 behaviour (as seen in *Apis* spp. [16]) (Figure 1) and regulation by the epigenetic processes which
193 limit behavioural plasticity [35].

194

195 *Mutual dependence in the life cycle - Reproductive control*

196 Reproductive control is determined through queen-worker communication across all levels of social
197 complexity (see [47,48] for reviews). It differs, however, significantly with group size, and by
198 extension, social complexity; in small, simple societies, queens retain dominance through
199 aggressive interactions and visual cues, while in large, complex societies, suppression of worker
200 reproduction is controlled by queen pheromones [49–52]. The hydrocarbons which act as a
201 sterility-inducing queen pheromone are highly conserved [53]. This same pheromone has been
202 found to also signal egg maternity, enabling workers to destroy worker-laid eggs in a selective

manner (worker policing) [54]. Worker policing occurs in species with multiply-mated queens because a worker force that is derived from two or more fathers is more closely related (on average) to the male offspring of their mother (brothers) than they are to the average worker-laid male [55]; worker policing can also evolve in singly-mated species if it increases colony efficiency [49,55]. Multiple mating and worker policing has evolved in many species of bees and ants [56,57]. To date, however, there have been no genomic studies examining the molecular processes by which these innovations have evolved. Among the wasps, the Vespinae are the only group known to be multiply-mated: within this clade, there are species that exhibit low (e.g. *Vespa crabro* [58]) and high (e.g. *Vespula* sp. [59]) levels of multiple mating, and associated levels of worker policing. This group is therefore an excellent model system for studying the molecular processes that regulate the innovations of multiple mating and worker policing.

214

215 **Conclusions**

216

The last few years have heralded significant theoretical and empirical progress in understanding the evolution of superorganismality [2,60], yet we still lack information on the mechanisms which precede the major transition. Here we summarise examples of life-history innovations, identified from behavioural studies, at which we would expect genomic changes (e.g. shifts in gene transcription networks; rewiring of genes; *de novo* evolution of genomic novelty) to be important; we put forward the Vespid wasps as a key group that would facilitate a molecular dissection of the process in a major evolutionary transition. These insights are likely to be of general relevance to social evolution at other levels of biological organisation, e.g. the evolution of multi-cellularity.

225

226 **Acknowledgements**

227 The authors are funded by NERC Grant NE/M012913/2.

228

229 **References**

- 230 1. Wheeler WM: **The ant-colony as an organism.** *J Morphol* 1911, **22**:307–325.
- 231 2. Boomsma JJ, Gawne R: **Superorganismality and caste differentiation as points of no**
232 **return: How the major evolutionary transitions were lost in translation.** *Biol Rev* 2017,
233 doi:10.1111/brv.12330.

- 234 3. Bank S, Sann M, Mayer C, Meusemann K, Donath A, Podsiadlowski L, Kozlov A, Petersen
235 M, Krogmann L, Meier R, et al.: **Molecular phylogenetics and evolution transcriptome**
236 **and target DNA enrichment sequence data provide new insights into the phylogeny of**
237 **vespid wasps (Hymenoptera: Aculeata: Vespidae).** *Mol Phylogenet Evol* 2017, **116**:213–
238 226.
- 239 4. Bell E, Sumner S: **Ecology and social organisation of wasps.** *eLS* 2013,
240 doi:10.1002/9780470015902.a0023597.
- 241 5. Hamilton WD: **The genetical evolution of social behaviour. {I & II}.** *J Theor Biol* 1964,
242 **7**:1.
- 243 6. Hamilton WD: **The evolution of altruistic behavior.** *Am Nat* 1963,
- 244 7. West SA, Gardner A: **Adaptation and inclusive fitness.** *Curr Biol* 2013, **23**:R577--R584.
- 245 8. West SA, Griffin AS, Gardner A: **Evolutionary explanations for cooperation.** *Curr Biol*
246 2007, **17**:R661--R672.
- 247 9. Szathmáry E, Maynard Smith J: **The major evolutionary transitions.** *Nature* 1995,
248 **374**:227–232.
- 249 10. West SA, Fisher RM, Gardner A, Kiers ET: **Major evolutionary transitions in**
250 **individuality.** *Proc Natl Acad Sci* 2015, **112**:10112–10119.
- 251 11. Bourke AFG: *Principles of social evolution.* Oxford University Press, Oxford; 2011.
- 252 12. Robinson GE, Grozinger CM, Whitfield CW: **Sociogenomics: Social life in molecular**
253 **terms.** *Nat Rev Genet* 2005, **6**:257–70.
- 254 13. Toth, AL Rehan S: **Climbing the social ladder: The molecular evolution of sociality.**
255 2015, doi:10.1111/j.1467-8330.1974.tb00606.x.
- 256 14. Sumner S, Bell E, Taylor D: **A molecular concept of caste in insect societies.** *Curr Opin*
257 *Insect Sci* 2017, doi:10.1016/j.cois.2017.11.010.
- 258 15. Toth AL, Rehan SM: **Molecular evolution of insect sociality: An eco-evo-devo**
259 **perspective.** *Annu Rev Entomol* 2017, **62**:419–442.

260 ** This comprehensive review defines levels of sociality and their importance in understanding the
261 evolution of social behaviour. It also summarises our current understanding of the molecular
262 mechanisms involved at each of these stages, as well as providing an inclusive overview of the
263 mechanistic hypotheses of social evolution.

- 264 16. Grozinger CM, Richards J, Mattila HR: **From molecules to societies: Mechanisms**
265 **regulating swarming behavior in honey bees (*Apis* spp.)**. *Apidologie* 2014, **45**:327–346.
- 266 17. Ashby R, Forêt S, Searle I, Maleszka R: **MicroRNAs in honey bee caste determination**.
267 *Nat Publ Gr* 2016, doi:10.1038/srep18794.
- 268 18. Warner MR, Mikheyev AS, Linksvayer TA: **Genomic signature of kin selection in an ant**
269 **with obligately sterile workers**. *Mol Biol Evol* 2017, **34**:1780–1787.
- 270 ** The authors used RNAseq in an ant species with an obligately sterile worker caste to identify
271 genomic signatures of kin selection and determine the relative contribution of ancient and young
272 genes to caste evolution.
- 273 19. Peters RS, Krogmann L, Mayer C, Rust J, Misof B, Niehuis O, Peters RS, Krogmann L,
274 Mayer C, Donath A, et al.: **Evolutionary history of the Hymenoptera**. *Curr Biol* 2017,
275 **27**:1–6.
- 276 ** This is the most comprehensive dataset on the phylogeny of the Hymenoptera, with 167 species
277 sequenced (and a further five annotated genomes included). This provides information for all the
278 major superfamilies, with the most in-depth phylogenetic relationships and divergence time
279 estimates.
- 280 20. Jandt JM, Toth AL: **Physiological and genomic mechanisms of social organization in**
281 **wasps (family: Vespidae)**. *Adv In Insect Phys* 2015, **48**:95–130.
- 282 21. West-Eberhard M: **Wasp societies as microcosms for the study of development and**
283 **evolution**. In *Natural History and Evolution of Paper Wasp*. Edited by Turillazzi, S.; West-
284 Eberhard M. Oxford University Press; 1996.
- 285 22. Field J, Foster W, Shreeves G, Sumner S: **Ecological constraints on independent**
286 **nesting in facultatively eusocial hover wasps**. *Proc R Soc B Biol Sci* 1998, **265**:973–977.
- 287 23. Baracchi D, Mazza G, Cini A, Petrocelli I, Bin Hashim R, Turillazzi S: **Social biology of**
288 ***Parischnogaster striatula* (Hymenoptera: Stenogastrinae)**. *Trop Zool* 2013, **26**:105–119.
- 289 24. Bolton A, Sumner S, Shreeves G, Casiraghi M, Field J: **Colony genetic structure in a**
290 **facultatively eusocial hover wasp**. *Behav Ecol* 2006, **17**:873–880.
- 291 25. Sumner S, Casiraghi M, Foster W, Field J: **High reproductive skew in tropical hover**
292 **wasps**. *Proc R Soc B Biol Sci* 2002, **269**:179–186.

- 293 26. Kocher SD, Li C, Yang W, Tan H, Yi S V, Yang X, Hoekstra HE, Zhang G, Pierce NE, Yu
294 DW: **The draft genome of a socially polymorphic halictid bee, *Lasioglossum albipes*.**
295 *Genome Biol* 2013, **14**.
- 296 27. Landi M, Queller DC, Turillazzi S, Strassmann JE: **Low relatedness and frequent queen**
297 **turnover in the stenogastrine wasp *Eustenogaster fraterna* favor the life insurance**
298 **over the haplodiploid hypothesis for the origin of eusociality.** *Insectes Soc* 2003,
299 **50:262–267.**
- 300 28. Sumner S, Kelstrup H, Fanelli D: **Reproductive constraints, direct fitness and indirect**
301 **fitness benefits explain helping behaviour in the primitively eusocial wasp, *Polistes***
302 ***canadensis*.** *Proc R Soc B Biol Sci* 2010, **277:1721–1728.**
- 303 29. Unnikrishnan S, Gadagkar R: **Dominance based reproductive queue in the primitively**
304 **eusocial wasp, *Ropalidia cyathiformis*.** *Insectes Soc* 2017, **64:495–503.**
- 305 30. Standage DS, Berens AJ, Glastad KM, Severin AJ, Brendel VP, Toth AL: **Genome,**
306 **transcriptome and methylome sequencing of a primitively eusocial wasp reveal a**
307 **greatly reduced DNA methylation system in a social insect.** *Mol Ecol* 2016, **25:1769–**
308 **1784.**
- 309 31. Bridge C, Field J: **Queuing for dominance: Gerontocracy and queue-jumping in the**
310 **hover wasp *Liostenogaster flavolineata*.** *Behav Ecol Sociobiol* 2007, **61:1253–1259.**
- 311 32. Thompson GJ, Hurd PL, Crespi BJ: **Genes underlying altruism.** *Biol Lett* 2013,
312 **9:20130395–20130395.**
- 313 33. Baracchi D: **The reproductive division of labour but not worker age affects spatial**
314 **sorting within the nest in a paper wasp.** *Insectes Soc* 2017, **64:379–385.**
- 315 34. West-Eberhard MJ: **Temporary queens in metapolybia wasps: Nonreproductive helpers**
316 **without altruism?** *Science (80-)* 1978, **200:441–443.**
- 317 35. Maleszka R: **Epigenetic code and insect behavioural plasticity.** *Curr Opin Insect Sci*
318 **2016, 15:45–52.**
- 319 36. Shukla S, Chandran S, Gadagkar R: **Ovarian developmental variation in the primitively**
320 **eusocial wasp *Ropalidia marginata* suggests a gateway to worker ontogeny and the**
321 **evolution of sociality.** *J Exp Biol* 2013, **216:181–187.**

- 322 37. Baio M V, Noll FB, Zucchi R: **Morphological caste differences and non-sterility of**
323 **workers in *Brachygastra augusti* (Hymenoptera, Vespidae, Epiponini), a neotropical**
324 **swarm-founding wasp.** *J New York Entomol Soc* 2004, **111**:242–252.
- 325 38. Hunt JH, O'Donnell S, Chernoff N, Brownie C: **Observations on two neotropical swarm-**
326 **founding wasps, *Agelaia yepocapa* and *A. panamaensis* (Hymenoptera: Vespidae).**
327 *Ann Entomol Soc Am* 2001, **94**:555–562.
- 328 39. Patnaik BB, Park SY, Kang SW, Hwang HJ, Wang TH, Park EB, Chung JM, Song DK, Kim
329 C, Kim S, et al.: **Transcriptome profile of the Asian giant hornet (*Vespa mandarinia*)**
330 **using Illumina HiSeq 4000 sequencing: De novo assembly, functional annotation, and**
331 **discovery of SSR markers.** *Int J Genomics* 2016, doi:10.1155/2016/4169587.
- 332 40. Kapheim KM: **Nutritional, endocrine, and social influences on reproductive physiology**
333 **at the origins of social behavior.** *Curr Opin Insect Sci* 2017, **22**:62–70.
- 334 41. Chen X, Hu Y, Zheng H, Cao L, Niu D, Yu D, Sun Y, Hu S, Hu F: **Transcriptome**
335 **comparison between honey bee queen- and worker-destined larvae.** *Insect Biochem*
336 *Mol Biol* 2012, **42**:665–673.
- 337 42. Guo X, Su S, Geir S, Li W, Li Z, Zhang S, Chen S, Chen R: **Differential expression of**
338 **miRNAs related to caste differentiation in the honey bee, *Apis mellifera*.** *Apidologie*
339 2016, **47**:495–508.
- 340 *This paper furthers our understanding of the mechanisms involved in caste development, by
341 showing that miRNAs show caste-specific expression in queen and worker honey bees.
- 342 43. Berens AJ, Hunt JH, Toth AL: **Comparative transcriptomics of convergent evolution:**
343 **Different genes but conserved pathways underlie caste phenotypes across lineages**
344 **of eusocial insects.** *Mol Biol Evol* 2015, **32**:690–703.
- 345 44. Chavarría L, Noll FB: **Age polyethism in the swarm-founding wasp *Metapolybia miltoni***
346 **(Andena & Carpenter) (Hymenoptera: Vespidae; Polistinae, Epiponini).** *Sociobiology*
347 2013, **60**:214–216.
- 348 45. Noll FB, Zucchi R: **Increasing caste differences related to life cycle progression in**
349 **some neotropical swarm-founding polygynic polistine wasps (Hymenoptera:**
350 **Vespidae: Epiponini).** *Ethol Ecol Evol* 1999, **12**:43–65.

- 351 46. Hughes CR, Queller DC, Strassmann JE, Soils CR, Negron-Sotomayor JA GK: *The*
352 *maintenance of high genetic related- ness in multi-queen colonies of social wasps. In:*
353 *Queen number and sociality in insects.* Oxford University Press; 1993.
- 354 47. Zhou X, Rokas A, Berger SL, Liebig J, Ray A, Zwiebel LJ: **Chemoreceptor evolution in**
355 **Hymenoptera and its implications for the evolution of eusociality.** *Genome Biol Evol*
356 2015, **7**:2407–2416.
- 357 48. Leonhardt SD, Menzel F, Nehring V, Schmitt T: **Ecology and evolution of communication**
358 **in social insects.** *Cell* 2016, **164**:1277–1287.
- 359 49. Olejarz J, Veller C, Nowak MA: **The evolution of queen control over worker**
360 **reproduction in the social Hymenoptera.** *Ecol Evol* 2017, **7**:8427–8441.
- 361 50. Mitra A: **Queen pheromone and monopoly of reproduction by the queen in the social**
362 **wasp *Ropalidia marginata*.** *Proc Indian Natl Sci Acad* 2014, **80**:1025–1044.
- 363 51. Kelstrup HC, Hartfelder K, Esterhuizen N, Wossler TC: **Juvenile hormone titers, ovarian**
364 **status and epicuticular hydrocarbons in gynes and workers of the paper wasp**
365 ***Belonogaster longitarsus*.** *J Insect Physiol* 2017, **98**:83–92.
- 366 52. Wen P, Cheng YN, Dong SH, Wang ZW, Tan K, Nieh JC: **The sex pheromone of a**
367 **globally invasive honey bee predator, the Asian eusocial hornet, *Vespa velutina*.** *Sci*
368 *Rep* 2017, **7**:1–11.
- 369 53. Oi CA, Millar JG, van Zweden JS, Wenseleers T: **Conservation of queen pheromones**
370 **across two species of Vespine wasps.** *J Chem Ecol* 2016, **42**:1175–1180.
- 371 54. Oi CA, Van Oystaeyen A, Caliari Oliveira R, Millar JG, Verstrepén KJ, Van Zweden JS,
372 Wenseleers T: **Dual effect of wasp queen pheromone in regulating insect sociality.**
373 *Curr Biol* 2015, **25**:1638–1640.
- 374 *The authors show that queen pheromones not only induce sterility in *V. vulgaris*, but also enables
375 the queen to communicate egg paternity, helping workers to recognise worker-laid eggs, and thus
376 facilitating policing.
- 377 55. Olejarz JW, Allen B, Veller C, Gadagkar R, Nowak MA: **Evolution of worker policing.** *J*
378 *Theor Biol* 2016, **399**:103–116.
- 379 56. Palmer KA, Oldroyd BP: **Evolution of multiple mating in the genus *Apis*.** *Apidologie*
380 2000, **31**:235–248.

- 381 57. Kronauer DJC, Johnson RA, Boomsma JJ: **The evolution of multiple mating in army**
382 **ants.** *Evolution (N Y)* 2007, **61**:413–422.
- 383 58. Foster KR, Gulliver J, Ratnieks FLW: **Worker policing in the European hornet *Vespa***
384 ***crabro*.** *Insectes Soc* 2002, **49**:41–44.
- 385 59. Loope KJ, Chien C, Juhl M: **Colony size is linked to paternity frequency and paternity**
386 **skew in yellowjacket wasps and hornets.** *BMC Evol Biol* 2014, **14**:1–12.
- 387 60. Kennedy P, Baron G, Qiu B, Freitak D, Helanterä H, Hunt ER, Manfredini F, O'Shea-Wheller
388 T, Patalano S, et al.: **Deconstructing superorganisms and societies to address big**
389 **questions in biology.** *Trends Ecol Evol* 2017, **32**:861–872.
- 390 61. Field J, Cronin A, Bridge C: **Future fitness and helping in social queues.** *Nature* 2006,
391 **441**:214–217.
- 392 62. Field J, Leadbeater E: **Cooperation between non-relatives in a primitively eusocial**
393 **paper wasp, *Polistes dominula*.** *Philos Trans R Soc B Biol Sci* 2016, **371**:20150093.
- 394 63. Patalano S, Vlasova A, Wyatt C, Ewels P, Camara F, Ferreira PG, Asher CL, Jurkowski TP,
395 Segonds-Pichon A, Bachman M, et al.: **Molecular signatures of plastic phenotypes in**
396 **two eusocial insect species with simple societies.** *Proc Natl Acad Sci* 2015, **112**:13970–
397 13975.
- 398 * This is the first published genome sequence for a Vespid wasp. It revealed how subtle
399 differences in gene expression, but not methylation, underpins the first stages of the major
400 transition in social evolution.
- 401 64. Nadolski J: **Structure of nests and colony sizes of the European Hornet (*Vespa crabro*)**
402 **and Saxon wasp (*Dolichovespula saxonica*) (Hymenoptera: Vespinae) in urban**
403 **conditions.** *Sociobiology* 2012, **59**:1075–1120.
- 404 65. Mateus S, Noll F, Zucchi R: **Morphological caste differences in the neotropical swarm-**
405 **founding Polistine wasps: *Parachartergus smithii* (Hymenoptera: Vespidae).** *J New*
406 *York Entomol Soc* 1997, **105**:129–139.
- 407 66. O'Donnell S: **Worker age, ovary development, and temporal polyethism in the swarm-**
408 **founding wasp *Polybia occidentalis* (Hymenoptera: Vespidae).** *J Insect Behav* 2001,
409 **14**:201–213.

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434