Blencowe, N., Waldon, R., & Vipond, M. N. (2018). Management of patients after laparoscopic procedures. *BMJ*, 360, [k120].
<https://doi.org/10.1136/bmj.k120>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1136/bmj.k120](https://doi.org/10.1136/bmj.k120)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via BMJ at <https://www.bmj.com/content/360/bmj.k120> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Management of patients following laparoscopic procedures, for the non-expert

Natalie S Blencowe¹, Richard Waldon², Mark N Vipond³

- 1 NIHR Clinical Lecturer and honorary specialty trainee, General Surgery, University of Bristol and the Severn Deanery
- 2 Senior Partner, Rowcroft Medical Centre, Stroud, Gloucestershire
- 3 Consultant Upper GI Surgeon, Gloucestershire Hospitals NHS Foundation Trust
Past President of the Association of Laparoscopic Surgeons of Great Britain and Ireland

Correspondence to: Mr M N Vipond

mark.vipond@nhs.net

Competing interests: None declared

Contribution statement: NB wrote the first draft of the article. All authors contributed to revisions and approved the final version for submission.

What you need to know

- The length of the recovery period depends upon the procedure that has been performed, and may be as short as 1-2 weeks
- Patients should refrain from driving until they are comfortably able to look in their blind spot and perform an emergency stop
- Worsening post-operative pain suggests a potential abdominal complication, and the patient requires urgent assessment
- Although most wound infections after laparoscopic surgery are superficial, some can involve the deeper tissues and should be re-assessed in hospital, particularly if prosthetic material has been inserted (e.g. mesh)

Introduction

Laparoscopy has revolutionized the field of abdominal surgery over the past thirty years. The total number of laparoscopic surgical operations performed annually worldwide is approximately 15 million, of which 10% are performed in the UK². This technique – also known as minimally invasive surgery - involves the use of cameras and specially designed instruments that allow surgeons to perform operations without the need for a large abdominal incision. It has gained worldwide popularity and acceptance by surgeons and patients alike. Better intra-operative visibility, minimal scarring, less pain, shorter hospital stay and faster recovery are the main reasons behind the appeal of laparoscopic surgery¹. This practice pointer discusses the important considerations around the management of patients following laparoscopic surgical procedures.

Laparoscopy – what's changed?

As well as cholecystectomy, appendicectomy and tubo-ovarian procedures, laparoscopy is now increasingly used for hysterectomy, bariatric procedures and many gastrointestinal and urological cancer resections, and it is also possible to perform aortic aneurysm repair laparoscopically^{3,4,5}. Further technological advances have resulted in the development of several variations on the 'standard' laparoscopic approach, which are described in Table 1.

Enhanced recovery programs are increasingly popular and are ideally suited to laparoscopic surgery. There is evidence to suggest that this approach reduces length of stay without increasing post-operative complications, when compared with standard care⁶. Enhanced recovery starts and ends many weeks before and after the surgical procedure itself; the main elements are outlined in Table 2.

When are patients typically discharged after laparoscopic surgery?

This depends upon the type of surgery that has been performed, and patient factors such as co-morbidities. Day case procedures are becoming more common: for example, over 60% of patients undergoing laparoscopic cholecystectomy are now discharged on the same day⁷. Discharge criteria include the ability to mobilise and control pain with simple analgesia. Although desirable, passing urine and oral intake are no longer seen as absolute requirements prior to discharge⁸. Patients are usually provided with information leaflets about expected recovery period, emergency contact details and follow-up schedules, although further appointments are often not required after routine procedures.

Which medications are usually prescribed?

Patients are usually discharged from hospital with simple analgesia. Sometimes it is appropriate to provide medication(s) to counteract the side effects of analgesia (such as laxatives, anti-emetics or proton pump inhibitors), and thromboprophylaxis (see below).

Analgesia

The duration of analgesia depends on the type of operation that has been performed. For operations such as appendicectomy or cholecystectomy, patients often require regular paracetamol, weak opioids (e.g. codeine), and/or NSAIDs for the first week after surgery, reducing to 'as required' after that. Following larger operations (e.g. for gastrointestinal cancer), patients may require regular analgesia for up to 2 months.

Thromboprophylaxis

Daily pharmacological prophylaxis is administered for the duration of all patients' hospital stay. Venous thromboembolism – i.e. deep venous thrombosis or pulmonary embolism – is not common following laparoscopic surgery, occurring in less than 1% of patients⁹. The

duration of thromboprophylaxis depends upon the procedure that has been performed. NICE advise the use of extended pharmacological VTE prophylaxis for 28 days following cancer surgery¹⁰. Although NICE does not specifically provide recommendations for bariatric procedures, obesity is a risk factor for VTE and such patients usually also receive extended VTE prophylaxis¹¹. There is no clear evidence to say which type of pharmacological prophylaxis is best, and individual units or surgeons may demonstrate a preference. Where appropriate, patients are taught the self-administration technique for heparin injections. If anticoagulant therapy was stopped pre-operatively, patients are usually able to recommence this before or at the point of discharge.

Wound care

Subcuticular absorbable sutures are the most commonly used materials for closing laparoscopic wounds. Simple adhesive dressings are typically applied to each wound, although there is no evidence to suggest that dressings reduce the rate of wound infection¹². Increasingly, skin glue is being used as a 'dressing', with the advantage that it renders the wound waterproof, meaning patients can shower immediately after surgery¹³. Patients with adhesive dressings are generally advised to avoid soaking the wounds in water for a few days after surgery. Unless there is excessive discharge of fluid from the wounds, there is generally no need to replace the dressings once they have fallen off.

Post-operative restrictions

Although the 'port site' wounds suggest a modest intervention, the underlying procedure is usually more extensive, and patients need to remember that a period of recovery will be required. The speed at which normal activity is resumed after laparoscopic surgery largely depends upon the extent and type of operation that has been performed. Many surgeons will use the phrase "if it hurts, don't do it" and explain to patients that they should notice an

almost daily increase in the activities they are able to undertake without experiencing pain or discomfort.

Driving

This is dependent on the type of laparoscopic procedure that has been performed. For example, a patient undergoing laparoscopic cholecystectomy or hernia repair may be able to drive after 1-2 weeks¹⁴, whereas those undergoing hernia repair or larger cancer procedures may need to wait longer. Patients should only be allowed to drive if they are comfortably able to apply an emergency brake, look in their blind spot and have sufficient reaction times. Drivers do not need to notify the DVLA of surgical recovery unless it is likely to persist for more than 3 months¹⁵.

Return to work

There are no clear guidelines on when patients should return to work. It is usually an individual decision, and depends on their occupation and the procedure performed. Patients may only require two weeks away from work after smaller procedures (such as cholecystectomy or appendicectomy)¹⁶. Following cancer surgery, a longer break, or gradual return to work may be advised, such as light duties or reduced hours¹⁶. "Sick notes" are provided by the hospital team to cover the anticipated period of absence.

Heavy lifting

Heavy lifting after laparoscopic surgery is controversial and under-researched. Whereas some sources state that lifting should be avoided for six weeks¹⁷, 1-2 and 4 weeks are generally thought to be sufficient for smaller and larger procedures, respectively¹⁶. During this time, patients are advised not to lift anything heavier than a kettle or a shopping bag.

They are also asked to avoid pushing and pulling activities, such as vacuuming, scrubbing the bath, hanging up heavy washing and mowing the lawn.

Eating and drinking

Restrictions on eating and drinking are dependent on the procedure that has been undertaken. For most, there are no restrictions. Operations for morbid obesity, anti-reflux procedures, and oesophago-gastric cancer, may initially require patients to follow a liquid diet¹⁸.

Travel

The risk of venous thromboembolism associated with air travel increases with the duration of travel and number of flights over a short time period¹⁹. The Civil Aviation Agency advise patients not to fly for 24 hours after laparoscopy, due to the potential for expansion of retained carbon dioxide in the abdominal cavity²⁰. A sensible precaution would be to use thromboembolic stockings if flying within one month of surgery.

What complications commonly present to primary care?

The majority of patients have very little pain after 'minor' laparoscopic surgery, and are often eager to mobilise soon afterwards. Appetite may be unaffected and apart from feeling tired, they may otherwise feel 'back to normal' quite quickly. It is important to bear these features in mind when assessing a patient following laparoscopic surgery, as a failure to meet these milestones may indicate an abdominal complication²¹.

Pain

Despite the use of small incisions, mild pain is common following laparoscopic surgery. Sometimes, the carbon dioxide used to inflate the abdomen can remain inside, causing

cramps, bloating and shoulder tip pain. These symptoms generally subside after 24 hours. If pain worsens after this time period, there should be a low threshold for suspecting that an abdominal complication has occurred. Table 3 (see below) outlines the other symptoms and signs that may be suggestive of an intra-abdominal complication²¹.

Wound infection

Any wound that is red, oozing, hot, swollen or tender may be infected and antibiotics are usually required. If there is a concern about deeper infection (and especially where prosthetic material, such as mesh, was inserted during the laparoscopic procedure), urgently refer the patient to hospital for review by the surgical team.

Distinguishing between haematoma, abscess and seroma

These complications all cause swelling at the operative site. There are various defining features of these complications that may help to distinguish one from another (Table 4). In cases of diagnostic doubt, ultrasound can be helpful. Whereas seromas are often treated conservatively, abscesses and (large) haematomas almost always require surgical intervention.

Other post-operative complications

Whilst laparoscopic surgery offers many advantages, abdominal complications may present more subtly than after a traditional 'open' operation²¹. The complications themselves are almost identical, and largely depend upon the nature of the operation that has been performed. For example, division of adhesions confers the risk of bowel damage, cholecystectomy may result in leakage of bile into the peritoneal cavity, and leakage from staple lines or anastomoses may occur after surgical procedures for morbid obesity or gastrointestinal cancer surgery. It is not the complication *per se* which is critical, rather the

time frame in which it is identified and treated. In the absence of clear evidence of continuing improvement, further investigation is urgently needed and prompt referral of patients to hospital is therefore crucial. Imaging and blood tests should be performed quickly at the point of hospital admission, rather than in the community, which may delay treatment. Table 3 outlines the symptoms and signs that may raise the suspicion of an intra-abdominal complication.

References

1. Agha R, Muir G. Does laparoscopic surgery spell the end of the open surgeon? *Journal of the Royal Society of Medicine*. 2003; 96(11):544-546.
2. Abbott TEF, Fowler AJ, Dobbs TD et al. Frequency of surgical treatment and related hospital procedures in the UK: a national ecological study using hospital episode statistics. *Br J Anaesthesia* 2017; 119(2):249–257
3. Green CJ, Maxwell R, Verne J, Martin RM, Blazeby JM. The influence of NICE guidance on the uptake of laparoscopic surgery for colorectal cancer. *Journal of Public Health*, 2009; 31(4):541-545
4. Cuesta MA, Biere SSAY, Henegouwen MI van B, van der Peet DL. Randomised trial, Minimally Invasive Oesophagectomy versus open oesophagectomy for patients with resectable oesophageal cancer. *Journal of Thoracic Disease*. 2012;4(5):462-464.
5. Janda M, Gebiski V, Davies LC et al. Effect of Total Laparoscopic Hysterectomy vs Total Abdominal Hysterectomy on Disease-Free Survival Among Women With Stage I Endometrial CancerA Randomized Clinical Trial. *JAMA*. 2017;317(12):1224–1233.
6. Varadhan KK, Neal KR, Dejong CH, Fearon KC, Ljungqvist O, Lobo DN. The Enhanced Recovery After Surgery (ERAS) pathway for patients undergoing major elective open colorectal surgery: a meta-analysis of randomized controlled trials. *Clin Nutr* 2010; 29(4):434–440.
7. CholeS Study Group. Population-based cohort study of outcomes following cholecystectomy for benign gallbladder diseases. *Br J Surg* 2-16; 103(12):1704–1715
8. The Association of Anaesthetists of Great Britain and Ireland, and the Association of Day Surgery. Day case and short stay surgery. 2011. Available at <https://www.aagbi.org/sites/default/files/Day%20Case%20for%20web.pdf>

9. Inderbitzin DT, Opitz I, Giger U, Kocher T, Krähenbühl L. Incidence of clinical pulmonary embolism after laparoscopic surgery. *Br J Surg* 2007; 94(5):599-603.
10. NICE. 2015. Venous thromboembolism: reducing the risk for patients in hospital. Available at <https://www.nice.org.uk/guidance/cg92/>
11. Bartlett MA, Mauck KF, Daniels PR. Prevention of venous thromboembolism in patients undergoing bariatric surgery. *Vascular Health and Risk Management*. 2015;11:461-477.
12. Blazeby JM on behalf of the Bluebelle study group. Do dressings prevent infection of closed primary wounds after surgery? *BMJ* 2016;353:i2270
13. SPARCS and the WMRC on behalf of the Bluebelle study group. Feasibility work to inform the design of a randomized clinical trial of wound dressings in elective and unplanned abdominal surgery. *Br J Surg* 2016;103(12):1738-1744.
14. Ismail W, Taylor SJC, Beddow E. Advice on driving after groin hernia surgery in the United Kingdom: questionnaire survey. *BMJ* 2000; 321(7268): 1056
15. Driver and Vehicle Licensing Agency. 2017. Assessing fitness to drive: a guide for medical professionals. Available at www.gov.uk/dvla/fitnesstodrive
16. Van Vliet DCR, van der Meij E, Bouwsma EVA, et al. A modified Delphi method toward multidisciplinary consensus on functional convalescence recommendations after abdominal surgery. *Surg Endosc*. 2016;30(12):5583-5595.
17. Medical Disability Advisor. Accessed 16th September 2017: <http://www.mdguidelines.com/medical-topics>
18. Mechanick JI, Youdim A, Jones DB, et al. Clinical Practice Guidelines for the Perioperative Nutritional, Metabolic, and Nonsurgical Support of the Bariatric Surgery Patient. *Obesity* 2013; 21(0 1):S1-27.
19. Philbrick JT, Shumate R, Siadat MS, Becker DM. Air Travel and Venous Thromboembolism: A Systematic Review. *Journal of General Internal Medicine*. 2007;22(1):107-114.
20. Civil Aviation Authority. Accessed 16th September 2017 www.caa.co.uk/Passengers/Before-you-fly/Am-I-fit-to-fly/Guidance-for-health-professionals/Surgical-conditions/
21. Association of Laparoscopic Surgeons. Recognition, management and prevention of abdominal complications of laparoscopic surgery. Accessed 1st September 2017: www.alsgbi.org/pdf/ALS_Complications_Management.pdf

Table 1 Variations on standard laparoscopic techniques

Name of laparoscopic technique		Description
SILS	Single incision laparoscopic surgery	The surgeon operates almost exclusively through a single entry point, typically the patient's umbilicus, leaving a single scar.
NOTES	Natural orifice endoscopic surgery	Performed with an endoscope passed through a natural orifice (e.g. mouth, anus) then through an internal incision (e.g. stomach, colon), thus avoiding any external incisions or scars.
Robotically-assisted laparoscopic surgery		Robotic systems, controlled by surgeons, are used to facilitate operations. The main perceived advantage is articulation beyond normal manipulation, resulting in improved ergonomics.
Laparo-endoscopic surgery		Combination of laparoscopic and endoscopic surgery (e.g. for removal of gastrointestinal polyps), eliminating the need for more major surgery (e.g. bowel or stomach resection).

Table 2 The main elements of enhanced recovery programmes

Time period	Element
Pre-operative care	Pre-operative assessment visit
	Lifestyle changes: stop smoking, lose weight, reduce alcohol, increase exercise
	Correct anaemia
	Optimise diabetic care
	Pre-operative carbohydrate drinks
Admission to hospital	Admission on the day of surgery
	Carbohydrate drinks and water up to 2 hours before surgery
Peri-operative care	Use of minimally invasive techniques
	Individualised fluid balance therapy
	Prevention of hypothermia
	Minimise use of drains and tubes
Post-operative care	Early mobilisation
	Early return to eating and drinking
	Early discharge planning

Table 3 Symptoms and signs suggestive of an intra-abdominal complication following laparoscopic surgery

Worsening abdominal pain
Anorexia or reluctance to drink
Reluctance to mobilise
Nausea or vomiting
Tachycardia
Abdominal tenderness or distension
Poor urine output
Pyrexia

Table 4 Distinguishing features of abscess, haematoma and seroma

	Haematoma	Abscess	Seroma
Time of onset	Usually within 24-48 hours of surgery	Usually 3-7 days after surgery	Usually later onset (after a week)
Overlying skin colour	Bruising may be present	Red, may be oozing pus	Normal
Overlying skin temperature	Normal	Hot	Normal
Degree of pain	May be painful	Often exquisitely tender	Usually only slight discomfort