Rakhra, D., Walker, T. W. M., Hall, S., Fleming, C. A., Thomas, S. J., Kerai, A., Horwood, J. P., & Waylen, A. E. (2018). Human papillomavirus (HPV) and its vaccine: Awareness and opinions of clinical dental students in a uk dental school. *British Dental Journal*, 225(10), 976–981. <https://doi.org/10.1038/sj.bdj.2018.1024>

Peer reviewed version

Link to published version (if available):
[10.1038/sj.bdj.2018.1024](https://doi.org/10.1038/sj.bdj.2018.1024)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Nature at <https://doi.org/10.1038/sj.bdj.2018.1024> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Human Papillomavirus (HPV) and its vaccine: Awareness and Opinions of Clinical Dental Students in a UK Dental School

Rakhra D, Walker T W M, Fleming CA, Kerai A, Hall S, Horwood J P, Waylen A E and Thomas S J.

In brief

- Students in the Bristol Dental Hospital understood the links between HPV and oropharyngeal carcinoma and its vaccination process.
- Students demonstrated that they believe oral healthcare practitioners have a role to play in health promotion regarding oral and oropharyngeal cancer but desired more teaching on the topic.
- Students felt unprepared to discuss the disease and associated risk factors with patients but further clarification as to why is needed.

Abstract

Introduction Dental clinicians are in a position to educate their patients on the subject of HPV as part of a primary healthcare multidisciplinary team and to detect HPV related disease. Attention needs to be paid to dental undergraduate teaching on the topic. This study aims to ascertain awareness and opinions of our dental undergraduate students about HPV, its relation to oral health and its vaccine. **Method** A questionnaire was distributed to undergraduate dental students in clinical years 3, 4 and 5 of the BDS programme at Bristol Dental School. **Results** Dental students demonstrated an understanding of HPV and oropharyngeal cancer and expressed that they felt dentists should play a role in health promotion in relation to oropharyngeal cancers. They also wished for more teaching on the subject and suggested topic areas that they wished to know more about. **Conclusion** New teaching strategies on the topic of HPV should be considered for inclusion into undergraduate dental programmes and encompass both knowledge and communication skills training to prepare future dentists for their role in confronting this preventable disease.

Introduction

The Human Papilloma Virus (HPV) is established as an aetiological risk factor for oropharyngeal cancer.¹ The number of cases of oropharyngeal cancer in the UK has risen significantly over the last two decades with the age-specific greatest increase in incidence seen in the 40-69 year old age group² and a similar increase in cases has been noted in other countries.³ HPV, in particular HPV-16, has seen a significant rise in association with oropharyngeal cancer over the last decade⁴, is identified in approximately 50-60% of cases.^{2,4-6} HPV is thought to explain the pattern of oropharyngeal cases in young, heterosexual male, non-smoking, low alcohol drinking patient cohort. Evidence suggests transmission of HPV is via high-risk sexual activity⁷ and is associated the number of sexual partners and oral sex partners over a lifetime.⁸⁻¹⁰ Tumour HPV status is considered to be a major prognostic factor for patients with oropharyngeal carcinoma¹¹ with survival for patients with HPV positive tumours being better.^{12,13} Management of patients with HPV-related oropharyngeal cancer is a high interest area of current research¹⁴ as effective treatment modalities differ from their non-HPV-related counterparts.

Oral health professionals can play a role in the prevention of HPV-related oropharyngeal cancers by educating patients about the risks of HPV transmission, instruct about lifestyle change and by encouraging patients and their relatives to participate in HPV vaccination programmes.¹⁵⁻¹⁷ There is a view in the United States that dental healthcare providers should be involved in education of their patients about HPV and its link to oropharyngeal cancers^{18,19} and some suggest oral health professionals should be at the forefront of confronting the 'HPV related oropharyngeal cancer epidemic'.¹⁸ Given that 28.9 million people accessed NHS dental care in the 24 months ending 31 Dec 2016²⁰, dental teams have a real opportunity to engage with a significant number of people about this escalating disease.

Since 2008, the current HPV vaccination programme in the UK has included girls aged 11-13, aimed at prevention of cervical cancers. Thus far it is reported to have decreased cancer rates by 86% in those women eligible for the vaccination as adolescents.²¹ Many organisations, including the British Dental Association²² are campaigning to include vaccinations for adolescent boys as well. The HPV vaccination became available for men who have sex with men in April 2018 after successful pilots in

the UK²³ and, most recently, the HPV vaccine has been proved to be cost effective in teenage boys and it will soon be offered to this cohort.²⁴

Articles on the subject of HPV have been presented more recently in dental literature in the UK²⁵⁻²⁸ and a recent study concluded that UK dental students lack confidence in communicating with patients about HPV.²⁹ It is clear that educational interventions are required to keep all dental professionals up to date with the rapidly changing landscape of oral and oropharyngeal cancer³⁰ in addition to training in communication skills to connect sensitively with their patients about potentially delicate subjects around sexual activity.

The present study has been conducted, as part of our teaching evaluation and development process, with an aim to ascertain the current awareness and opinions of HPV and its vaccine among our dental undergraduate students. The outcome of the study will inform the undergraduate dental teaching programme review process to ensure our students graduate to carry out effective health promotion on this subject.

Methods

A cross-sectional study was performed among undergraduate dental students in years 3, 4 and 5 of the five-year Bachelor of Dental Surgery programme at Bristol Dental School. A questionnaire was developed using previous literature^{8,29} and validated by means of a round table discussion (Appendix 1). Ethical approval was obtained via the University of Bristol, Faculty of Medicine & Dentistry Ethics Committee.

The clinical dental undergraduate students were selected because all these students had received some teaching about HPV, the extent of which depending on which year of the programme they were enrolled in. The students were advised by e-mail one week ahead of questionnaire distribution that they would be surveyed using a hard copy of the questionnaire at the end of a specified lecture. Students in all years were given the opportunity to leave without completing the survey. Questionnaires were deposited by students in to a drop box after they had completed them. Completed questionnaires were immediately entered in to a secure password protected University accessed database. Descriptive analysis was applied to the data.

Questions were developed to acquire qualitative data to assess dental students':

- Current awareness about HPV and its link to oropharyngeal cancer (yes/no answers),
- Knowledge of the methods of HPV transmission (single best answer).
- Knowledge of the current UK vaccination schedule (single best answer).
- Opinions on the importance of the role that general dental practitioners play in health promotion related to oral and oropharyngeal cancer (4 point Likert scale).
- Views on how much more information regarding HPV and its vaccine they would like to be taught (4 point Likert scale)

In addition, a free text area was included to gain further information regarding what else students would like to know about HPV.

Results

Response rates

A total of 167 questionnaires were returned from the 211 students across the 3 clinical years, giving a response rate across all years of 79%. Students from third year responded highest with a 92% response rate, followed by year 5 at 74% and then year 4 with a response rate of 72%. The absolute number of responses follows this trend with 61, 55 and 51 responses respectively.

Responses to questions regarding HPV awareness

The results to the questions relating to students' awareness of HPV and its link to oropharyngeal cancer, awareness of HPV transmission and the HPV vaccine are illustrated in Figures 1 to 3. There was a slight decrease in responses to individual questions. The questions represented by Figure 1 had two respondents not answer and Figure 3 lacked one answer. In summary: 93% of students knew HPV and cause oropharyngeal cancer, 71% knew it could be contracted via oral-genital sex and 83% correctly identified the cohort eligible for the vaccine under the NHS at the time. With regard to the latter point, half of those who responded incorrectly, chose answers that included the correct cohort.

Role of the dentist

Students were asked how important they thought the role of the dentist is in health promotion related to oropharyngeal cancer. In total 143 (86%) of students responded that the role of the dentist was either *important* or *very important*. Only two students, both from year 3, responded that the dentist's role was *not important at all*. The results are illustrated in Figure 4.

Dental education

When the students were asked how much more they felt they needed to know about HPV and its vaccine, 139 (83%) of 167 responders answered 'some more' or 'lots more' as illustrated in Figure 5. Only three students (2%) felt they knew enough about HPV and its vaccine.

The free text answers to '*What more would you like to know about HPV and its relation to oral health?*' were categorised into 'knowledge' and 'communication' themed comments. The responses are summarised in Table 1. Across all three years, there is a rough 2:1 ratio between students' desire for more knowledge of HPV and its vaccine and teaching in communication of these topics with their patients.

Respondents tended to show a desire for more knowledge in prevention, vaccination and risk management. With regards to communications, students desired teaching about how and what to be advising to their patients about HPV and its vaccination.

Discussion

The position that the oral healthcare professional has a role to play in health promotion regarding HPV, oral and oropharyngeal cancers and its vaccine clearly resonates with our undergraduate dental students; all but one registered that it has importance. It follows that for this to be possible, knowledge of the virus and its sequelae should be taught to students so they can adopt this role when they begin to practice. It is worth noting that the question did not have an answer for equally important (i.e. a shared responsibility) between dental and medical team.

The question arises as to what dentists need to know about HPV, how they are expected to use that information to care for their patients, and indeed how much time is dedicated to this topic in a busy

undergraduate dental curriculum. A learning outcome relating to health promotion and disease prevention in the GDC's dental undergraduate curriculum document 'Preparing for practice' states that dentists are expected to *'provide patients with comprehensive and accurate preventive education'* and *"underpin all patient care with a preventive approach that contributes to the patient's long-term oral and general health"*³¹. With no specific guidance available, some authors have attempted to propose a level of knowledge about HPV that dentists should aim for, focussing on understanding the pathophysiology of HPV and oropharyngeal cancer, associated risk factors and on available vaccination programmes.^{1,7,8} Others advocate that national organisations representing oral health professionals should develop campaigns to offer education on HPV as part of continuing professional development.^{18,30}

In light of no reliable screening tools for oral HPV, prevention and early detection of disease is paramount. Education about lifestyle risk factors, vaccination programmes and the virus itself should be available to patients if they enquire, especially if the patient falls within the ever-expanding cohort of those who are offered the vaccination on the NHS. With respect to detection, practitioners should be able to identify oral and oropharyngeal carcinomas at their early stage and provide appropriate referral and council.

Most of the students (93%) surveyed had an understanding that HPV and oropharyngeal cancers are causally linked, however, just under a third – across all year groups - did not know that it could be contracted via oral-genital sex. This is significant as lack of knowledge about transmission may lead to patients being misinformed about their actual risk of the disease. Van der Valk and Walker¹⁷ found similar results in 2014, finding that two-thirds of newly qualified dentists in the Berkshire area felt poorly trained on discussing HPV with their patients. It is apparent, then, that there is a need for further education around this topic. A wider-reaching study including multiple institutions would be beneficial and may highlight places where students are very well prepared to deal with HPV.

Fortunately, the vaccination programme in the UK was better understood with 91% of respondents choosing an answer including the correct cohort. This is important because the danger with poor knowledge of vaccination programmes is lack of referral and education, not over-education or over-referral. However, this should not lead to complacency as new developments in the eligible vaccination cohort mean continued education is needed.

It is not just the author's opinion, however, that more teaching is needed here. The majority of our students expressed that they felt they needed to know 'some more' or 'lots more' about HPV and its vaccine (Figure 5) to inform patients. There were many knowledge themed free text comments supporting this (Table 1) such as what are the '*risks and prevalence?*' of HPV and '*what can we do to reduce the risk?*'. Acquiring this data about gaps in our students' knowledge is key to improve our teaching programme so that students are given the information they need to help educate their patients in the future.

Dental students are conventionally well trained in discussing risk factors related to oral cancer with patients, such as tobacco use and alcohol consumption²⁹ however, HPV and its relation to oropharyngeal cancer is a relatively new topic in the dental programme. There were several communication themed free text comments that demonstrated students may not be confident in communicating with patients about HPV, for example '*how do you talk to patients about HPV?*' and '*when appropriate to bring this up with patients?*' (Table 1). A recent study found that only 14% of dental students felt confident in discussing the risks of HPV in relation to oral health and 16% felt confident in advising on HPV vaccination programmes.²⁹ It is not clear, however, that this is a function of students' lack of knowledge more than the fact that discussion of oral-genital sex, cancer and high-risk lifestyles is inherently difficult. Particularly when those patients are young girls, and soon boys, under the age of 18. The ability to broach these topics professionally and with confidence is unlikely to come from the classroom and more from dedicated communications teaching and experience in practice.

It is beneficial for patients and practitioners alike that dental education does not lag behind its practice, as some authors have alluded to in other areas of dentistry.³² Given that this is a quickly changing field, there is potential here for more modern and alternative teaching methods, such as online courses or seminars, that can be easily spread, digested at leisure and updated with new developments remotely. Further research into why students and dentists are not comfortable discussing the above topics, and indeed whether this feeling persists with veteran oral healthcare professionals, would help guide both content and teaching methods and delivery.

Conclusion

The rise in HPV-related oropharyngeal cancer is recognised by the dental profession and attention must be given to training in this topic in contemporary dental undergraduate programmes. The study highlights that, despite a good acknowledgment of the disease and its sequelae, students from the Bristol Dental Hospital would like further teaching regarding the communication and prevention of HPV. Undergraduate dental programmes should aim to incorporate training in knowledge about HPV, as well as communication skills teaching, focussed on prevention, referring patients appropriately and answering patients' questions in order to educate them as part of a multidisciplinary primary healthcare approach to tackling this preventable disease. Future studies should assess novel ways to improve dental students' knowledge and awareness of oral and oropharyngeal cancer, including HPV as a risk factor, and examine methods to develop dental students' confidence in communicating with patients on the subject.

References

1. Khode SR, Dwivedi RC, Rhys-Evans P, Kazi R. Exploring the link between human papilloma virus and oral and oropharyngeal cancers. *J Cancer Res Ther* [Internet]. 2014;**10**(3):492–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25313727>
2. Marur S, D'Souza G, Westra WH, Forastiere AA. HPV-associated head and neck cancer: A virus-related cancer epidemic. Vol. 11, *The Lancet Oncology*. 2010. p. 781–9.
3. Hocking JS, Stein A, Conway EL, et al. Head and neck cancer in Australia between 1982 and 2005 show increasing incidence of potentially HPV-associated oropharyngeal cancers. *Br J Cancer*. 2011;**104**(5):886–91.
4. Evans M, Newcombe R, Fiander A, et al. Human Papillomavirus-associated oropharyngeal cancer: An observational study of diagnosis, prevalence and prognosis in a UK population. *BMC Cancer*. 2013;**13**.
5. Heath S, Willis V, Allan K, et al. Clinically significant human papilloma virus in squamous cell carcinoma of the head and neck in UK practice. *Clin Oncol* [Internet]. 2012 Feb [cited 2018 Jun 28];**24**(1):e18–23. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21752613>
6. Schache AG, Powell NG, Cuschieri KS, et al. HPV-related oropharynx cancer in the United Kingdom: An evolution in the understanding of disease etiology. *Cancer Res*. 2016;**76**(22):6598–606.
7. D'Souza G, Kreimer AR, Viscidi R, et al. Case–Control Study of Human Papillomavirus and Oropharyngeal Cancer. *N Engl J Med* [Internet]. 2007;**356**(19):1944–56. Available from: <http://www.nejm.org/doi/abs/10.1056/NEJMoa065497>
8. Heck JE, Berthiller J, Vaccarella S, et al. Sexual behaviours and the risk of head and neck cancers: A pooled analysis in the International Head and Neck Cancer Epidemiology (INHANCE) consortium. *Int J Epidemiol*. 2010;**39**(1):166–81.
9. Smith EM, Ritchie JM, Summersgill KF, et al. Age, sexual behavior and human papillomavirus infection in oral cavity and oropharyngeal cancers. *Int J Cancer*. 2004;**108**(5):766–72.
10. D'Souza G, Agrawal Y, Halpern J, Bodison S, Gillison ML. Oral Sexual Behaviors

- Associated with Prevalent Oral Human Papillomavirus Infection. *J Infect Dis* [Internet]. 2009;**199**(9):1263–9. Available from: <https://academic.oup.com/jid/article-lookup/doi/10.1086/597755>
11. Ang KK, Harris J, Wheeler R, et al. Human Papillomavirus and Survival of Patients with Oropharyngeal Cancer. *N Engl J Med* [Internet]. 2010;**363**(1):24–35. Available from: <http://www.nejm.org/doi/abs/10.1056/NEJMoa0912217>
 12. Gillison ML. Human papillomavirus-associated head and neck cancer is a distinct epidemiologic, clinical, and molecular entity. Vol. 31, *Seminars in Oncology*. 2004. p. 744–54.
 13. Sudhoff HH, Schwarze HP, Winder D, et al. Evidence for a causal association for HPV in head and neck cancers. Vol. 268, *European Archives of Oto-Rhino-Laryngology*. 2011. p. 1541–7.
 14. Masterson L, Moualed D, Liu ZW, et al. De-escalation treatment protocols for human papillomavirus-associated oropharyngeal squamous cell carcinoma: A systematic review and meta-analysis of current clinical trials. *Eur J Cancer*. 2014;**50**(15):2636–48.
 15. McCann PJ, Sweeney MP, Gibson J, Bagg J. Training in oral disease, diagnosis and treatment for medical students and doctors in the United Kingdom. *Br J Oral Maxillofac Surg*. 2005;**43**(1):61–4.
 16. Shanks LA, Walker TWM, McCann PJ, Kerin MJ. Oral cavity examination: Beyond the core curriculum? *Br J Oral Maxillofac Surg*. 2011;**49**(8):640–2.
 17. Valk R a. Van Der, Walker TWM. Newly qualified dentists and the Human Papilloma Virus (HPV). *Br J Oral Maxillofac Surg* [Internet]. 2014 Oct 1 [cited 2018 Jun 28];**52**(8):e121. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0266435614005026>
 18. Fried JL. Confronting human papilloma virus/oropharyngeal cancer: A model for interprofessional collaboration. *J Evid Based Dent Pract*. 2014;**14**(SUPPL.).
 19. Daley E, Debate R, Dodd V, et al. Exploring awareness, attitudes, and perceived role among oral health providers regarding HPV-related oral cancers. *J Public Health Dent*. 2011;**71**(2):136–42.
 20. NHS Dental Statistics. NHS Dental Statistics for England - 2016-17, Second Quarterly Report [Internet]. 2017 [cited 2018 Jun 28]. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/nhs-dental->

- statistics/nhs-dental-statistics-for-england-2016-17-second-quarterly-report
21. Mesher D, Panwar K, Thomas SL, et al. The Impact of the National HPV Vaccination Program in England Using the Bivalent HPV Vaccine: Surveillance of Type-Specific HPV in Young Females, 2010–2016. *J Infect Dis* [Internet]. 2018 Jun 18 [cited 2018 Jun 28];**jiy249**. Available from: <https://academic.oup.com/jid/article/doi/10.1093/infdis/jiy249/5025126>
 22. British Dental Association. Overwhelming support from dentists for gender neutral HPV vaccinations [Internet]. British Dental Association. 2017 [cited 2018 Jun 28]. Available from: <https://bda.org/news-centre/press-releases/overwhelming-support-from-dentists-for-gender-neutral-hpv-vaccinations>
 23. Department of Health and Social Care. HPV vaccination programme for men who have sex with men - GOV.UK [Internet]. gov.uk. 2018 [cited 2018 Jun 28]. Available from: <https://www.gov.uk/government/news/hpv-vaccination-programme-for-men-who-have-sex-with-men>
 24. Rose D. HPV jabs will be offered to thousands of teen boys on the NHS [Internet]. The Mail on Sunday. 2018 [cited 2018 Jun 28]. Available from: <http://www.dailymail.co.uk/news/article-5852705/HPV-jabs-offered-thousands-teen-boys-NHS.html>
 25. Thavaraj S, Stokes A, Odell E. Is head and neck cancer infectious? Human papilloma virus in oral and pharyngeal cancer. *Fac Dent J* [Internet]. 2010 [cited 2018 Jun 28];**1**(3):108–13. Available from: [https://kclpure.kcl.ac.uk/portal/en/publications/is-head-and-neck-cancer-infectious-human-papilloma-virus-in-oral-and-pharyngeal-cancer\(eddc6d9-3c5e-4a8e-adce-8506ac8b7541\).html](https://kclpure.kcl.ac.uk/portal/en/publications/is-head-and-neck-cancer-infectious-human-papilloma-virus-in-oral-and-pharyngeal-cancer(eddc6d9-3c5e-4a8e-adce-8506ac8b7541).html)
 26. Mitchell DA. HPV vaccination and the prevention of oropharyngeal cancer: two steps forward and one to the side? *Fac Dent J* [Internet]. 2015 Jan 8 [cited 2018 Jun 28];**6**(1):14–7. Available from: <http://publishing.rcseng.ac.uk/doi/10.1308/204268515X14174408395768>
 27. Hancocks S. HPV vaccine for boys. *British Dental Journal*. 2014;**216**(1):1.
 28. Baker P. Vaccinating boys against HPV. Vol. 215, *British Dental Journal*. 2013. p. 543–4.
 29. Walker TWM, Fleming C, Kerai A, et al. Are dental students well-equipped to deal

- with difficult communication situations? *Br Dent J*. 2018;**224**(3):163–8.
30. Rowan SD, Hu SL, Brotzman JS, et al. Knowledge Assessment of the Dental Community in Texas on the Role of Human Papilloma Virus in Oropharyngeal Cancer. *Tex Dent J* [Internet]. 2015 Aug [cited 2018 Jun 28];**132**(8):528–36. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26489292>
 31. General Dental Council. Preparing for practice. 2012;1–104.
 32. Walker TWM, Gately F, Stagnell S, et al. Can UK undergraduate dental programmes provide training in non-surgical facial aesthetics? *Br Dent J*. 2017;**222**(12):949–53.

Tables

Table 1: Tabulated free text responses by year group

Year 3	
Knowledge	<i>Risk factors/signs/symptoms</i>
	<i>I don't really know anything, just that it can occur</i>
	<i>Likelihood of HPV contraction, chances of Ca development, mode of transmission</i>
	<i>Everything. The only health link I was aware of was HPV and cervical cancer</i>
	<i>How to prevent infection. The exact implications of HPV</i>
	<i>How to prevent infection. Treatments</i>
	<i>Prevalence.</i>
	<i>What can we do to reduce the risk?</i>
	<i>Signs and symptoms, consequences, action that needs to be taken</i>
	<i>Risks and prevalence</i>
Communication	<i>Where to advise patients to gain more info/vaccination</i>
	<i>A lecture on it and how to give advice</i>
	<i>What to tell patients</i>
	<i>How to approach subject with patient</i>
	<i>How can we make people more aware without worrying them?</i>
Year 4	
Knowledge	<i>Vaccines available</i>
	<i>How to identify risks to patients and clinical signs of symptoms in patients</i>
	<i>Be able to confidently diagnose oral warts</i>
	<i>The risks / benefits of HPV vaccine. How it acts. Who it's available to.</i>
	<i>Causes. How caught. Consequences. Treatment</i>
	<i>Causes, signs and symptoms, prognosis, when to refer</i>
	<i>Exactly how it impacts oral health and how big the risk is for oropharyngeal cancer</i>
	<i>We have not had any practical teaching</i>
	<i>Its aetiology, effects, public health info and programmes currently in place</i>
	<i>Key presentation signs.</i>
Communication	<i>Risks of vaccine. Stats.</i>
	<i>How to communicate this risk to patients with reference to HPV. Communication training to extract this info from patients</i>
	<i>We haven't had much teaching on HPV communication skills,</i>
	<i>How do you talk to patients about HPV?</i>
	<i>Advice to give patients. Who to target the advice towards. ?screening programme</i>
	<i>Ways to discuss with patients their risk of oropharyngeal cancer.</i>

	<i>When appropriate to bring this up with patients</i>
	<i>Who you should tell and when</i>
	<i>How to discuss the issue sensitively with patients</i>
Year 5	
Knowledge	<i>About the possible vaccines - when they are available and what protection they convey the patients</i>
	<i>About the vaccine, Epidemiology of HPV and oral cancer</i>
	<i>Exact effects</i>
	<i>The vaccine</i>
	<i>Management</i>
	<i>What to do if you have a problem</i>
	<i>HPV prevalence, treatment, % of people that develop cancer, screening/detection</i>
	<i>How to prevent. Who to talk to about it. Consequences and risk rates</i>
Communication	<i>Statistics. Can an oral 'wart' progress to cancer?</i>
	<i>What to tell patients</i>
	<i>Advising patients</i>
	<i>What we should be advising to patients</i>
	<i>How to advise patients</i>
	<i>What should be advise patients to reduce risk? Who should we give this advice?</i>