McNally, K. E., & Cullen, P. J. (2018). Endosomal Retrieval of Cargo: Retromer Is Not Alone. *Trends in Cell Biology*, 28(10), 807-822.
<https://doi.org/10.1016/j.tcb.2018.06.005>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.tcb.2018.06.005](https://doi.org/10.1016/j.tcb.2018.06.005)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cell Press at <https://www.sciencedirect.com/science/article/pii/S0962892418301065>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Endosomal retrieval of cargo: retromer is not alone

Kerrie E. McNally¹ and Peter J. Cullen^{1*}

¹ School of Biochemistry, Biomedical Sciences Building, University of Bristol, Bristol
BS8 1TD, UK.

*Correspondence: Pete.cullen@bristol.ac.uk

Keywords: endosomes, retromer, retriever, WASH complex, CCC complex, sorting
nexin (SNX)

Abstract

Endosomes are major protein sorting stations in cells. Endosomal localized multi-protein complexes sort integral proteins, including signaling receptors, nutrient transporters, adhesion molecules and lysosomal hydrolase receptors, for lysosomal degradation or conversely for retrieval and subsequent recycling to various membrane compartments. Correct endosomal sorting of these proteins is essential for maintaining cellular homeostasis, with defects in endosomal sorting implicated in various human pathologies including neurodegenerative disorders. Retromer, an ancient multi-protein complex, is essential for the retrieval and recycling of hundreds of transmembrane proteins. Whilst retromer is a major player in endosomal retrieval and recycling, several studies have recently identified retrieval mechanisms that are independent of retromer. Here we review endosomal retrieval complexes, with a focus on recently discovered retromer-independent mechanisms.

24 **Endosomal sorting of cargo is achieved by multi-protein complexes**

25 The endocytic pathway consists of a series of membrane trafficking steps which act
26 together to regulate cell surface levels of lipids and integral proteins (referred to as
27 cargo). Following endocytosis cargo enter the endosomal network and are sorted for
28 one of two fates; degradation as mature endosomes fuse with lysosomes or retrieval
29 from this fate. Following retrieval, cargo are recycled back to the plasma membrane,
30 the TGN (*trans*-Golgi Network) or to other organelles such as melanosomes (**Figure**
31 **1**) [1, 2]. Thus endosomal sorting of cargo for degradation or retrieval and recycling
32 tightly regulates the composition of the cell surface and hence the ability of the cell to
33 sense and respond to its environment. While endocytosis is a key entry point into the
34 endosomal network, cargo such as the lysosomal hydrolase receptors CI-MPR
35 (cation-independent mannose 6-phosphate receptor) and sortilin undergo iterative
36 rounds of trafficking between the biosynthetic pathway and endosomes, in order to
37 deliver lysosomal hydrolases to the endosomal lumen (**Figure 1**) [2]. Therefore,
38 endosomal retrieval and recycling is essential not only for plasma membrane
39 homeostasis but also for maintaining lysosomal health [3]. Consistent with this,
40 defective endosomal sorting is associated with many human diseases, including
41 Parkinson's disease [4, 5].

42 Endosomal sorting requires a series of spatially and temporally regulated multi-protein
43 complexes to facilitate cargo enrichment and membrane deformation to generate
44 cargo-enriched budding profiles that undergo scission to generate cargo-enriched
45 transport carriers [6]. Whilst ESCRT (endosomal sorting complexes required for
46 transport) complexes specifically recognize ubiquitinated cargo and sort these into
47 **ILVs (intra-luminal vesicles; see Glossary)** for lysosomal degradation [7], it was
48 long thought that retrieval and recycling of cargo was achieved through **geometric-**

49 **based principles** and bulk membrane flow [8]. However, many studies have
50 established that endosomal retrieval and recycling is achieved by sequence-
51 dependent sorting of cargo by endosomal localized multi-protein machineries and
52 branched actin [9, 10]. One of the best characterized endosomal retrieval complexes
53 is the evolutionary conserved retromer complex [2, 11]. In human cultured cells,
54 retromer regulates cell surface levels of over 100 integral plasma membrane proteins
55 as well as mediating recycling of cargo from the endosome to the TGN and to
56 lysosomal-related organelles [1, 2, 12]. However, not all cargo require retromer for
57 their retrieval and recycling [13-16]. Recently retriever and the CCC complex have
58 been identified as additional evolutionary conserved endosomal complexes [14, 17].
59 Importantly, together these complexes provide a mechanism for retromer-independent
60 endosomal sorting of additional cargo [14]. The WASH complex, which promotes
61 endosomal branched actin polymerization, plays essential roles in both retromer-
62 dependent and retromer-independent pathways, demonstrating that the WASH
63 complex is fundamental for endosomal cargo sorting [14, 17-19]. Here, we review the
64 recent advancements in our understanding of endosomal cargo recycling, with a focus
65 on the first step of this process - retrieval of cargo from a lysosomal degradative fate
66 **(Figure 1).**

67

68 **Lysosomal degradation vs retrieval and recycling**

69 Once cargo enter the endosomal network, a major fate decision is made; are cargo
70 targeted for lysosomal degradation or will they be retrieved and subsequently recycled
71 back to the relevant membrane compartment [2]? Integral membrane cargo to be
72 degraded are marked through lysine-63 (K-63)-linked ubiquitination [20]. The

73 evolutionary conserved ESCRT complexes sequentially localize to the cytosolic face
74 of the endosome and drive cargo enrichment, membrane deformation, and vesicle
75 scission to generate cargo-enriched ILVs. Upon fusion of late endosomes with
76 lysosomes, cargo within the ILVs are degraded (**Figure 1**) [7]. Whilst ubiquitination is
77 a key marker of cargo to be degraded, it should be noted that ubiquitin-independent
78 cargo incorporation into ILVs has been reported. ALIX (apoptosis-linked gene 2-
79 interacting protein X) interacts with sequence-specific motifs within cargo and acts as
80 an adaptor for the ESCRT pathway, bypassing the requirement of ubiquitination [21].

81 Interestingly, upon suppression or knock-out of retromer, retriever, the WASH or CCC
82 complexes (see subsequent discussions), cargo are degraded in lysosomes
83 suggesting that in the absence of these sorting complexes, cargo degradation may be
84 the default pathway [12, 14]. Conversely, suppression or deletion of proteins involved
85 in membrane remodelling, **carrier scission** and **carrier transport** and fusion often
86 results in the accumulation of cargo in endosomal or vesicular compartments. For
87 example, depletion of retromer results in lysosomal degradation of the glucose
88 transporter GLUT1, whereas suppression of SNX-BARs, which are important for
89 carrier formation, results in endosomal accumulation of the transporter [12]. Retromer,
90 retriever, the CCC and WASH complexes will therefore be referred to as retrieval
91 complexes because they are required for retrieval of cargo away from lysosomal
92 degradation.

93

94 **Retrieval and degradative functions of the endosome are spatially segregated**

95 The antagonistic roles of ESCRT and retrieval complexes in cargo sorting are spatially
96 segregated on the cytosolic leaflet of the endosome (**Figure 1**). ESCRT and clathrin

co-localize in endosomal sub-domains which are distinct from sub-domains containing endosomal retrieval complexes such as retriever, retromer and the WASH complex [14, 22, 23]. The formation and maintenance of these distinct subdomains may spatially segregate cargo into either degradative or retrieval sub-domains based upon recognition of ubiquitinated cargo by ESCRT or sequence-specific recognition of cargo by retrieval complexes. Whilst the precise mechanism by which cargo leak into the lysosomal pathway in the absence of retrieval complexes is not established, it could be speculated that cargo are no longer spatially segregated away from ESCRT sub-domains and instead become incorporated into ILVs. The dependence of this process on cargo ubiquitin status or recognition by ALIX remains to be investigated.

The WASH complex organises endosomal retrieval sub-domains

Flat clathrin lattices are proposed to scaffold the ESCRT subdomain whereas the formation of endosomal retrieval sub-domains, in which cargo and their respective retrieval complexes are enriched, is driven by branched actin polymerization by the **Arp2/3** complex [10, 22-25]. WASH1 (WASHC1) is an endosomally localized nucleation promoting factor for Arp2/3 and it exists as part of the pentameric WASH complex along with (new nomenclature for the subunits are shown in brackets): FAM21A/B/C (WASHC2A/B/C), CCDC53 (WASHC3), SWIP also known as KIAA1033 (WASHC4) and strumpellin also known as KIAA0196 (WASHC5) (**Figure 1**) [26, 27]. The WASH complex is of ancient origin but has not been retained by Fungi [28]. Depolymerization of endosomal actin causes distinct WASH labelled retrieval subdomains to merge indicating that endosomal actin is important for the formation and maintenance of retrieval subdomains [23]. However WASH labelled subdomains

remain spatially distinct to degradative sub-domains upon actin depolymerization [23]. Depletion of the WASH interactor RME-8 (receptor-mediated endocytosis-8, also known as DNAJC13) or the **sorting nexin** SNX1 (see later discussion) however results in the mixing of degradative and retrieval subdomains in *Caenorhabditis elegans* and HeLa cells, a process speculated to occur due to defects in removing clathrin from the retrieval subdomain [22, 29].

Moreover the WASH complex also organizes the retrieval sub-domain by acting as an endosomal scaffold for the recruitment of additional multi-protein retrieval complexes and regulators of endosomal sorting to the retrieval sub-domain. The long unstructured tail of FAM21 interacts with and recruits the CCC complex along with retriever, to the endosomal membrane (see subsequent discussions) [14, 17]. Moreover, the tail of FAM21 interacts with FKBP15 (FK506-binding protein 15) and mediates its localization to endosomes, although a functional role for FKBP15 is yet to be reported [30-32].

Strikingly, in fibroblasts derived from WASH knock-out mice and melanocytes from strumpellin knock-out mice, the endosomal and lysosomal networks become collapsed to the peri-nuclear region suggesting that the WASH complex is also important for the spatial organization of the endo-lysosomal system although the underlying mechanisms remain unknown [33, 34].

The WASH complex is essential for the retrieval and recycling of many cargo

Consistent with the notion that the WASH complex is essential for retrieval sub-domain formation and maintenance as well as endosomal positioning, the WASH complex is necessary for endosomal retrieval and recycling of many cargo to both the TGN and the plasma membrane, including: CI-MPR [35], the retromer-independent cargo $\alpha_5\beta_1$

integrin [14, 16, 36], GLUT1 [34], TCR (T cell receptor) [37], TfnR (transferrin receptor) [26], β 2-adrenergic receptor [10, 38] and LDLR (low-density lipoprotein receptor) [18]. In *Dictyostelium discoideum*, the WASH complex is required for recycling of phagocytic receptors from phagosomes to the plasma membrane [39]. Moreover, WASH and its ability to polymerize actin was required for the removal of **v-ATPase** from lysosomal membranes, to allow lysosome neutralization prior to exocytosis of indigestible material [40]. The role for WASH in recycling of the v-ATPase was also demonstrated in WASH knock-out *Drosophila melanogaster* [41]. In addition hemocytes from WASH knock-out *Drosophila* display defects in cell spreading and cell migration, indicative of impaired integrin recycling [41].

Besides the role of the WASH complex and actin in endosomal organization, it was reported that depletion of components of the WASH complex resulted in elongated endosomal recycling tubules, suggesting that the WASH complex is required for tubule scission [26, 35]. However WASH knock-out MEFs (mouse embryonic fibroblasts) did not display this phenotype [34]. Endosomal actin also helps to generate and stabilize endosomal tubules into which cargo are sorted in a sequence-dependent manner [10]. Interestingly, β 2-adrenergic receptors that lack a PDZbm (**PDZ binding motif**), required for retrieval and recycling, but instead harbor an actin-binding domain can recycle normally, indicating that interactions between endosomal actin and cargo may be sufficient for recycling [10].

Recruitment of the WASH complex to endosomes

Retromer directly binds to the multiple **LFa** (leucine-phenylalanine-a series of acidic residues) repeats in the carboxy-terminal tail of FAM21 and this was reported to be

sufficient and necessary for endosomal recruitment of the WASH complex [31, 32, 42, 43]. However, in a retromer knock-out HeLa cell line, a significant proportion of FAM21 remains associated with the endosomal network, establishing that in human cells the endosomal association of the WASH complex is mediated through both retromer-dependent and retromer-independent mechanisms [14]. This agrees with work in *Dictyostelium* describing the retromer-independent recruitment of the WASH complex [39]. Importantly, retromer-independent recruitment of the WASH complex may explain how endosomal retrieval of some cargo requires the WASH complex but occurs independently of retromer (see subsequent discussions) [14].

The mechanism of retromer-independent recruitment of the WASH complex to endosomal membranes remains poorly characterized. However, purified WASH complex displayed the ability to associate with liposomes and the carboxy-terminal tail of FAM21 was shown to bind to PI3P and PI(3,5)P₂, phosphoinositides found enriched on the cytosolic leaflet of early and late endosomes, suggesting that intrinsic lipid-association of the WASH complex may be important for retromer-independent recruitment to endosomal membranes [26, 27, 44]. In addition, interactions between the WASH complex and endosomally localized proteins including the sorting nexin SNX27 (see subsequent discussion) and RME-8 may also underlie retromer-independent recruitment of the WASH complex [12, 29].

Retromer

Retromer is an ancient endosomal sorting complex, conserved in all eukaryotes [45]. Retromer was first discovered in the budding yeast *Saccharomyces cerevisiae*, where retromer exists as a pentameric complex, composed of a dimer of Vps5p and Vps17p

193 and a trimer of Vps26p, Vps35p and Vps29p [11, 46]. In mammals the genes encoding
 194 retromer have undergone duplication and divergence [45]. SNX1 and SNX2 are
 195 orthologues of the yeast *Vps5* gene, whilst SNX5, SNX6 and SNX32 are orthologues
 196 of *Vps17* [45]. These proteins contain a **BAR domain** (Bin/amphiphysin/Rvs) which
 197 can sense and induce curvature, generating endosomal tubules for cargo recycling
 198 [47, 48]. SNX1 or SNX2 dimerize with SNX5, SNX6 or SNX32 [49]. In mammals
 199 VPS26 exists as three paralogues: VPS26A, VPS26B and DSCR3 [45]. DSCR3 is not
 200 a sub-unit of retromer and instead incorporates into the retriever complex (see
 201 subsequent discussion) [14]. It is important to note that whilst retromer is highly
 202 evolutionary conserved, mammalian retromer is not an obligatory pentameric
 203 complex. The mammalian VPS26, VPS35 and VPS29 proteins exist as a hetero-trimer
 204 which does not form a stable complex in solution with the SNX-BAR dimer [50]. When
 205 discussing mammalian retromer, the membrane deformation SNX-BAR hetero-dimer
 206 will subsequently be referred to as the 'retromer linked SNX-BAR' complex whereas
 207 the VPS26:VPS35:VPS29 hetero-trimer will be referred to simply as 'retromer' [13].

208 Retromer was originally identified as being required for endosome to TGN retrograde
 209 transport of multiple cargo including Vps10p, a receptor in *S. cerevisiae* required for
 210 delivery of hydrolases such as carboxypeptidase Y, from the Golgi to the vacuole (the
 211 yeast equivalent of the lysosome) and its mammalian homologues sorLA and sortilin,
 212 the lysosomal hydrolase receptor CI-MPR and the mammalian iron transporter DMT1-
 213 II [11, 46, 51-57]. Retromer engages cargo containing a ØX(L/M) (where Ø represents
 214 an aromatic amino acid and x represents any amino acid) motif in their cytosolic tail
 215 and VPS26 directly binds a FANSHY motif within the cytoplasmic tail of sorLA [52, 57-
 216 59].

217 Whilst some cargo have been reported to directly bind to retromer, it was discovered
218 that a 'cargo adaptor', SNX3, could directly engage cargo such as the yeast iron
219 transporter Fet3p-Ftr1p and Wntless whilst simultaneously associating with retromer
220 for retrieval of these cargo away from degradation and for their recycling back to the
221 TGN [59-62]. In addition, retromer can retrieve and recycle cargo back to the plasma
222 membrane and this often occurs through association with a distinct retromer cargo
223 adaptor, SNX27, which recognises a PDZbm within the cytosolic domain of integral
224 proteins [12, 38, 63, 64].

225 Retromer is suggested to function by directly engaging specific sequences in the
226 cytosolic tails of cargo through its VPS26:VPS35:VPS29 hetero-trimer or through
227 association with SNX3 or SNX27 [38, 59, 60]. Cargo are enriched and corralled, with
228 the aid of WASH-dependent actin polymerization, into endosomal tubules generated
229 by the retromer-linked SNX-BAR hetero-dimers for transport to target membranes [1].
230 Recently super-resolution imaging established that retromer cargo destined for the
231 plasma membrane and the TGN are sorted into the same tubular carriers, suggesting
232 that recycling to either membrane occurs downstream of retromer [65].

233

234 **Organization and structure of retromer**

235 VPS35 forms an extended central scaffold with VPS26 and VPS29 independently
236 engaging VPS35 at the amino- and carboxy-terminals respectively (**Figure 2 A**) [50,
237 59, 66]. VPS35 is composed of 33 helices, which form 16 pairs of antiparallel α -helices,
238 a domain structure referred to as a HEAT (Huntington/EF3/PP2A/TOR1) repeat.
239 These HEAT repeats form an extended α -helical solenoid structure which is slightly
240 curved [59, 66]. At the carboxy-terminus of VPS35, the solenoid structure of VPS35

wraps itself around VPS29 which forms a fold similar to that found in phosphoesterases, although no phosphoesterase activity of VPS29 is observed *in vitro* (**Figure 2 A**) [66, 67]. VPS26A and VPS26B adopt arrestin-like folds, which consist of two beta-sandwich domains connected by a flexible linker and a polar core (**Figure 2 A**) [68, 69].

SNX3 binds to the VPS26-VPS35 interface through its amino terminal flexible region and its **PX domain**. The binding of retromer to SNX3 causes a conformational change in VPS26 which opens up a hydrophobic, cargo-binding pocket in the interface between the carboxy-terminal lobe of VPS26 and the PX domain of SNX3 [59]. SNX27 directly binds to VPS26 through SNX27's PDZ (Postsynaptic density 95 – Disc large – ZO1) domain [12, 70]. The **PDZ domain** of SNX27 also mediates the interaction with cargo, although through a distinct binding site within the PDZ domain [70, 71]. SNX27's affinity for cargo is increased when SNX27 is bound to VPS26 thus coupling cargo recognition with retromer association [70].

Not all cargo are retrieved away from lysosomal degradation by retromer

The ability of retromer to directly interact with cargo and via cargo adaptors such as SNX3 and SNX27 increases the repertoire of cargo retromer can retrieve and recycle in a sequence-dependent manner, thus increasing the plethora of cellular functions for which retromer mediated retrieval and recycling is required [72]. Quantitative proteomic analysis of plasma membrane integral proteins under retromer or SNX27 suppression indicated that retromer is required for the endosome to plasma membrane retrieval and recycling of over one hundred cargo including proteins involved in cell adhesion, ion transport and amino acid transport [12]. It is therefore not surprising that

mutations in retromer and retromer-associated proteins have been associated with various human diseases including Parkinson's disease (see [3, 73] for recent reviews). Yet, even multiple SNX-retromer complexes are unlikely to account for the sequence-dependent retrieval and recycling of all cargo transiting through the endosomal system, suggesting that additional retrieval and recycling pathways exist. Consistent with this, two papers have recently reported that the retromer linked SNX-BAR proteins (SNX1/SNX2:SNX5/SNX6/SNX32) can bind to cargo such as CI-MPR and IGF1R (insulin-like growth factor 1 receptor) and facilitate their retrieval and recycling back to the TGN or the plasma membrane respectively, independently of the retromer VPS26:VPS35:VPS29 trimer [13, 15]. Further support for the existence of retromer-independent retrieval pathways came from studies on the SNX27-related sorting nexin, SNX17. SNX17 and SNX27 both contain a **FERM** (4.1/ezrin/radixin/moesin) **domain** but SNX17 lacks the amino-terminal PDZ domain found in SNX27 required for binding to PDZbm in cargo and for interactions with retromer [63, 74, 75]. The FERM domain of SNX17 mediates the binding to NPx(Y/F) motifs within the cytosolic tails of cargo such as LRP1 (Low density lipoprotein receptor-related protein 1), LDLR and $\alpha_5\beta_1$ integrin and upon suppression of SNX17, these cargo are degraded in lysosomes [16, 76-78]. A global proteomic approach in HeLa cells revealed that SNX17 regulates the plasma membrane levels of over 200 cargo [14]. Consistent with the inability of SNX17 to engage retromer, the recycling of the majority of these cargo were not affected by SNX27 or retromer suppression [12, 14, 16]. SNX17-dependent, retromer-independent retrieval of cargo instead relies upon the recently discovered ancient endosomal retrieval complex, retriever [14].

Retriever is a 'retromer-like' retrieval complex

Retriever is a hetero-trimer consisting of DSCR3, C16orf62 (chromosome 16 open reading frame 62) and the retromer sub-unit VPS29 (**Figure 2 A**). Suppression or knock-out of retriever sub-units or perturbations in the ability of retriever to interact with SNX17 results in the lysosomal degradation of SNX17 cargo such as $\alpha_5\beta_1$ integrin. Importantly, knock-out of VPS35, the core component of retromer, does not give this phenotype demonstrating that retriever-dependent sorting of cargo is independent of retromer [14].

Retriever shares several similarities with retromer, for example, both complexes are ubiquitously expressed and are found in the last common eukaryotic ancestor [28, 45]. However, DSCR3 and C16orf62 have been selectively lost, along with the WASH complex and the CCC complex (see later discussion) in all Fungi [14, 28, 45] (**Figure 2 B**) In addition, in HeLa cells the estimated protein copy number of retromer sub-units is much higher than the retriever sub-units DSCR3 and C16orf62 (**Figure 2 B**) [79].

Retriever and retromer share similarities in their structural composition. The most obvious similarity is that VPS29 is a sub-unit in both retromer and retriever. DSCR3, as previously discussed, is a known paralogue of VPS26 [45, 80]. Furthermore, although C16orf62 and VPS35 share little residue conservation, C16orf62 is predicted to contain HEAT repeats, similar to those found in VPS35. Therefore retriever and retromer are hetero-trimers which contain a VPS29 subunit, a protein with an arrestin-like fold (either DSCR3 or VPS26A/B respectively) and a protein containing a series of HEAT-repeats (C16orf62 or VPS35 respectively) (**Figure 2 A**) [14].

312 Immuno-precipitations in DSCR3 knock-out HeLa cells indicate that SNX17 may
313 interact with retriever via DSCR3, although evidence of a direct interaction is currently
314 lacking [14]. This is reminiscent of SNX27 engagement with the equivalent subunit of
315 retromer, VPS26 [12, 70]. SNX31, which is expressed mainly in the urinary tract, is
316 closely related to SNX17 and is also important for the recycling of various integrins
317 [81]. Both SNX17 and SNX31 possess the conserved carboxy-terminal motif that is
318 necessary and sufficient to bind to retriever, indicating that SNX31 may play an
319 important role in retrieving cell-type specific cargo [14]. It has yet to be established
320 whether retriever can bind to cargo directly, as has been reported for retromer [52,
321 59].

322 Retriever and retromer undertake distinct cargo retrieval itineraries but are localized
323 to the same endosomal retrieval sub-domain [14, 17]. Retromer does not contain any
324 membrane binding domains and instead relies upon interactions with SNX3 and the
325 late endosomal/lysosomal Rab GTPase Rab7 for membrane association to early and
326 early-to-late transitioning endosomes (**Figure 3**) [82-84]. Like retromer, retriever is not
327 predicted to bind membranes and therefore retriever also relies upon protein: protein
328 interactions for its recruitment to endosomes. In contrast to retromer, the recruitment
329 of retriever to endosomal sub-domains does not depend upon SNX3 and Rab7 but
330 through interactions with an additional endosomal sorting complex, the CCC complex
331 (**Figure 3**) [14].

332

333 **The CCC complex**

334 The CCC complex colocalizes with retromer, retriever and the WASH complex on
335 endosomes and is required for the endosomal retrieval and recycling of LDLR, Notch2

and ATP7A to the plasma membrane [14, 17, 18, 85]. The CCC complex comprises of a heterodimer of CCDC22 (coiled-coil domain-containing protein 22) and CCDC93 (coiled-coil domain-containing protein 93) and the association of COMMD (copper metabolism MURR1 domain-containing) proteins [17, 86].

The COMMDs are a family of 10 members (named COMMD 1-10), largely conserved throughout evolution [87, 88]. Whilst vertebrate genomes contain all 10 COMMD genes, *Drosophila* only have 5 COMMD genes (COMMD 2, 3, 4, 5 and 10), *C. elegans* only encodes COMMD4 whereas *Dicytostelium* possesses all 10 genes [87-89]. The defining characteristic of COMMDs is the presence of a unique carboxy-terminal COMM domain. The COMM domain is responsible for homo- and hetero-dimerization of COMMD proteins and there is evidence for preferential dimers existing in cells [85, 87]. In addition, the COMM domain has been shown to be essential for interactions between COMMD1 and the CCC complex subunits CCDC22 and CCDC93, a mechanism which probably underlies how every COMMD protein has the ability to associate with CCDC22 and CCDC93 [17, 86]. It should be noted that whilst the CCC complex has recently been implicated in endosomal retrieval and recycling of cargo, roles for these proteins, especially the COMMDs, in other cellular functions such as the hypoxia response and NF- κ B activation have been reported (see [89, 90] for recent reviews on COMMD1 function).

The retriever-CCC-WASH complex pathway

A full mechanistic understanding of how the CCC complex engages cargo and promotes its retrieval from degradation for subsequent recycling has not been achieved. However, like retromer, components of the CCC complex have been shown

to directly interact with the tail of FAM21, an interaction which is necessary for CCC complex localization [17, 31]. In agreement with the WASH complex being important for the function of the CCC complex, *Wash1* knock-out MEFs display endosomal accumulation of LDLR and increased degradation of the receptor, which can be rescued through treatment with **bafilomycin**, which prevents lysosomal acidification and thus pH-dependent degradation [18]. Furthermore, the reintroduction of a WASH mutant, that cannot activate Arp2/3, into *Wash1* knock-out MEFS did not rescue LDLR from lysosomal degradation, suggesting that branched actin polymerization is essential for CCC complex-dependent recycling of cargo [18].

Several proteomics studies have identified that the CCC complex interacts with components of retriever (**see Box 1**) [17, 28, 91, 92]. Moreover, components of the CCC complex were also identified as interactors of SNX17 [14]. SNX17 interactions with the CCC complex are dependent upon retriever, suggesting that SNX17 interacts indirectly with the CCC complex through retriever [14]. Furthermore, suppression of CCDC22 or CCDC93 results in the lysosomal degradation of $\alpha_5\beta_1$ integrin, indicating that SNX17, retriever and the CCC complex are functionally linked.

A model for how these complexes function together in retrieval of $\alpha_5\beta_1$ integrin was proposed when it was observed that suppression of the CCC complex results in the loss of endosomal association of retriever [14, 17]. Extending this further, knock-out of FAM21, which recruits the CCC complex to endosomes, also results in the loss of endosomal association of retriever and consequently lysosomal degradation of $\alpha_5\beta_1$ integrin [14]. It has previously been shown that the WASH complex is essential for the retrieval and recycling of $\alpha_5\beta_1$ integrin (or their homologues) in invasive ovarian cancer cells, mouse fibroblast cells, *Dictyostelium* and *Drosophila* [36, 39, 41, 93]. Similarly, the retrieval of LFA-1 (lymphocyte function-associated antigen 1) and TCR (T cell

385 receptor), two other SNX17-dependent cargo, from lysosomal degradation in T cells,
386 also depends upon the WASH complex [37, 94]. However, until recently the
387 mechanism linking SNX17 and the WASH complex, in a retromer-independent
388 manner, had not been described. The current model suggests that the WASH complex
389 is recruited to endosomes in retromer-dependent and retromer-independent
390 mechanisms, where it recruits the CCC complex through interactions with FAM21. The
391 CCC complex then recruits retriever to the retrieval sub-domain of endosomes where
392 it can engage SNX17 and its bound cargo, driving retrieval of cargo away from
393 degradation (**Figure 3**). Evolutionary conservation analysis infers that the WASH,
394 CCC and retriever complexes co-evolved, consistent with the notion that these
395 complexes depend on each other for localization and function [14, 28, 89].

396 Whilst this model suggests that the CCC complex is forming a bridge between retriever
397 and the WASH complex it must be stressed that it is highly unlikely that this is the only
398 function of the CCC complex. Indeed it has been proposed that the COMMDs
399 themselves may act as cargo-binding adaptors [18, 85]. Whilst many of the COMMD
400 proteins are ubiquitously expressed in tissues, some COMMDs such as COMMD9
401 have preferential expression in myeloid cells and the central nervous system [88].
402 Moreover, it has recently been shown that the CCC complex containing COMMD9
403 dimerized to COMMD5 or COMMD10 was involved in the retrieval and recycling of
404 Notch2 to the plasma membrane [85], whereas COMMD1 is involved in ATP7A and
405 LDLR recycling [17, 18]. This raises the possibility that distinct CCC complexes
406 containing different COMMD hetero-dimers exist, analogous to the SNX-retromers [2].
407 Consistent with this idea, COMMD9 and COMMD1 knock-out mice, whilst both
408 embryonic lethal, display different phenotypes indicating that they are not completely
409 redundant with one another [85, 95].

Furthermore, the COMM domain of COMMDs shares structural homology with the lipid binding **pleckstrin homology (PH) domains** [89, 96]. Recombinant COMMD1 was found to bind to liposomes containing PI(4,5)P₂, an interaction which required the COMM domain and promoted COMMD1 oligomerisation [96]. It is unknown whether the proposed lipid-binding ability of COMMDs plays a role in endosomal retrieval and recycling.

Concluding remarks and future perspectives

At the time of writing, four major, evolutionary conserved, endosomal retrieval multi-protein complexes (retromer, retriever, WASH and the CCC complexes) are thought to drive retrieval of cargo in a sequence-dependent manner away from a lysosomal degradative fate. These complexes are summarized in **Table 1**. The identification of retromer-independent pathways has expanded our knowledge of sequence-dependent cargo retrieval and recycling, with retriever likely to be responsible for retrieval and recycling of at least an additional 200 integral proteins back to the plasma membrane [14]. As yet unidentified cargo adaptors could feed into the retriever-CCC-WASH pathway and these complexes may directly engage sequence motifs in cargo, accounting for additional cargo retrieval events.

The importance of endosomal retrieval complexes in maintaining cellular homeostasis is highlighted by patients carrying mutations in genes encoding for these retrieval complexes (**Table 2**). Interestingly, mutations in CCC and WASH complex subunits present very similar patient phenotypes, such as intellectual disability, suggesting a shared mechanism of pathology (**Table 2**). Moreover, homozygous knock-out mouse models of COMMD1, COMMD6, COMMD9, COMMD10, WASH1, strumpellin are

434 embryonic lethal [34, 85, 95, 97, 98] and a point mutation in SNX17 in mice causes
435 congenital abnormalities such as an atrioventricular septal heart defect and duplex
436 kidneys [99, 100], demonstrating that these complexes play critical roles in
437 developmental processes.

438 Whilst the discovery of retromer-independent endosomal retrieval pathways have
439 allowed a greater understanding of the molecular mechanisms of endosomal sorting,
440 many outstanding questions remain (**see Outstanding Questions**). Future research
441 investigating the molecular mechanisms of endosomal cargo retrieval and recycling
442 and how they are regulated will answer these important questions.

443

Glossary

Arp2/3 (Actin-related protein 2/3) complex: A complex composed of seven subunits which nucleates branched actin polymerisation. However, Arp2/3 itself is not intrinsically active and requires activation by nucleation promoting factors such as the WASH complex.

Bafilomycin: An inhibitor of the vacuolar-H⁺-ATPase, preventing acidification of lysosomes.

BAR domain: BAR domains are a protein dimerization interface that dimerise to form a rigid banana-shaped structure. BAR domains are membrane curvature sensing domains and can also induce membrane curvature and tubulation of liposomes.

Carrier scission: Detachment of the nascent cargo enriched carrier from the donor membrane.

Carrier transport: Transport of the cargo enriched carrier from the donor to the acceptor membrane. Usually occurs via microtubule and actin based motors.

FERM domain (4.1/ezrin/radixin/moesin): A domain with a cloverleaf structure and binds to NPx(Y/F) motifs within proteins.

Geometric based principles: The recycling tubules display a high surface area to low volume ratio, thus transmembrane proteins entering the endosome are much more likely to be present in the endosomal tubules than the vesicular portion of the endosome which has a high volume to low surface area ratio. Therefore, unless cargo are specifically recognized by ESCRT, they are likely to be recycled back to the plasma membrane or the TGN based upon bulk membrane recycling.

Intra-luminal vesicles (ILVs): Vesicles formed from inward-budding of the endosomal membrane.

LFa motifs: Leucine-Phenylalanine-a series of acidic residues. Multiple motifs found in the tail of FAM21 that mediate binding to VPS35.

472 **PDZ binding motif:** Protein:protein interaction motif found in the cytosolic tails of
473 transmembrane cargo, usually at the extreme carboxy-termini. The consensus
474 sequence of a class I PDZbm is [T/S]-x- ϕ , where x denotes any amino acid and ϕ
475 represents an amino acid with a bulky hydrophobic group.

476 **PDZ domain (Postsynaptic density 95 – Disc large – ZO1):** PDZ domains interact
477 with PDZ binding motifs (PDZbm).

478 **Pleckstrin homology (PH) domain:** A phosphoinositide binding domain, interacts
479 with PI(3,4)P₂ and PI(3,4,5)P₃ and various other phosphoinositide species.

480 **PX domain (Phox homology):** A phosphoinositide binding domain, interacting
481 primarily with PI3P.

482 **Sorting nexin (SNX):** Sorting nexins are a family of proteins defined by the
483 presence of a PX domain with over 50% sequence similarity to the first member of
484 the family, SNX1. Some members of this protein family also contain extra functional
485 domains such as PDZ and FERM domains.

486 **v-ATPase:** vacuolar H⁺-ATPase. Proton pump that utilises ATP hydrolysis to pump
487 protons from the cytosol into the lumen of endosomes and lysosomes, generating
488 and maintaining an acidic pH in the lumen of these organelles.

489

490

491

492

493

494

495

496

497 **BOX 1:**

498 **The COMMander complex**

499 Genome-wide human interaction networks and phylogenetic analyses have
500 suggested the presence of a large, ancient, multi-protein complex named
501 'COMMander' because of the presence of a large number of COMMD proteins [28,
502 92, 101, 102]. The current evidence for COMMander has been recently reviewed,
503 suggesting that that the core COMMander complex consists of 14 proteins:
504 CCDC22, CCDC93, COMMD 1-10, C16orf62 and DSCR3 [89]. It could be
505 speculated that COMMander is a super-complex, composed of retriever and the
506 CCC complex. Whilst retriever can be purified from insect cells as a stable hetero-
507 trimer, the CCC complex is required for retriever localization, protein stability and
508 function [14, 103]. Moreover, their shared evolution, along with the WASH complex,
509 suggests a fundamental role for the assembly in cellular function [28]. Further
510 biochemical evidence will be needed to clarify the organization, definition and
511 functional role of COMMander.

512

513

514

515

516

517

518 **Table 1 Summary of endosomal retrieval complexes**

	Retromer	Retriever	CCC complex	WASH complex
Subunits	<ul style="list-style-type: none"> VPS26A/B VPS35 VPS29 	<ul style="list-style-type: none"> DSCR3 (VPS26C) C16orf62 (VPS35L) VPS29 	<ul style="list-style-type: none"> CCDC22 CCDC93 COMMD 1-10 	<ul style="list-style-type: none"> WASH (WASHC1) FAM21A/C (WASHC2A/C) CCDC53 (WASHC3) SWIP (WASHC4, KIAA1033) Strumpellin (WASHC5, KIAA0196)
Recruitment to endosomes	<ul style="list-style-type: none"> SNX3 Rab7 	<ul style="list-style-type: none"> CCC complex 	<ul style="list-style-type: none"> WASH complex 	<ul style="list-style-type: none"> Retromer Unknown retromer-independent mechanism Intrinsic lipid binding?
Direct cargo binding	<ul style="list-style-type: none"> VPS26 binds to FANSHY motifs Øx(L/M) (where Ø represents any hydrophobic amino acid and x represents any amino acid) motif 	<ul style="list-style-type: none"> Unknown 	<ul style="list-style-type: none"> Unknown Cargo possibly binding to COMMDs 	<ul style="list-style-type: none"> Unknown Cargo may bind actin
Cargo adaptors	<ul style="list-style-type: none"> SNX3 SNX27 	<ul style="list-style-type: none"> SNX17 SNX31 	<ul style="list-style-type: none"> COMMDs? 	

519

520 C16orf62; Chromosome 16 open reading frame 62, DSCR3; Downs syndrome critical
521 region 3, VPS35L; VPS35-like, CCDC; coiled-coil domain containing, WASH; Wiskott-
522 Aldrich syndrome protein and SCAR homologue, SWIP; Strumpellin and WASH

523 interacting protein, CCC complex; CCDC22, CCDC93 and COMMD, SNX; sorting
 524 nexin.

525

526 **Table 2 Human diseases associated with mutations in the retriever-CCC-WASH**
 527 **pathway**

Protein	Mutation	Disease	Examples of phenotypes	References
CCDC22	p.T17A	X-linked intellectual disability	<ul style="list-style-type: none"> Intellectual disability Ventricular and atrial septal defects Craniofacial dysmorphisms Hypercholesterolemia 	[18, 86, 104]
	p.Y557C	X-linked intellectual disability with symptoms of Ritscher/Schinzel syndrome	<ul style="list-style-type: none"> Intellectual disability Dandy-walker malformations Camptodactyly Ventricular septal defects 	[105]
Strumpellin	p.I226T	Hereditary spastic paraplegia (SPG8)	<ul style="list-style-type: none"> Lower limb weakness 	[106]
	p.N471D		<ul style="list-style-type: none"> Spasticity 	[107]
	p.R583S			[108]
	p.S591P			[109]
	p.L619F			[107]
	p.V620A			[110, 111]
	p.V626F			[107]
	p.G696A			[112]
	p.E713K			[113]
	g.ex11-15del			[108]
	c.3335+2T>A c.3335+4C>A	Ritscher–Schinzel/3C syndrome	<ul style="list-style-type: none"> Intellectual disability Hypertelorism 	[114]

	c.3335+8A>G		<ul style="list-style-type: none"> Atrial and ventricular septal defects vary within the patient cohort 	
SWIP	p.P1019R	Autosomal recessive intellectual disease	<ul style="list-style-type: none"> Intellectual disability Short stature Severe delays in motor development 	[115]

528

529 CCDC; coiled-coil domain containing, SWIP; Strumpellin and WASH interacting
530 protein

531

532

533 References

- 534 1. Gallon, M. and Cullen, P.J. (2015) Retromer and sorting nexins in endosomal
535 sorting. *Biochem Soc Trans* 43 (1), 33-47.
- 536 2. Burd, C. and Cullen, P.J. (2014) Retromer: a master conductor of endosome sorting.
537 *Cold Spring Harb Perspect Biol* 6 (2).
- 538 3. McMillan, K.J. et al. (2017) The emerging role of retromer in neuroprotection. *Curr*
539 *Opin Cell Biol* 47, 72-82.
- 540 4. Maxfield, F.R. (2014) Role of endosomes and lysosomes in human disease. *Cold*
541 *Spring Harb Perspect Biol* 6 (5), a016931.
- 542 5. Schreij, A.M. et al. (2016) Endocytic membrane trafficking and neurodegenerative
543 disease. *Cell Mol Life Sci* 73 (8), 1529-45.
- 544 6. McGough, I.J. and Cullen, P.J. (2011) Recent advances in retromer biology. *Traffic*
545 12 (8), 963-71.
- 546 7. Christ, L. et al. (2017) Cellular Functions and Molecular Mechanisms of the ESCRT
547 Membrane-Scission Machinery. *Trends Biochem Sci* 42 (1), 42-56.
- 548 8. Maxfield, F.R. and McGraw, T.E. (2004) Endocytic recycling. *Nat Rev Mol Cell Biol*
549 5 (2), 121-32.
- 550 9. Hsu, V.W. et al. (2012) Getting active: protein sorting in endocytic recycling. *Nat*
551 *Rev Mol Cell Biol* 13 (5), 323-8.
- 552 10. Puthenveedu, M.A. et al. (2010) Sequence-dependent sorting of recycling proteins
553 by actin-stabilized endosomal microdomains. *Cell* 143 (5), 761-73.
- 554 11. Seaman, M.N. et al. (1998) A membrane coat complex essential for endosome-to-
555 Golgi retrograde transport in yeast. *J Cell Biol* 142 (3), 665-81.

- 556 12. Steinberg, F. et al. (2013) A global analysis of SNX27-retromer assembly and
557 cargo specificity reveals a function in glucose and metal ion transport. *Nat Cell Biol* 15
558 (5), 461-71.
- 559 13. Simonetti, B. et al. (2017) Sequence-dependent cargo recognition by SNX-BARs
560 mediates retromer-independent transport of CI-MPR. *J Cell Biol* 216 (11), 3695-3712.
- 561 14. McNally, K.E. et al. (2017) Retriever is a multiprotein complex for retromer-
562 independent endosomal cargo recycling. *Nat Cell Biol*.
- 563 15. Kvainickas, A. et al. (2017) Cargo-selective SNX-BAR proteins mediate retromer
564 trimer independent retrograde transport. *J Cell Biol* 216 (11), 3677-3693.
- 565 16. Steinberg, F. et al. (2012) SNX17 protects integrins from degradation by sorting
566 between lysosomal and recycling pathways. *J Cell Biol* 197 (2), 219-30.
- 567 17. Phillips-Krawczak, C.A. et al. (2015) COMMD1 is linked to the WASH complex and
568 regulates endosomal trafficking of the copper transporter ATP7A. *Mol Biol Cell* 26 (1),
569 91-103.
- 570 18. Bartuzi, P. et al. (2016) CCC- and WASH-mediated endosomal sorting of LDLR is
571 required for normal clearance of circulating LDL. *Nat Commun* 7, 10961.
- 572 19. Seaman, M.N. et al. (2013) Retromer-mediated endosomal protein sorting: all
573 WASHed up! *Trends Cell Biol* 23 (11), 522-8.
- 574 20. Clague, M.J. et al. (2012) Governance of endocytic trafficking and signaling by
575 reversible ubiquitylation. *Dev Cell* 23 (3), 457-67.
- 576 21. Bissig, C. and Gruenberg, J. (2014) ALIX and the multivesicular endosome: ALIX
577 in Wonderland. *Trends Cell Biol* 24 (1), 19-25.
- 578 22. Norris, A. et al. (2017) SNX-1 and RME-8 oppose the assembly of HGRS-
579 1/ESCRT-0 degradative microdomains on endosomes. *Proc Natl Acad Sci U S A* 114
580 (3), E307-E316.
- 581 23. Derivery, E. et al. (2012) Actin polymerization controls the organization of WASH
582 domains at the surface of endosomes. *PLoS One* 7 (6), e39774.
- 583 24. Raiborg, C. et al. (2002) Hrs sorts ubiquitinated proteins into clathrin-coated
584 microdomains of early endosomes. *Nature Cell Biology* 4 (5), 394-398.
- 585 25. Raiborg, C. (2006) Flat clathrin coats on endosomes mediate degradative protein
586 sorting by scaffolding Hrs in dynamic microdomains. *Journal of Cell Science* 119 (12),
587 2414-2424.
- 588 26. Derivery, E. et al. (2009) The Arp2/3 activator WASH controls the fission of
589 endosomes through a large multiprotein complex. *Dev Cell* 17 (5), 712-23.
- 590 27. Jia, D. et al. (2010) WASH and WAVE actin regulators of the Wiskott-Aldrich
591 syndrome protein (WASP) family are controlled by analogous structurally related
592 complexes. *Proc Natl Acad Sci U S A* 107 (23), 10442-7.
- 593 28. Li, Y. et al. (2014) Expansion of biological pathways based on evolutionary
594 inference. *Cell* 158 (1), 213-25.
- 595 29. Freeman, C.L. et al. (2014) RME-8 coordinates the activity of the WASH complex
596 with the function of the retromer SNX dimer to control endosomal tubulation. *J Cell Sci*
597 127 (Pt 9), 2053-70.
- 598 30. Harbour, M.E. et al. (2010) The cargo-selective retromer complex is a recruiting
599 hub for protein complexes that regulate endosomal tubule dynamics. *J Cell Sci* 123
600 (Pt 21), 3703-17.
- 601 31. Harbour, M.E. et al. (2012) Recruitment of the endosomal WASH complex is
602 mediated by the extended 'tail' of Fam21 binding to the retromer protein Vps35.
603 *Biochem J* 442 (1), 209-20.
- 604 32. Jia, D. et al. (2012) Multiple repeat elements within the FAM21 tail link the WASH
605 actin regulatory complex to the retromer. *Mol Biol Cell* 23 (12), 2352-61.

606 33. Tyrrell, B.J. et al. (2016) Loss of strumpellin in the melanocytic lineage impairs the
607 WASH Complex but does not affect coat colour. *Pigment Cell Melanoma Res* 29 (5),
608 559-71.

609 34. Gomez, T.S. et al. (2012) Trafficking defects in WASH-knockout fibroblasts
610 originate from collapsed endosomal and lysosomal networks. *Mol Biol Cell* 23 (16),
611 3215-28.

612 35. Gomez, T.S. and Billadeau, D.D. (2009) A FAM21-containing WASH complex
613 regulates retromer-dependent sorting. *Dev Cell* 17 (5), 699-711.

614 36. Zech, T. et al. (2011) The Arp2/3 activator WASH regulates alpha5beta1-integrin-
615 mediated invasive migration. *J Cell Sci* 124 (Pt 22), 3753-9.

616 37. Piotrowski, J.T. et al. (2013) WASH knockout T cells demonstrate defective
617 receptor trafficking, proliferation, and effector function. *Mol Cell Biol* 33 (5), 958-73.

618 38. Temkin, P. et al. (2011) SNX27 mediates retromer tubule entry and endosome-to-
619 plasma membrane trafficking of signalling receptors. *Nat Cell Biol* 13 (6), 715-21.

620 39. Buckley, C.M. et al. (2016) WASH drives early recycling from macropinosomes
621 and phagosomes to maintain surface phagocytic receptors. *Proc Natl Acad Sci U S A*
622 113 (40), E5906-E5915.

623 40. Carnell, M. et al. (2011) Actin polymerization driven by WASH causes V-ATPase
624 retrieval and vesicle neutralization before exocytosis. *J Cell Biol* 193 (5), 831-9.

625 41. Nagel, B.M. et al. (2016) Drosophila WASH is required for integrin-mediated cell
626 adhesion, cell motility and lysosomal neutralization. *J Cell Sci*.

627 42. Helfer, E. et al. (2013) Endosomal recruitment of the WASH complex: active
628 sequences and mutations impairing interaction with the retromer. *Biol Cell* 105 (5),
629 191-207.

630 43. McGough, I.J. et al. (2014) Retromer Binding to FAM21 and the WASH Complex
631 Is Perturbed by the Parkinson Disease-Linked VPS35(D620N) Mutation. *Curr Biol* 24
632 (14), 1670-6.

633 44. Cullen, P.J. (2011) Phosphoinositides and the regulation of tubular-based
634 endosomal sorting. *Biochem Soc Trans* 39 (4), 839-50.

635 45. Koumandou, V.L. et al. (2011) Evolutionary reconstruction of the retromer complex
636 and its function in *Trypanosoma brucei*. *J Cell Sci* 124 (Pt 9), 1496-509.

637 46. Horazdovsky, B.F. et al. (1997) A sorting nexin-1 homologue, Vps5p, forms a
638 complex with Vps17p and is required for recycling the vacuolar protein-sorting
639 receptor. *Mol Biol Cell* 8 (8), 1529-41.

640 47. Peter, B.J. et al. (2004) BAR domains as sensors of membrane curvature: the
641 amphiphysin BAR structure. *Science* 303 (5657), 495-9.

642 48. Carlton, J. et al. (2004) Sorting nexin-1 mediates tubular endosome-to-TGN
643 transport through coincidence sensing of high- curvature membranes and 3-
644 phosphoinositides. *Curr Biol* 14 (20), 1791-800.

645 49. van Weering, J.R. et al. (2012) Molecular basis for SNX-BAR-mediated assembly
646 of distinct endosomal sorting tubules. *EMBO J* 31 (23), 4466-80.

647 50. Norwood, S.J. et al. (2011) Assembly and solution structure of the core retromer
648 protein complex. *Traffic* 12 (1), 56-71.

649 51. Seaman, M.N. (2004) Cargo-selective endosomal sorting for retrieval to the Golgi
650 requires retromer. *J Cell Biol* 165 (1), 111-22.

651 52. Fjorback, A.W. et al. (2012) Retromer binds the FANSHY sorting motif in SorLA to
652 regulate amyloid precursor protein sorting and processing. *J Neurosci* 32 (4), 1467-
653 80.

654 53. Nielsen, M.S. et al. (2007) Sorting by the cytoplasmic domain of the amyloid
655 precursor protein binding receptor SorLA. *Mol Cell Biol* 27 (19), 6842-51.

656 54. Seaman, M.N. et al. (1997) Endosome to Golgi retrieval of the vacuolar protein
657 sorting receptor, Vps10p, requires the function of the VPS29, VPS30, and VPS35
658 gene products. *J Cell Biol* 137 (1), 79-92.

659 55. Marcusson, E.G. et al. (1994) The sorting receptor for yeast vacuolar
660 carboxypeptidase Y is encoded by the VPS10 gene. *Cell* 77 (4), 579-86.

661 56. Arighi, C.N. et al. (2004) Role of the mammalian retromer in sorting of the cation-
662 independent mannose 6-phosphate receptor. *J Cell Biol* 165 (1), 123-33.

663 57. Tabuchi, M. et al. (2010) Retromer-mediated direct sorting is required for proper
664 endosomal recycling of the mammalian iron transporter DMT1. *J Cell Sci* 123 (Pt 5),
665 756-66.

666 58. Seaman, M.N. (2007) Identification of a novel conserved sorting motif required for
667 retromer-mediated endosome-to-TGN retrieval. *J Cell Sci* 120 (Pt 14), 2378-89.

668 59. Lucas, M. et al. (2016) Structural Mechanism for Cargo Recognition by the
669 Retromer Complex. *Cell* 167 (6), 1623-1635 e14.

670 60. Strohlic, T.I. et al. (2007) Grd19/Snx3p functions as a cargo-specific adapter for
671 retromer-dependent endocytic recycling. *J Cell Biol* 177 (1), 115-25.

672 61. Harterink, M. et al. (2011) A SNX3-dependent retromer pathway mediates
673 retrograde transport of the Wnt sorting receptor Wntless and is required for Wnt
674 secretion. *Nat Cell Biol* 13 (8), 914-23.

675 62. Zhang, P. et al. (2011) SNX3 controls Wingless/Wnt secretion through regulating
676 retromer-dependent recycling of Wntless. *Cell Res* 21 (12), 1677-90.

677 63. Lauffer, B.E. et al. (2010) SNX27 mediates PDZ-directed sorting from endosomes
678 to the plasma membrane. *J Cell Biol* 190 (4), 565-74.

679 64. Chen, D. et al. (2010) Retromer is required for apoptotic cell clearance by
680 phagocytic receptor recycling. *Science* 327 (5970), 1261-4.

681 65. Varandas, K.C. et al. (2016) Retromer Endosome Exit Domains Serve Multiple
682 Trafficking Destinations and Regulate Local G Protein Activation by GPCRs. *Curr Biol*
683 26 (23), 3129-3142.

684 66. Hierro, A. et al. (2007) Functional architecture of the retromer cargo-recognition
685 complex. *Nature* 449 (7165), 1063-7.

686 67. Collins, B.M. et al. (2005) Vps29 has a phosphoesterase fold that acts as a protein
687 interaction scaffold for retromer assembly. *Nat Struct Mol Biol* 12 (7), 594-602.

688 68. Collins, B.M. et al. (2008) Structure of Vps26B and mapping of its interaction with
689 the retromer protein complex. *Traffic* 9 (3), 366-79.

690 69. Shi, H. et al. (2006) The retromer subunit Vps26 has an arrestin fold and binds
691 Vps35 through its C-terminal domain. *Nat Struct Mol Biol* 13 (6), 540-8.

692 70. Gallon, M. et al. (2014) A unique PDZ domain and arrestin-like fold interaction
693 reveals mechanistic details of endocytic recycling by SNX27-retromer. *Proc Natl Acad*
694 *Sci U S A*.

695 71. Balana, B. et al. (2011) Mechanism underlying selective regulation of G protein-
696 gated inwardly rectifying potassium channels by the psychostimulant-sensitive sorting
697 nexin 27. *Proc Natl Acad Sci U S A* 108 (14), 5831-6.

698 72. Cullen, P.J. and Korswagen, H.C. (2012) Sorting nexins provide diversity for
699 retromer-dependent trafficking events. *Nat Cell Biol* 14 (1), 29-37.

700 73. Cui, Y. et al. (2018) The functional roles of retromer in Parkinson's disease. *FEBS*
701 *Lett* 592 (7), 1096-1112.

702 74. Gallon, M. et al. (2014) A unique PDZ domain and arrestin-like fold interaction
703 reveals mechanistic details of endocytic recycling by SNX27-retromer. *Proc Natl Acad*
704 *Sci U S A* 111 (35), E3604-13.

705 75. Ghai, R. et al. (2011) Phox homology band 4.1/ezrin/radixin/moesin-like proteins
706 function as molecular scaffolds that interact with cargo receptors and Ras GTPases.
707 Proc Natl Acad Sci U S A 108 (19), 7763-8.

708 76. Böttcher, R.T. et al. (2012) Sorting nexin 17 prevents lysosomal degradation of β 1
709 integrins by binding to the β 1-integrin tail. Nat Cell Biol 14 (6), 584-92.

710 77. van Kerkhof, P. et al. (2005) Sorting nexin 17 facilitates LRP recycling in the early
711 endosome. EMBO J 24 (16), 2851-61.

712 78. Burden, J.J. et al. (2004) Sorting motifs in the intracellular domain of the low
713 density lipoprotein receptor interact with a novel domain of sorting nexin-17. J Biol
714 Chem 279 (16), 16237-45.

715 79. Kulak, N.A. et al. (2014) Minimal, encapsulated proteomic-sample processing
716 applied to copy-number estimation in eukaryotic cells. Nature Methods 11 (3), 319-
717 324.

718 80. Aubry, L. et al. (2009) The arrestin fold: variations on a theme. Curr Genomics 10
719 (2), 133-42.

720 81. Tseng, H.Y. et al. (2014) Sorting nexin 31 binds multiple β integrin cytoplasmic
721 domains and regulates β 1 integrin surface levels and stability. J Mol Biol 426 (18),
722 3180-94.

723 82. Rojas, R. et al. (2008) Regulation of retromer recruitment to endosomes by
724 sequential action of Rab5 and Rab7. J Cell Biol 183 (3), 513-26.

725 83. Seaman, M.N. et al. (2009) Membrane recruitment of the cargo-selective retromer
726 subcomplex is catalysed by the small GTPase Rab7 and inhibited by the Rab-GAP
727 TBC1D5. J Cell Sci 122 (Pt 14), 2371-82.

728 84. van Weering, J.R. et al. (2012) SNX-BAR-mediated endosome tubulation is co-
729 ordinated with endosome maturation. Traffic 13 (1), 94-107.

730 85. Li, H. et al. (2015) Endosomal sorting of Notch receptors through COMMD9-
731 dependent pathways modulates Notch signaling. J Cell Biol 211 (3), 605-17.

732 86. Starokadomskyy, P. et al. (2013) CCDC22 deficiency in humans blunts activation
733 of proinflammatory NF-kappaB signaling. J Clin Invest 123 (5), 2244-56.

734 87. Burstein, E. et al. (2005) COMMD proteins, a novel family of structural and
735 functional homologs of MURR1. J Biol Chem 280 (23), 22222-32.

736 88. Maine, G.N. and Burstein, E. (2007) COMMD proteins: COMMing to the scene.
737 Cell Mol Life Sci 64 (15), 1997-2005.

738 89. Mallam, A.L. and Marcotte, E.M. (2017) Systems-wide Studies Uncover
739 Commander, a Multiprotein Complex Essential to Human Development. Cell Syst 4
740 (5), 483-494.

741 90. Riera-Romo, M. (2018) COMMD1: A Multifunctional Regulatory Protein. J Cell
742 Biochem 119 (1), 34-51.

743 91. Huttlin, E.L. et al. (2017) Architecture of the human interactome defines protein
744 communities and disease networks. Nature 545 (7655), 505-509.

745 92. Wan, C. et al. (2015) Panorama of ancient metazoan macromolecular complexes.
746 Nature 525 (7569), 339-44.

747 93. Duleh, S.N. and Welch, M.D. (2012) Regulation of integrin trafficking, cell
748 adhesion, and cell migration by WASH and the Arp2/3 complex. Cytoskeleton
749 (Hoboken) 69 (12), 1047-58.

750 94. Osborne, D.G. et al. (2015) SNX17 affects T cell activation by regulating TCR and
751 integrin recycling. J Immunol 194 (9), 4555-66.

752 95. van de Sluis, B. et al. (2007) Increased activity of hypoxia-inducible factor 1 is
753 associated with early embryonic lethality in Commd1 null mice. Mol Cell Biol 27 (11),
754 4142-56.

96. Burkhead, J.L. et al. (2009) COMMD1 forms oligomeric complexes targeted to the endocytic membranes via specific interactions with phosphatidylinositol 4,5-bisphosphate. *J Biol Chem* 284 (1), 696-707.

97. Jahic, A. et al. (2015) The spectrum of KIAA0196 variants, and characterization of a murine knockout: implications for the mutational mechanism in hereditary spastic paraplegia type SPG8. *Orphanet J Rare Dis* 10, 147.

98. Bartuzi, P. et al. (2013) Tuning NF-kappaB activity: a touch of COMMD proteins. *Biochim Biophys Acta* 1832 (12), 2315-21.

99. San Agustin, J.T. et al. (2016) Genetic link between renal birth defects and congenital heart disease. *Nat Commun* 7, 11103.

100. Li, Y. et al. (2015) Global genetic analysis in mice unveils central role for cilia in congenital heart disease. *Nature* 521 (7553), 520-4.

101. Huttlin, E.L. et al. (2015) The BioPlex Network: A Systematic Exploration of the Human Interactome. *Cell* 162 (2), 425-40.

102. Hein, M.Y. et al. (2015) A human interactome in three quantitative dimensions organized by stoichiometries and abundances. *Cell* 163 (3), 712-23.

103. Fedoseienko, A. et al. (2018) COMMD Family Regulates Plasma LDL Levels and Attenuates Atherosclerosis Through Stabilizing the CCC Complex in Endosomal LDLR Trafficking. *Circ Res*.

104. Voineagu, I. et al. (2012) CCDC22: a novel candidate gene for syndromic X-linked intellectual disability. *Mol Psychiatry* 17 (1), 4-7.

105. Kolanczyk, M. et al. (2014) Missense variant in CCDC22 causes X-linked recessive intellectual disability with features of Ritscher-Schinzel/3C syndrome. *Eur J Hum Genet*.

106. Bettencourt, C. et al. (2013) Exome sequencing expands the mutational spectrum of SPG8 in a family with spasticity responsive to L-DOPA treatment. *J Neurol* 260 (9), 2414-6.

107. Valdmann, P.N. et al. (2007) Mutations in the KIAA0196 gene at the SPG8 locus cause hereditary spastic paraplegia. *Am J Hum Genet* 80 (1), 152-61.

108. Ishiura, H. et al. (2014) Molecular epidemiology and clinical spectrum of hereditary spastic paraplegia in the Japanese population based on comprehensive mutational analyses. *J Hum Genet* 59 (3), 163-72.

109. Wang, X. et al. (2014) A novel KIAA0196 (SPG8) mutation in a Chinese family with spastic paraplegia. *Chin Med J (Engl)* 127 (10), 1987-9.

110. Bogucki, P. and Sobczynska-Tomaszewska, A. (2017) First patient with hereditary spastic paraplegia type 8 in Poland. *Clin Case Rep* 5 (9), 1468-1470.

111. Jahic, A. et al. (2014) A novel strumpellin mutation and potential pitfalls in the molecular diagnosis of hereditary spastic paraplegia type SPG8. *J Neurol Sci* 347 (1-2), 372-4.

112. de Bot, S.T. et al. (2013) Pure adult-onset spastic paraplegia caused by a novel mutation in the KIAA0196 (SPG8) gene. *J Neurol* 260 (7), 1765-9.

113. Ichinose, Y. et al. (2016) Exome sequencing reveals a novel missense mutation in the KIAA0196 gene in a Japanese patient with SPG8. *Clin Neurol Neurosurg* 144, 36-8.

114. Elliott, A.M. et al. (2013) A novel mutation in KIAA0196: identification of a gene involved in Ritscher-Schinzel/3C syndrome in a First Nations cohort. *J Med Genet* 50 (12), 819-22.

115. Ropers, F. et al. (2011) Identification of a novel candidate gene for non-syndromic autosomal recessive intellectual disability: the WASH complex member SWIP. *Hum Mol Genet* 20 (13), 2585-90.

116. Gershlick, D.C. and Lucas, M. (2017) Endosomal Trafficking: Retromer and Retriever Are Relatives in Recycling. *Curr Biol* 27 (22), R1233-R1236.

Figure 1 Overview of endosomal retrieval and recycling of cargo

Cargo from the biosynthetic pathway (e.g. CI-MPR) or the plasma membrane (e.g. β_2 adrenergic receptor, $\alpha_5\beta_1$ integrin, EGFR) enter the endosomal system where they are either sorted for degradation or are retrieved away from a degradative fate and recycled to the plasma membrane, to the TGN or to other organelles such as melanosomes. Retrieval complexes such as retriever, retromer, the CCC and WASH complexes localize to retrieval sub-domains, which are spatially distinct from the degradative sub-domain in which ESCRT complexes reside. Cargo destined for lysosomal degradation, such as activated EGFR, are recognized by the ESCRT complexes which package the cargo into intraluminal vesicles (ILVs) that bud into the endosomal lumen and eventually pinch off (represented as black circles inside the endosome). As endosomes mature, the number of ILVs in the lumen increases and eventually late endosomes fuse with lysosomes and cargo is degraded. Hydrolase receptors, such as CI-MPR, deliver hydrolases from the TGN to the endosome. Upon delivery to endosomes, the change in pH results in the dissociation of the hydrolase from the receptor into the endosomal lumen. The hydrolase is ultimately delivered to the lysosome upon fusion of endosomes and lysosomes. The hydrolase receptor and other cargo destined for recycling are conversely retrieved away from lysosomal degradation through the sequence-dependent recognition of cytosolic facing motifs by cargo retrieval complexes. After retrieval, cargo are recycled back to various membrane compartments.

830 C16orf62; Chromosome 16 open reading frame 62, DSCR3; Downs syndrome critical
831 region 3, VPS35L; VPS35-like, CCDC; coiled-coil domain containing, WASH; Wiskott-
832 Aldrich syndrome protein and SCAR homologue, SWIP; Strumpellin and WASH
833 interacting protein, CCC complex; CCDC22, CCDC93 and COMMD, SNX; sorting
834 nexin, ESCRT ; endosomal sorting complex required for transport, EGFR; epidermal
835 growth factor receptor, CI-MPR; cation independent mannose-6 phosphate receptor,
836 TGN; trans-Golgi network, ILVs; intraluminal vesicles

837

838 **Figure 2 Retriever is predicted to be a retromer-like complex**

839 A) The mammalian retromer hetero-trimer consists of VPS26A/B, VPS35 and
840 VPS29. VPS35 forms an extended central scaffold with VPS26 and VPS29
841 independently engaging VPS35 at the amino- and carboxy-terminals
842 respectively. Retriever is a hetero-trimer consisting of DSCR3 (VPS26C),
843 C16orf62 (VPS35L) and VPS29, with predicted structural similarity with
844 retromer.

845 B) Estimated protein copy numbers of retrieval complexes in *S. cerevisiae* and
846 HeLa cells. Subunits of retriever, the CCC complex and the WASH complex are
847 absent in the yeast *S. cerevisiae*. Interestingly, the retriever subunits C16orf62
848 and DSCR3 are expressed at much lower levels than the equivalent VPS35
849 and VPS26A/B subunits in retromer. Copy number data from [79].

850 [The predicted structures of retromer and retriever, adapted from \[116\].](#)

851 C16orf62; Chromosome 16 open reading frame 62, DSCR3; Downs syndrome
852 critical region 3, VPS35L; VPS35-like, CCDC; coiled-coil domain containing,
853 WASH; Wiskott-Aldrich syndrome protein and SCAR homologue, SWIP;

854 Strumpellin and WASH interacting protein, CCC complex; CCDC22, CCDC93
855 and COMMD

856 **Figure 3 The endosomal retrieval sub-domain**

857 'Degradative' sub-domains containing ESCRT complexes and 'retrieval' sub-domains
858 containing retriever, retromer and the WASH and CCC complexes are spatially
859 segregated on endosomal membranes. 1) The pentameric WASH complex (individual
860 subunits are not shown for simplicity) activates Arp 2/3, which polymerizes branched
861 actin on the endosomal membrane. The WASH complex is recruited to endosomal
862 membranes through interactions of the FAM21 tail (represented as a black wavy line)
863 with retromer and also through unidentified retromer-independent mechanisms. 2)
864 Retromer is recruited to the endosomal membrane through interactions with SNX3 and
865 GTP-bound Rab7a. Sequence motifs within cytosolic domains of cargo are recognized
866 by retromer, either directly via a $\phi x(L/M)$ motif (where ϕ represents an aromatic residue
867 and x represents any residue) or indirectly through the cargo adaptors SNX3 and
868 SNX27, the latter of which binds to cargo containing a PDZbm. VPS29 can also bind
869 to regulatory factors such as TBC1D5, a Rab7 GAP and VARP. 3) The tail of FAM21
870 can interact with and recruit the CCC complex to endosomal membranes. The CCC
871 complex interacts with and recruits retriever to the retrieval subdomain. Precise
872 interactions between the CCC complex and retriever have not been elucidated. 4)
873 SNX17 associates with endosomal membranes through its PX domain and engages
874 cargo, such as $\alpha_5\beta_1$ integrin, containing NPxY or NPxF motifs. SNX17 is a cargo
875 adaptor for retriever, associating with retriever through its flexible carboxy-terminal tail.
876 Retriever may be able to directly engage cargo. Other sequence-dependent cargo
877 entry points into this pathway could occur via the CCC complex. Adapted from [14].

878 C16orf62; Chromosome 16 open reading frame 62, DSCR3; Downs syndrome critical
879 region 3, VPS35L; VPS35-like, CCDC; coiled-coil domain containing, WASH; Wiskott-
880 Aldrich syndrome protein and SCAR homologue, SWIP; Strumpellin and WASH
881 interacting protein, CCC complex; CCDC22, CCDC93 and COMMD, SNX; sorting
882 nexin, ESCRT; endosomal sorting complex required for transport, TBC1D5; TBC1
883 domain family member 5, PDZbm; PDZ binding motif, GAP: GTPase-activating
884 protein, VARP: VPS9-ankyrin-repeat protein.

885

886

887

Outstanding questions

- **What is the precise role of retromer in endosome-to-TGN transport?**
- **Is retriever also involved in endosome-to-TGN trafficking of cargo and if so, which and how many?**
- **Can retriever bind to cargo directly?**
- **Are retromer and retriever cargo corralled into the same cargo carriers?**
- **What complexes act downstream of retriever and the CCC complex to mediate cargo carrier formation, scission and transport?**
- **How is sequence-dependent cargo retrieval and recycling co-ordinated with RAB switching to allow for pathway progression?**
- **Is there redundancy in cargo retrieval and is this a point of regulation and/or therapeutic intervention?**
- **What are the regulators of endosomal retrieval complexes?**
- **Are there other undiscovered multi-protein complexes which regulate endosomal retrieval of cargo?**

Highlights

- Endosomal cargo are retrieved from degradation by multi-protein complexes prior to recycling
- Retromer, retriever, the CCC and WASH complexes are required for sequence-specific retrieval of cargo
- Retromer is a key orchestrator of endosomal sorting but retromer-independent pathways exist
- Retriever is a retromer-like complex that functions independently of retromer
- Retriever requires the CCC and WASH complexes for endosomal localisation and function
- Mutations in these multi-protein complexes are increasingly associated with human pathologies including neurodegeneration and developmental disorders.

Glossary:

Arp2/3 (Actin-related protein 2/3) complex: A complex composed of seven subunits which nucleates branched actin polymerisation. However, Arp2/3 itself is not intrinsically active and requires activation by nucleation promoting factors such as the WASH complex.

Bafilomycin: An inhibitor of the vacuolar-H⁺-ATPase, preventing acidification of lysosomes.

BAR domain: BAR domains are a protein dimerization interface that dimerise to form a rigid banana-shaped structure. BAR domains are membrane curvature sensing domains and can also induce membrane curvature and tubulation of liposomes.

Cargo: Transmembrane proteins, and their associated proteins and lipids, that are trafficked through the endosomal network.

Carrier scission: Detachment of the nascent cargo enriched carrier from the donor membrane.

Carrier transport: Transport of the cargo enriched carrier from the donor to the acceptor membrane. Usually occurs via microtubule and actin based motors.

FERM domain (4.1/ezrin/radixin/moesin): A domain with a cloverleaf structure and binds to NPx(Y/F) motifs within proteins.

Geometric based principles: The recycling tubules display a high surface area to low volume ratio, thus transmembrane proteins entering the endosome are much more likely to be present in the endosomal tubules than the vesicular portion of the endosome which has a high volume to low surface area ratio. Therefore, unless cargo are specifically recognized by ESCRT, they are likely to be recycled back to the plasma membrane or the TGN based upon bulk membrane recycling.

Intra-luminal vesicles (ILVs): Vesicles formed from inward-budding of the endosomal membrane.

LFa motifs: Leucine-Phenylalanine-a series of acidic residues. Multiple motifs found in the tail of FAM21 that mediate binding to VPS35.

PDZ binding motif: Protein:protein interaction motif found in the cytosolic tails of transmembrane cargo, usually at the extreme carboxy-termini. The consensus sequence of a class I PDZbm is [T/S]-x- ϕ , where x denotes any amino acid and ϕ represents an amino acid with a bulky hydrophobic group.

PDZ domain (Postsynaptic density 95 – Disc large – ZO1): PDZ domains interact with PDZ binding motifs (PDZbm).

Pleckstrin homology (PH) domain: A phosphoinositide binding domain, interacts with PI(3,4)P₂ and PI(3,4,5)P₃ and various other phosphoinositide species.

PX domain (Phox homology): A phosphoinositide binding domain, interacting primarily with PI3P.

Sorting nexin (SNX): Sorting nexins are a family of proteins defined by the presence of a PX domain with over 50% sequence similarity to the first member of the family, SNX1. Some members of this protein family also contain extra functional domains such as PDZ and FERM domains.

v-ATPase: vacuolar H⁺-ATPase. Proton pump that utilises ATP hydrolysis to pump protons from the cytosol into the lumen of endosomes and lysosomes, generating and maintaining an acidic pH in the lumen of these organelles.

Retrieval complexes

Retriever

Retromer

CCC complex

WASH complex

A

Retromer

Retriever

B

Complex	Protein	Copy Number in <i>S. cerevisiae</i>	Copy Number in HeLa cells
Retromer	VPS26A	1140.8	826147.4
	VPS26B	0.0	76397.8
	VPS35	571.3	780462.6
Retromer/Retriever	VPS29	4009.1	1424734.0
Retriever	C16orf62 (VPS35L)	0.0	286.3
	DSCR3 (VPS26C)	0.0	2751.5
CCC complex	CCDC93	0.0	33942.3
	CCDC22	0.0	55222.3
	COMMD1	0.0	83103.2
WASH complex	WASH1 (WASHC1)	0.0	43936.1
	FAM21A (WASHC2A)	0.0	58981.4
	FAM21B (WASHC2B)	0.0	558.1
	FAM21C (WASHC2C)	0.0	1135.1
	CCDC53 (WASHC3)	0.0	99617.0
	SWIP (WASHC4)	0.0	65245.8
	Strumpellin (WASHC5)	0.0	141599.6