Tailor, S., Manzotti, M., Asghar, S., Rowsell, B., Luckham, S., Sparkes, H., & Bedford, R. (2019). Revisiting Claims of the Iron-, Cobalt-, Nickel-, and Copper-Catalyzed Suzuki Biaryl Cross-Coupling of Aryl Halides with Aryl Boronic Acids. *Organometallics*, 38(8), 1770-1777. <https://doi.org/10.1021/acs.organomet.9b00083>

Peer reviewed version

License (if available):
Other

Link to published version (if available):
[10.1021/acs.organomet.9b00083](https://doi.org/10.1021/acs.organomet.9b00083)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via ACS Publications at <https://doi.org/10.1021/acs.organomet.9b00083>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Revisiting claims of the iron-, cobalt-, nickel- and copper-catalyzed Suzuki biaryl cross-coupling of aryl halides with aryl boronic acids.

Sanita B. Tailor,^a Mattia Manzotti,^a Soneela Asghar,^{a,b} Benjamin J. S. Rowsell,^a Stephen L. J. Luckham,^a Hazel Sparkes^a and Robin B. Bedford.^{*a}

a. School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, U.K.

b. Department of Chemistry and Chemical Engineering, SBA School of Science & Engineering Lahore University of Management Sciences Lahore 54792, Pakistan.

ABSTRACT: Intrigued by recent reports on the surprisingly excellent activity of a range of cobalt, iron, copper and nickel-based catalysts in the Suzuki biaryl cross-coupling of simple arylboronic acids with aryl halides, we undertook a reexamination of the syntheses of representative examples of the reported pre-catalysts and their application to the catalytic reaction. A reported PNP-Fe pincer complex, in our hands, proved to be a mixture of starting materials; a mono-Schiff base cobalt complex in fact the bis-ligated adduct and a monomeric copper (II) PNP pincer complex a di- or oligomeric copper(I) species. In our hands, neither these complexes, nor any other of the selected pre-catalysts investigated, showed any activity in a Suzuki cross-coupling reaction of an electronically-activated aryl bromide with phenyl boronic acid. Meanwhile, switching the nucleophile to the BuLi-activated phenyl boronic pinacol ester gave some promising activity with cobalt pre-catalysts.

INTRODUCTION

The Suzuki biaryl cross-coupling reaction,¹ (Scheme 1) is a robust and widely used method for the construction of aryl-aryl bonds. Commercial applications that have exploited Suzuki biaryl bond formation include the synthesis of OTBN, an intermediate in the production of a range of the Sartan class of drugs for the treatment of hypertension, and Boscalid, a broad-spectrum fungicide used in crop protection.²

Scheme 1. Suzuki biaryl cross-coupling and selected commercial applications.

The vast majority of Suzuki biaryl cross-coupling reactions rely on palladium-based catalysts. However, the high cost of this metal, combined with the need to remove it to the low ppm level from active pharmaceutical

intermediates³ has inspired the hunt for more sustainable catalysts for Suzuki cross-coupling based on cheap, readily available first-row transition metals. Of these, nickel⁴⁻⁶ and copper-catalyzed^{7,8} Suzuki cross-coupling reactions are arguably the most well developed. Of particular note is the coupling of aryl halides with aryl boronic acids, as this represents the simplest, yet perhaps most synthetically valuable variant of the cross-coupling reaction.^{4-6a,e,7}

Cobalt-catalyzed Suzuki biaryl cross-coupling is rapidly developing,⁹ but is currently largely limited to the coupling of arylboronic esters, activated with either an organolithium reagent or an alkoxide base, rather than free arylboronic acids. Meanwhile, despite growing success in iron-catalyzed Suzuki couplings,¹⁰ simple biaryl cross-coupling under mild conditions¹¹ remains limited to either highly activated 2-pyridyl or 2-pyrimidyl halide substrates,^{10a} or aryl halide substrates that can engender directed aryl halide activation.¹² Again, in all these cases, the free boronic acid cannot be exploited, rather a tetraorganoborate or an activated aryl boronic ester is required.

A series of apparent exceptions to the failure to exploit simple aryl boronic acids using Fe and Co (as well as Cu and Ni) pre-catalysts have recently been published across five papers by Bhat and co-workers.¹³ The authors reported that attractively simple pre-catalysts based on the easily produced ligands **1** - **3** and closely related ligands all show good to excellent activity in the highly desirable Suzuki

1

2

3a: E = CH; Z = OMe; EWG = C(O)Me
b: E = N; Z = OMe; EWG = Br
c: E = CH; Z = H; EWG = C(O)Me

Attempted syntheses of the pre-catalysts. The Co and Ni complexes of ligand **2** (salophen), complexes **4** and **5** respectively, have been reported many times, indeed they are commercially available, and were readily prepared by Bhat's procedures.^{13d} The IR data we obtained for the complexes were in agreement with the data reported in the literature,¹⁵ while the NMR spectroscopic data of the diamagnetic Ni(II) complex **5** were in accord with literature values.^{15b} Bhat reported a value of $1.23 \mu_B$ for the solid state magnetic moment of complex **4**,^{13d} which we note is far too low for a pure sample; low-spin square planar cobalt(II) complexes typically show 'normal' room temperature magnetic moments in the range $2.1 - 2.9 \mu_B$,¹⁶ while the room-temperature magnetic moment of closely related cobalt(II) salen has variously been reported in the range $2.2 - 2.7 \mu_B$.¹⁷ We obtained a room-temperature magnetic moment for **4** of $3.02 \mu_B$, essentially identical to a previously reported value.^{15d}

Perhaps the most important data in support of the proposed formulation, is the ESI mass-spectrum of **6a** presented by Bhat,^{13c} which shows a single mass peak at $m/z = 568.10$, as well as single mass peaks consistent with the loss of chloride, and the appearance of free Schiff-base ligand. While this mass is consistent with $[\text{MH}]^+$, the complete

Taken together, the data are inconsistent with the formulation of **6a**. Structurally characterized examples of bis-Schiff base Fe(III) chloride complexes of the general type **6** are surprisingly rare with, to the best of our knowledge, only two reported in the literature,²⁰ while the products of reactions of FeCl₃ with Schiff base ligands is highly dependent on the presence or absence of water (note here water is used as a co-solvent in ethanol at reflux temperature) and the presence or absence of base.¹⁸ Regardless, the purple solid “**6a***” was used, as is, in subsequent Suzuki cross-coupling reactions (*vide infra*). Meanwhile, we prepared the previously reported, closely related complex **6b**,^{20b} and used the same synthetic approach to prepare what we believe is a genuine sample of **6a**. The HR-ESI-MS of **6a** prepared this way showed a peak at $m/z = 568.0820$ corresponding to [MH]⁺ (calculated $m/z = 568.0847$), with the expected isotope distribution pattern (see Figure S4, supplementary information). Both of the complexes **6a** and **b** were investigated in the Suzuki reaction (*vide infra*).

From the outset, the reported formulation of the cobalt complex **7**^{3c} appeared extremely unlikely to us: firstly because the proposed $\kappa^3\text{-N,O,O}$ -chelation appeared impossibly strained, as is clearly apparent from the distorted tetrahedral and square-planar geometries obtained from DFT calculations (B3LYP-D3BJ/cc-pVDZ) of both the high- and low-spin ground-state configurations respectively (Figure 1). Secondly, while many structurally-characterized cobalt bis-Schiff base complexes are known, mono-adducts of simple aryl Schiff bases are rare.²¹ Furthermore, Bhat reported that complex **7** has a low spin ($S = \frac{1}{2}$) ground state electronic configuration,^{13c} based on a (presumably ambient temperature) solid state magnetic moment measurement of $1.26 \mu_B$,^{13c} a value far too low for a typical Co(II) low spin complex. In addition, the low spin assignment is perhaps unexpected for a five-coordinate Co(II) with only a single Schiff-base ligand and moderate to weak-field coligands. Indeed, DFT calculations (B3LYP-D3BJ/cc-pVTZ//cc-pVDZ) of the putative complex **7** revealed a significant preference for a high-spin ground-state configuration at room temperature of around 12.2 kcal/mol, suggesting that the reported low-spin configuration for this mononuclear complex is unlikely, at least at room-temperature.²²

Figure 1. DFT (B₃LYP-D₃BJ/cc-pVDZ)-calculated high-spin (left) and low-spin ground-state geometries for the putative complex **7**. Single point energy calculations at the B₃LYP-D₃BJ/cc-pVTZ//cc-pVDZ level of theory indicate the high-spin structure is preferred by around 12.2 kcal/mol.

Following the synthetic procedure reported by Bhat for complex **7**,^{13c} namely heating one equivalent of ligand **3a** with hydrated cobalt(II) chloride in ethanol gave a red-brown solid; the same solid was produced on reaction of cobalt chloride with two equivalents of **3a** (see Figure S10, supplementary information for UV-Vis spectroscopic data). The IR spectroscopic data for the sample produced from a 2 : 1 ratio of **3a** : CoCl₂·6H₂O matched the data reported by Bhat.^{13c} Taken together, the data do not support the reported formulation of **7**, but indicate that the reaction instead yields a bis-Schiff base adduct, **8a**, irrespective of the ratio of ligand to metal used. The high-resolution spectrum supported the formulation of **8a**, with a peak at $m/z = 596.1338$ corresponding to the [MH]⁺ ion (calculated $m/z = 596.1358$), with the anticipated isotope distribution pattern (Figure S11, Supplementary Information).²³ The room-temperature magnetic moment recorded of 4.02 μ_B is consistent with a tetrahedral $S = 3/2$ Co(II) complex. Finally, it should be noted that Bhat reported separately that, in sharp contrast, the equivalent reaction with the analogous Schiff base derived from salicylaldehyde gives the bis-Schiff base complex **8b**,^{13c} a finding with which we concur.

We next reexamined the reported syntheses of iron, cobalt and copper complexes of the PNP-pincer ligand **1**. The synthesis of the iron-containing complex **9** was attempted following Bhat's reported procedure,^{13b} which consisted of heating ligand **1** with [Fe(OH₂)₆]SO₄·H₂O in THF at reflux temperature for 4 hours. In our hands, this gave a pale brown precipitate that proved insoluble in THF, dichloromethane, acetonitrile and acetone. An IR spectrum of the precipitate was essentially identical to that of the starting material, hydrated iron(II) sulfate (see Figure S15), dominated by a large, broad peak at 1081 cm⁻¹ corresponding to the sulfate ion. Removal of the solvent from the supernatant from the synthetic reaction gave a pale brown residue which proved to be predominantly the free ligand **1**, as determined by ¹H and ³¹P NMR spectroscopy. Summarizing, in our hands, no reaction was obtained between ligand **1** and hydrated ferrous sulfate. The same results were obtained on repeated attempts.

The reaction of ligand **1** with cobalt(II) acetate, employing the method of Bhat,^{13a} gave a brown solid whose IR spectrum (Figure S19) had some similar peaks to those reported for complex **10**, most notably those associated with the acetate ligands. The ESI-MS spectrum of a methanol solution (Figure S20) showed, along with higher mass peaks, a peak of $m/z = 627.1$ consistent with loss of an acetate ligand from **10** and coordination of a solvent molecule. Attempts to produce crystals of **10** suitable for X-ray analysis were unsuccessful, and at this stage, we remain hesitant about the assignment of the structure.

9

10: M = Co
11: M = Cu

Our attempts to produce the reported copper complex **11** by the method described by Bhat from Cu(OAc)₂·H₂O proved fruitless.^{13b} The NMR spectroscopic data obtained (see Figures S21 – S24, Supporting Information) for the product mixture was entirely inconsistent with a monometallic Cu(II) species of the formulation **11**. Firstly, the ¹H NMR spectrum showed peaks in the standard range for a diamagnetic complex, whereas, as formulated, the monomeric complex **11** should be paramagnetic, with $S = 1/2$. Secondly, the ³¹P spectrum showed, in addition to some free ligand **1**, an AB system with broad doublets observed at δ 24.1 and 20.3 ppm with a mutual coupling of around 166 Hz and a very broad, featureless peak centered at around -17 ppm. Interestingly, the same AB pattern was observed when the reaction was repeated with copper(I) acetate as the precursor, along with other peaks indicative of di- or polymetallic products. In addition, an ESI-MS of the product mixture indicated some oxidation of the ligand **1** to a diphosphine oxide, which in turn coordinated to a copper

center.²⁴ Taken together, the data indicate that if the mononuclear copper(II) complex **11** is indeed produced at any stage, it undergoes very facile reduction to copper(I) with concomitant oxidation of the phosphine residues.²⁵ This is perhaps not surprising in view of the well-known instability of Cu(II)-phosphine adducts with respect to facile reduction to Cu(I), especially in the presence of even trace amounts of adventitious water,²⁶ such as the water of crystallization in the copper (II) acetate used in the reported synthesis of **11**.^{13b} Furthermore, reported copper(I) often consist of dimetallic or higher order copper clusters rather than monometallic complexes, indeed, dinuclear Cu(I) complex formation has been demonstrated previously for ligand **1**.²⁷

In view of the facile reduction of the putative intermediate **11** to a mixture of dimeric (or higher nuclearity) Cu(I) species, and in light of the proven catalytic activity of copper(I) complexes in the Suzuki coupling of aryl iodides with arylboronic esters,^{8f} we decided instead to focus our efforts on a well-defined Cu(I) complex of ligand **1**, the previously reported dimeric copper(I) iodide species, complex **12** which could be readily prepared and isolated according to a literature method.^{27a}

Similarly, our lack of success in preparing complex **9** as reported, or indeed any tractable material other than starting materials, led us to investigate another closely related iron-based analogue containing ligand **1**. The best catalytic activity reported for the material described as complex **9** was obtained in acetonitrile.¹³ Therefore we reasoned that if **9** were indeed accessible, when dissolved in acetonitrile, as per the reported catalytic conditions, it would be highly likely to form the adduct $[\text{Fe}(\text{1})(\text{NCMe})_3]^{2+}$, **13**, which has previously been reported and structurally characterized as the tetrafluoroborate, $[\text{13}][\text{BF}_4]$, by Kirchner and co-workers.²⁸ This complex was therefore synthesized and examined as a well-defined pre-catalyst (*vide infra*).

While the data we obtained for complex **10** was not entirely inconsistent with those reported by Bhat,^{13a} in

the absence of definitive structural characterization, we were keen to also target a well-defined cobalt pre-catalyst. During catalysis it is highly likely that bromide ligands from the aryl bromide substrates will coordinate to the metal center, suggesting the bromide-containing adducts of the PNP-Co motif should be catalytically competent. Indeed, Bhat proposes that the acetate ligands in **10** are lost during pre-catalyst activation to be replaced by a halide on oxidative addition.^{13a} Accordingly, we synthesized the cobalt dibromide adduct of ligand **1**, complex **14**. The single crystal X-ray structure of **14** is shown in Figure 2(a) and the structure is broadly similar to that for the previously reported chloride analogue.²⁹ Recrystallization of **14** from acetonitrile/acetone gave the cationic complex **15**, the crystal structure of which is shown in Figure 2(b), supporting the assertion made above that acetonitrile is likely to replace anionic ligands during catalysis. In both cases the distorted square-based pyramidal structures obtained are consistent with low-spin ($S = \frac{1}{2}$) five-coordinate Co(II) complexes and this ground state is further supported by a value of $2.22 \mu_B$ for the effective magnetic moment of **14** recorded in solution (Evans' method) at room temperature, a value which is in line with the data published for a range of Co(II)-PNP complexes, including the dichloride analogue of **14**.²⁹

(a)

(b)

Figure 2. Single crystal X-ray structures of (a) complex **14** and (b) complex **15**. Hydrogen atoms (except for H-bond in **15**) and solvent molecules omitted for clarity, thermal ellipsoids set at 50% probability.

Summarizing our synthetic efforts, we were able to reproduce Bhat's syntheses of the previously very well-known Co- and Ni-salophen complexes **4** and **5**, although we note that Bhat's reported value for the magnetic susceptibility of **4** is far too low for a pure sample, but were unable to conclusively reproduce most of the otherwise previously unreported claims. The reaction of ligand **3c** under the conditions described yields a purple complex with the same IR and UV-Vis spectroscopic data, reported by Bhat, but *in toto* the data obtained by Bhat and us do not support the formulation of **6a**. Of particular note the MS data presented by Bhat are remarkably devoid of any isotope patterns, and the $[MH]^+$ peak appears to be in the wrong place in the spectrum shown in their supplementary information. The complex **6a** could, however, be isolated by an alternative synthetic procedure. Similarly, ligand **3a** does not, in our hands, give a mono-ligated Co(II) complex **7**, but rather the bis-ligated adduct **8a**, the structural analogue of **8b**, a complex also reported by Bhat, irrespective of the ligand to metal ratio used in the synthesis. Furthermore, the magnetic data for "**7**" presented by Bhat and co-workers does not support their formulation, and the proposed chelating tridentate nature of the Schiff-base ligand is highly improbable, indeed a search of the CCDC data base revealed no such motif in any crystallographically characterized complex.³⁰ Turning to the PNP ligand **1**, this may possibly give the cobalt (II) acetate adduct **10** as reported (we believe there is insufficient data across Bhat's and our work to state this definitively), but no reaction was obtained between **1** and ferric sulfate, while the reaction with copper (II) acetate led to the formation of di- or oligomeric copper (I) species, rather than the monomeric copper(II) adduct claimed. Accordingly we also synthesized well-defined Co(II), Fe(II) and Cu(I) adducts of the ligand **1** for the catalytic testing described in the following section.

Suzuki biaryl cross-coupling reactions. Across the papers reported by Bhat and co-workers,¹³ particularly good results were claimed in the coupling of the electronically activated aryl bromide substrate, 4-bromocyanobenzene, **16**, with phenylboronic acid, **17** to give 4-cyanobiphenyl **18**, using a variety of bases and solvents (Scheme 2(a)). Indeed, replotting the catalytic data obtained by Bhat in this reaction, using acetonitrile as the solvent at 80 °C and a selection of pre-catalysts types^{13a-c} (Figure 3) shows a remarkable insensitivity of catalyst performance to variations in either the metal (Co, Fe or Cu) or the ligand (PNP pincer or Schiff base).³¹ Unfortunately, under these conditions, we were unable to produce any of the desired biaryl **18** using the salophen or Schiff-base Co, Ni or Fe-based pre-catalysts **4**, **5**, "**6a**" (the purple solid incorrectly formulated as **6a**), a

genuine sample of **6a** prepared by an alternative method, the closely related analogue **6b**, **8a** (the actual formulation for the reported pre-catalyst **7**), or **8b**.³²

Iron-salen-type complexes have been successfully exploited in both the coupling of secondary alkyl Grignard reagents with activated aryl chlorides,³³ and the coupling of aryl Grignard reagents with primary and secondary alkyl halides,³⁴ while the latter reactions can also be catalysed by an iron Schiff base complex,^{35,36} however it appears that here the use of phenylboronic acid and an aryl bromide as substrates is beyond the scope of these catalyst types. The lack of activity with the cobalt and nickel complexes **4**, **5** and **8a** is in accord with the previous observation that salen complexes of Co(II) and Ni(II), in contrast with the Fe(III) analogue, give only traces of cross-coupled product in the coupling of alkyl halides with aryl Grignards.³⁴

Scheme 2. (a) Unsuccessful and (b) successful Suzuki biaryl cross-coupling reactions. Full conditions for the reactions are given in the Supplementary Information. All attempted couplings of **16** with **17** were repeated independently by (at least) two separate authors. ^a1,4-dioxane, K₂CO₃, 110 °C. ^bMeCN, NEt₃, 80 °C. ^c **6a***, **9*** and **11*** refer to the solids obtained according to Bhat's synthetic procedures, incorrectly formulated by Bhat as **6a**, **9** and **11** respectively. ^dToluene, K₂CO₃, 110 °C. ^eMeCN, CS₂CO₃, 80 °C. ^fMeCN, K₂CO₃, 80 °C. ^gLack of conversion to product demonstrated by GC-MS (see Figure S33). ^hSpectroscopic yield, determined by ¹H NMR spectroscopy (1,3,5-(MeO)₃C₆H₃, internal standard).

Figure 3. Replotting of Bhat's reported data^{13a-c} for the claimed coupling of **16** with **17** in acetonitrile at 80 °C, catalyzed by the PNP-Fe, -Co and -Cu complexes (**9** – **11**) and the Fe(II) and Co(II) bis-schiff base complexes **6a** and **8b**.

Similarly, no cross-coupled product **18** was obtained using the cobalt PNP-pincer complex tentatively ascribed as **10**, nor with the structurally authenticated analogue, complex **14**. Neither was any catalytic activity observed with the pre-formed iron complex of ligand **1**, complex **13**, nor with the intractable residue "**9**" obtained under Bhat's synthetic conditions (most likely the starting material, [Fe(OH₂)₆]SO₄, *vide supra*). Likewise, neither the ill-defined solid "**11**", obtained under Bhat's conditions, nor the well-defined copper(I) adduct of **1**, complex **12**, showed any activity. Instead, in all cases, we observed peaks in the GC-MS corresponding with unreacted **16** and the trimeric cyclic ester of phenyl boronic acid, (PhBO)₃ (see Figure S33, Supporting Information for a representative example).

Following our complete failure to reproduce any of the catalysis reported by Bhat using a simple aryl boronic acid substrate with an electronically-activated aryl bromide, we decided to briefly investigate the application of selected cobalt pre-catalysts in the coupling of the electronically deactivated aryl bromide **19** with the activated nucleophilic substrate, **20**, under conditions that we have previously reported (Scheme 2(b)).^{9b} While the salophen complex **4** and the PNP-pincer complex **14** gave only low spectroscopic yields of the desired cross-coupled product **21**, a modest spectroscopic yield (47%) was obtained when the bis-Schiff base complex **8a** was employed as a pre-catalyst.

CONCLUSIONS

In conclusion, while we were able to substantiate some of Bhat's simpler synthetic claims, notably of complexes that have previously been reported multiple times, the majority of the complexes reported for the first time by Bhat proved to be either starting materials or to have structures very

different from those claimed. An examination of some of the reported mass spectra pivotal to Bhat's reported formulations raise intriguing questions regarding lack of observable isotope patterns and positions of peaks. Contrary to Bhat's findings, none of the Fe, Co, Cu or Ni complexes, nor structurally closely related examples, proved, in our hands at least, to be viable catalysts in the Suzuki biaryl cross-coupling of a simple aryl boronic acid with an electronically-activated aryl bromide.

While we did not screen all of the reported complexes, we believe the examples chosen reflect the wider set of catalysts reported and we are confident that the selected examples represent a fair cross-section. Interestingly, while we observed no reaction with phenylboronic acid, some activity was observed with cobalt catalysts when a BuLi-activated aryl pinacol boronic ester was employed under our previously reported conditions;^{9b} we are currently examining this promising lead further and these results will be published in due course.

The coupling of simple aryl boronic acids with aryl halides catalyzed by cobalt or, ideally, iron, remains, in our opinion, a highly desirable, yet unmet goal.³⁷

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental details; spectroscopic data; DFT calculations (PDF).

xyz coordinate file of the DFT-calculated structures (XYZ).

AUTHOR INFORMATION

Corresponding Author

* E-mail: r.bedford@bristol.ac.uk

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We thank the EPSRC for a studentship (S.B.T.) and the provision of studentships through the EPSRC Centre for Doctoral Training in Catalysis (M.M. and S.L.J.L) and the Bristol Chemical Synthesis Centre for Doctoral Training (B.J.S.R) and the International Research Support Initiative Program, HEC, Pakistan for funding (S.A.).

REFERENCES

- (a) Miyaura, N.; Suzuki, A., Palladium-Catalyzed Cross-Coupling Reactions of Organoboron Compounds, *Chem. Rev.* **1995**, 95, 2457-2483. (b) Valente, C.; Organ M. G. in *Boronic Acids* (Ed.: D. G. Hall), Wiley- VCH, Weinheim, **2011**, pp. 213 – 262
- For a review, see: Torborg, C.; Beller, M., Recent Applications of Palladium-Catalyzed Coupling Reactions in the Pharmaceutical,

Agrochemical, and Fine Chemical Industries, *Adv. Synth. Catal.* **2009**, *351*, 3027–3043.

3. (a) Garrett, C. E.; Prasad, K., The Art of Meeting Palladium Specifications in Active Pharmaceutical Ingredients Produced by Pd-Catalyzed Reactions, *Adv. Synth. Catal.* **2004**, *346*, 889–900. (b) Guideline on the specification limits for residues of metal catalysts or metal reagents, (Doc.Ref. EMEA/CHMP/SWP/4446/2000). Committee for medicinal products for human use (CHMP). European Medicines Agency, London, February **2008**, pp. 1–34.

4. Han, F.-S., Transition-metal-catalyzed Suzuki–Miyaura cross-coupling reactions: a remarkable advance from palladium to nickel catalysts, *Chem. Soc. Rev.* **2013**, *42*, 5270–5298.

5. Early examples: (a) Saito, S.; Sakai, M.; Miyaura, N., A synthesis of biaryls via nickel(o)-catalyzed cross-coupling reaction of chloroarenes with phenylboronic acids, *Tetrahedron Lett.*, **1996**, *37*, 2993–2996. (b) Saito, S.; Oh-tani, S.; Miyaura, N., Synthesis of Biaryls via a Nickel(o)-Catalyzed Cross-Coupling Reaction of Chloroarenes with Arylboronic Acids, *J. Org. Chem.*, **1997**, *62*, 8024–8030.

6. For selected recent examples, see: (a) Mastalir, M.; Stöger, B.; Pittenauer, E.; Allmaier, G.; Kirchner, K., Air-Stable Triazine-Based Ni(II) PNP Pincer Complexes As Catalysts for the Suzuki–Miyaura Cross-Coupling, *Org. Lett.* **2016**, *18*, 3186–3189; (b) Zhou, J.; Berthel, J. H. J.; Kuntze-Fechner, M. W.; Friedrich, A.; Marder, T. B.; Radius, U., NHC Nickel-Catalyzed Suzuki–Miyaura Cross-Coupling Reactions of Aryl Boronate Esters with Perfluorobenzenes, *J. Org. Chem.* **2016**, *81*, 5789. (c) Shi, S.; Meng, G.; Szostak, M., Synthesis of Biaryls through Nickel-Catalyzed Suzuki–Miyaura Coupling of Amides by Carbon–Nitrogen Bond Cleavage, *Angew. Chem. Int. Ed.* **2016**, *55*, 6959. (d) Malan, F. P.; Singleton, E.; van Rooyen, P. H.; Landman, M., Facile Suzuki–Miyaura coupling of activated aryl halides using new CpNiBr(NHC) complexes, *J. Organomet. Chem.* **2016**, *813*, 7. (e) Shields, J. D.; Gray, E. E.; Doyle, A. G., A Modular, Air-Stable Nickel Precatalyst, *Org. Lett.* **2015**, *17*, 2166.

7. Review: Thapa, S.; Shrestha, B.; Gurung, S. K.; Giri, R., Copper-catalysed cross-coupling: an untapped potential, *Org. Biomol. Chem.* **2015**, *13*, 4816–4827.

8. Selected recent examples: (a) Basnet, P.; Thapa, S.; Dickie, D. A.; Giri, R., The copper-catalysed Suzuki–Miyaura coupling of alkylboron reagents: disproportionation of anionic (alkyl)(alkoxy)borates to anionic dialkylborates prior to transmetalation, *Chem. Commun.* **2016**, *52*, 11072–11075. (b) Gurung, S. K.; Thapa, S.; Shrestha, B.; Giri, R., Copper-catalysed cross-couplings of arylboronate esters with aryl and heteroaryl iodides and bromides, *Org. Chem. Front.* **2015**, *2*, 649–653. (c) Zhang, Z.-Q.; Yang, C.-T.; Liang, L.-J.; Xiao, B.; Lu, X.; Liu, J.-H.; Sun, Y.-Y.; Marder, T. B.; Fu, Y., Copper-Catalyzed/Promoted Cross-coupling of gem-Diborylalkanes with Nonactivated Primary Alkyl Halides: An Alternative Route to Alkylboronic Esters, *Org. Lett.* **2014**, *16*, 6342–6345. (d) Sun, Y.-Y.; Yi, J.; Lu, X.; Zhang, Z.-Q.; Xiao, B.; Fu, Y., Cu-Catalyzed Suzuki–Miyaura reactions of primary and secondary benzyl halides with arylboronates, *Chem. Commun.* **2014**, *50*, 11060–11062. (e) Zhou, Y.; You, W.; Smith, K. B.; Brown, M. K., Copper-Catalyzed Cross-Coupling of Boronic Esters with Aryl Iodides and Application to the Carboboration of Alkynes and Allenes, *Angew. Chem., Int. Ed.* **2014**, *53*, 3475–3479. (f) Gurung, S. K.; Thapa, S.; Kafle, A.; Dickie, D. A.; Giri, R., Copper-Catalyzed Suzuki–Miyaura Coupling of Arylboronate Esters: Transmetalation with (PN)CuF and Identification of Intermediates, *Org. Lett.* **2014**, *16*, 1264–1267.

9. (a) Neely, J. M.; Bezdek, M. J.; Chirik, P. J., Insight into Transmetalation Enables Cobalt-Catalyzed Suzuki–Miyaura Cross Coupling, *ACS Cent. Sci.*, **2016**, *2*, 935–942. (b) Asghar, S.; Tailor,

S. B.; Elorriaga, D.; Bedford, R. B., Cobalt-Catalyzed Suzuki Biaryl Coupling of Aryl Halides, *Angew. Chem. Int. Ed.* **2017**, *56*, 16367–16370. (c) Duong, H. A.; Wu, W.; Teo, Y.-Y., Cobalt-Catalyzed Cross-Coupling Reactions of Arylboronic Esters and Aryl Halides, *Organometallics* **2017**, *36*, 4363–4366.

10. (a) Bedford, R. B.; Hall, M. A.; Hodges, G. R.; Huwe, M.; Wilkinson, M. C., Simple mixed Fe–Zn catalysts for the Suzuki couplings of tetraarylborates with benzyl halides and 2-halopyridines, *Chem. Commun.* **2009**, 6430–6432. (b) Hatakeyama, T.; Hashimoto, T.; Kondo, Y.; Fujiwara, Y.; Seike, H.; Takaya, H.; Tamada, Y.; Ono, T.; Nakamura, M., Iron-Catalyzed Suzuki–Miyaura Coupling of Alkyl Halides, *J. Am. Chem. Soc.* **2010**, *132*, 10674–10676. (c) Hashimoto, T.; Hatakeyama, T.; Nakamura, M., Stereospecific Cross-Coupling between Alkenylboronates and Alkyl Halides Catalyzed by Iron–Bisphosphine Complexes, *J. Org. Chem.* **2012**, *77*, 1168–1173. (d) Hatakeyama, T.; Hashimoto, T.; Kathiriarachchi, K. K.; Zenmyo, T.; Seike, H.; Nakamura, M., Iron-Catalyzed Alkyl–Alkyl Suzuki–Miyaura Coupling, *Angew. Chem. Int. Ed.* **2012**, *51*, 8834–8837. (e) Bedford, R. B.; Brenner, P. B.; Carter, E.; Carvell, T. W.; Cogswell, P. M.; Gallagher, T.; Harvey, J. N.; Murphy, D. M.; Neeve, E. C.; Nunn, J.; Pye, D. R., Expedient Iron-Catalyzed Coupling of Alkyl, Benzyl and Allyl Halides with Arylboronic Esters, *Chem. Eur. J.* **2014**, *20*, 7935–7938. (f) Bedford, R. B.; Brenner, P. B.; Carter, E.; Clifton, J.; Cogswell, P. M.; Gower, N. J.; Haddow, M. F.; Harvey, J. N.; Kehl, J. A.; Murphy, D. M.; Neeve, E. C.; Neidig, M. L.; Nunn, J.; Snyder, B. E. R.; Taylor, J., Iron Phosphine Catalyzed Cross-Coupling of Tetraorganoborates and Related Group 13 Nucleophiles with Alkyl Halides, *Organometallics* **2014**, *33*, 5767–5780. (g) Crockett, M. P.; Tyrol, C. C.; Wong, A. S.; Li, B.; Byers, J. A., Iron-Catalyzed Suzuki–Miyaura Cross-Coupling Reactions between Alkyl Halides and Unactivated Arylboronic Esters, *Org. Lett.* **2018**, *20*, 5233–5237.

11. For an example of a reported Co- or Fe-catalyzed Suzuki biaryl cross-coupling performed under extremely high pressure, see: Guo, Y.; Young, D. J.; Hor, T. S. A., Palladium-free Suzuki–Miyaura cross-coupling at elevated pressures, *Tetrahedron Lett.* **2008**, *49*, 5620.

12. (a) Bedford, R. B.; Gallagher, T.; Pye, D. R.; Savage, W., Towards Iron-Catalysed Suzuki Biaryl Cross-Coupling: Unusual Reactivity of 2-Halobenzyl Halides, *Synthesis* **2015**, *47*, 1761–1765. (b) O'Brien, H. M.; Manzotti, M.; Abrams, R. D.; Elorriaga, D.; Sparkes, H. A.; Davis, S. A.; Bedford, R. B., Iron-catalysed substrate-directed Suzuki biaryl cross-coupling, *Nat. Catal.*, **2018**, *1*, 429–437.

13. (a) Kumar, L. M.; Bhat, B. R., Cobalt pincer complex catalyzed Suzuki–Miyaura cross coupling – A green approach, *J. Organomet. Chem.*, **2017**, *827*, 41–48. (b) Kumar, L. M.; Ansari, R. M.; Bhat, B. R., Catalytic activity of Fe(II) and Cu(II) PNP pincer complexes for Suzuki coupling reaction, *Appl. Organometal. Chem.* **2017**, *32*, e4054. (c) Ansari, R. M.; Bhat, B. R., Schiff base transition metal complexes for Suzuki–Miyaura cross-coupling reaction, *J. Chem. Sci.* **2017**, *129*, 1483–1490. (d) Ansari, R. M.; Kumar, L. M.; Bhat, B. R., Air-Stable Cobalt(II) and Nickel(II) Complexes with Schiff Base Ligand for Catalyzing Suzuki–Miyaura Cross-Coupling Reaction, *Russian J. Coord. Chem.*, **2018**, *44*, 1–8. (e) Ansari, R. M.; Mahesh, L. K.; Bhat, B. R., Cobalt Schiff base Complexes: Synthesis Characterization and Catalytic Application in Suzuki–Miyaura Reaction, *Chin. J. Chem. Eng.* **2018**, <https://doi.org/10.1016/j.cjche.2018.05.002>

14. Professor Bhat was contacted several times regarding our findings, the first time over eight months before submission of this manuscript, and he stands by his original claims. Meanwhile the editors of the five journals in question were contacted four months before submission. We thank the editors-in-chief of *J. Organomet. Chem.* and *J. Chem. Sci.* for responding.

15. (a) Pradhan, K.; Selvaraj, K.; Nanda, A. K., A Convenient Approach to the Synthesis of Different Types of Schiff's Bases and Their Metal Complexes, *Chem. Lett.* **2010**, 39, 1078-1079. (b) Chong, J. H.; Ardakani, S. J.; Smith, K. J.; MacLachlan, M. J., Triptycene-Based Metal Salphens—Exploiting Intrinsic Molecular Porosity for Gas Storage, *Chem. Eur. J.* **2009**, 15, 11824–11828. (c) Escudero-Adán, E. C.; Benet-Buchholz, J.; Kleij, A. W., Expedient Method for the Transmetalation of Zn(II)-Centered Salphen Complexes, *Inorg. Chem.* **2007**, 46, 7265–7267. (d) Mehta, B. H.; Joishar, D., X-Ray Crystallography and Biological Studies of Iron(II), Cobalt(II) and Nickel(II) Complexes Derived from Bidentate Schiff Bases, *Asian J. Chem.*, **2005**, 17, 1041-1049.
16. Kent Baretield, E.; Busch, D. H.; Nelson, S. M., Iron, cobalt, and nickel complexes having anomalous magnetic moments, *Quart. Rev., Chem. Soc.* **1968**, 22, 457.
17. Hobday, M. D.; Smith, T. D., N,N'-ethylenebis(salicylideneiminato) transition metal ion chelates, *Coord. Chem. Rev.* **1972-1973**, 9, 311-337.
18. van den Bergen, A.; Murray, K. S.; O'Connor, M. J.; Rehak, N.; West, B. O., N-Substituted salicylaldimine complexes of iron(III). I. Synthesis, infrared and mass spectra, *Aust. J. Chem.*, **1968**, 21, 1505-15.
19. Similarly intriguing is the complete lack of isotope patterns observed in the mass spectra of the analogous ML₂ complexes of Co(II), Ni(II) and Cu(II) reported in the same paper; the lack of any peak corresponding to ⁶³Cu in the latter case is particularly noteworthy.
20. (a) Elerman Y.; Paulus, H., A Five-Coordinate Bis(p-bromophenyl)-salicylaldimino)chloroiron(III) Complex, *Acta Cryst.* **1996**, C52, 1971-1973. (b) Fazekas, E.; Nichol, G. S.; Garden J. A.; Shaver, M. P., Iron(III) Half Salen Catalysts for Atom Transfer Radical and Ring-Opening Polymerizations, *ACS Omega* **2018**, 3, 16945–16953.
21. Examples: (a) Wang, L.; Sun, W.-H.; Han, L.; Li, Z.; Hu, Y.; He, C.; Yan, C., Cobalt and nickel complexes bearing 2,6-bis(imino)phenoxy ligands: syntheses, structures and oligomerization studies, *J. Organomet. Chem.* **2002**, 650, 59–64. (b) Du, J.; Han, L.; Cui, Y.; Li, J.; Li, Y.; Sun, W.-H., Synthesis, Characterization, and Ethylene Oligomerization of 2,6-Bis(imino)phenoxy Cobalt Complexes, *Aust. J. Chem.* **2003**, 56, 703–706. (c) Salehi, M.; Hasan-zadeh, M., Characterization, crystal structures, electrochemical and antibacterial studies of four new binuclear cobalt(III) complexes derived from o-aminobenzyl alcohol, *Inorg. Chim. Acta* **2015**, 426, 6–14.
22. The B3LYP functional has been shown to correctly predict the ground state of various Co(II) complexes, see, for example: Pietrzyk, P.; Srebro, M.; Radon, M.; Sojka, Z.; Michalak, A., Spin Ground State and Magnetic Properties of Cobalt(II): Relativistic DFT Calculations Guided by EPR Measurements of Bis(2,4-acetylacetonate)cobalt(II)-Based Complexes, *J. Phys. Chem. A* **2011**, 115, 2316–2324.
23. This contrasts with the ESI spectrum reported by Bhat for the supposed complex **7**, which shows a peak at *m/z* = 380.10, again notably devoid of an isotope pattern. See reference 13(e).
24. See Supplementary Information for details.
25. It is also worth noting that related pyridyl-NH-P(O)Ph₂ adducts of copper(II) are known, see: Yeh, C.-W.; Chang, K.-H.; Hu, C.-Y.; Hsu, W.; Chen, J.-D., Syntheses, structures and ligand conformations of Cu(II), Co(II) and Ag(I) complexes containing the phosphinic amide ligands, *Polyhedron* **2012**, 31, 657.
26. (a) Tisato, F.; Vallotto, F.; Pilloni, G.; Refosco, F.; Corvaja, C.; Corain, B., [Cu(H₂dped)](BF₄)₂(H₂dped = N,N'-bis(2-(diphenylphosphino)phenyl)ethane-1,2-diamine): a stable phosphine complex of copper(II), *J. Chem. Soc., Chem. Commun.*, **1994**, 2397.
- (b) Adner, D.; Möckel, S.; Korb, M.; Buschbeck, R.; Rüffer, T.; Schulze, S.; Mertens, L.; Hietschold, M.; Mehring, M.; Lang, H., Copper(II) and triphenylphosphine copper(I) ethylene glycol carboxylates: synthesis, characterisation and copper nanoparticle generation, *Dalton Trans.*, **2013**, 42, 15599-15609.
27. (a) Pan, Z.; Gamer, M. T.; Roesky, P. W., Copper(I) and Gold(I) Complexes with N,N'-Bis(diphenylphosphino)-2,6-diaminopyridine as Ligand: Synthesis and Molecular Structures, *Z. Anorg. Allg. Chem.* **2006**, 632, 744. (b) Arce, P.; Vera, C.; Escudero, D.; Guerrero, J.; Lappin, A.; Oliver, A.; Jara, D. H.; Ferraudi, G.; Lemus, L., Structural diversity in Cu(I) complexes of the PNP ligand: from pincer to binuclear coordination modes and their effects upon the electrochemical and photophysical properties, *Dalton Trans.*, **2017**, 46, 13432.
28. Benito-Garagorri, D.; Becker, E.; Wiedermann, J.; Lackner, W.; Pollak, M.; Mereiter, K.; Kisala, J.; Kirchner, K., Achiral and Chiral Transition Metal Complexes with Modularly Designed Tridentate PNP Pincer-Type Ligands Based on N-Heterocyclic Diamines, *Organometallics* **2006**, 25, 1900-1913.
29. Rösler, S.; Obenauf, J.; Kempe, R., A Highly Active and Easily Accessible Cobalt Catalyst for Selective Hydrogenation of C=O Bonds, *J. Am. Chem. Soc.* **2015**, 137, 7998–8001.
30. Searches performed 20th December 2018, using a phenolate ligand, with an ortho-OMe and an ortho C-N (with 'any bond' selected between the C and N) and bonds between the three heteroatom donors and a single metal (set as both 'any transition metal' and 'any metal').
31. An examination of Bhat's papers (ref. 13) shows that this remarkable insensitivity extends to variations in other parameters as well. For instance, replotting the results from the variation in solvents in the same reaction, catalysed by **9** or **11** (ref 13b) shows that while there is a difference claimed between the performance of the two catalysts, the slope of the linear trend line applied to the five data points in each solvent is constant (4.7(1) with **9**, 3.89(4) with **11**) across toluene, ethanol, 1,4-dioxane and acetonitrile, which suggests a notable insensitivity to the rate across a wide range of solvent types. See Figure S34, supplementary information for details.
32. Each catalytic reaction was tried independently by at least two authors.
33. Fürstner, A.; Leitner, A.; Méndez, M.; Krause, H., Iron-Catalyzed Cross-Coupling Reactions, *J. Am. Chem. Soc.* **2002**, 124, 13856–13863.
34. Bedford, R. B.; Bruce, D. W.; Frost, R. M.; Goodby, J. W.; Hird, M., Iron(III) salen-type catalysts for the cross-coupling of aryl Grignards with alkyl halides bearing β-hydrogens, *Chem. Commun.*, **2004**, 2822-2823.
35. Bedford, R. B.; Betham, M.; Bruce, D. W.; Davis, S. A.; Frost, R. M.; Hird, M., Iron nanoparticles in the coupling of alkyl halides with aryl Grignard reagents, *Chem. Commun.*, **2006**, 1398–1400.
36. In all these cases it is likely that the salen and Schiff-base complexes are converted to iron nanoparticles under the reaction conditions, see reference 35 for details.
37. For an earlier discussion on the problems of reproducibility in claimed iron-catalysed Suzuki biaryl cross-coupling, see: Bedford, R. B.; Nakamura, M.; Gower, N. J.; Haddow, M. F.; Hall, M. A.; Huwe, M.; Hashimoto, T.; Okopie, R. A., Iron-catalysed Suzuki coupling? A cautionary tale, *Tetrahedron Lett.* **2009**, 50, 6110–6111.

TOC Graphic: