Rosansky, S. J., Thamer, M., Caskey, F., Couchoud, C., McDonald, S. P., & Moist, L. (2019). A comparison of predialysis estimated glomerular filtration rate in the US, Canada, France, Australia, and the UK between 2005 and 2015. *Kidney International*, 95(5).
<https://doi.org/10.1016/j.kint.2019.02.008>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.kint.2019.02.008](https://doi.org/10.1016/j.kint.2019.02.008)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.kint.2019.02.008>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

A comparison of predialysis eGFR in the US, Canada, France, Australia and the UK between 2005-2015

Steven J Rosansky MD, Dorn Research Institute, WJBD VA, Columbia SC

Mae Thamer PhD Research Director Medical Technology & Practice
Patterns Institute
Bethesda, MD

Fergus Caskey MD Consultant Nephrologist, North Bristol NHS Trust
Honorary Senior Clinical Lecturer, School of Social and Community
Medicine, University of Bristol, Medical Director, UK Renal Registry,
Bristol

Cécile Couchoud MD Coordination Nationale de REIN
Agence de la Biomédecine 1 avenue du Stade de France

Stephen P McDonald MBBS(Hons) PhD FRACP Executive Officer,
ANZDATA Registry, Adelaide Australia

Louise Moist MD Division of Nephrology, Department of Medicine,
Western University, London, Ont.; Kidney Clinical Research Unit London
Health Sciences Centre, London, Ont.;

Correspondence
Steven Rosansky
8034225427
sjrcra@yahoo.com
526 N Trenholm Rd
Columbia S.C.29206

Running head
Predialysis eGFR between 2005-2015

Evidence from international observational studies and one randomized controlled trial found no survival benefit of "early start "of dialysis, generally defined as predialysis eGFR >10 ml/min/1.73m² (ES)(1). We compared the frequency of ES as reported to dialysis registries for the US, Canada, France, Australia and the UK between 2005- 2015. In 2015, ES was more common in the US, Canada and France (38.8%, 37.2%, 31.8%, respectively) than Australia and the UK (21.7% and 19.1% respectively), It has been suggested that early start rates may be influenced by financial incentives, but this alone does not explain these differences (2,3).

For dialysis patients aged ≥ 75 ES was approximately half as frequent in the UK and Australia versus the US and Canada and France and appears to have plateaued for most countries by 2010 (figure1). Predialysis eGFR >15 ml/min/1.73m² for ≥ 75 new dialysis starts with comorbidity congestive heart failure in the US, Canada and France was 17.0%, 17.6% and 18.0%, respectively, a finding that deserves further exploration.

The first eGFR related dialysis initiation guidelines appeared in the US in 1997(1). These guidelines might have had the greatest impact on ES in the US. For African Americans ES in 2015 was 26.7% versus 39.4% for whites. Thus, the US may have the highest rate of early start after controlling for race. Since ES has not been shown to provide a comorbidity adjusted survival or quality of life benefit one could argue that all five countries should find ways to decrease the rates of ES. This may be especially important for the 75+ populations where conservative management may be the preferred approach (4)

1. Rosansky SJ, Cancarini G, Clark WF et al. Dialysis initiation: what's the rush? *Semin Dial.* 2013;26:650–7
2. Vanholder R, Davenport A, Hannedouche T et al. Reimbursement of Dialysis: A Comparison of Seven Countries *J Am Soc Nephrol.*2012; 23: 1291–1298
3. Berns JS, Saffer TL, LinE. Addressing Financial Disincentives to Improve CKD Care *J Am Soc Nephrol.* : 2018, 29
4. Rosansky SJ, Schell J, Shega J et al. Treatment decisions for older adults with advanced chronic kidney disease *BMC Nephrology* .2017;18:200.

Figure 1 legend:

Proportion of the population aged ≥ 75 years initiating dialysis “early”, defined by starting eGFR > 10 ml/min/1.73m² shown in two categories: eGFR 10 - 14.9; and eGFR ≥ 15 ml/min/1.73m² in 2005, 2010 and 2015

across five countries. The incident average dialysis ≥ 75 aged population during this time period was 28,214 for the US, 1,330 for Canada, 3,310 for France, 492 for Australia, and 1,652 for U.K. Missing starting eGFR measurements were removed from all calculations and measured on average between 2005-2015-- 2.9% in the US, 9.3% in Canada, 17.8% in France, 1.9% in Australia, and 51.8% in the U.K. France used data from as many regions as were available for each study period.