Saitta, E. T., Kaye, T. G., & Vinther, J. (2019). Sediment-encased maturation: a novel method for simulating diagenesis in organic fossil preservation. *Palaeontology*, 62(1), 135-150.
<https://doi.org/10.1111/pala.12386>

Peer reviewed version

License (if available):
Other

Link to published version (if available):
[10.1111/pala.12386](https://doi.org/10.1111/pala.12386)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://doi.org/10.1111/pala.12386> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Sediment-encased maturation: a novel method for simulating diagenesis in organic fossil preservation

Evan T. Saitta¹, Thomas G. Kaye², & Jakob Vinther^{1,3}

¹School of Earth Sciences, University of Bristol, Bristol, United Kingdom; e-mails: evansaitta@gmail.com (ORCID ID: orcid.org/0000-0002-9306-9060), jakob.vinther@bristol.ac.uk (ORCID ID: orcid.org/0000-0002-3584-9616)

²Foundation for Scientific Advancement, Sierra Vista, Arizona, USA; e-mail: tom@tomkaye.com (ORCID ID: orcid.org/0000-0001-7996-618X)

³School of Biological Sciences, University of Bristol, Bristol, United Kingdom

ABSTRACT: Exceptional fossils can preserve diagenetically-altered biomolecules, and understanding the pathways to such preservation is vital to utilising fossil information in evolutionary and palaeoecological studies. Experimental taphonomy explores the stability of tissues during microbial/autolytic decay or their molecular stability through maturation under high pressure and temperature. Maturation experiments are often hampered by the fact that maturation occurs inside sealed containers, which does not allow for the loss of labile, mobile, or volatile molecules. On the other hand, maturation of tissues wrapped inside aluminium foil can sometimes be too open of a system, leading to loss of both labile and recalcitrant materials. Here we present a novel experimental procedure for maturing tissues under elevated pressure/temperature inside compacted sediment. In this procedure, porous sediment allows for labile maturation breakdown products to escape from the sample into the sediment and maturation chamber while recalcitrant,

immobile components are contained, in a manner expected to be similar to the natural conditions of fossilisation. To test the efficacy of this novel procedure with respect to simulating fossil diagenesis in a simple case study, we investigate the differential survival of melanosomes relative to proteinaceous tissues through maturation of fresh lizard body parts and feathers. Macro- and ultrastructures are then compared to fossils. Similar to many carbonaceous exceptional fossils, the resulting organic components are thin, dark films composed mainly of exposed melanosomes resting on the sediment in association darkened bones. Keratinous, muscle, collagenous, and adipose tissues appear to be lost. Such results are consistent with predictions derived from non-sediment-encased maturation experiments and our understanding of biomolecular stability. These experiments might also suggest that organic preservation is largely driven by the original molecular composition of the tissue and the diagenetic stability of those molecules, rather than the tissue's decay resistance alone, and this should be experimentally explored in the future.

KEY WORDS: maturation, diagenesis, fossils, melanosomes, protein, taphonomy

TAPHONOMY examines the physical (e.g., transport or burial) and chemical (e.g., microbial/autolytic decay or diagenesis) changes involved in fossilisation, from death to discovery (Parry *et al.* 2018). Determining the relative impact of processes that alter organisms during fossilisation has benefitted from advances in analytical techniques and experimental frameworks (Briggs & McMahon 2016; Parry *et al.* 2018; Purnell *et al.* 2018). Experiments may help elucidate the chemical and physical factors influencing decay (Briggs & Kear 1993, Clements *et al.* 2017), patterns of decay (Sansom *et al.* 2010a, 2010b, 2013), and authigenic mineralisation of decaying tissues (Briggs *et al.* 1993; Briggs & Kear 1994). Decay resistance, however, may not always

correlate to fossilisation potential (Wilson *et al.* 2016; Parry *et al.* 2018). Some decay-resistant tissues have low fossilisation potential, such as the keratin component of integument (Saitta *et al.* 2017a) or the collagenous notochord (Parry *et al.* 2018), and decay-labile tissues can contain recalcitrant components, such as melanosomes within the retinal pigment epithelium (Clements *et al.* 2016). For example, eyes decay relatively quickly (Sansom *et al.* 2010a, 2010b, 2013), but preserve commonly in carbonaceous exceptional fossils (Vinther *et al.* 2008; Sallan and Coates 2014; Clements *et al.* 2016).

The role and challenges of maturation experiments

In order to understand chemical changes to organic molecules, taphonomists have adopted methodologies from fossil fuel petrology and organic geochemistry (Rhead *et al.* 1971; Peters *et al.* 1981; Lewan 1983; Monthieux *et al.* 1985; Lewan *et al.* 1986; Rullkötter & Marzi 1988; Teerman & Hwang 1991; Behar *et al.* 1995, 1997, 2003; Koopmans *et al.* 1996; Versteegh *et al.* 2004; Schimmelmann *et al.* 2006). These maturation experiments, as they are called, illuminate the alterations of organic molecules through exposure to elevated temperatures and pressures (typically ~200–300 °C/several hundred bars/hours–days for paleontological studies) and have proven valuable in simulating the processes taking place during late diagenesis (Stankiewicz *et al.* 2000; Gupta *et al.* 2006a, 2007a; Schiffbauer *et al.* 2012; Glass *et al.* 2013; Colleary *et al.* 2015; Briggs & McMahon 2016; Saitta *et al.* 2017a). The underlying concept behind maturation experiments is kinetics; by subjecting samples to elevated temperatures and pressures over short timescales, chemical degradation reactions that normally occur at relatively lower diagenetic temperatures over much longer, geologic durations are accelerated and can be observed.

Maturation experiments can help determine which organics have high fossilisation potential and can produce results that approach fossils in chemical signature (Rhead *et al.* 1971; Stankiewicz *et al.* 2000; Gupta *et al.* 2006a, 2007a; Glass *et al.* 2013; Colleary *et al.* 2015; Saitta *et al.* 2017a). The stability of organic molecules over geologic timescales is related to their chemical structure, which influences the thermodynamics and kinetics of chemical reactions. Hydrophobic, water-insoluble organic molecules with a carbon backbone can show high longevity with minimal alteration, while nucleic acids and proteins show the lowest longevity of biomacromolecules due to their thermodynamically unstable phosphodiester and peptide bonds and the varying thermal stability of amino acids (Eglinton & Logan 1991).

Maturation experiments typically take place within small, sealed capsules composed of noble metals (e.g., ~2 mm diameter in some water-pressurised autoclaves); labile or volatile molecules produced during maturation that might normally be lost during fossilisation are trapped in the capsule (Saitta *et al.* 2017a) and will be analysed alongside more stable molecules, unless subsequent treatment of the product with solvents is performed (Stankiewicz *et al.* 2000). Furthermore, macrostructural observations are often difficult, and decay and maturation experiments are rarely integrated (Gupta *et al.* 2009). It is possible to mature larger tissues within aluminium foil using an Ar gas pressurised autoclave (McNamara *et al.* 2013), but this can allow for leaking of maturation products. Within common timeframes used in maturation experiments (e.g., ~24 hours or more), entire tissue samples can leak from such a foil setup and be lost (Saitta *et al.* 2017a). Note that these previous foil experiments (McNamara *et al.* 2013) only reported on one-hour maturation treatments (McNamara *et al.* 2017) with 24 hour durations resulting in sample loss (Saitta *et al.* 2017a).

94
95 We have designed an experimental protocol that allows for maturing small tissue samples within
96 compacted sediment. Since tissues consist of a multitude of molecules with different stabilities,
97 maturation experiments inside small, sealed containers will generate a consortium of recalcitrant
98 and labile products. This consortium is typically resulting in a solid or fluid with some degassing
99 upon opening the container, as evidenced by a strong odour (Saitta *et al.* 2017a). Analysing this
100 mixture and comparing it to fossils may be complicated, as volatile and soluble molecules would
101 be expected to dissipate within porous rock matrix. Stankiewicz *et al.* (2000) treated matured
102 samples with chloroform to yield the recalcitrant fraction only. However, this approach would be
103 expected to result in the loss of molecules that are soluble in nonpolar organic solvents that may
104 be of interest as geochemical biomarkers, such as steroid lipids (Mackenzie *et al.* 1982; Dreier *et*
105 *al.* 1988; Stevens *et al.* 2010; Melendez *et al.* 2013; Plet *et al.* 2017). During fossilisation, tissues
106 are held within a matrix of sediment, which is sometimes, but not always, cemented. Sediment is
107 predicted to hold the more stable and less mobile components of tissues in place, while fine pore
108 spaces allow for labile fluids and gasses to leach away. Therefore, fossilisation is often expected
109 to occur, at least partly, in a fairly open system.

110 To overcome these difficulties, with implications for testing a wide variety of taphonomic
111 hypotheses, our novel experimental procedure involves initial compaction in sediment, followed
112 by maturation. In order to monitor tissue macrostructure with heterogeneous composition (e.g.,
113 tissues with colour patterns or multiple organs), the maturation chamber is relatively large (19 mm
114 inner diameter as opposed to ~2 mm diameter for some water-pressurised autoclaves). We
115 hypothesise that in our experiments, the encasing, compacted sediment will establish an open-

system filter whereby labile degradation products escape out into the sediment and maturation chamber. Recalcitrant, immobile components with preservation potential, on the other hand, are expected to remain trapped within the compacted specimen. This experimental platform would then allow for testing of the relative stability and mobility of the various organic and inorganic components in complex tissues and for predicting what components of heterogeneous tissues are likely to be lost or retained in fossils. As discussed in the following section, previous analyses of fossils and non-sediment-encased maturation experiments, as well as an understanding of biomolecular stability, allow us to predict how a sediment filtration mechanism, similar to that expected during fossilisation, might appear.

Testing the protocol: melanosomes as key variable in vertebrate soft tissue preservation

To test this hypothesis of sediment filtration, we chose to mature two types of vertebrate specimens that occasionally preserve carbonaceous soft tissue fossils: feathers and scale-clad diapsids, here represented by lizards. Feathers are mainly composed of keratin protein and can be pigmented, most commonly by melanin. Freshly deceased lizards will have various proteins (mostly keratin, collagen, and muscle), adipose tissue and other lipids, as well as pigments like melanin. The results of our sediment-encased maturation experiments were then compared to fossils macro- and ultrastructurally through microscopy to judge the efficiency of the procedure at establishing a sediment filter and improving upon previous maturation experiments in simulating fossilisation.

According to a conservative interpretation of organic stability, amniote integument, including feathers and diapsid skin, can preserve in fossils as a film composed of melanosomes (Vinther *et al.* 2008, 2016; Lindgren *et al.* 2014; Vinther 2015). Non-pigmented feathers and scales

139 leave no organic residue, but can still preserve through residues of calcium phosphate, known to
140 harden keratinous tissues *in vivo* (Blakely *et al.* 1963; Pautard 1963, 1964), which can be detected
141 through fluorescence imaging (Mayr *et al.* 2016; Vinther *et al.* 2016). However, we focus here on
142 organic preservation pathways.

143 The discovery that melanosomes preserve commonly in exceptional fossils (Vinther *et al.*
144 2008; Colleary *et al.* 2016) has opened new avenues of palaeontological research, but their study
145 has not been without controversy. These structures have alternatively been identified as fossil
146 bacteria (Wuttke 1983; Davis & Briggs 1995), a position still maintained by some (Moyer *et al.*
147 2014; Lindgren *et al.* 2015; Schweitzer *et al.* 2015). However, this stance has been countered by
148 the fact that microbodies commonly found in fossil skin, hair, and feathers conform to
149 melanosomes in distribution, size, and organisation (Vinther *et al.* 2008; Vinther 2015, 2016).
150 Furthermore, chemical analyses show that these structures contain melanin (Glass *et al.* 2012,
151 2013; Lindgren *et al.* 2012, 2014; Colleary *et al.* 2015; Clements *et al.* 2016; Gabbott *et al.* 2016;
152 Brown *et al.* 2017).

153 In addition, it has been suggested that keratin protein can preserve organically in fossils
154 (Schweitzer *et al.* 1999, 2015; Edwards *et al.* 2011; Moyer *et al.* 2016a, 2016b) and, thus,
155 melanosomes might be obscured by a keratin protein matrix (Zhang *et al.* 2010; Moyer *et al.* 2014;
156 Pan *et al.* 2016). However, studies have described melanosomes as being exposed on the sediment
157 while non-pigmented regions yield only rock matrix (Vinther *et al.* 2008; Colleary *et al.* 2015;
158 Vinther 2015). More recently, studies have argued that the only constituents of keratinous tissues
159 preserved deep in the geologic record are calcium phosphate and pigments (Mayr *et al.* 2016;
160 Vinther *et al.* 2016; Saitta *et al.* 2017a).

Experimental taphonomy echoes the poor preservation potential of keratin protein. Previous, non-sediment-encased maturation experiments turned keratin into a volatile-rich, water-soluble fluid (Saitta *et al.* 2017a) while extracted melanosomes can survive largely intact (Colleary *et al.* 2015). This contrasting behaviour between keratin protein and melanin during maturation suggests that diagenetic degradation and loss of keratin protein is to be expected in fossils, leaving behind melanosomes. However, this keratin-melanin dynamic has yet to be experimentally observed simultaneously.

Similarly, decay-resistant collagenous tissue (Sansom *et al.* 2010a, 2010b, 2013) is expected to be diagenetically unstable (Parry *et al.* 2018), like other proteinaceous organics (Bada *et al.* 1999). However, epidermal collagen protein has been proposed in Mesozoic fossils (Lingham-Soliar *et al.* 2007), although such claims are debated (Smith *et al.* 2015; Smithwick *et al.* 2017).

Regarding the specific experiments described here, we hypothesise that tissues containing diagenetically-unstable organics like proteins and labile lipids (e.g., keratinous, collagenous, muscle, and adipose tissues) will largely become degraded and lost into the sediment, while diagenetically-stable organics like melanin (i.e., melanosomes) will tend to remain with the specimen. If the hypothesised filtering effect of porous sediment on diagenetically-altered organic materials is correct, then experimental results from sediment-encased maturation should be expected to resemble fossils such that preserved organic stains consist largely of exposed melanosomes resting on the sediment with a loss of surrounding tissues.

A statement on the relative role of diagenesis versus decay in molecular preservation?

A second aspect of our experiments involves investigating the relative importance of diagenesis and decay in carbonaceous exceptional fossils. None of our experimental samples were subjected to a decay treatment prior to sediment-encased maturation. In many natural settings, these tissues are often rapidly decayed through autolytic or microbial mechanisms. We therefore specifically test the survivability of pristine (i.e., non-decayed) tissues during maturation. We predict that if molecular thermal stability is a major factor in determining the composition of carbonaceous exceptional fossils, then our experimentally matured samples will experience dramatic volume loss and largely consist of a film of melanosomes similar to fossil observations without requiring a prior decay treatment.

METHODS

Fresh feathers (*Gallus gallus* and *Meleagris gallopavo* from UK farms) and lizards (*Anolis* captured from the wild in Arizona, USA) were matured shortly after acquisition/humane euthanasia via CO₂ asphyxiation with their full range of tissue composition present (see Saitta *et al.* (2018) for additional, preliminary results on non-vertebrates). Specimens were buried in easily-compacted bentonitic clay (purchased from Clay Terra) inside a metal piston and compacted using a hydraulic press (9–18 tonnes over 126.7 mm²), producing a consolidated tablet (Fig. 1A). Previous attempts with loose sediment did not produce results macrostructurally comparable to fossils, or amenable to easy structural analysis, indicating that compaction is important in establishing the desired pore space filtration. Tablets were loaded into a welded metal tube (19 mm inner diameter), forming an airtight chamber tapped for a high-pressure airline, with the goal of providing space for the escape of maturation products from the sediment (Fig. 1B). The chamber

resided inside a ceramic-lined laboratory oven. The airline exited a hole in the oven and connected a pressure-regulated air compressor. Experiments ran at 210–250 °C/225–300 bars/12–23 hours (Saitta *et al.* 2018), consistent with other maturation studies of fossilisation. After maturation, the chamber was cut open using a lathe. Tablets were split into a part and counterpart and analysed with light microscopy (Nikon SMZ-U stereo microscope, Nikon D810 DSLR camera) and scanning electron microscopy (SEM, gold coated specimens, Zeiss Evo10 SEM, Lab6 emitter).

Institutional abbreviations. BHI, Black Hills Institute Museum, Hill City, South Dakota, USA; BSP, Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany; IVPP, Institute of Vertebrate Palaeontology and Palaeoanthropology, Beijing, China; LEIUG, Leicester University Geology Department, Leicester, UK; MB, Humboldt Museum für Naturkunde, Berlin, Germany; MU, National Museum of Ethiopia, Authority for Research and Conservation of Cultural Heritage, Addis Ababa, Ethiopia.

RESULTS

Matured sediment tablets consistently split to reveal organic-containing parts and counterparts amenable to being easily manipulated and analysed. Lizard body parts lost significant volume (Fig. 2A–I). Only flattened and darkened skeletal material remained (greater temperature produced darker hues in the bones), surrounded by a thin, dark film where soft tissues existed previously, resulting in a body outline. SEM revealed that these organic stains lacked original tissue structure and histology, were associated with sediment impressions, and contained exposed oblate melanosomes (about 0.4 and 0.3 µm in length and width, respectively) resting on the sediment. No

ligaments or other clearly identifiable collagenous, keratinous, adipose, or muscle tissues were discernable.

Feathers lost volume, producing thin, dark stains (Fig. 3A–J). SEM revealed that these organic stains contained free-lying oblong melanosomes (about 1.1 and 0.2 μm in length and width, respectively) in regions that were pigmented *in vivo*, resting on the sediment. The melanosomes lost their original alignment. Regions that were non-pigmented *in vivo* lacked melanosomes and showed barb and barbule impressions in the sediment.

Variation in the amount of preserved organics between the part and counterpart was sometimes apparent (Saitta *et al.* 2018). The samples appear to be stable after maturation with no noticeable changes having occurred over more than a year of storage at dry, room temperature conditions inside a plastic SEM stub box.

DISCUSSION

In the following discussion, the feather and lizard sediment-encased maturation experiments' ability to discriminate stable from unstable organics as well as their similarity to fossils is judged. Then, the experiments are discussed in relation to how they might elucidate our understanding of the importance of diagenesis in fossilisation and of the correlation between molecular stability and preservation. Next, the usefulness of sediment-encased maturation in understanding fossilisation is compared to that of previous taphonomic experiments and methodologies, with particular discussion of the importance and implications of the sediment filtration mechanism. Finally, the potential and versatility of sediment-encased maturation is discussed, particularly in relation to

preservational modes beyond that of the melanin and bone preservation similar to some carbonaceous exceptional fossils as demonstrated by the experiments in this study.

Feather and lizard experiments: selective preservation in a manner resembling some carbonaceous exceptional fossils

The observed changes in our sediment-encased maturation experiments are very consistent with observations from carbonaceous vertebrate tissues in exceptional fossils. Matured samples resemble fossils in many ways, and over a range of scales and specimens, evidenced by ultrastructural features observed through microscopic structural analysis. Melanosomes are the sole unequivocally identifiable component of the dark soft tissue films. Tissues underwent extensive volume loss, and proteinaceous and adipose tissues appear to have, at least largely, degraded and escaped into the sediment. Bones and teeth, of course, easily survived, as in the fossil record. Sediment-encased maturation is therefore expected to allow for comparisons of the survivability of organics by simulating deep-burial diagenesis. As is the case for any scientific model, including the physical model used to simulate fossilisation through diagenesis here, results bearing resemblance to the natural system are a testament to the accuracy of the model and, by extension, our understanding of the underlying processes involved.

Like the experimentally matured lizard skin observed here, fossil reptile skin preserves melanosomes (Li *et al.* 2014; Lindgren *et al.* 2014; Vinther *et al.* 2016; Lindgren *et al.* 2017), as is the case for other tetrapod skin fossils (Fig. 2J–L; Colleary *et al.* 2015). Connective, muscle, and adipose tissues appear to have largely escaped as breakdown products into the sediment and chamber so as to be unrecognisable ultrastructurally. Although some of these tissues can quickly

decay, these experiments suggest that if they were to survive decay, subsequent significant diagenetic alteration likely eliminates them. As Parry *et al.* (2018) discussed, decay resistance does not always correlate with preservation potential, ultimately.

Similarly, liberated melanosomes are the only unequivocally identifiable structure in the matured feathers, as in fossil feathers (Vinther *et al.* 2008; Vinther 2015) (Fig. 3K–L). Original melanosome alignment is lost, potentially due to a lack of continuous, concurrent compaction during keratin degradation whereby encasing sediment might help to lock melanosomes in place and regulate the release of keratin degradation products away from the specimen. An optional second compaction after maturation can superficially remove void spaces but lacks this concurrent dynamic likely required to produce more realistic results.

Variation in the degree of organic staining preserved on the part compared to the counterpart is likely a result of recalcitrant organics settling within the sediment under gravity as tissues undergo volume loss. Again, the lack of continuous, concurrent compaction of the sediment as would happen in nature allowed the resulting residues to easily accumulate at the bottom of void spaces. However, different organic staining patterns between parts and counterparts due to uneven splitting occur in fossils as well (e.g., MB.Av.100 and BSP 1869 VIII 1, counterpart and part of the isolated feather assigned to *Archaeopteryx*).

The importance of diagenesis in carbonaceous exceptional fossils

In addition to discriminating between different organismal components in a manner similar to some carbonaceous exceptional fossils, these sediment-encased maturation experiments may provide insight into the relative importance of diagenesis within the overall taphonomic process.

It must first be kept in mind that no samples underwent decay treatments prior to experimentation, yet produced macro- and microscopic results comparable to fossils. The observed microbodies in regions of the samples that were pigmented *in vivo* can therefore be reliably identified as melanosomes as opposed to bacteria, providing yet another line of evidence within the growing consensus that such structures in fossils are indeed melanosomes (Vinther 2016).

More importantly, however, these non-decayed samples highlight the influence of the variables simulated with maturation (i.e., the time, heat, and pressure of diagenesis) as well as the effect of encasing sediment during fossilisation, particularly regarding carbonaceous exceptional fossils. While decay can result in tissue loss and obscure tissue identification in fossils (Briggs and Kear 1993; Sansom *et al.* 2010a, 2010b, 2013), our results appear to demonstrate that sediment-encased maturation on its own can discriminate between tissues with low and high organic preservation potential based on the thermodynamic stability of their molecular components and the mobility of their breakdown products. Unstable, mobile organics and their breakdown products appear to be lost through sediment pores while stable, immobile organics remain *in situ*. Experimentally-produced soft tissue carbonaceous stains resemble preservational modes of some Konservat-Lagerstätten like the Yixian Formation (see Benton *et al.* 2008 for a review of such Jehol Biota). Although carbonaceous fossils exhibit inter-locality variation, they might generally result from decay-limited environments, with late diagenesis ultimately impacting which organics are preserved or lost. Minimising early taphonomic processes (e.g., scavenging, transport, decay) might generally favour whole, articulated, organically-preserved, soft tissue-containing fossils. Maturation experiments would therefore be important to understand many Konservat-Lagerstätten.

Consideration of biomolecular stability informs predictions of preservation potential. Thinking in terms of diagenetic influences on fossilisation, one would benefit from first considering molecular thermal stability when attempting to ascertain which molecules might survive these sediment-encased maturation experiments and, we suspect, in fossils themselves. Proteins are not predicted to be diagenetically stable in sediments (Bada *et al.* 1999) and appear to have been largely lost during the maturation treatment. Similarly, adipose tissues have largely been lost during maturation and are composed of triglycerides (Björntorp 1996) that are also predicted to be thermally unstable due to hydrolytic cleavage of ester bonds between fatty acids and the hydrophilic glycerol group (Wardell *et al.* 1981; Eglinton & Logan 1991). Cell membranes are composed mainly of phospholipids, which again are expected to be thermally unstable due to hydrolytic cleavage of ester bonds between fatty acids and the hydrophilic glycerol-phosphate polar heads (Eglinton & Logan 1991; Zuidam & Crommelin 1995). The free fatty acid chains of labile lipids like triglycerides and phospholipids can, however, undergo *in situ* polymerisation to preserve in fossils as aliphatic hydrocarbons (Stankiewicz *et al.* 2000; Gupta *et al.* 2006a, 2006b, 2007a, 2007b, 2007c, 2008, 2009). Although chemical analysis was not performed in this study, it is possible that such kerogen-like aliphatic molecules are present in experimental samples. Indeed, occasionally an unidentifiable amorphous material was observed in the matured lizards under SEM, consistent with a kerogen-like substance but requiring further investigation (Saitta *et al.* 2018). Similarly, sterols are thermally stable and often have low solubility in water with an extremely high preservation potential (Mackenzie *et al.* 1982; Dreier *et al.* 1988; Stevens *et al.* 2010; Melendez *et al.* 2013; Plet *et al.* 2017), and it is possible that sterols such as cholesterol remained in the experimental samples, again requiring further chemical analysis. However, before any further discussion of potential avenues for future research into sediment-encased maturation,

the observed results of the experiments described herein should be used to inform a decision about the utility of the novel protocol in studying taphonomy.

Sediment-encased maturation: improved results over other taphonomic experiments

How does the novel protocol presented here compare to other methods in experimental taphonomy at realistically modelling fossilisation? Does it represent a significant methodological improvement?

It seems that sediment-encased, pressure-temperature maturation produces results comparable to carbonaceous fossils and, potentially, sub-fossil resins (see Saitta *et al.* (2018) for supplemental, preliminary results showing hardened tree resin with evidence of escaped maturation products, leaves producing dark stains with associated volume loss, and arthropod cuticles showing browning and relatively high stability through maturation). In comparison, decay experiments can sometimes authigenically mineralise certain tissues through phosphatisation (Briggs & Kear 1993) or pyritisation (Grimes *et al.* 2001), but do not result in the sort of carbonaceous preservation observed here. While acknowledging the ability for decay to induce authigenic mineralisation in some circumstances, Parry *et al.* (2018) discussed how decay resistance does not always correlate to organic preservation. Although decay can expose melanosomes from a keratin matrix (Saitta *et al.* 2017b), the results here show that this can be achieved by maturation alone, consistent with the idea that decay is more of a factor in instances of unexceptional preservation rather than exceptional carbonaceous preservation (Parry *et al.* 2018). Acidic-thermal experiments, on the other hand, produce compounds unlike fossils, such as SiC ceramics from wood (Shin *et al.* 2005), while attempts to produce highly-precise silicified or

carbonised 3-dimensional structural replicas of whole organisms through to cells and extra-cellular proteins via replacement of original organics (Townson *et al.* 2014) similarly appear to produce results largely incomparable to fossils, and the experimental conditions of such treatments are likely inappropriate for modelling natural environments of fossilisation.

Non-sediment-encased maturation experiments do reveal quite a bit about organic preservation, such as the origin of sedimentary organic matter and fossil fuels (Rhead *et al.* 1971; Peters *et al.* 1981; Lewan 1983; Monthioux *et al.* 1985; Lewan *et al.* 1986; Rullkötter & Marzi 1988; Teerman & Hwang 1991; Behar *et al.* 1995, 1997, 2003; Koopmans *et al.* 1996; Versteegh *et al.* 2004; Schimmelmann *et al.* 2006), *in situ* polymerisation of lipids (Stankiewicz *et al.* 2000; Gupta *et al.* 2006a, 2007a), or the fossilisation of melanin (Glass *et al.* 2013; Colleary *et al.* 2015). However, some maturation experiments suffer from methodological shortcomings that result in unrealistic simulation of diagenesis through time. For example, some maturation experiments only elevate temperature (Moyer *et al.* 2016a), allowing water to boil off. This is problematic for simulating fossilisation since no fossilisation pathway is entirely anhydrous, and hydrolysis, a major biomacromolecule degradation reaction, requires water, meaning that the conclusion of keratin preservation in those experiments by Moyer *et al.* (2016a) might be questionable. Furthermore, maturation duration should be of sufficient duration and temperature in order to mimic fossil chemistries of the appropriate age/thermal maturity (Saitta *et al.* 2017a, 2017b). For example, maturing feathers for only 1 hour does not completely degrade keratin (McNamara *et al.* 2013, 2017) and does not give comparable results to fossils of appreciable age or thermal maturity (Saitta *et al.* 2017a). However, in addition to careful consideration of maturation conditions such as duration and pressure, might our sediment-encased maturation experiments benefit from another improvement over these previous experiments, one that is more fundamental to the technique itself

rather than simply conditional? To answer this, one must consider how sediment encasing fundamentally changes the observed results of maturation experiments.

Sediment pore filtration: the key process. Maturation studies of organic molecules inside sealed, noble metal capsules produce an amalgam of recalcitrant and labile maturation products. However, by allowing labile molecules to escape, any subsequent analysis will only involve recalcitrant and immobile molecules, as one would expect to find in fossils. Here we show that sediment-encased maturation appears to allow degraded molecules like protein fragments to escape through pore spaces, while recalcitrant organelles and tissues like melanosomes and bones remain, as might be expected in fossilisation. In contrast, non-sediment-encased feather maturation under similar conditions hydrolysed keratin into a thick, water-soluble fluid not seen in fossils (Saitta *et al.* 2017a) nor in sediment-encased feather maturation. This non-sediment-encased feather maturation did not reveal melanosomes under SEM (Fig. 4A–B; Saitta *et al.* 2017a, 2017b), presumably obscured by the fluid.

We hypothesise that sediment-encased maturation might, therefore, yield chemical signatures more comparable to fossils than non-sediment-encased maturation. This hypothesis is consistent with the patterns of degradation observed here: the colour changes and loss of tissue structure are evidence that chemical alteration has occurred, while the volume loss and loss of certain tissues are consistent with the leaching of these degradation products, such as the hydrolysis of peptides and the leaching of their fragments from the specimen. For example, previous, non-sediment-encased maturation on melanin inside gold capsules showed that the resulting negative secondary ion spectra from time-of-flight secondary ion mass spectrometry (TOF-SIMS) began to converge upon those of fossil melanin when examined using principal component analysis

(Colleary *et al.* 2015). However, at best, the artificially matured melanin spectra only partially overlapped with those of fossil melanin. We hypothesise that this discrepancy could be due to trapped labile molecules in the non-sediment-matured melanin that are not present in the fossil melanin analysed alongside recalcitrant molecules, meaning that sediment-encased matured melanin might show improved chemical overlap with fossil melanin. Thus, precise chemical differences between sediment-encased and non-sediment-encased matured organics should be further investigated and compared to fossils in future experiments using various chemical analyses (e.g., TOF-SIMS). Based on our understanding of the underlying processes of maturation, particularly hydrolysis, chemical signatures of proteins and labile lipids, for example, are expected to decrease in more highly matured samples. In a similar manner, compacted sediment may even improve upon maturation experiments examining fossil fuel or sedimentary organic matter formation (Rhead *et al.* 1971; Peters *et al.* 1981; Lewan 1983; Monthioux *et al.* 1985; Lewan *et al.* 1986; Rullkötter & Marzi 1988; Teerman & Hwang 1991; Behar *et al.* 1995, 1997, 2003; Koopmans *et al.* 1996; Versteegh *et al.* 2004; Schimmelmann *et al.* 2006) or allow us to study differential migration patterns of various organics through sediment. In the meantime, however, the structural data presented here is sufficient to illustrate the mechanism of sediment filtration and the differing survivability of various organics.

But is establishing such a filtration mechanism indeed a worthwhile endeavour? Others in the past have attempted to separate out maturation products using organic solvents, such as chloroform (Stankiewics *et al.* 2000). While attempting to eliminate mobile maturation products from any subsequent chemical analysis might help produce results more comparable to fossils, the use of an organic solvent is not a proper simulation of the loss of mobile molecules that occurs in fossilisation, and some organic molecules that are capable of surviving in fossils might be organic-

soluble and lost from the experiment. Instead, our method accomplishes this filtration using a physical, rather than chemical mechanism, and this mechanism is directly comparable to what is expected to occur during fossilisation. The advantage of this filter is the ability to study variation in the preservation potential of different components within heterogeneous samples, such as whole bodies and tissues with colour patterns. In another example, Behar *et al.* (1997) used multiple cold trap pyrolysis-gas chromatography to mature kerogen in an open system with elevated temperature (300 °C), but for short durations (6 min) and without elevated pressure, and such open system attempts to artificially mature kerogen and other sedimentary organic matter while simultaneously allowing for fluid products to escape have been recognised as an established method (Behar *et al.* 2003; Schimmelmann *et al.* 2006). Again, compacted sediment likely represents a more realistic maturation environment for fossilisation without creating an overly open system that might result in the loss of stable organics naturally found in fossils, as occurred in the attempts to replicate the maturation experiments on feathers in foil of McNamara *et al.* (2013) by Saitta *et al.* (2017a). Furthermore, large sediment tablets make for easy downstream analyses of matured samples using various techniques beyond those such as pyrolysis-gas chromatography-mass spectrometry (Py-GC-MS), including surface analyses like TOF-SIMS.

Overall, based on the current experiments, our sediment-encased maturation appears to more closely model carbonaceous fossilisation than previous experimental attempts. Knowing this, what are the potential uses for such a protocol, and might it help to inform our understanding of other modes of fossil preservation?

The versatility and potential of sediment-encased maturation: new ways to test taphonomic hypotheses

458

459 Future experimentation with sediment-encased maturation will allow for examining the relative
460 impact of specific taphonomic variables. Sediment type, moisture content and fluid species/pH,
461 compaction magnitude/number, and maturation temperature, pressure, duration, and pressuring
462 gas species could all be controlled to test many taphonomic hypotheses. While the individual effect
463 of each of these variables in isolation has yet to be fully disentangled, given that the maturation
464 conditions used here are comparable to those of earlier non-sediment-encased maturation
465 experiments (particularly Saitta *et al.* 2017a), it can be safely concluded that it is largely the effect
466 of encasing sediment responsible for the dramatically improved results under our protocol.
467 Depending on the age and thermal maturity of the fossils one is interested in, further changes to
468 the experimental variables of our method might produce results more comparable to those fossils.

469 Furthermore, additional consideration can be made as to exactly what organics one wishes
470 to examine for potential selective preservation of certain components. Here, we focus mainly on
471 relative survivability of vertebrate melanin and proteins, but such studies could be done on any
472 type of heterogeneous tissue containing varying organics (e.g., those containing chitin, cellulose,
473 various lipids, etc.).

474 Large sample sizes allow for easier observations of many tissue types simultaneously, and
475 the parts and counterparts produced are easily manipulated by hand and amenable to various types
476 of downstream structural and chemical analyses commonly run on fossils (e.g., placement onto
477 stubs for SEM or TOF-SIMS).

478

479 *Combining decay and maturation.* A decay treatment could be applied in order to understand the
480 impact of tissue loss prior to maturation (Sansom *et al.* 2010a, 2010b, 2013; Briggs & McMahon

2016; Parry *et al.* 2018). Subsequent maturation of specimens decayed within sediment is often complicated due to difficulties in extricating the tissues from the sediment without affecting structural observations. Our method can help to circumvent this issue, since the buried, decayed tissue can first be compacted into a sediment tablet that can then be easily split apart after maturation. Attempting to seriously couple decay and maturation experiments in the future is important since decay inside sediment may lead to novel chemical reactions that are of significance in fossilisation (Wilson and Butterfield 2014). The hypothesis of ‘stemward slippage’ observed through decay (Sansom *et al.* 2010a, 2010b) can be easily tested all the way through diagenesis using sediment-encased maturation to determine if the stemward pattern of character loss from decay is still present after samples at various decay stages have been matured.

Authigenic mineralisation can preserve decay-prone tissues, such as muscles and the digestive tract through phosphatisation (Briggs 2003; Parry *et al.* 2015) or arthropod cuticles through pyritisation (Briggs *et al.* 1991; Parry *et al.* 2018). By maturing specimens encased in sediment subsequent to decay, authigenic mineralisation can be easily studied in conjunction with organic diagenesis using our method. Decay is also thought to be important in the formation of carbonate concretions, which can lead to exceptionally preserved soft tissues (McCoy *et al.* 2015), although experimental concretion formation has yet to be achieved (Briggs & McMahon 2016).

Other preservational modes leading to organic stabilisation. Sediment-encased maturation will likely permit the easy testing of hypotheses regarding organic stabilisation mechanisms during decay and early diagenesis (e.g., kerogenisation, adipocere formation, sulfurisation, Maillard reactions, sub-fossil and fossil resin formation) as well as organic-inorganic interactions and how these affect the resulting stability of the organics through late diagenesis.

Decay can sometimes convert labile organic molecules into more recalcitrant forms. Studies have highlighted the potential for labile lipids such as fatty acids, endogenous to the organism within tissues like arthropod cuticle, graptolite periderm, or leaves, to produce kerogen-like, long aliphatic molecules in a process termed *in situ* polymerisation during both decay and maturation, which may be widely occurring in exceptional fossil preservation (Stankiewicz *et al.* 2000; Gupta *et al.* 2006a, 2006b, 2007a, 2007b, 2007c, 2008, 2009). For example, experiments with scorpion cuticle subjected to confined, non-sediment-encased maturation with subsequent chloroform washing to remove organic-soluble maturation products found Py-GC-MS profiles that closely resembled Carboniferous scorpion cuticle, largely consisting of aliphatic molecules (Stankiewicz *et al.* 2000). Preliminary, supplemental results of our experiments are promising with respect to possible kerogen-like preservation: angiosperm leaves lost volume and produced dark stains, and arthropod cuticles survived well through maturation and became browned (Saitta *et al.* 2018). Therefore, sediment-encased maturation may show promise regarding experiments on lipid *in situ* polymerisation to produce kerogen-like residues.

Sulphurisation of organic molecules is known to take place (Sinninghe Damste 1989; Sinninghe Damste & De Leeuw 1990), whereby S crosslinking stabilises molecules in a manner that leads one to draw similarities to the vulcanisation of rubber (Ding & Leonov 1996). While melanin has been shown to react with S (Glass *et al.* 2012; McNamara *et al.* 2016), it is already recalcitrant in nature. Organically-preserved muscle has been reported in a salamander specimen from Miocene lacustrine sediments in Spain (McNamara *et al.* 2010), which is remarkable for its organic preservation of an originally proteinaceous tissue due to extensive sulfurisation. The same deposits are also reported to yield what appears to be organically-preserved, sulphurised bone marrow (McNamara *et al.* 2006). Therefore, decay and maturation experiments under euxinic

conditions should be conducted in a sediment-encased protocol in order to examine the potential for organic preservation of labile tissue via S crosslinking.

Beyond the establishment of S crosslinking, other mechanisms of organic stabilisation through covalently-bonded crosslinks between structural biomolecules exist via tanning-like processes (Wilson & Butterfield 2014). Such crosslinking can derive from various agents including multivalent Fe, Ti, Zr, and Si, aluminosilicates, polyphosphates, and mineral salts such as aluminium sulphate or chromium sulphate. A common form of crosslinking involves polyphenolic tannins from plants that can bind to organic molecules like proteins, forming the foundation of the early leather tanning industry (Kanagy & Kronstadt 1943; Covington 1997; Hueffer *et al.* 2010). However, crosslinking might be more applicable to the archeological, rather than fossil, record, such as the cold, acidic, anoxic, reducing peat bog environment that preserves ‘bog bodies’ with sphagnum, which contains a carbonyl-rich polyssacharide (Painter 1991; Evershed 1992; Stankiewicz *et al.* 1997).

While the above examples of crosslinking greatly slow down the rate of organic degradation, it might not entirely halt the process, as evidenced by the limitations of tissue fixation with formaldehyde (Helander 1994; Kiernan 2000). Recently though, Wilson and Butterfield (2014) proposed that tissues decaying in kaolin, as opposed to sand, lime, or smectitic clays, may result in protein crosslinking, similar to some forms of leather tanning (Plapper *et al.* 1981), which might impact the quality of the resulting carbonaceous fossil. Even more applicable to the fossil record are chemical reactions involving protein degradation products, either through Maillard reactions between the amino group of amino acids (normally involved in peptide bonds prior to hydrolytic cleavage) and the carbonyl group of reducing sugars to form melanoidins that result in browning as in the cooking of food (Stankiewicz *et al.* 1997; Martins *et al.* 2000), or between

oxidised fatty acids and amino acids that also produce browning (Hidalgo *et al.* 1999). Sugars can also react with lipids, in addition to proteins, to form advanced glycation end products (AGEs). These reactions between sugars, lipids, and proteins are non-enzymatic and are accelerated under elevated temperatures (Uribarri *et al.* 2010). The darkened bones observed in our lizard experiments may derive from such browning processes and further chemical analyses should be carried out (Jasmina Wiemann, personal communication); similarly, browning in any under-matured samples should consider these reactions as potential explanations (e.g., the predominantly white region of the feather resulting in discoloured sediment in Fig. 3B might be due to slightly insufficient maturation conditions leading to incomplete protein degradation or from the low melanin concentrations in some places on this predominantly white region as visible in Fig. 3A). Sediment-encased maturation can begin to address what crosslinking processes are likely to occur in various taphonomic settings, what types of bonds survive subsequent diagenesis, whether or not initial crosslinking ultimately impacts the quality of the carbonaceous fossil, and whether any crosslinks can stabilise polymers such as peptides through diagenesis or if the only surviving derivatives of complex, labile biomolecules are simpler AGE or melanoidin-like molecules derived from previously hydrolysed polymers.

Perhaps the most remarkable form of organic preservation is in amber (e.g., Xing *et al.* 2017). Resin fossilisation involves water and monoterpene loss followed by polymerisation, isomerisation, and crosslinking of the remaining recalcitrant diterpenoids (Langenheim 1969, 1990; Lambert & Frye 1982; Mills *et al.* 1984; Pike 1993; Ragazzi *et al.* 1993; Villanueva-García *et al.* 2005). Preliminary, supplemental results here suggest that sediment-encased maturation can harden tree resin into a translucent mass with a surrounding escape front of maturation products leaching into the sediment (Saitta *et al.* 2018). Whether this hardening is only a result of the escape

of unstable components into the sediment, as in copal, or if subsequent polymerisation and crosslinking of the more stable components has also occurred, as in amber, should be examined in more detail using chemical analyses. Ultimately, sediment-encased maturation might provide an interesting platform by which to study the organic degradation of the inclusions within an organic matrix of stabilised resin. Thus, the method might be able to examine organic geochemistry in settings other than inorganic sediment matrices.

Mimicking the taphonomic histories and lithologies of specific Konservat-Lagerstätten using sediment-encased maturation may produce various preservational modes beyond those described here, especially if a decay treatment is performed prior to maturation in order to induce authigenic mineralisation alongside carbonaceous stains. The effect of sediment type or other abiotic factors of the burial environment on observed organic preservation can easily be examined using our method.

CONCLUSION

We propose that this novel experimental procedure of maturation in compacted sediment allows for observations of the macro- and ultrastructural changes in co-occurring tissues simultaneously. Sediment filtration lets labile molecules to escape while retaining recalcitrant ones. For the first time, we can use maturation to illustrate the dynamic of simultaneous loss versus retention of different organics in tissues (e.g., melanin vs. proteins), owing to the mechanism of sediment filtration as shown here. The results match with those predicted based on analyses of fossils and previous non-sediment-encased maturation experiments and are consistent with our understanding of biomolecular stability (Eglinton & Logan 1991). Sediment-encased matured samples are

comparable to carbonaceous exceptional fossils down to the ultrastructural level, highlighting the importance of diagenesis in such preservational modes in determining which tissues are likely to preserve as well as the differential stability of the molecular components of such tissues. Promising supplemental, preliminary results on non-vertebrate tissues (Saitta *et al.* 2018) and the ability to control many variables prior to and during maturation vouch for the versatility and potential of the protocol, providing a novel framework for hypothesis testing in taphonomy. These include examining the interplay between decay and maturation, which may potentially produce further preservational modes (e.g., authigenic mineralisation alongside carbonaceous preservation as well as crosslinking of more labile molecules). Sediment-encased maturation may one day improve upon the earliest maturation experiments studying fossil fuel formation.

Acknowledgments. Many thanks to reviewers James D. Schiffbauer, Imran Rahman, and Robert Sansom and to Nicholas Edwards and Wang Yuan for providing images. We also thank Matthew J. Collins and Jasmina Wiemann for helpful discussion. Work was funded by the Geological Society of London's Daniel Pidgeon Fund.

DATA ARCHIVING STATEMENT

Data for this study are available in the [Dryad Digital Repository]:
<http://datadryad.org/review?doi=doi:10.5061/dryad.0t67n>

Ethics. Chicken and turkey feathers were taken from purchased, deceased specimens from UK farms. Lizards, beetle, and plant tissues were captured/collected from the wild in Arizona.

Euthanasia adhered to UK Schedule One of the Animals (Scientific Procedures) Act of 1986.

REFERENCES

BADA, J. L., WANG, X. S. and HAMILTON, H. 1999. Preservation of key biomolecules in the fossil record: current knowledge and future challenges. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **354**, 77–87.

BEHAR, F., VANDENBROUCKE, M., TEERMANN, S. C., HATCHER, P. G., LEBLOND, C. and LERAT, O. 1995. Experimental simulation of gas generation from coals and a marine kerogen. *Chemical Geology*, **126**, 247–260.

— TANG, Y. and LIU, J. 1997. Comparison of rate constants for some molecular tracers generated during artificial maturation of kerogens: influence of kerogen type. *Organic Geochemistry*, **26**, 281–287.

— LEWAN, M. D., LORANT, F. and VANDENBROUCKE, M. 2003. Comparison of artificial maturation of lignite in hydrous and nonhydrous conditions. *Organic Geochemistry*, **34**, 575–600.

BENTON, M. J., ZHONGHE, Z., ORR, P. J., FUCHENG, Z. and KEARNS, S. L. 2008. The remarkable fossils from the Early Cretaceous Jehol Biota of China and how they have changed our knowledge of Mesozoic life: Presidential Address, delivered 2nd May 2008. *Proceedings of the Geologists' Association*, **119**, 209–228.

BJÖRNTORP, P. 1996. The regulation of adipose tissue distribution in humans. *International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*, **20**, 291–302.

642 BLAKEY, P. R., EARLAND, C. and STELL, J. G. P. 1963. Calcification of keratin. *Nature*, **198**,
643 481–481.

644 BRIGGS, D. E. G. 2003. The role of decay and mineralization in the preservation of soft-bodied
645 fossils. *Annual Review of Earth and Planetary Sciences*, **31**, 275–301.

646 — and KEAR, A. J. 1993 Fossilization of soft tissue in the laboratory. *Science*, **259**, 1439–1442.

647 — and KEAR, A. J. 1994. Decay and mineralization of shrimps. *Palaios*, **9**, 431–456.

648 — and MCMAHON, S. 2016. The role of experiments in investigating the taphonomy of
649 exceptional preservation. *Palaeontology*, **59**, 1–11.

650 — BOTTRELL, S. H. and RAISWELL, R. 1991. Pyritization of soft-bodied fossils: Beecher's
651 trilobite bed, Upper Ordovician, New York State. *Geology*, **19**, 1221–1224.

652 — KEAR, A. J., MARTILL, D. M. and WILBY, P. R., 1993. Phosphatization of soft-tissue in
653 experiments and fossils. *Journal of the Geological Society*, **150**, 1035–1038.

654 BROWN, C. M., HENDERSON, D. M., VINTHER, J., FLETCHER, I., SISTIAGA, A.,
655 HERRERA, J. and SUMMONS, R. E. 2017. An Exceptionally Preserved Three-
656 Dimensional Armored Dinosaur Reveals Insights into Coloration and Cretaceous Predator-
657 Prey Dynamics. *Current Biology*, **27**, 2514–2521.

658 CLEMENTS, T., DOLOCAN, A., MARTIN, P., PURNELL, M. A., VINTHER, J. and
659 GABBOTT, S. E. 2016 The eyes of *Tullimonstrum* reveal a vertebrate affinity. *Nature*,
660 **532**, 500–503.

661 CLEMENTS, T., COLLEARY, C., DE BAETS, K. and VINTHER, J. 2017. Buoyancy
662 mechanisms limit preservation of coleoid cephalopod soft tissues in Mesozoic
663 Lagerstätten. *Palaeontology*, **60**, 1–14.

664 COLLEARY, C., DOLOCAN, A., GARDNER, J., SINGH, S., WUTTKE, M., RABENSTEIN,

665 R., HABERSETZER, J., SCHAAL, S., FESEHA, M., CLEMENS, M., JACOBS, B. F.,
 666 CURRANO, E. D., JACOBS, L. L., SYLVESTERSEN, R. L., GABBOTT, S. E. and
 667 VINTHER, J. 2015 Chemical, experimental, and morphological evidence for
 668 diagenetically altered melanin in exceptionally preserved fossils. *Proceedings of the*
 669 *National Academy of Sciences*, **112**, 12592–12597.

670 COVINGTON, A. D. 1997. Modern tanning chemistry. *Chemical Society Reviews*, **26**, 111–126.

671 DAVIS, P. G. and BRIGGS, D. E. G. 1995. Fossilization of feathers. *Geology*, **23**, 783–786.

672 DING, R. and LEONOV, A. I. 1996. A kinetic model for sulfur accelerated vulcanization of a
 673 natural rubber compound. *Journal of Applied Polymer Science*, **61**, 455–463.

674 DREIER, F., BUCHS, A. and GÜLAÇAR, F. O. 1988. The degradation rates of some sterols in a
 675 heated sediment. *Geochimica et Cosmochimica Acta*, **52**, 1663–1666.

676 EDWARDS, N. P., BARDEN, H. E., VAN DONGEN, B. E., MANNING, P. L., LARSON, P.
 677 L., BERGMANN, U., SELLERS, W. I. and WOGELIUS, R. A. 2011. Infrared mapping
 678 resolves soft tissue preservation in 50 million year-old reptile skin. *Proceedings of the*
 679 *Royal Society B*, **278**, 3209–3218.

680 EGLINTON, G. and LOGAN, G. A. 1991. Molecular preservation. *Philosophical Transactions of*
 681 *the Royal Society of London. Series B, Biological Sciences*, **333**, 315–328.

682 EVERSLED, R. P. 1992. Chemical composition of a bog body adipocere. *Archaeometry*, **34**, 253–
 683 265.

684 GABBOTT, S. E., DONOGHUE, P. C., SANSOM, R. S., VINTHER, J., DOLOCAN, A. and
 685 PURNELL, M. A. 2016. Pigmented anatomy in Carboniferous cyclostomes and the
 686 evolution of the vertebrate eye. *Proceedings of the Royal Society B*, **283**, 20161151.

687 GLASS, K., ITO, S., WILBY, P. R., SOTA, T., NAKAMURA, A., BOWERS, C. R., VINTHER,
688 J., DUTTA, S., SUMMONS, R., BRIGGS, D. E. G. and WAKAMATSU, K. 2012. Direct
689 chemical evidence for eumelanin pigment from the Jurassic period. *Proceedings of the*
690 *National Academy of Sciences*, **109**, 10218–10223.

691 — ITO, S., WILBY, P. R., SOTA, T., NAKAMURA, A., BOWERS, C. R., MILLER, K. E.,
692 DUTTA, S., SUMMONS, R. E., BRIGGS, D. E. and WAKAMATSU, K. 2013. Impact of
693 diagenesis and maturation on the survival of eumelanin in the fossil record. *Organic*
694 *Geochemistry*, **64**, 29–37.

695 GRIMES, S. T., BROCK, F., RICKARD, D., DAVIES, K. L., EDWARDS, D., BRIGGS, D. E.
696 and PARKES, R. J. 2001. Understanding fossilization: experimental pyritization of plants.
697 *Geology*, **29**, 123–126.

698 GUPTA, N. S., BRIGGS, D. E. G. and PANCOST, R. D. 2006b. Molecular taphonomy of
699 graptolites. *Journal of the Geological Society*, **163**, 897–900.

700 — MICHELS, R., BRIGGS, D. E. G., EVERSLED, R. P. and PANCOST, R. D., 2006a. The
701 organic preservation of fossil arthropods: an experimental study. *Proceedings of the Royal*
702 *Society of London B: Biological Sciences*, **273**, 2777–2783.

703 — BRIGGS, D. E. G., COLLINSON, M. E., EVERSLED, R. P., MICHELS, R., JACK, K. S.
704 and PANCOST, R. D. 2007b. Evidence for the in situ polymerisation of labile aliphatic
705 organic compounds during the preservation of fossil leaves: implications for organic matter
706 preservation. *Organic Geochemistry*, **38**, 499–522.

707 — MICHELS, R., BRIGGS, D. E., COLLINSON, M. E., EVERSLED, R. P. and PANCOST, R.
708 D. 2007a. Experimental evidence for the formation of geomacromolecules from plant leaf
709 lipids. *Organic Geochemistry*, **38**, 28–36.

710 — TETLIE, O. E., BRIGGS, D. E. and PANCOST, R. D. 2007c. The fossilization of eurypterids:
711 a result of molecular transformation. *Palaios*, **22**, 439–447.

712 — BRIGGS, D. E. G., LANDMAN, N. H., TANABE, K. and SUMMONS, R. E. 2008.
713 Molecular structure of organic components in cephalopods: Evidence for oxidative cross
714 linking in fossil marine invertebrates. *Organic Geochemistry*, **39**, 1405–1414.

715 — CODY, G. D., TETLIE, O. E., BRIGGS, D. E. and SUMMONS, R. E. 2009. Rapid
716 incorporation of lipids into macromolecules during experimental decay of invertebrates:
717 Initiation of geopolymer formation. *Organic Geochemistry*, **40**, 589–594.

718 HELANDER, K. G. 1994. Kinetic studies of formaldehyde binding in tissue. *Biotechnic &*
719 *Histochemistry*, **69**, 177–179.

720 HIDALGO, F. J., ALAIZ, M. and ZAMORA, R. 1999. Effect of pH and temperature on
721 comparative nonenzymatic browning of proteins produced by oxidized lipids and
722 carbohydrates. *Journal of Agricultural and Food Chemistry*, **47**, 742–747.

723 HUEFFER, S., SCHROEDER, S., WAGNER, E., RAEDLER, T. and VILL, K. 2010. Formulation
724 for use in chrome or chrome-free tannage. *United States Patent*, 7,771,489.

725 KANAGY, J. R. and KRONSTADT, R. A. 1943. Iron as a tanning agent. *Journal of Research of*
726 *the National Bureau of Standards*, **31**, 279–289.

727 KIERNAN, J. A. 2000. Formaldehyde, formalin, paraformaldehyde and glutaraldehyde: what they
728 are and what they do. *Microscopy today*, **1**, 8–12.

729 KOOPMANS, M. P., DE LEEUW, J. W., LEWAN, M. D. and SINNINGHE DAMSTE, J. S.
730 1996. Impact of dia- and catagenesis on sulphur and oxygen sequestration of biomarkers
731 as revealed by artificial maturation of an immature sedimentary rock. *Organic*
732 *Geochemistry*, **25**, 391–426.

733 LAMBERT, J. B. and FRYE, J. S. 1982. Carbon functionalities in amber. *Science*, **217**, 55–57.

734 LANGENHEIM, J. H. 1969. Amber: a botanical inquiry. *Science*, **163**, 1157–1169.

735 — 1990. Plant resins. *American Scientist*, **78**, 16–24.

736 LEWAN, M. D. 1983. Effects of thermal maturation on stable organic carbon isotopes as
737 determined by hydrous pyrolysis of Woodford Shale. *Geochimica et Cosmochimica*
738 *Acta*, **47**, 1471–1479.

739 — BJORØY, M. and DOLCATER, D. L. 1986. Effects of thermal maturation on steroid
740 hydrocarbons as determined by hydrous pyrolysis of Phosphoria Retort Shale. *Geochimica*
741 *et Cosmochimica Acta*, **50**, 1977–1987.

742 LI, Q., CLARKE, J. A., GAO, K. Q., ZHOU, C. F., MENG, Q., LI, D., D’ALBA, L. and
743 SHAWKEY, M. D. 2014. Melanosome evolution indicates a key physiological shift within
744 feathered dinosaurs. *Nature*, **507**, 350–353.

745 LINDGREN, J., UVDAL, P., SJÖVALL, P., NILSSON, D. E., ENGDAHL, A., SCHULTZ, B. P.
746 and THIEL, V. 2012. Molecular preservation of the pigment melanin in fossil
747 melanosomes. *Nature Communications*, **3**, 824.

748 — SJÖVALL, P., CARNEY, R. M., UVDAL, P., GREN, J. A., DYKE, G.,
749 SCHULTZ, B. P., SHAWKEY, M. D., BARNES, K. R. and POLCYN, M. J. 2014. Skin
750 pigmentation provides evidence of convergent melanism in extinct marine reptiles. *Nature*,
751 **506**, 484–488.

752 — MOYER, A., SCHWEITZER, M. H., SJÖVALL, P., UVDAL, P., NILSSON, D. E.,
753 HEIMDAL, J., ENGDAHL, A., GREN, J. A., SCHULTZ, B. P. and KEAR, B. P. 2015.
754 Interpreting melanin-based coloration through deep time: a critical review. *Proceedings*
755 *of the Royal Society B*, **282**, 20150614.

756 — KURIYAMA, T., MADSEN, H., SJÖVALL, P., ZHENG, W., UVDAL, P., ENGDAHL, A.,
 757 MOYER, A. E., GREN, J. A., KAMEZAKI, N. and UENO, S. 2017. Biochemistry and
 758 adaptive colouration of an exceptionally preserved juvenile fossil sea turtle. *Scientific*
 759 *Reports*, **7**, 13324.

760 LINGHAM-SOLIAR, T., FEDUCCIA, A. and WANG, X. 2007. A new Chinese specimen
 761 indicates that ‘protofeathers’ in the Early Cretaceous theropod dinosaur *Sinosauropteryx*
 762 are degraded collagen fibres. *Proceedings of the Royal Society B*, **274**, 1823–1829.

763 MACKENZIE, A. S., BRASSELL, S. C., EGLINTON, G. and MAXWELL, J. R., 1982. Chemical
 764 fossils: the geological fate of steroids. *Science*, **217**, 491–504.

765 MARTINS, S. I., JONGEN, W. M. and VAN BOEKEL, M. A. 2000. A review of Maillard reaction
 766 in food and implications to kinetic modelling. *Trends in Food Science & Technology*, **11**,
 767 364–373.

768 MAYR, G., PITTMAN, M., SAITTA, E., KAYE, T. G. and VINTHER, J. 2016. Structure and
 769 homology of Psittacosaurus tail bristles. *Palaeontology*, **59**, 793–802.

770 MCCOY, V. E., YOUNG, R. T. and BRIGGS, D. E. G. 2015. Factors controlling exceptional
 771 preservation in concretions. *Palaios*, **30**, 272–280.

772 MCNAMARA, M. E., ORR, P. J., KEARNS, S. L., ALCALÁ, L., ANADÓN, P. and
 773 PEÑALVER-MOLLÁ, E. 2006. High-fidelity organic preservation of bone marrow in ca.
 774 10 Ma amphibians. *Geology*, **34**, 641–644.

775 — ORR, P. J., KEARNS, S. L., ALCALÁ, L., ANADÓN, P. and PEÑALVER-MOLLÁ, E.
 776 2010. Organic preservation of fossil musculature with ultracellular detail. *Proceedings of*
 777 *the Royal Society of London B: Biological Sciences*, **277**, 423–427.

- 778 — BRIGGS, D. E. G., ORR, P. J., FIELD, D. J. and WANG, Z. 2013. Experimental maturation
779 of feathers: implications for reconstructions of fossil feather colour. *Biology Letters*, **9**,
780 20130184.
- 781 — DONGEN, B. E., LOCKYER, N. P., BULL, I. D. and ORR, P. J. 2016. Fossilization of
782 melanosomes via sulfurization. *Palaeontology*, **59**, 337–350.
- 783 — BRIGGS, D. E. G., ORR, P. J., FIELD, D. J. and WANG, Z. 2017. Correction to
784 ‘Experimentalmaturation of feathers: implications for reconstructions of fossil feather
785 colour’. *Biology Letters*, **13**, 20170128.
- 786 MELENDEZ, I., GRICE, K. and SCHWARK, L. 2013. Exceptional preservation of Palaeozoic
787 steroids in a diagenetic continuum. *Scientific Reports*, **3**, 2768.
- 788 MILLS, J.S., WHITE, R. and GOUGH, L.J. 1984. The chemical composition of Baltic
789 amber. *Chemical Geology*, **47**, 15–39.
- 790 MONTHIOUX, M., LANDAIS, P. and MONIN, J. C. 1985. Comparison between natural and
791 artificial maturation series of humic coals from the Mahakam delta, Indonesia. *Organic*
792 *Geochemistry*, **8**, 275–292.
- 793 MOYER, A. E., ZHENG, W., JOHNSON, E. A., LAMANNA, M. C., LI, D. Q., LACOVARA,
794 K. J. and SCHWEITZER, M. H. 2014. Melanosomes or microbes: testing an alternative
795 hypothesis for the origin of microbodies in fossil feathers. *Scientific Reports*, **4**, 4233.
- 796 — ZHENG, W. and SCHWEITZER, M. H. 2016a. Keratin durability has implications for the
797 fossil record: results from a 10 year feather degradation experiment. *PLoS One*, **11**,
798 e0157699.
- 799 — ZHENG, W. and SCHWEITZER, M. H. 2016b. Microscopic and immunohistochemical

800 analyses of the claw of the nesting dinosaur, *Citipati osmolskae*. *Proceedings of the Royal*
801 *Society B*, **283**, 20161997.

802 PAINTER, T. J. 1991. Lindow man, tollund man and other peat-bog bodies: the preservative and
803 antimicrobial action of Sphagnum, a reactive glycuronoglycan with tanning and
804 sequestering properties. *Carbohydrate Polymers*, **15**, 123–142.

805 PAN, Y., ZHENG, W., MOYER, A. E., O’CONNOR, J. K., WANG, M., ZHENG, X., WANG,
806 X., SCHROETER, E. R., ZHOU, Z. and SCHWEITZER, M. H. 2016. Molecular evidence
807 of keratin and melanosomes in feathers of the Early Cretaceous bird *Eoconfuciusornis*.
808 *Proceedings of the National Academy of Sciences*, **113**, E7900–E7907.

809 PARRY, L. A., WILSON, P., SYKES, D., EDGECOMBE, G. D. and VINTHER, J. 2015. A new
810 fireworm (Amphinomidae) from the Cretaceous of Lebanon identified from three-
811 dimensionally preserved myoanatomy. *BMC evolutionary biology*, **15**, 256.

812 — SMITHWICK, F., NORDÉN, K. K., SAIITA, E. T., LOZANO-FERNANDEZ, J., TANNER,
813 A. R., CARON, J. B., EDGECOMBE, G. D., BRIGGS, D. E. and VINTHER, J. 2018.
814 Soft-Bodied Fossils Are Not Simply Rotten Carcasses—Toward a Holistic Understanding
815 of Exceptional Fossil Preservation. *BioEssays*, **40**, 1700167.

816 PAUTARD, F. G. E. 1963. Mineralization of keratin and its comparison with the enamel
817 matrix. *Nature*, **199**, 531–535.

818 — 1964. Calcification of keratin. *Progress in the Biological Sciences in Relation to*
819 *Dermatology*, **2**, 227.

820 PETERS, K. E., ROHRBACK, B. G. and KAPLAN, I. R. 1981. Carbon and hydrogen stable
821 isotope variations in kerogen during laboratory-simulated thermal maturation. *American*
822 *Association of Petroleum Geologists Bulletin*, **65**, 501–508.

823 PIKE, E. M. 1993. Amber taphonomy and collecting bias. *Palaios*, **8**, 411–419.

824 PLAPPER, J., SCHUMANN, K., SCHWUGER, M. J., SMOLKA, H.
825 G., ARNDT, E., RUSCHEINSKY, E. and JANSEN, H. 1981, Use of water-insoluble
826 aluminosilicates in tanning process for the production of leather. *United States Patent*,
827 4,272,242.

828 PLET, C., GRICE, K., PAGÈS, A., VERRALL, M., COOLEN, M. J., RUEBSAM, W.,
829 RICKARD, W. D. and SCHWARK, L. 2017. Palaeobiology of red and white blood cell-
830 like structures, collagen and cholesterol in an ichthyosaur bone. *Scientific reports*, **7**,
831 13776.

832 PURNELL, M. A., DONOGHUE, P. J., GABBOTT, S. E., MCNAMARA, M. E., MURDOCK,
833 D. J. and SANSOM, R. S. 2018. Experimental analysis of soft-tissue fossilization: opening
834 the black box. *Palaeontology*, **61**, 317–323.

835 RAGAZZI, E., ROGHI, G., GIARETTA, A. and GIANOLLA, P. 2003. Classification of amber
836 based on thermal analysis. *Thermochimica Acta*, **404**, 43–54.

837 RHEAD, M. M., EGLINTON, G. and DRAFFAN, G. H. 1971 Hydrocarbons produced by the
838 thermal alteration of cholesterol under conditions simulating the maturation of sediments.
839 *Chemical Geology*, **8**, 277–297.

840 RULLKÖTTER, J. and MARZI, R. 1988. Natural and artificial maturation of biological markers
841 in a Toarcian shale from northern Germany. *Organic Geochemistry*, **13**, 639–645.

842 SAITTA, E. T., ROGERS, C. S., BROOKER, R. A., ABBOTT, G. D., KUMAR, S., O'REILLY,
843 S. S., DONOHUE, P., DUTTA, S., SUMMONS, R. E. and VINTHER, J. 2017a Low
844 fossilization potential of keratin protein revealed by experimental taphonomy.
845 *Palaeontology*, **60**, 547–556.

846 — ROGERS, C. S., BROOKER, R. A. and VINTHER, J. 2017b. Experimental taphonomy of
847 keratin: a structural analysis of early taphonomic changes. *Palaios*, **32**, 647–657.

848 — KAYE T. G. and VINTHER, J. 2018. Data from: Sediment-encased maturation: a method for
849 simulating diagenesis in organic fossil preservation. Dryad Digital Repository.
850 <http://datadryad.org/review?doi=doi:10.5061/dryad.0t67n>

851 SALLAN, L. C. and COATES, M. I. 2014. The long-rostrumed elasmobranch *Bandringa* Zangerl,
852 1969, and taphonomy within a Carboniferous shark nursery. *Journal of Vertebrate*
853 *Paleontology*, **34**, 22–33.

854 SANSOM, R. S., GABBOTT, S. E. and PURNELL, M. A. 2010a. Decay of vertebrate characters
855 in hagfish and lamprey (Cyclostomata) and the implications for the vertebrate fossil
856 record. *Proceedings of the Royal Society of London B: Biological Sciences*, **278**, 1150–
857 1157.

858 — GABBOTT, S. E. and PURNELL, M. A. 2010b. Non-random decay of chordate
859 characters causes bias in fossil interpretation. *Nature*, **463**, 797–800.

860 — GABBOTT, S. E. and PURNELL, M.A. 2013. Atlas of vertebrate decay: a visual and
861 taphonomic guide to fossil interpretation. *Palaeontology*, **56**, 457–474.

862 SCHIFFBAUER, J. D., WALLACE, A. F., HUNTER, J. L., KOWALEWSKI, M., BODNAR, R.
863 J. and XIAO, S. 2012 Thermally-induced structural and chemical alteration of organic-
864 walled microfossils: an experimental approach to understanding fossil preservation in
865 metasediments. *Geobiology*, **10**, 402–423.

866 SCHIMMELMANN, A., SESSIONS, A. L. and MASTALERZ, M. 2006. Hydrogen isotopic
867 (D/H) composition of organic matter during diagenesis and thermal maturation. *Annual*
868 *Review of Earth and Planetary Sciences*, **34**, 501–533.

869 SCHWEITZER, M. H., WATT, J. A., AVCI, R., FORSTER, C. A., KRAUSE, D. W., KNAPP,
 870 L., ROGERS, R. R., BEECH, I. and MARSHALL, M. 1999. Keratin immunoreactivity in
 871 the Late Cretaceous bird *Rahonavis ostromi*. *Journal of Vertebrate Paleontology*, **19**, 712–
 872 722.

873 — LINDGREN, J. and MOYER, A. E. 2015 Melanosomes and ancient
 874 coloration re-examined: A response to Vinther. *BioEssays*, **37**, 1174–1183.

875 SHIN, Y., WANG, C. and EXARHOS, G. J. 2005. Synthesis of SiC ceramics by the
 876 carbothermal reduction of mineralized wood with silica. *Advanced Materials*, **17**, 73–77.

877 SINNINGHE DAMSTE, J. S. and DE LEEUW, J. W. 1990. Analysis, structure and geochemical
 878 significance of organically-bound sulphur in the geosphere: state of the art and future
 879 research. *Organic Geochemistry*, **16**, 1077–1101.

880 — EGLINTON, T.I., DE LEEUW, J.W. and SCHENCK, P.A. 1989. Organic sulphur in
 881 macromolecular sedimentary organic matter: I. Structure and origin of sulphur-containing
 882 moieties in kerogen, asphaltenes and coal as revealed by flash pyrolysis. *Geochimica et*
 883 *Cosmochimica Acta*, **53**, 873–889.

884 SMITH, N. A., CHIAPPE, L. M., CLARKE, J. A., EDWARDS, S. V., NESBITT, S. J., NORELL,
 885 M. A., STIDHAM, T. A., TURNER, A., VAN TUINEN, M., VINTHER, J. and XU, X.
 886 2015. Rhetoric vs. reality: a commentary on “Bird Origins Anew” by A. Feduccia. *The*
 887 *Auk*, **132**, 467–480.

888 SMITHWICK, F. M., MAYR, G., SAITTA, E. T., BENTON, M. J. and VINTHER, J. 2017.
 889 On the purported presence of fossilized collagen fibres in an ichthyosaur and a theropod
 890 dinosaur. *Palaeontology*, **60**, 409–422.

891 STANKIEWICZ, B. A., HUTCHINS, J. C., THOMSON, R., BRIGGS, D. E. and EVERSHED,
892 R. P. 1997. Assessment of bog-body tissue preservation by pyrolysis-gas
893 chromatography/mass spectrometry. *Rapid Communications in Mass Spectrometry*, **11**,
894 1884–1890.

895 — BRIGGS, D. E. G., MICHELS, R., COLLINSON, M. E., FLANNERY, M. B. and
896 EVERSHED, R. P. 2000. Alternative origin of aliphatic polymer in kerogen. *Geology*, **28**,
897 559–562.

898 STEVENS, M. M., HONERKAMP-SMITH, A. R. and KELLER, S. L., 2010. Solubility
899 limits of cholesterol, lanosterol, ergosterol, stigmasterol, and β -sitosterol in
900 electroformed lipid vesicles. *Soft Matter*, **6**, 5882–5890.

901 TEERMAN, S. C. and HWANG, R. J. 1991. Evaluation of the liquid hydrocarbon potential of coal
902 by artificial maturation techniques. *Organic Geochemistry*, **17**, 749–764.

903 TOWNSON, J. L., LIN, Y. S., CHOU, S. S., AWAD, Y. H., COKER, E. N., BRINKER, C. J. and
904 KAEHR, B. 2014. Synthetic fossilization of soft biological tissues and their shape-
905 preserving transformation into silica or electron-conductive replicas. *Nature*
906 *Communications*, **5**, 5665.

907 URIBARRI, J., WOODRUFF, S., GOODMAN, S., CAI, W., CHEN, X., PYZIK, R., YONG, A.,
908 STRIKER, G. E. and VLASSARA, H. 2010. Advanced glycation end products in foods
909 and a practical guide to their reduction in the diet. *Journal of the American Dietetic*
910 *Association*, **110**, 911–916.

911 VERSTEEGH, G. J., BLOKKER, P., WOOD, G. D., COLLINSON, M. E., DAMSTÉ, J. S. S. and
912 DE LEEUW, J. W. 2004. An example of oxidative polymerization of unsaturated fatty

913 acids as a preservation pathway for dinoflagellate organic matter. *Organic*
914 *Geochemistry*, **35**, 1129–1139.

915 VILLANUEVA-GARCÍA, M., MARTÍNEZ-RICHA, A. and ROBLES, J. 2005. Assignment of
916 vibrational spectra of labdatriene derivatives and ambers: a combined experimental and
917 density functional theoretical study. *Arkivoc*, **6**, 449–458.

918 VINTHER, J., BRIGGS, D. E. G., PRUM, R. O., and SARANATHAN, V. 2008. The colour of
919 fossil feathers. *Biology Letters*, **4**, 522–525.

920 — 2015. A guide to the field of palaeo colour. *BioEssays*, **37**, 643–656.

921 — 2016. Fossil melanosomes or bacteria? A wealth of findings favours
922 melanosomes. *BioEssays*, **38**(3), pp.220-225.

923 — NICHOLLS, R., LAUTENSCHLAGER, S., PITTMAN, M., KAYE, T. G., RAYFIELD, E.,
924 MAYR, G. and CUTHILL, I. C. 2016. 3D camouflage in an ornithischian dinosaur.
925 *Current Biology*, **26**, 2456–2462.

926 WARDELL, J. M., HILL, C. M. and SOUZA, S. 1981. Effect of pasteurization and of freezing
927 and thawing human milk on its triglyceride content. *Acta Paediatrica*, **70**, 467–471.

928 WILSON, L. A. and BUTTERFIELD, N. J. 2014. Sediment effects on the preservation of Burgess
929 Shale-type compression fossils. *Palaios*, **29**, 145–154.

930 WILSON, P., PARRY, L. A., VINTHER, J. and EDGECOMBE, G. D. 2016 Unveiling biases in
931 soft-tissue phosphatization: extensive preservation of musculature in the Cretaceous
932 (Cenomanian) polychaete *Rollinschaeta myoplana* (Annelida: Amphinomidae).
933 *Palaeontology*, **59**, 463–479.

934 WUTTKE, M. 1983. Weichteil-erhaltung durch lithifizierte mikroorganismen bei mittel-eozänen

vertebraten aus den ölschiefern der ‘Grube Messel’ bei Darmstadt. *Senckenbergiana
lethaea*, **64**, 509–527.

XING, L., O’CONNOR, J. K., MCKELLAR, R. C., CHIAPPE, L. M., TSENG, K., LI, G. and
BAI, M. 2017. A mid-Cretaceous enantiornithine (Aves) hatchling preserved in Burmese
amber with unusual plumage. *Gondwana Research*, **49**, 264–277.

ZHANG, F., KEARNS, S. L., ORR, P. J., BENTON, M. J., ZHOU, Z., JOHNSON, D., XU, X.
and WANG, X. 2010. Fossilized melanosomes and the colour of Cretaceous dinosaurs and
birds. *Nature*, **463**, 1075–1078.

ZUIDAM, N. J. and CROMMELIN, D. J. A. 1995. Chemical hydrolysis of phospholipids. *Journal
of Pharmaceutical Sciences*, **84**, 1113–1119.

FIGURE CAPTIONS

Fig. 1. Experimental setup. A, compaction phase. B, maturation phase.

Fig. 2. Experimental lizards compared to fossils. Lizard foot, A, before burial and, B–C,
counterpart after maturation showing browned bones, organic stains, and scale imprints. Lizard
hand, D, before burial and, E, after maturation showing scale imprints, some organic staining, and
discoloured bones. Lizard head, F, before burial and, G, after maturation revealing browned,
crushed bones and organic stain. H–I, SEM images of exposed oblate melanosomes from the
matured lizard foot in A–C. J, Fossil oblate melanosomes from a pipidid frog epidermis, Early
Miocene, Mush Valley, Ethiopia (MU 32-2A/B). Fossil lizards, K, Eocene, Green River Formation
(BHI-102B, image courtesy of Nicholas Edwards) and, L, Early Cretaceous, Yixian Formation

(IVPP V14715, image courtesy of Wang Yuan). Scale bars represent 5 mm (A–B, D–G, K); 1 mm (C), 2 μm (H), 1 μm (I–J), 1 cm (L). Frog silhouette in J was obtained through phylopic.org (credit: Sarah Werning, Creative Commons Attribution 3.0 Unported (CC BY 3.0), <https://creativecommons.org/licenses/by/3.0/legalcode>).

Fig. 3. Experimental feathers compared to fossils. Black and white feather with indications of the side of the feather from which the SEM images are derived, A, before burial, B, counterpart after treatment (in this case, involving a second compaction step after maturation) showing preferential preservation of more heavily melanised regions and sediment infilling of rachis, C, barb imprints in sediment from the pigmented region after treatment with textures resulting from regions of melanosomes, D, barb and barbule imprints in sediment from the predominantly white region lacking melanosomes after treatment as evidenced by the scarcity of textures from melanosomes in void spaces, E, void space in pigmented region left behind due to organic loss from a barb after treatment with textures resulting from regions of melanosomes, and F–G, oblong melanosomes from pigmented region after treatment. Darkly pigmented feather, H, after treatment, I, barb and barbule imprints in sediment after treatment, J, SEM image of exposed oblong melanosomes after treatment. K, Fossil oblong melanosomes in the organic stains of, L, fossil feather, Early Cretaceous, Crato Formation (LEIUG 115562), showing colour patterning and preferential preservation correlated to melanin pigmentation. Scale bars represent 5 mm (A–B, H); 200 μm (C); 100 μm (D); 10 μm (E); 2 μm (F–G, J); 30 μm (I); 1 μm (K); 1 mm (L).

Fig. 4. Non-sediment-encased matured (250 °C/250 bars/24 hours) dark feathers as described by Saitta *et al.* (2017a, 2017b) where maturation was performed inside a small, sealed, noble metal

capsule. A, fluid produced from maturation observed adhering to the inside wall of a glass vial after extruding from the opened maturation capsule. B–D, SEM images of, B, fluid extruding onto the surface of the opened maturation capsule, C, the topography of the fluid (likely influenced by surface charging) and, D, the finer ultrastructure of the fluid. No melanosomes are visible. Scale bars represent 2 mm (A); 1 mm (B); 200 μm (C); 10 μm (D).

A

B

988

989

A

B

C

D

990