Chang, Y., Chen, Z., Zhou, J., Luo, Z., He, Z., Wu, G., Ashfold, M. N. R., Yuan, K., & Yang, X. (2019). Striking Isotopologue-Dependent Photodissociation Dynamics of Water Molecules: The Signature of an Accidental Resonance. *Journal of Physical Chemistry Letters*, 10, 4209-4214. [15]. <https://doi.org/10.1021/acs.jpcllett.9b01710>

Peer reviewed version

Link to published version (if available):
[10.1021/acs.jpcllett.9b01710](https://doi.org/10.1021/acs.jpcllett.9b01710)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via American Chemical Society at <https://pubs.acs.org/doi/10.1021/acs.jpcllett.9b01710> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Striking Isotopologue Dependent Photodissociation Dynamics of Water Molecules: The Signature of an Accidental Resonance

Yao Chang^{a#}, Zhichao Chen^{a#}, Jiami Zhou^a, Zijie Luo^a, Zhigang He^a, Guorong Wu^a,
Michael N.R. Ashfold^b, Kaijun Yuan^{a*}, Xueming Yang^{a*}

a. *State Key Laboratory of Molecular Reaction Dynamics, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, 457 Zhongshan Road, Dalian, 116023, China*

b. *School of Chemistry, University of Bristol, Bristol, BS8 1TS, U.K.*

These authors have equal contributions

*) To whom correspondence should be addressed. Email address: kjyuan@dicp.ac.cn, xmyang@dicp.ac.cn

Abstract

Investigations of the photofragmentation patterns of both light and heavy water at the state-to-state level are a pre-requisite for any thorough understanding of chemical processing and isotope heterogeneity in the interstellar medium (ISM). Here we reveal dynamical features of the dissociation of water molecules following excitation to the $\tilde{C}(010)$ state using a tunable vacuum ultraviolet source in combination with the high resolution H(D)-atom Rydberg tagging time-of-flight technique. The action spectra for forming H(D) atoms and the OH(OD) product state distributions resulting from excitation to the $\tilde{C}(010)$ states of H₂O and D₂O both show striking differences, which are attributable to the effects of an isotopologue-specific accidental resonance. Such accidental resonance induced state mixing may contribute to the D/H isotope heterogeneity in the Solar System. The present study provides an excellent example of competitive state-to-state non-adiabatic decay pathways involving at least five electronic states.

Introduction

Water, ubiquitous in the universe, absorbs light at all wavelengths (λ) in the vacuum ultraviolet (VUV) region below 190 nm,¹ resulting in fragmentation leading to a hydrogen (H) atom and a hydroxyl (OH) radical and/or, at shorter wavelengths, the three constituent atoms. The single bond fission process is an important source of OH radicals in the interstellar medium (ISM).^{2,3} An in-depth understanding of the OH product quantum state population distributions from the photodissociation of water is thus an essential prerequisite for interstellar related chemistry modelling. Many distinctive features of the OH product state distributions arising from water photodissociation have already been identified, and explained as exemplars of a rich gamut of non-adiabatic dynamics, *e.g.*, passage through conical intersections (CIs),^{4,5} Coriolis couplings,^{6,7} vibronic couplings,⁸ etc. Here we demonstrate another novel feature of the OH product state distribution arising in the VUV photodissociation of water, which is isotope-specific and can be attributed to an accidental vibronic resonance. Accidental resonances are often observed in molecular spectra – as perturbations in line positions, intensities or widths.⁹ However, the effects of accidental resonances in the parent absorption upon the subsequent fragmentation dynamics have rarely been recognized hitherto.

The photodissociation of H₂O has been the subject of extensive experimental and theoretical investigations over the past decades, so much so that H₂O (and its various isotopomers) is now viewed as a benchmark system for illustrating and understanding

non-adiabatic coupling pathways between potential energy surfaces (PESs). Excitation of H₂O at $\lambda \sim 160$ nm populates the lowest excited electronic ($\tilde{A}^1B_1$) state; the subsequent direct dissociation from this state yields an H atom plus a ground state OH($X^2\Pi$) radical with little internal excitation.¹⁰⁻¹⁶ Absorption to the second excited singlet ($\tilde{B}^1A_1$) state maximizes at $\lambda \sim 128$ nm. This state displays a minor direct dissociation channel to H + OH($A^2\Sigma^+$) products, and major non-adiabatic pathways via two CIs with the ground ($\tilde{X}^1A_1$) state PES at linear HOH and HHO configurations that favor formation of highly rotationally excited OH(X , $v=0$) products.¹⁷⁻²⁹ The next two excited states identified in absorption have predominant Rydberg character, and predissociate by non-adiabatic coupling to those lower (predominantly valence) states. The $\tilde{C}^1B_1$ state shows resolvable rotational structure at $\lambda \sim 124$ nm,³⁰⁻³³ and previous studies have identified dramatic variations in the OH product state distributions and angular distributions that arise following excitation to different rotational levels of this state.³⁴ Dissociation from $\tilde{C}$ state rotational levels with $K_a' = 0$ (where K_a describes the projection of the parent rotational angular momentum (J) onto the a inertial axis) occurs exclusively by electronic coupling to the $\tilde{A}$ state and yields highly vibrationally excited OH(X) products with vibrational quantum number $v'' \leq 13$ (henceforth termed “*high v*” products). A rival decay pathway opens for H₂O ($\tilde{C}$) molecules in levels with $K_a' > 0$. This relies on Coriolis-type (*i.e.* rotationally induced) coupling to the $\tilde{B}$ state PES, and results in both ground (X) and electronically excited (A) state OH fragments with very high rotational but minimal vibrational excitation (henceforth termed “*high N*” products).

Replacing H by deuterium (D) often leads to significant isotopic effects in processes that are dynamically controlled, particularly in cases where non-adiabatic dynamics are involved. Indeed, investigations of the photodissociation of heavy water (D_2O) following excitation to the same excited states have revealed qualitative variations from the H_2O data but, in all cases studied to date, the same dissociation mechanisms appear to prevail for both H_2O and D_2O .³⁵⁻³⁷ However, a recent ultrafast time-resolved photoelectron spectroscopy study revealed quantitatively different predissociation behaviors for the $\tilde{\text{C}}(010)$ states (i.e. the states carrying one quantum of bending vibrational excitation, ν_2') of H_2O and D_2O .³⁸ The present article unravels the state-to-state dynamics associated with this unusual isotopic effect, using tunable VUV pump and probe laser sources and the high resolution H/D atom Rydberg tagging technique. The distinctive OH and OD quantum state distributions determined in this work provide further exquisitely detailed illustrations of the sensitivity of the various possible non-adiabatic decay pathways to details of the excited state PESs and the inter-state coupling matrix elements.

The first experiment involved scanning the VUV photolysis wavelength while recording the total H (D) atom yield (see Experimental Methods in *supporting information*), thereby obtaining action spectra for forming H (D) atoms from VUV photodissociation of H_2O (D_2O). **Figure 1A** shows the action spectrum from photolysis of H_2O in the wavelength range 121.7-122.4 nm, which spans transitions to the $\tilde{\text{C}}(010)$ state and to the origin (000) level of the $\tilde{\text{D}}^1\text{A}_1$ state. The broad peak centered at $\lambda \sim 122.1$ nm and extending across this entire spectral window is

attributable to the $\tilde{D} \leftarrow \tilde{X}$ origin band; the width of this feature reflects the short predissociation lifetime of the $\tilde{D}(000)$ state.³⁸ Note that this spectrum is distorted by an obvious dip in the VUV tuning curve at $\lambda \sim 122.09$ nm, which is also apparent in the action spectra for forming D atoms from both D₂O and CD₄ photolysis shown in **Figure 1C**. Analogy with the rotational fine structures of the $\tilde{C} \leftarrow \tilde{X}$ origin band allows the partially resolved peaks at $\lambda \sim 121.85$ nm to be assigned to the three strongest rotational lines in the jet-cooled $\tilde{C}(010) \leftarrow \tilde{X}(000)$ transition – as shown in **Figure 1B**. This analysis provides an experimental estimate of the bending vibrational wavenumber for the $\tilde{C}$ state of H₂O, $\nu_2' = 1402 \pm 3$ cm⁻¹, which is in good accord with the values determined in previous multiphoton spectroscopy studies³⁰ and theoretical predictions³⁹.

In contrast, the action spectrum from the photodissociation of D₂O shows well resolved rotational structure associated with the $\tilde{C}(010) \leftarrow \tilde{X}(000)$ transition (**Figure 1C**). The striking similarity between this spectrum and that of the $\tilde{C} \leftarrow \tilde{X}$ origin band (**Figure 1D**) allows straightforward assignment of the rotational fine structure in the $\tilde{C}(010) \leftarrow \tilde{X}(000)$ spectrum and yields a value of $\nu_2' = 1040 \pm 2$ cm⁻¹ for the $\tilde{C}$ state of D₂O. The widths of individual lines in the $\tilde{C}(010) \leftarrow \tilde{X}(000)$ action spectra (**Figures. 1B** and **1D**) are ~ 15 cm⁻¹ and 1.2 - 1.7 cm⁻¹ for H₂O and D₂O, respectively. These widths correspond to predissociation lifetimes of ~ 330 fs for H₂O and 3.0 - 4.2 ps for D₂O, in good accord with the time resolved measurements of He *et al.*³⁸ and with the conclusions from earlier analyses of the respective resonance enhanced multiphoton ionization spectra of H₂O and D₂O³⁰, wherein the striking

isotope effect was attributed to the effects of an accidental resonance between the $\tilde{C}$ (010) and $\tilde{D}$ (000) vibronic levels of H₂O. The $\tilde{C}$ (010) and $\tilde{D}$ (000) states of D₂O, in contrast, are ~ 270 cm⁻¹ out of resonance³⁰ and the mixing between the two states is thus negligible.

We have measured TOF spectra of the H (D) atom products formed with the VUV photolysis laser tuned to each of the more intense rotational lines in the $\tilde{C}$ (010) $\leftarrow \tilde{X}$ (000) bands of H₂O and D₂O. Knowing the distance travelled from the photodissociation region to the detector and the fragment masses, such TOF spectra can be converted into spectra of the total kinetic energy release (TKER). The TKER distributions obtained with the polarization vector $\mathbf{\epsilon}_{\text{phot}}$ aligned, respectively, parallel and perpendicular to the detection axis can then be used to construct 3-dimensional (3-D) flux diagrams of the H + OH (D + OD) fragments. **Figure 2** shows such 3-D flux diagrams for the products formed following excitation to the same $\tilde{C}$ (010) $\leftarrow \tilde{X}$ (000), $1_{10} \leftarrow 0_{00}$ transitions of H₂O and D₂O. Each diagram shows two groups of features: an inner group, associated with formation of levels of electronically excited A state of OH (OD) fragments, and an outer group associated with ground X state OH (OD) products. The obvious differences in the 3-D distributions displayed in **Figure 2** clearly illustrate that H₂O and D₂O molecules exhibit different predissociation dynamics, even when excited via the same transition to the same J_{KaKc} level of the respective $\tilde{C}$ (010) state.

Figure 3 displays TKER distributions from D₂O photolysis at 122.282, 122.235 and 121.954 nm with $\mathbf{\epsilon}_{\text{phot}}$ aligned parallel and perpendicular to the detection axis. These wavelengths excite, respectively, the $1_{01} \leftarrow 1_{11}$ and $1_{10} \leftarrow 0_{00}$ transitions of the $\tilde{C}$

(010) $\leftarrow \tilde{X}$ (000) band, and near the center of the $\tilde{D} \leftarrow \tilde{X}$ origin band. All of these TKER distributions show progressions of sharp peaks. Given that the total energy and linear momentum must be conserved in the photodissociation process, we can write

$$h\nu + E_{\text{int}}(\text{D}_2\text{O}) - D_0(\text{D-OD}) = E_T(\text{D} + \text{OD}) + E_{\text{int}}(\text{OD}), \quad (1)$$

where the internal energy (E_{int}) distribution of the OD fragment can be determined from the TKER (E_T) distribution, $h\nu$ is the photolysis photon energy, h is Planck's constant and $D_0(\text{D-OD})$ is the bond dissociation energy. All of the sharp peaks in **Figure 3** can be assigned to population of specific rovibrational levels of the A and X states of OD.

The excited level populated at the first of these wavelengths has $K_a' = 0$ and, as **Figure 3A** shows, the TKER spectrum is invariant to the alignment of ϵ_{phot} (i.e. the recoil velocity distribution is isotropic) and the fine structure indicates that the OD products are formed in their ground (X) state, in a very wide spread of vibrational states (up to $v'' = 18$), and with little rotational excitation. Both the energy disposal and the isotropy of the product recoil are very similar to those found when exciting the same rotational transition within the $\tilde{C}(000) \leftarrow \tilde{X}(000)$ origin band,³⁷ but totally different to that observed when exciting to the $\tilde{D}(000)$ state – which yields ‘high N ’ OD products (**Figure 3C**). The TKER spectra obtained following excitation of the $1_{10} \leftarrow 0_{00}$ transition (i.e. populating a $\tilde{C}(010)$ state level with $K_a' = 1$) are different again. As **Fig. 3B** shows, the spectrum recorded in the perpendicular direction is dominated by ‘high v ’ OD(X) products, whereas ‘high N ’ OD (A) and OD(X) products dominate the TKER spectrum recorded in the parallel direction. Again, these very

obvious differences are reminiscent of those reported previously when exciting the same rotational transition in the $\tilde{C} \leftarrow \tilde{X}$ origin band of D₂O (and H₂O).³⁴ These parallels suggest that the predissociation of $\tilde{C}(010)$ and $\tilde{C}(000)$ state D₂O molecules follow essentially the same mechanisms – namely a Coriolis-induced pathway to the $\tilde{B}$ state PES and an electronic coupling to the $\tilde{A}$ state PES as summarized in **scheme 1**. The former yields ‘*high N*’ OD(X) and OD(A) products whereas the latter yields ‘*high v*’ OD(X) products. The products of these two pathways show opposite recoil anisotropies and, for $\tilde{C}(010)$ levels with $K_a' = 1$, the respective signals suggest that the two processes occur with comparable efficiencies.

Scheme 1

The situation with H₂O is very different. **Figure 4** shows the TKER distributions obtained when exciting each of the three $\tilde{C}(010) \leftarrow \tilde{X}(000)$ transitions identified in **Figure 1B**. TKER distributions were also measured with the photolysis wavelength tuned to the valley between the peaks at 121.851 and 121.814 nm in the action spectrum, and to 122.117 nm (where absorption is ascribed exclusively to the $\tilde{D} \leftarrow \tilde{X}$ origin transition). These are shown in the supplementary information (**Figures S1** and **S2**). The TKER distributions obtained at all five wavelengths are essentially identical, showing progressions of sharp peaks that can be assigned to population of *high N* levels of the A and X states of OH.

The TKER distributions and the recoil anisotropies appear insensitive to whether the initial excitation is to any of the three $\tilde{C}(010)$ state rotational levels or to the $\tilde{D}(000)$ state. Note that each of the probed $\tilde{C}(010)$ levels of H₂O has K_a' (and thus $\langle J_a^2 \rangle$) = 1. Unfortunately, the linestrengths of the $0_{00} \leftarrow 1_{10}$ and $1_{01} \leftarrow 1_{11}$ transitions (which populate levels with $\langle J_a^2 \rangle = 0$) are too weak to be identified under the stronger $\tilde{D} \leftarrow \tilde{X}$ origin band, and any transitions to levels with $K_a' = 2$ are too heavily predissociated to be resolved.³⁰ What is immediately obvious, however, is that – in contrast to D₂O – the fragmentation of H₂O $\tilde{C}(010)$ molecules with $K_a' = 1$ yields *high N* products, i.e. the fragmentation dynamics are indicative of eventual nuclear motion on the $\tilde{B}$ state PES, which are different from that when exciting the same rotational transition within the $\tilde{C}(000) \leftarrow \tilde{X}(000)$ origin band (**Figures S3**). The TKER spectra in the perpendicular direction show no indication of any ‘*high v*’ OH(X) products such as would be expected if non-adiabatic coupling from these levels to the $\tilde{A}$ state PES was a competitive process.

The striking isotope-dependent dynamics of the $\tilde{C}(010)$, $K_a' = 1$ levels of H₂O and D₂O are directly attributable to the accidental resonance between the $\tilde{C}(010)$ and $\tilde{D}(000)$ states of H₂O. As noted above, the $\tilde{D}$ and $\tilde{B}$ states share a common (¹A₁) symmetry, and the $\tilde{D}(000)$ states of both isotopomers are efficiently predissociated by electronic mixing with the $\tilde{B}$ state continuum. The $\tilde{C}$ state, in contrast, has ¹B₁ symmetry. As shown in **Scheme 1**, $\tilde{C}$ state molecules can decay by electronic (same symmetry) coupling to the $\tilde{A}$ state continuum or, if prepared with appropriate nuclear motion (a-axis rotation), by coupling to the $\tilde{B}$ state continuum. The two decay pathways have comparable probabilities for D₂O molecules with $K_a' = 1$ in

either the $\tilde{C}(000)$ or $\tilde{C}(010)$ states, and for H_2O , $\tilde{C}(000)$ molecules with $K_a' = 1$. However, the accidental resonance with the $\tilde{D}(000)$ state greatly boosts the efficiency of Coriolis-coupling from the $\tilde{C}(010)$ state, $K_a' = 1$ levels of H_2O to the $\tilde{B}$ state continuum, as illustrated in **Scheme 2**, to the extent that this pathway to ‘*high N*’ products overwhelms the potential rival electronic coupling route to ‘*high v*’ $\text{OH}(X)$ products.

Scheme 2

A further feature of the $\text{OH}(X, v=0)$ rotational distributions displayed in **Figure 4** is the clear intensity alternation in the populations of successive N levels. These alternations are reminiscent of those observed in the $\text{OH}(X)$ fragments arising in the 121.6 nm photolysis of H_2O_5 and, as in that case, can be attributed to dynamical interferences between dissociation pathways via the CIs between the $\tilde{B}$ and $\tilde{X}$ state PESs at linear H–O–H and H–H–O geometries.

The dissociation mechanisms shown in schemes 1 and 2 highlight a clear yet unprecedented isotope dependence in the photodissociation of $\tilde{C}(010)$ state water molecules. Previous studies have revealed qualitatively different dissociation behaviors for H_2O and D_2O molecules following excitation to the same electronic states, most of which effects could be attributed to mass changes.³⁵⁻³⁷ Here, we reveal a much more dramatic isotope effect, caused by accidental resonance, which leads to different dissociation rates, different dissociation mechanisms, and different product

energy disposals following excitation to a common excited state of H₂O and D₂O. Given the high density of excited vibronic levels of H₂O and D₂O accessible at shorter excitation wavelengths⁴⁰ it would be surprising if the accidental resonance – and the striking isotope dependent fragmentation dynamics – identified in the present study were to be the only example available following VUV photoexcitation of water molecules.

The photodissociation of water is invoked as an important photochemical process in interstellar molecular clouds.⁴¹ Thus the isotope-specific dissociation process identified in the present study may have implications in modelling chemistry prevailing in the interstellar medium. For example, self-shielding has been proposed to account for the observed isotope heterogeneity within the Solar System,^{42,43} but recent investigations of the VUV photodissociation of CO support the view that perturbation-dominated state mixing dynamics will also introduce isotopologue-dependent dissociation probabilities.^{44,45} The strikingly different fragmentation dynamics of H₂O and D₂O observed in the present work represent an extreme example of such accidental resonance induced state mixing and may be expected to contribute to the D/H isotope heterogeneity in the Solar System.

In summary, detailed dynamical features of the photodissociation of $\tilde{C}(010)$ state H₂O and D₂O molecules have been determined using a tunable VUV source in combination with the high resolution H/D atom Rydberg tagging technique. The predissociation mechanisms of H₂O and D₂O following excitation to the $\tilde{C}(010)$ state show a striking isotopologue dependence. The OH product state distributions are

determined by the accidental resonance mediated non-adiabatic coupling pathway, but this pathway has negligible influence on the OD product state distributions. The present study provides an excellent example of competitive state-to-state non-adiabatic couplings and decay pathways involving at least five electronic PESs.

Acknowledgements

The experimental work is supported by the National Natural Science Foundation of China (NSFC Nos. 21873099, 21673232), the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB 17000000), and the Chemical Dynamics Research Center (Grant No. 21688102). MNRA gratefully acknowledges funding from the Engineering and Physical Sciences Research Council (EPSRC, EP/L005913) and the NSFC Center for Chemical Dynamics for the award of a Visiting Fellowship.

Figures and Captions

Figure 1 One-photon action spectra for forming (A) H atoms from H₂O and (C) D atoms from D₂O following excitation at wavelengths around 122 nm. The resolved structures at $\lambda \sim 121.8$ nm for H₂O and $\lambda \sim 122.2$ nm for D₂O are attributable to rotational lines of the respective $\tilde{C}(010) - \tilde{X}(000)$ transitions. The broad features centered at $\lambda \sim 122.1$ nm for H₂O and ~ 121.9 nm for D₂O are the respective $\tilde{D} - \tilde{X}$ origin bands, the profiles of which are distorted by a dip in the VUV tuning curve at $\lambda \sim 122.09$ nm (which is also evident in the action spectrum for forming D atoms from one photon photolysis of CD₄ shown in panel (C)). The extended rotational structures and line assignments of the $\tilde{C}(010) - \tilde{X}(000)$ transitions of H₂O and D₂O are shown in panels (B) and (D), along with comparisons with the respective $\tilde{C} - \tilde{X}$ origin bands. The red arrows in panels (A) and (B) indicate the excitation wavelengths used when measuring the TKER spectra reported in Figures 4, S1 and S2.

Figure 2 3D contour plots of the (A) H + OH products from the photodissociation of H₂O and (B) the D + OD products from the photodissociation of D₂O via the $1_{10} \leftarrow 0_{00}$ lines of the respective $\tilde{C}(010) - \tilde{X}(000)$ transitions. The double headed arrow in (A) shows the alignment of $\mathbf{\epsilon}_{\text{phot}}$. The outer rings in both plots are associated with formation of rovibrational states of the ground (X) state OH radical products, while the inner structures are primarily due to OH/OD(A) products.

Figure 3 TKER spectra of the D + OD products from the photodissociation of D₂O at (A) 122.282, (B) 122.235 and (C) 121.954 nm, resonant with the $1_{01} \leftarrow 1_{11}$ and $1_{10} \leftarrow 0_{00}$ lines of the $\tilde{C}(010) - \tilde{X}(000)$ transition and the center of the $\tilde{D} - \tilde{X}$ origin band, respectively, with ϵ_{phot} aligned parallel (black) and perpendicular (red) to the detection axis.

Figure 4 TKER spectra of the H + OH products from photodissociation of H₂O at (A) 121.881, (B) 121.851 and (C) 121.814 nm, resonant with the $1_{11} \leftarrow 1_{01}$, $1_{10} \leftarrow 0_{00}$ and $2_{11} \leftarrow 1_{01}$ rotational lines of the $\tilde{C}(010) - \tilde{X}(000)$ transition, respectively, with ϵ_{phot} aligned parallel (black) and perpendicular (red) to the detection axis.

References

- (1) Lee, L. C.; Suto, M. Quantitative Photoabsorption and Fluorescence Study of H₂O and D₂O at 50-190 nm. *Chem. Phys.* **1986**, *110*, 161-169.
- (2) Carr, J. S.; Najita, J. R. The OH Rotational Population and Photodissociation of H₂O in DG Tauri. *Astrophys. J.* **2014**, *788*, 1-15.
- (3) Chang, Y.; Yu, Y.; Wang, H. L.; Hu, X. X.; Li, Q. M.; Yang, J. Y.; Su, S.; He, Z. G.; Chen, Z. C.; Che, L.; Wang, X. N.; Zhang, W. Q.; Wu, G. R.; Xie, D. Q.; Ashfold, M. N. R.; Yuan, K. J.; Yang, X. M. Hydroxyl Super Rotors from Vacuum Ultraviolet Photodissociation of Water. *Nat. Commun.* **2019**, *10*, 1025.
- (4) Dixon, R. N.; Hwang, D. W.; Yang, X. F.; Harich, S.; Lin, J. J.; Yang, X. Chemical "Double Slits": Dynamical Interference of Photodissociation Pathways in Water. *Science* **1999**, *285*, 1249-1253.
- (5) Harich, S. A.; Yang, X. F.; Hwang, D. W. H.; Lin, J. J.; Yang, X. M.; Dixon, R. N. Photodissociation of D₂O at 121.6 nm: A State-to-state Dynamical Picture. *J. Chem. Phys.* **2001**, *114*, 7830-7837.
- (6) Yuan, K. J.; Dixon, R. N.; Yang, X. M. Photochemistry of the Water Molecule: Adiabatic versus Nonadiabatic Dynamics. *Acc. Chem. Res.* **2011**, *44*, 369-378.
- (7) Hwang, D. W.; Yang, X. F.; Harich, S.; Lin, J. J.; Yang, X. Photodissociation Dynamics of H₂O at 121.6 nm: Effect of Parent Rotational Excitation on Reaction Pathways. *J. Chem. Phys.* **1999**, *110*, 4123-4126.
- (8) Dixon, R. N.; Oliver, T. A. A.; Cheng, L.; Cheng, Y.; Yuan, K. J.; Yang, X. M. Vibronically Induced Decay Paths from the $\tilde{C}^1B_1$ -state of Water and Its Isotopomers. *J. Chem. Phys.* **2013**, *138*, 104306.
- (9) Herzberg, G. *Electronic Spectra of Polyatomic Molecules*, Van Nostrand, Princeton, 1966.
- (10) Engel, V.; Schinke, R.; Staemmler, V. Photodissociation Dynamics of H₂O and D₂O in the First Absorption Band - A Complete Ab Initio Treatment. *J. Chem. Phys.* **1988**, *88*, 129-148.
- (11) Guo, H.; Murrell, J. N. Dynamics of the A-state Photodissociation of H₂O at 193 nm. *Mol. Phys.* **1988**, *65*, 821-827.
- (12) Engel, V.; Staemmler, V.; Vanderwal, R. L.; Crim, F. F.; Sension, R. J.; Hudson, B.; Andresen, P.; Hennig, S.; Weide, K.; Schinke, R. Photodissociation of Water in the

First Absorption Band - A Prototype for Dissociation on a Repulsive Potential Energy Surface. *J. Phys. Chem.* **1992**, *96*, 3201-3213.

(13) Yang, X. F.; Hwang, D. W.; Lin, J. J.; Ying, X. Dissociation Dynamics of the Water Molecule on the $\tilde{A}^1B_1$ Electronic Surface. *J. Chem. Phys.* **2000**, *113*, 10597-10604.

(14) van Harrevelt, R.; van Hemert, M. C. Photodissociation of Water in the $\tilde{A}$ Band Revisited with New Potential Energy Surfaces. *J. Chem. Phys.* **2001**, *114*, 9453-9462.

(15) Zhou, L. S.; Xie, D. Q.; Sun, Z. G.; Guo, H. Product Fine-structure Resolved Photodissociation Dynamics: The A Band of H₂O. *J. Chem. Phys.* **2014**, *140*, 024310.

(16) Lu, I. C.; Wang, F. Y.; Yuan, K. J.; Cheng, Y.; Yang, X. M. Nonstatistical Spin Dynamics in Photodissociation of H₂O at 157 nm. *J. Chem. Phys.* **2008**, *128*, 066101.

(17) van Harrevelt, R.; van Hemert, M. C. Photodissociation of Water. I. Electronic Structure Calculations for the Excited States. *J. Chem. Phys.* **2000**, *112*, 5777-5786.

(18) van Harrevelt, R.; van Hemert, M. C. Photodissociation of Water. II. Wave Packet Calculations for the Photofragmentation of H₂O and D₂O in the $\tilde{B}$ Band. *J. Chem. Phys.* **2000**, *112*, 5787-5808.

(19) Fillion, J. H.; van Harrevelt, R.; Ruiz, J.; Castillejo, N.; Zanganeh, A. H.; Lemaire, J. L.; van Hemert, M. C.; Rostas, F. Photodissociation of H₂O and D₂O in $\tilde{B}$, $\tilde{C}$, and $\tilde{D}$ States (134-119 nm). Comparison between Experiment and Ab Initio Calculations. *J. Phys. Chem. A* **2001**, *105*, 11414-11424.

(20) Mordaunt, D. H.; Ashfold, M. N. R.; Dixon, R. N. Dissociation Dynamics of H₂O (D₂O) Following Photoexcitation at the Lyman- α Wavelength (121.6 nm). *J. Chem. Phys.* **1994**, *100*, 7360-7375.

(21) Cheng, Y.; Yuan, K. J.; Cheng, L. N.; Guo, Q.; Dai, D. X.; Yang, X. M. Photodissociation Dynamics of H₂O: Effect of Unstable Resonances on the $\tilde{B}^1A_1$ Electronic State. *J. Chem. Phys.* **2011**, *134*, 064301.

(22) Vondirke, M.; Heumann, B.; Kuhl, K.; Schroder, T.; Schinke, R. Fluctuations in Absorption Spectra and Final Product State Distributions following Photodissociation Processes. *J. Chem. Phys.* **1994**, *101*, 2051-2068.

(23) Zhou, L. S.; Jiang, B.; Xie, D. Q.; Guo, H. State-to-state Photodissociation Dynamics of H₂O in the B Band: Competition between Two Coexisting Nonadiabatic Pathways. *J. Phys. Chem. A* **2013**, *117*, 6940-6947.

-
- (24) Zhou, L. S.; Lin, G. S. M.; Xie, D. Q. State to State Photodissociation Dynamics of D₂O in the B Band. *J. Chem. Phys.* **2013**, *139*, 114303.
- (25) Bin, J.; Xie, D. Q.; Guo, H. State-to-state Photodissociation Dynamics of Triatomic Molecules: H₂O in the B band. *J. Chem. Phys.* **2012**, *136*, 034302.
- (26) Hu, X. X.; Zhou, L. S.; Xie, D. Q. State-to-state Photodissociation Dynamics of the Water Molecule. *Wires Comput. Mol. Sci.* **2018**, *8*, 1350.
- (27) Su, S.; Wang, H. Z.; Chen, Z. C.; Yu, S. R.; Dai, D. X.; Yuan, K. J.; Yang, X. M. Photodissociation Dynamics of HOD via the $\tilde{B}^1A_1$ Electronic State. *J. Chem. Phys.* **2015**, *143*, 184302.
- (28) Yi, W. K.; Park, J.; Lee, J. Photodissociation Dynamics of Water at Lyman- α (121.6 nm). *Chem. Phys. Lett.* **2007**, *439*, 46-49.
- (29) Zanganeh, A. H.; Fillion, J. H.; Ruiz, J.; Castillejo, M.; Lemaire, J. L.; Shafizadeh, N.; Rostas, F. Photodissociation of H₂O and D₂O below 132 nm. *J. Chem. Phys.* **2000**, *112*, 5660-5671.
- (30) Ashfold, M. N. R.; Bayley, J. M.; Dixon, R. N. Molecular Predissociation Dynamics Revealed through Multiphoton Ionization Spectroscopy.1. The $\tilde{C}^1B_1$ States of H₂O and D₂O. *Chem. Phys.* **1984**, *84*, 35-50.
- (31) Kuge, H. H.; Kleinermmanns, K. Rotational Predissociation of H₂O ($\tilde{C}^1B_1$) Studied by Multiphoton Ionization Spectroscopy in a Supersonic Free Jet. *J. Chem. Phys.* **1989**, *90*, 46-52.
- (32) Hodgson, A.; Simons, J. P.; Ashfold, M. N. R.; Bayley, J. M.; Dixon, R. N. Quantum State-selected Photodissociation Dynamics in H₂O and D₂O. *Mol. Phys.* **1985**, *54*, 351-368.
- (33) Yang, C. H.; Sarma, G.; ter Meulen, J. J.; Parker, D. H.; Western, C. M. REMPI Spectroscopy and Predissociation of the $\tilde{C}^1B_1$ ($v=0$) Rotational Levels of H₂O, HOD and D₂O. *Phys. Chem. Chem. Phys.* **2010**, *12*, 13983-13991.
- (34) Yuan, K. J.; Cheng, Y.; Cheng, L.; Guo, Q.; Dai, D. X.; Wang, X. Y.; Yang, X. M.; Dixon, R. N. Nonadiabatic Dissociation Dynamics in H₂O: Competition between Rotationally and Nonrotationally Mediated Pathways. *Pro. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 19148-19153.
- (35) Harich, S. A.; Yang, X. F.; Hwang, D. W. H.; Lin, J. J.; Yang, X. M.; Dixon, R. N. Photodissociation of D₂O at 121.6 nm: A State-to-state Dynamical Picture. *J. Chem.*

Phys. **2001**, *114*, 7830-7837.

(36) Cheng, Y.; Cheng, L. N.; Guo, Q.; Yuan, K. J.; Dai, D. X.; Wang, X. Y.; Dixon, R. N.; Yang, X. M. Rotational State Specific Dissociation Dynamics of D₂O via the $\tilde{C}$ Electronic State. *J. Chem. Phys.* **2010**, *133*, 034307.

(37) Cheng, Y.; Cheng, L. N.; Guo, Q.; Yuan, K. J.; Dai, D. X.; Yang, X. M. Photodissociation Dynamics of D₂O via the $\tilde{B}^1A_1$ Electronic State. *J. Chem. Phys.* **2011**, *134*, 104305.

(38) He, Z. G.; Yang, D. Y.; Chen, Z. C.; Yuan, K. J.; Dai, D. X.; Wu, G. R.; Yang, X. M. An Accidental Resonance Mediated Predissociation Pathway of Water Molecules Excited to the Electronic $\tilde{C}$ State. *Phys. Chem. Chem. Phys.* **2017**, *19*, 29795-29800.

(39) Johns, J. W. C. On the Absorption Spectrum of H₂O and D₂O in Vacuum Ultraviolet. *Can. J. Phys.* **1963**, *41*, 209-219.

(40) Fillion, J. H.; Ruiz, J.; Yang, X. F.; Castillejo, M.; Rostas, F.; Lemaire, J. L. High resolution photoabsorption and photofragment fluorescence spectroscopy of water between 10.9 and 12 eV. *J. Chem. Phys.* **2004**, *120*, 6531-6541.

(41) Van Dishoeck, E. F.; Herbst, E.; Neufeld, D. A. Interstellar Water Chemistry: From Laboratory to Observations. *Chem. Rev.* **2013**, *113*, 9043-9085.

(42) Yurimoto, H.; Kuramoto, K. Molecular Cloud Origin for the Oxygen Isotope Heterogeneity in the Solar System. *Science* **2004**, *305*, 1763-1766.

(43) Lyons, J. R.; Young, E. D. CO Self-shielding as the Origin of Oxygen Isotope Anomalies in the Early Solar Nebula. *Nature* **2005**, *435*, 317-320.

(44) Chakraborty, S.; Rude, B.; Ahmed, M.; Thieme, M. H. Carbon and Oxygen Isotopic Fractionation in the Products of Low-temperature VUV Photodissociation of Carbon Monoxide. *Chem. Phys.* **2018**, *514*, 78-86.

(45) Chakraborty, S.; Ahmed, M.; Jackson, T. L.; Thieme, M. H. Experimental Test of Self-shielding in Vacuum Ultraviolet Photodissociation of CO. *Science* **2008**, *321*, 1328-1331.