Leoni, F., Shi, R., Tanaka, H., & Russo, J. (2019). Crystalline clusters in mW water: Stability, growth, and grain boundaries. *Journal of Chemical Physics*, 151(4), [044505].
<https://doi.org/10.1063/1.5100812>

Publisher's PDF, also known as Version of record

License (if available):
Other

Link to published version (if available):
[10.1063/1.5100812](https://doi.org/10.1063/1.5100812)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via AIP at <https://doi.org/10.1063/1.5100812> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Crystalline clusters in mW water: Stability, growth, and grain boundaries

Cite as: J. Chem. Phys. **151**, 044505 (2019); <https://doi.org/10.1063/1.5100812>

Submitted: 20 April 2019 . Accepted: 01 July 2019 . Published Online: 29 July 2019

Fabio Leoni, Rui Shi , Hajime Tanaka , and John Russo

View Online

Export Citation

CrossMark

ARTICLES YOU MAY BE INTERESTED IN

[Some considerations on the water polymorphism and the liquid-liquid transition by the density behavior in the liquid phase](#)

The Journal of Chemical Physics **151**, 044504 (2019); <https://doi.org/10.1063/1.5095687>

[Ice growth rate: Temperature dependence and effect of heat dissipation](#)

The Journal of Chemical Physics **151**, 044509 (2019); <https://doi.org/10.1063/1.5103273>

[On the link between polyamorphism and liquid-liquid transition: The case of salty water](#)

The Journal of Chemical Physics **151**, 044503 (2019); <https://doi.org/10.1063/1.5100959>

Lock-in Amplifiers up to 600 MHz

starting at

\$6,210

Zurich
Instruments

Watch the Video

Crystalline clusters in mW water: Stability, growth, and grain boundaries

Cite as: J. Chem. Phys. 151, 044505 (2019); doi: 10.1063/1.5100812

Submitted: 20 April 2019 • Accepted: 1 July 2019 •

Published Online: 29 July 2019

Fabio Leoni,^{1,a)} Rui Shi,^{2,b)} Hajime Tanaka,^{2,c)} and John Russo^{1,2,d)}

AFFILIATIONS

¹School of Mathematics, University of Bristol, Bristol BS8 1TW, United Kingdom

²Department of Fundamental Engineering, Institute of Industrial Science, University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo 153-8505, Japan

Note: This paper is part of a JCP Special Topic on Chemical Physics of Supercooled Water.

^{a)}fabio.leoni@bristol.ac.uk

^{b)}rshi@iis.u-tokyo.ac.jp

^{c)}tanaka@iis.u-tokyo.ac.jp

^{d)}john.russo@bristol.ac.uk

ABSTRACT

With numerical simulations of the mW model of water, we investigate the energetic stability of crystalline clusters both for Ice *I* (cubic and hexagonal ice) and for the metastable Ice 0 phase as a function of the cluster size. Under a large variety of forming conditions, we find that the most stable cluster changes as a function of size: at small sizes, the Ice 0 phase produces the most stable clusters, while at large sizes, there is a crossover to Ice *I* clusters. We further investigate the growth of crystalline clusters with the seeding technique and study the growth patterns of different crystalline clusters. While energetically stable at small sizes, the growth of metastable phases (cubic and Ice 0) is hindered by the formation of *coherent* grain boundaries. A fivefold symmetric twin boundary for cubic ice, and a newly discovered coherent grain boundary in Ice 0, promotes cross nucleation of cubic ice. Our work reveals that different local structures can compete with the stable phase in mW water and that the low energy cost of particular grain boundaries might play an important role in polymorph selection.

Published under license by AIP Publishing. <https://doi.org/10.1063/1.5100812>

I. INTRODUCTION

The pioneering work of Frank¹ revealed that the lowest energy local structure can differ from the arrangement found in the stable crystalline phases. In the Lennard-Jones fluid, for example, Frank found that the 13-atom icosahedral cluster has an energy which is 8.4% lower than 13-atom clusters corresponding to nearest-neighbor environments in close-packed crystals (both the fcc and hcp crystalline phases). The idea of competing local structures has had a profound influence on our understanding of supercooled liquid states and of their nonequilibrium extension, i.e., glasses.^{2–8}

The study of locally favored structures has also played an important role in the understanding of the behavior of water.⁹ Differently from simple liquids with isotropic interactions, water is characterized by strong directional interactions that favor a local

tetrahedral arrangement of particles. It is possible to classify these tetrahedral environments in two different groups: amorphous and crystalline local structures.

The first group is composed of amorphous local structures. Amorphous structures are characterized by a lack of full translational and rotational symmetry, meaning that there is a high degree of degeneracy with respect to the relative position and/or orientation of the tetrahedral units. One of the key properties of water, that sets it apart from nonanomalous liquids, is that its amorphous structures can be further grouped into two states.¹⁰ The first state, which we call the *S*-state, is a low density/energy configuration in which a central water molecule is surrounded by four tetrahedra that are translationally ordered (i.e., they form a well-defined second shell of nearest neighbors), but orientationally disordered (meaning that the relative orientation of the tetrahedra is disordered).¹¹ The second state, which we call the *p*-state, is a high-density/energy configuration with

a collapsed second shell of nearest neighbors. The definition of an appropriate order parameter has revealed bimodality in liquid structures up to ambient conditions and has allowed the definition of a microscopic two-state model that accounts for all water thermodynamic anomalies.¹¹ Locally favored structures also impose strong constraints on the dynamics of water molecules: more specifically, a water molecule has a slower diffusive dynamics the more S-states surround it. A hierarchical two-state model was built with this idea in mind and was found to be able to account for water's dynamic anomalies¹² while also predicting a strong-to-strong crossover at supercooled conditions.¹³

The idea of different states playing a pivotal role in determining water's behavior is further supported by the behavior of water in supercooled and out-of-equilibrium conditions. In particular, several simulation works have shown that water can exist in two distinct liquid forms: the LDL (low density liquid) and HDL (high density liquid) phases, which become indistinguishable at a liquid-liquid critical point.^{14–17} A nonequilibrium counterpart of this transition is also found in water glasses, where low-density (LDA) and high density (HDA) amorphous glasses have been found (with the proposal of an additional very-high density VHDA amorphous glass^{18,19}).

The second group of local structures is composed of crystalline local structures. While the anomalous behavior of the liquid phase appears to be strongly linked with the competition of two different amorphous local structures, the process of ice nucleation is associated with a different type of local structure, i.e., crystalline local structures. These structures are defined as clusters of molecules whose nearest-neighbor environment is the same as those found in bulk crystals. They differ from the amorphous local structures for their high degree of local translational and rotational order. They can thus extend to arbitrarily large sizes, and their growth becomes irreversible above the critical nucleus size. The relationship of these crystalline local structures to amorphous local structures formed in a supercooled state is one of the important factors controlling the ease of crystallization.^{3,8,9,20} These clusters originate from fluctuations that are very localized in space and time. For this reason, computer simulations are giving a fundamental contribution to our understanding of this important process.^{8,9,17,21,22}

Homogeneous ice nucleation occurs by the growth of the stable Ice I phase from a few nucleation sites within the supercooled liquid state. Ice I has two different polytypes: *hexagonal* ice (space group $P6_3/mmc$) and *cubic* ice (space group $Fd\bar{3}m$). Hexagonal ice is the thermodynamic stable form, but nucleation at supercooled conditions often proceeds through metastable phases, and deviations from sixfold rotational symmetry have been found in diffraction patterns for a vast array of crystallization scenarios (see Ref. 23 for a review). These include recrystallization from either amorphous ices^{24–28} or high-pressure ices.^{29–32} This metastable phase was long identified with the cubic phase, but it has recently been pointed out that due to the high amount of stacking disorder,^{33–40} this phase should be more appropriately called stacking disordered ice, ice I_{sd} , which retains a threefold rotational symmetry and can be identified with the space group $P3m1$.^{31,41}

The formation of ice I_{sd} can be understood in terms of cross-nucleation,^{42,43} which occurs when two polymorphs share a common crystallographic plane, i.e., the (111) plane for cubic ice and the basal plane (0001) of hexagonal ice. In this case, the two crystals

match perfectly at the interface plane (both in symmetry and lattice constant), without discontinuities across the two phases. This causes a great reduction of the interface free energy. In the language of interphase interfaces, the two Ice I polytypes form a *coherent* interface. Very recently, computer simulations of the mW model have shown that ice crystals with stacking disorder are about 14 kJ mol^{-1} more stable than hexagonal crystals for sizes up to 100 000 molecules at $T = 230 \text{ K}$,⁴⁴ nicely showing that the driving force for crystallization is size dependent.

Cubic ice and ice I_{sd} are the only metastable phases found in laboratory experiments at ambient pressure. However, there are different candidate crystalline structures, which retain the local tetrahedral arrangement of water molecules and have additional properties not found in the known forms of ice.⁴⁵ We focus here on the so-called Ice 0 form, which is currently the only known phase (of the mW potential) that is metastable in the regime of ambient pressure and low temperatures that is relevant for homogeneous ice nucleation.²⁰ The unit cell of Ice 0 contains 12 oxygen atoms and is tetragonal where the ratio between the long and the short sides of the cell is $c/a = 1.81$.

Several simulation studies have addressed the problem of homogeneous nucleation of water, both in molecular models, such as TIP4P,^{38,46,47} and the coarse grained mW model,^{22,48–50} which sacrifices orientational degrees of freedom in exchange of a considerable faster crystallization dynamics. To enhance the sampling of the crystalline cluster, several techniques such as umbrella sampling, metadynamics, and forward flux sampling have been employed.^{51–54} An understanding of ice nucleation requires accurate studies of interfacial properties^{55–60} which are commonly used within classical nucleation theory parameterizations.^{61–63} The behavior of water at interfaces is central in understanding heterogeneous nucleation,^{64–68} nucleation with free surfaces,^{69–74} in nanoconfinement,^{75,76} and crystallization in solutions.^{77–79}

In the present work, we focus on crystalline clusters for the mW model of water.⁴⁸ We consider both polytypes of Ice I, i.e., the cubic and hexagonal ice phases, and the metastable Ice 0 phase. We first examine their energetic stability under a wide variety of conditions (temperature and density). We will show that for small clusters, the Ice I crystalline clusters are not the most energetically stable. We instead show that small clusters with the nearest neighbor environment of Ice 0 have a lower total energy due to the abundance of 5-membered rings. While Ice I clusters (both hexagonal and cubic forms) grow isotropically in all directions, Ice 0 grows along layers composed of 5-membered rings. We then perform seeding simulations at supercooled conditions and examine the growth behavior of the different clusters. In particular, we highlight the importance of grain boundaries in the kinetic pathway of growth of the different crystalline clusters. We focus here on three coherent grain boundaries, whose low free energy penalties make them particularly abundant. We will estimate these free energy penalties and try to rationalize their impact on ice nucleation.

II. METHODS

Here, we perform Monte Carlo simulations of the mW model of water.⁴⁸ We employ isobaric simulations at ambient pressure and focus on highly supercooled conditions, $T = 218 \text{ K}$, which is low enough to have accessible nucleation barriers but high enough that

nucleation is a rare process (not spontaneously occurring in unbiased simulations) and that dynamics can still be observed over 1000 s of Brownian times. Ice crystals are then grown both via biased simulations, with the CNT-US method, and unbiased simulations, with the *seeding* technique.

The CNT-US method is a variant of umbrella sampling introduced in Ref. 20, where the expression for the free-energy barrier of classical nucleation theory is used as a biasing potential in simulations. The new Hamiltonian is then written as

$$\mathcal{H}' = \mathcal{H} + |\Delta\mu|n^{2/3}(n^{1/3} - 3n_c^{1/3}/2), \quad (1)$$

where $\mathcal{H}$ is the given by the mW potential, $\Delta\mu$ is the chemical potential difference between Ice I and supercooled water, n is the size of the largest nucleus in the system, and n_c is the critical nucleus size. The advantage of this method over standard umbrella sampling is that the whole free energy landscape can be explored in equilibrium in one independent simulation, avoiding kinetic traps that are often incurred when restricting simulations to small sampling windows.

To distinguish the nucleation barriers between different phases, e.g., Ice I_h , I_c , and nuclei with stacking disorder, we perform standard umbrella sampling simulations in which we initialize the simulations with a nuclei of the desired phase. These simulations only preserve the biasing potential of the CNT-US scheme, but, as in traditional US schemes, the range of the order parameter (the size of the largest nucleus) is divided into many small windows, each one allowing fluctuations of the order parameter between N_i and $N_i + \Delta N$. Since the initialization of each window occurs with a seed of the correct type, fluctuations to a seed with a different composition occur very rarely. We thus monitor the trajectory in each window and perform a check of the nucleus composition every 1000 Monte Carlo (MC) cycles to ensure that the nucleus has preserved the correct composition. If not, we perform a rejection move of the trajectory of the last 1000 MC time steps. We typically set $\Delta N = 5$, but we have checked that the results are insensitive to this choice for $\Delta N = 1$ (where the technique is formally equivalent to the successive umbrella sampling scheme⁸⁰) and $\Delta N = 10$.

We then monitor the different simulations to make sure that the original crystal structure is preserved during the sampling and ensure that no additional stacking faults have originated.

For unbiased simulations, we use the *seeding* technique,^{81–84} which consists in the insertion of perfect crystalline nuclei in the melt in order to determine at which conditions the nucleus is critical, and then use classical nucleation theory to estimate parameters such as the surface tension and the crystallization rate. While being successful in the estimation of the crystallization rate of water at moderate supercooling, with critical nuclei composed of several thousand molecules and where the approximations of classical nucleation theory are most likely to hold, potential problems with the extension of this technique to deeply supercooled conditions have been recently highlighted.⁸⁵ On the contrary, for systems seeded with stable Ice I crystals at deeply supercooled conditions, we will find that the critical nucleus sizes are in excellent agreement with the ones we obtain independently from umbrella sampling simulations.

To distinguish the different crystalline phases, and to bias the umbrella sampling simulations, we use the standard

bond-orientational order parameters. In particular, we employ the Q_{12} bond orientational order parameter, where a molecule is identified as having a crystal-like environment if it has more than 12 connected neighbors among its first 16 nearest neighbors. A connected neighbor of particle i is defined as a neighbor j for which the normalized product satisfies $Q_{12}(i) \cdot Q_{12}(j) > 0.75$. See Ref. 2 for a recent review of this methodology.

Where not differently specified, the system size of seeding simulations is $N = 10\,000$ water molecules, while umbrella sampling simulations are run with $N = 4000$ molecules. Since we only focus on small nuclei (up to $N = 100$ molecules), no strong system size dependence of our results is expected.

III. RESULTS

A. Energy of crystalline clusters

In order to understand crystal cluster competition in supercooled water, we compare the energy of clusters of perfect crystals in three different ways to account for different scenarios of cluster formation: (i) at the equilibrium density ρ , corresponding to a specific temperature T , which depends on the structure under consideration (see Fig. 1); (ii) at fixed density corresponding to that of the liquid phase at equilibrium; (iii) at fixed bond distance, in analogy with the work of Frank.¹ Afterward, (iv) we extend the approach (i) to account for cluster equilibration in the liquid phase and (v) we also consider the energetic contribution coming from the hydration layer.

In Fig. 1, we show the ρ vs T relation at constant pressure $P = 0$ atm for the liquid phase (L), cubic ice (I_c), hexagonal ice (I_h), and ice 0 (Ice 0) as obtained from Monte Carlo simulations of the mW model in the NPT ensemble for $N = 5376$ particles with periodic boundary conditions. We can notice the well-known temperature of maximum density (TMD) anomaly of liquid water found at around $T = 250$ K in the mW model at normal conditions.^{48,86} We also note crystal densities being lower than the liquid density above 207 K (see Fig. 1).

FIG. 1. ρ vs T relation at constant pressure $P = 0$ atm, as described in the main text, for cubic ice (I_c , black circles), hexagonal ice (I_h , red squares), ice 0 (Ice 0, green diamonds), and liquid phase (L, blue triangles).

Clusters of mW particles are built by adding particles to a crystalline lattice (or from a configuration of the liquid at equilibrium in the case of disordered clusters) considering the following energetic approach: starting from any position in the crystalline lattice, we add one particle at each step choosing among the lattice positions that result in the minimum energy cluster. This involves computing the bond energy of every particle on the surface of the cluster with its potential neighbors and choosing to add the particle that generates the cluster with minimum energy. In the case of degenerate lattice positions, i.e., different clusters with the same size and same energy, the choice of which lattice position to fill is done randomly. We repeat this procedure until we get a cluster of the desired

FIG. 2. Energy per particle of clusters E/n as a function of cluster size n for cubic ice (I_c , black circles), hexagonal ice (I_h , red squares), ice 0 (Ice 0, green diamonds and dark green left-pointing triangles), and liquid phase (L, blue triangles). The vertical orange band marks the crossing region between Ice 0 and I_c/I_h curves. In (a), every cluster is considered at the same density of the liquid phase at equilibrium at $T = 200$ K, which is $\rho = 0.976$ g/cm³ [method (ii) described in the main text]. The shortest bond distance in each structure is $d = 2.71$, 2.71 , and 2.67 Å for I_c , I_h , and Ice 0, respectively, while the shortest neighboring distance in the case of the liquid phase L is the position of the first peak of the liquid two-point correlation function. In order to compare the different clusters following the method (iii), we also show Ice 0 at the density $\rho = 0.937$ g/cm³ (dark green left-pointing triangles) which corresponds to $d = 2.71$ Å, the same of I_c/I_h . In (b), every cluster is at its own equilibrium density corresponding to $T = 235$ K [method (i)], which is $\rho = 0.982$, 0.982 , 0.951 , and 1.002 g/cm³ for I_c , I_h , Ice 0, and L, respectively. Inset in (a): energy per particle of clusters E/n as a function of cluster size n using the spherical symmetric growth of clusters based on the distance from the seed, as described in the main text.

size. Since clusters will have different morphologies (i.e., different numbers of interparticle bonds), we compute the average energy of clusters over 50 independent growth processes.

In Figs. 2(a) and 2(b), we show the energy per particle of clusters, E/n , as a function of the cluster size n for the crystals and liquid phases at the conditions described above (ii), (iii), and (i), respectively.

From Figs. 2(a) and 2(b), we can notice that clusters of I_c and I_h have always the same energy for any size n , as expected. We also note that Ice 0 has a lower energy with respect to Ice I for small cluster sizes up to the critical value $n^* \sim 40$ for the three conditions (i), (ii), and (iii) [we computed the value of $n = n^*$ at which curves associated with Ice 0 and I_c/I_h cross for other values of ρ and T following methods (ii) and (iii) and found always $n^* \sim 40$].

To extend the previous results, we account for cluster equilibration in the liquid phase [method (iv)]. In order to do so, we perform Monte Carlo simulations sampling the dynamics according to a fixed topology structure, where the energetic neighbors of each particle of a cluster are constrained to be the same during the dynamics and their bond elongation bounded by a cutoff set to ≈ 1.3 Å. In Fig. 3, we show the energy per particle of clusters, E/n , as a function of the cluster size n for the crystals obtained following method (iv). The curves are obtained averaging for each size n over 10 different uncorrelated simulation times and 50 independent growth processes. From Fig. 3, we can see that clusters of Ice 0 are more stable with respect to clusters of ice I up to the size $n_{eq}^* \approx 30$.

Considering also the contribution to the energy of clusters coming from the hydration layer [method (v)], we observe (inset of Fig. 3) that the maximum size up to which clusters of Ice 0 are more stable with respect to ice I is reduced further with respect to the previous cases up to the value $n_{eq-hy}^* \approx 24$. This behavior is similar to what has been observed for the stability of the 13-particle icosahedron with respect to 13-particle fcc and hcp clusters interacting through the Lennard-Jones potential when embedding clusters in the liquidlike environment.⁸⁷

FIG. 3. Energy per particle of equilibrated clusters E/n as a function of cluster size n for cubic ice (I_c , black circles), hexagonal ice (I_h , red squares), and ice 0 (Ice 0, green diamonds). The vertical orange band marks the crossing region between Ice 0 and I_c/I_h curves. Inset: energy per particle (including the hydration layer) of equilibrated and hydrated clusters E/n_{eq-hy} as a function of cluster size n .

1. Cluster morphology

In Fig. 4, we show typical clusters of ices I_c , I_h , and Ice 0 projected in the xy (left panels) and xz (right panels) planes for clusters of size $n = 40$ that is the typical value at which the curves of energy per particle associated with Ice 0 and I_c/I_h cross, for methods (i), (ii), and (iii). Particles are colored according to the growth sequence such that the seeding particle is the darkest one, and the last particle attached to the cluster is the lightest one. For I_c , the xy plane corresponds to the direction $[111]$, while xz to $[\bar{1}10]$. For I_h , xy is the basal face corresponding to the Miller indices $\{0001\}$, while xz is the secondary prism face corresponding to the Miller indices $\{11-20\}$. For Ice 0, the long side of the unit cell c is aligned along the z axis.

The first important difference between Ice 0 and I_c/I_h is that Ice 0 grows preferentially along the (100) plane, while other clusters grow with spherical symmetry. We understand the planar symmetry of growing clusters of Ice 0 in terms of its neighboring bond lengths. While I_c and I_h are characterized by a unique length, which is the distance between two neighboring particles, Ice 0 is characterized by three different neighboring distances (d_s , d_i , and d_l), very close to each other. Two of these three lengths (d_s and d_l) are involved in the formation of planar structures. In the snapshots of Ice 0 in Fig. 4, white bonds are associated with the short distance ($d_s \approx 2.67$ Å for $\rho = 0.976$ g/cm³), while blue bonds to the long distance ($d_l \approx 2.73$ Å for $\rho = 0.976$ g/cm³, which is about 2% longer than d_s). The intermediate distance ($d_i \approx 2.68$ Å for $\rho = 0.976$ g/cm³) connects particles along the z direction forming distorted hexagonal rings, and

FIG. 4. Snapshots of typical clusters of size $n = 40$ projected in the xy (left panels) and xz (right panels) planes, for the crystalline phases I_c , I_h , and Ice 0. Bonds connect pairs of neighboring particles. Particles are colored according to the growth sequence such that the seeding particle is the darkest one, and the last particle attached to the cluster is the lightest one.

then it does not play any role in the formation of clusters built following the minimum energy rule, as discussed below. From Fig. 4, we can see that clusters of Ice 0 grow forming rings of pentagons composed of d_s (white) and d_l (blue) bonds. For this reason, the energy of Ice 0 is lower than I_c/I_h already for clusters of $n = 5$ particles which form pentagons in Ice 0, that is a ring, while I_c/I_h can form the first ring only for $n = 6$. Each ring forming in the cluster gives an extra term in the calculation of its total energy. Indeed, the number of bonds N_b forming in a cluster is $N_b = n + N_r - 1$, where N_r is the number of rings. Counting the number of bonds for the different growing clusters, we find that on average I_c and I_h can form more bonds than Ice 0 only for $n \geq 40$, which corresponds to the typical value at which the curves of energy per particle associated with Ice 0 and I_c/I_h cross (highlighted by the orange vertical band in Fig. 2).

Instead of adding the strongest bonded particle to a cluster, we can add particles with increasing radial distance from a central particle. In this case, we get for Ice 0 clusters that are similar to those obtained following the minimum energy rule up to a size of $n = 13$, while for $n > 13$, we get a spherical symmetric structure whose energy becomes in average larger than the energy of clusters of I_c/I_h (computed with this same rule) for $n \gtrsim 22$ [see inset of Fig. 2(a)]. This value of n is smaller than the critical value of $n (\gtrsim 40)$ found for clusters of Ice 0 growing according to the minimum energy rule because for equal number of particles, a spherical symmetric cluster forms a smaller number of rings than a planar symmetric cluster for the Ice 0, while for I_c/I_h , the growth is always spherically symmetric following both growing rules.

The reason why the number of bonds N_b can be used to compute the difference in the energy content between clusters of I_c/I_h and Ice 0 is related to the two characteristic lengths in Ice 0, d_s and d_l , being very close to each other [i.e., $(d_l - d_s)/d_s \ll 1$]. If the characteristic length of neighboring particles forming a cluster is only one, as in the case of I_c and I_h , this length will be the minimum of the potential energy (note that for perfect crystals the three-body term of the mW potential is zero). If, on the other hand, the characteristic lengths of neighboring particles forming a cluster are more than one, in general, they will be different from the length corresponding to the minimum of the potential energy and then the eventual increase in the number of bonds will not balance the increase in energy of each individual bond. In the case of Ice 0, since the two characteristic lengths d_s and d_l forming clusters are close to each other and close to the distance corresponding to the minimum of the potential energy (which is 2.71 Å for $\rho = 0.976$ g/cm³), the energy of each bond is very close to that of bonds in I_c/I_h clusters such that the value of N_b is a clear indication of the energy content of a cluster.

The mechanism responsible for the lower energy content of small clusters of Ice 0 with respect to clusters of I_c/I_h now described is the same causing the icosahedron to have a lower energy content than fcc and hcp clusters in simple one-component liquids with particles interacting through spherically symmetric pair-potentials, like the Lennard-Jones. Indeed, fcc and hcp structures are characterized by a single length, which is the minimum of the potential. The icosahedron is characterized by two different lengths: the distance between the central particle and its 12 neighbors, and the edge of the 20 equilateral triangles forming the surface of the icosahedron where the vertices are the 12 neighboring particles, which is about 5% longer. The total number of bonds forming

the icosahedron is 42. In the case of fcc or hcp structures, the 13 particles (central particle plus its 12 first near neighbors) composing them can form only 36 bonds. Although the 42 bonds formed in the icosahedron have a slightly higher energy with respect to the 36 bonds formed in fcc and hcp, the total energy of a 13-particle cluster of icosahedron is smaller (by 8.4% in the case of a Lennard-Jones liquid¹) than that of a 13-particle cluster of fcc or hcp. Another analogy between Ice 0 and the icosahedron is their tendency to form more clusters as the temperature is lowered.^{87,88} On the other hand, an important difference between Ice 0 and the icosahedron is that the former can form a crystal, even if it is metastable, while the latter, due to its fivefold rotational symmetry which is not compatible with translational periodicity, cannot tile the space and form a crystal so that it leads to frustration in simple one-component liquids.⁸⁹

The stability range in size of clusters of Ice 0 with respect to ice I_c/I_h will depend on the particular water model adopted. For example, the mW model with respect to molecular models underestimates the lattice energy of Ice 0 relative to ice I_c/I_h .⁹⁰ However, since in our study the stability of Ice 0 with respect to ice I_c/I_h originates from topology (that is the number of bonds formed in a cluster), we expect that a different water model should give a similar result for the energy per particle difference between Ice 0 and ice I_c/I_h (that is, Ice 0 is more stable than ice I_c/I_h up to a specific cluster size), while the energy per particle of each ice could change with respect to the mW model.

Another aspect to take into account when considering fully atomistic realizations of clusters is the configuration of the hydrogen bond network. In the case of small clusters, the HB network rearrangements take place in the hydration layer, as verified for small aggregated domains in the case of an atomistic model of water in Ref. 91.

In conclusion, we find that even a tetrahedral liquidlike water can be characterized by competing crystalline clusters. In our case, clusters of Ice 0 are even more stable than isolated clusters of I_c/I_h up to a critical size of $n^* \sim 40$ for isolated clusters growing following the minimum energy rule (see Fig. 2), up to a critical size of $n_{eq}^* \sim 30$ and $n_{eq-hy}^* \sim 24$ following the same rule for equilibrated clusters and equilibrated and hydrated clusters, respectively (see Fig. 3), and up to a critical size of $n^* \sim 22$ for clusters growing following the closest distance rule (see Fig. 2).

B. Cluster growth and grain boundaries

In this section, we address the growth of different crystalline seeds, both from critical nuclei (with the seeding technique and umbrella sampling simulations) and as planar interfaces. We will show how the growth of each polymorph is associated with a specific coherent grain boundary. Ice I_h is hindered by stacking disorder formed via a twin boundary between its basal plane and the (111) surface of Ice I_c , giving rise to stacking disordered Ice I_{sd} . Ice I_c growth is instead hindered by a fivefold coherent grain boundary. Ice 0 has a coherent grain boundary between its (100) face and the (310) plane of cubic ice.

1. Growth of hexagonal ice

Nucleation of the stable Ice I phase in the mW model has been extensively addressed in previous studies.^{40,44,50,52,83,92–94} The

free energy difference between the two polytypes, cubic and hexagonal ice, was found to be very small, $\Delta G_{ch} = 2 \pm 1.5 \text{ J mol}^{-1}$ at $T = 200 \text{ K}$.⁴⁰ No free energy difference was indeed found in similar studies.^{20,95} The difference in nucleation barriers between an hexagonal and cubic nucleus at $T = 240 \text{ K}$ was also found to be negligibly small, $2 \pm 2 \text{ J mol}^{-1}$,⁹⁴ which is equivalent to a difference of about $1 k_B T$.

Figure 5 shows the result for our calculations of the nucleation barriers at $T = 218 \text{ K}$ with the umbrella sampling method detailed in Sec. II. The height of the nucleation barrier is the same ($\beta\Delta G = 27.2 \pm 0.25$, where $\beta = 1/k_B T$) for both a nucleus of Ice I_h and I_c . We see that the nucleation barrier difference between the two ices is indeed negligible at lower temperatures, and within the umbrella sampling method's precision.

Both Ice I_h and I_c are made of layers composed of six-membered rings and differ only in the way these layers are stacked. In Ice I_h , the oxygen positions are the same between all layers (parallel to the basal plane), while in Ice I_c , each layer is shifted by a distance equal to half the diameter of a ring. Given the low free energy difference between the cubic and hexagonal polytypes, the growth of ice I_h along the direction perpendicular to the basal plane is controlled by the free energy cost of introducing a twin grain boundary that changes the order of the stacking. It was recently pointed out that the presence of stacking disorder gives rise to a new crystal, ice I_{sd} which retains only a threefold rotational symmetry instead of the full sixfold symmetry of ice I_h .³⁸ The signature of stacking disorder was found in many X-ray and neutron diffraction patterns which were previously attributed to cubic ice,³⁸ leading to the idea that at low temperatures ice always nucleates as Ice I_{sd} rather than cubic ice.^{38,41} We will challenge this idea in Sec. III B 2.

The free energy cost of forming a stacking fault is offset by an increase in configurational entropy coming from the different ways to arrange the layers. Previous studies have addressed both these terms.^{40,44,94} The most recent estimates⁹⁴ put the stacking fault free

FIG. 5. Free energy barriers scaled by thermal energy $\beta\Delta G$ as a function of nucleus size n at $T = 218 \text{ K}$ for the nucleation of Ice I_h (red squares), I_c (black circles), and a nucleus of Ice I_{sd} with alternating layers with cubic and hexagonal stacking (green diamonds). The inset shows the barrier around its maximum.

energy per unit area between $\gamma_{sf} = 0 \text{ mJ m}^{-2}$ at $T = 180 \text{ K}$ and $\gamma_{sf} = 0.11 \text{ mJ m}^{-2}$ at the melting point.

We estimate the value of γ_{sf} from umbrella sampling simulations for an Ice I_{sd} nucleus in which the cubic and hexagonal layers alternate between each other (see, e.g., Fig. 9). The barrier is shown with green diamond symbols in Fig. 5. We emphasize that this is not the nucleation barrier for Ice I_{sd} , as it does not take into account the entropy contribution coming from different realizations of the stacking disorder. The curve in Fig. 5 is instead the nucleation barrier for a single realization of stacking disorder, from which we can estimate the cost of forming the twin boundary defect. The free energy difference between the stacking disordered nucleus and the hexagonal (or cubic) nucleus is shown in the inset of Fig. 5 and corresponds to a difference of $0.3 \pm 0.1 k_B T$. The estimation of the difference in free energies between stacking disordered and cubic and hexagonal ice was done by fitting the nucleation barriers with the classical nucleation theory expression in the range $60 < N < 120$ around the top of the barrier. The expression is $\Delta G = ax + bx^{2/3}$. We obtain maxima at $\Delta G = 27.6$ for the stacking disordered seed and $\Delta G = 27.3$ for both cubic and hexagonal seeds. The propagated error from fitting the classical nucleation theory expression is low (1%), but to account for systematic effects coming from the umbrella sampling method, we have increased this error to be of the same order of the deviation between barriers computed from independent trajectories, which is of the order of $0.1 k_B T$.

To the value $0.3 \pm 0.1 k_B T$, one would need to subtract the bulk free energy difference between the cubic and hexagonal crystals, but as one can see from the inset of Fig. 5, this difference falls below our resolution at $T = 218 \text{ K}$, so we simply set it to zero. We then measure the surface area between the hexagonal and cubic layers during the simulation and estimate the surface tension as $\gamma_{sf} = 0.16 \pm 0.05 \text{ mJ m}^{-2}$. Ignoring the bulk free energy difference makes our value an upper bound to the true surface penalty, so overall our estimate is in agreement with the results of Ref. 94.

2. Growth of cubic ice

We first consider unbiased simulations from cubic ice seeds following the seeding technique. We first determine the equilibrium densities of both the crystal and supercooled liquid phase and prepare crystalline nuclei at different system sizes. We then run isobaric simulations of $N = 10\,000$ water molecules at $T = 218 \text{ K}$ and find the size that is critical, where the nucleus has the same probability to either grow or shrink. The inset of Fig. 6 shows the time evolution of different runs for seeds of size $n = 110$ molecules, where half the trajectories (continuous blue lines) see the nucleus melting, and half the trajectories (lines with symbols) see the nucleus growing. We thus determine the size $n \sim 110$ to be critical, in excellent agreement with the umbrella sampling simulations shown in Fig. 5, and study the growth process of the initial ice I_c seed. Via bond orientational order parameters, we determine the fraction of cubic ice present in the nucleus and compute the *cubicity* parameter, defined as $f_{\text{cubic}} = n_{\text{cubic}} / (n_{\text{cubic}} + n_{\text{hexagonal}})$. The main panel in Fig. 6 plots the cubicity parameter as a function of the growth steps for the three trajectories that experience crystal growth, showing that cubicity steadily decreases during growth in all cases. These results confirm that at these sizes, crystal growth involves the formation of stacking faults.

FIG. 6. Fraction of molecules in a cubic ice environment as a function of simulation steps, for three independent runs at $T = 218 \text{ K}$. The inset shows the total amount of molecules in the largest nucleus as a function of time, both for three runs with growing nuclei (lines with symbols), and for three runs where the initial seed entirely melts (lines).

In order to study how the process of stacking disorder proceeds, in Fig. 7, we plot the $\langle 110 \rangle$ projection of the crystal growth process from an initial cubic seed [Fig. 7(a)]. In the growth stages [Figs. 7(b)–7(d)], we notice the development of stacking disorder. A crucial difference between Ice I_{sd} and Ice I_c is that the former has stacking disorder only in one direction, while the latter, due to its cubic symmetry, has a family of equivalent planes $\{1, 1, 1\} = (111), (\bar{1}\bar{1}\bar{1}), (1\bar{1}\bar{1}), (\bar{1}1\bar{1})$. The growth along two of such planes is highlighted in Fig. 7(d), where the order of the stacking is indicated with “c” for cubic arrangement and “h” for hexagonal arrangement.

The appearance of random stacking along different crystallographic directions poses a problem with cubic seeds trying to grow. As shown in Fig. 7, the interface between the two stacking directions is incompatible with crystalline order, i.e., a grain boundary forms. We observe the formation of such grain boundaries in all 10 trajectories that are started with a cubic seed.

While Ice I_{sd} only pays a small free energy penalty for stacking, ice I_c must additionally incur in the free energy penalty associated with the formation of grain boundaries. For incoherent boundaries, this penalty would be too high and cubic ice would then play no role in crystal nucleation. Instead, we always observe the formation of a coherent grain boundary with fivefold symmetry. This particular structure was first noted in Ref. 52. We can already get a hint of this process by considering the angle θ at which the $\{111\}$ planes meet in Fig. 7(d), which is $\theta = \cos^{-1}(1/3)$. We note that this angle, $\theta \sim 70.5^\circ$, is very close to the value 72° that corresponds to fivefold symmetry. The fivefold grain boundary is depicted in Fig. 8. It is a defect in which five cubic ice grains with hexagonal stacking on coplanar $\{111\}$ planes merge together in a coherent way.

It is important to note that this particular grain only appears as a consequence of stacking disorder of ice I_c and cannot originate when nucleation of Ice I_h or I_{sd} occurs instead (since there is only one stacking direction in these forms of ice). The formation

FIG. 7. $\langle 110 \rangle$ projection of a cubic ice seed composed of 48 water molecules (a). Oxygen atoms are connected with bonds and colored according to their phase: blue for cubic ice and orange for hexagonal ice. The different stages of growth are shown in (b) (115 molecules), (c) (484 molecules), and (d) (1092 molecules). The growth occurs by random stacking of new $\{111\}$ planes (dashed lines) from the original seed. The symmetry of the stacking along two directions forming an angle of $\cos^{-1}(1/3)$ is indicated with c for cubic and h for hexagonal.

of this grain boundary is thus an indirect piece of evidence toward nucleation of ice I_c compared to Ice I_h or I_{sd} . In the following, we will show that nucleation simulations indicate the presence of this structure in at least 50% of trajectories.

FIG. 8. Fivefold symmetric coherent grain boundary formed from cubic ice (blue atoms) with hexagonal stacking (orange atoms) on coplanar $\{111\}$ planes.

To study nucleation events without seeding, we use CNT-US simulations at $T = 218$ K. According to Eq. (1), the two unknown parameters that need to be determined are the driving force for crystallization, $\Delta\mu$, and the critical nucleus size, n_c . The first one is obtained via thermodynamic integration from the melting point of mW ice I, $T = 275.6$ K. The parameter n_c is instead determined as follows: several simulations at different values of n_c are run independently, and the fluctuations of the largest nucleus size are monitored. When n_c is close to the true value of the critical size, one observes large fluctuations of the size n , with the system sampling from small to large crystalline sizes in the course of the same simulation. From these simulations, one can reconstruct the free energy barrier and thus obtain the true value of n_c . We run 10 independent trajectories at the predetermined values of $\beta\Delta\mu$ and n_c .

Figure 9 shows representative configurations in which only nuclei of size $n \sim 300$ are plotted and colored according to their phase (orange for the hexagonal crystal and blue for the cubic crystal). The results reveal three possible outcomes of nucleation. In the top row, we see the formation of a fivefold symmetric grain boundary of Fig. 8. In the bottom row (left panel), we see nuclei with stacking disorder in at least two directions, which also originates from the cubic phase. Finally, the bottom row (right panel) shows nuclei with stacking in only one direction, which corresponds to Ice I_{sd} nucleation.

The abundance of stacking disorder in two or more directions, and, in particular, the presence of the fivefold symmetric grain, provides indirect evidence that nucleation indeed involves also the cubic

FIG. 9. Results from 10 independent runs, for ice nuclei grown in CNT-US simulations at $T = 218$ K. Nuclei correspond to sizes $n \sim 300$. The output is classified into three groups: (1) nuclei with fivefold symmetry grain; (2) cubic ice nuclei, in which stacking occurs in at least two-different directions; and (3) ice I_{sd} nuclei, in which stacking occurs only in one direction (i.e., the basal plane).

phase. The cubic phase is able to avoid paying the full free-energy cost of an incoherent grain boundary by forming coherent fivefold symmetric grain boundaries, in which the hydrogen bond network is only distorted and not disrupted. The fivefold symmetric grain boundary forms a coherent interface at the expense of some elastic energy. We will now attempt a first free energy estimate of the grain boundary cost.

We tried obtaining a free energy barrier for crystals containing the fivefold grain boundary, but the nucleus fluctuations always favored growth on one of the cubic sides, until eventually the grain boundary disappeared. This could be remedied by introducing additional biasing fields that stabilize the grain boundary, but here instead we use a simpler approach and leave a more detailed estimate for future studies.

We decrease the temperature to $T = 204$ K, in a regime where nucleation occurs spontaneously in the system. We then estimate the free energy barrier of nucleation via mean first passage theory.^{96,97} The mean first passage time $t_{fp}(n)$ is defined as the average time elapsed until the appearance of a nucleus of size n in the system. For homogeneous nucleation in the steady-state, the mean first passage time is given by

$$t_{fp}(n) = \frac{1}{2kV} \{1 + \text{erf}[c(n - n_c)]\}, \quad (2)$$

where k is the nucleation rate, n_c is the critical nucleus size, erf is the error function, and c is related to the curvature of the free energy barrier at its maximum (i.e., the Zeldovich factor), $c = \sqrt{\Delta G''(n_c)/k_B T}$.

We run 100 homogeneous nucleation trajectories at $T = 204$ K until the appearance of a supercritical nucleus. We then divide the trajectories into two sets depending on whether the nucleus has formed a fivefold symmetric grain boundary or not. To identify

the grain boundary, we use bond orientational order parameters to detect the atoms in the inner five-membered ring of the defect (depicted in cyan in Fig. 8). These atoms are identified when the following two conditions are satisfied: $Q_4 < 0.1$ and $Q_{12} > 0.2$. Both Q_4 and Q_{12} refer to coarse-grained bond orientational parameters⁹⁸ (see Ref. 2 for a review of these methods).

The computed mean first passage times for nuclei of size up to $n = 100$ are plotted in Fig. 10, distinguishing between nuclei with the fivefold grain (red squares) and without it (black circles). We notice a very clear distinction between the two different sets, with nuclei

FIG. 10. Mean first passage time as a function of nucleus size, n . The time is in arbitrary Monte Carlo sweeps. Symbols are measured averaging mean first passage times over 100 trajectories of homogeneous nucleation at $T = 204$ K, divided into two groups depending on whether they form the fivefold grain boundary or not. The lines are fits according to Eq. (2).

with the fivefold grain having shorter mean first passage times. This already reveals that the nucleation barrier for these states is lower compared to the nuclei without the grain. By using Eq. (2), we fit both curves and display the results as continuous lines in Fig. 10. The results for the fit are (the superscript 5f refers to structures with the fivefold grain boundary): $n_c^{5f} = 41$, $n_c = 44$, $k^{5f}/k = 1.44$, $c^{5f} = 0.040$, and $c = 0.038$. We note that both the critical size and the free energy curvature are about the same in both cases, while the main difference is in the nucleation rate. We can thus estimate the free energy difference between the nuclei with and without a fivefold grain boundary ΔG^{5f} via the following relation: $\beta \Delta G^{5f} = -\log(k^{5f}/k) = -0.37$ or $\Delta G^{5f} = -0.6$ KJ/mol. As discussed above, for small nuclei (here $n_c \sim 40$), the energetic cost of forming the fivefold grain boundary is very small, so the free energy advantage comes from the entropic gain in the coherent stacking in multiple directions. For much bigger nuclei, we expect the strain on the bonds far from the grain core to make ΔG^{5f} big and positive.

Overall, the abundance of fivefold grain boundaries in both direct nucleation and umbrella sampling simulations, and the absence of a significant free-energy penalty for small nuclei, shows that the nucleation of the cubic phase occurs distinctly and at least with the same probability of the nucleation of Ice I_h and I_{sd} . The free-energy cost of the fivefold grain boundary is expected to grow considerably with increasing nucleus size, due to the strain in the bonds far from the grain core, and this is likely to stop the stacking of cubic ice in multiple directions. It should thus be more advantageous, for cubic seeds above a certain size, to have stacking disorder in only one direction, thus converting them into Ice I_{sd} . This suggests another mechanism, besides just the difference in bulk free energy between the cubic and hexagonal phases, for the conversion of cubic ice in Ice I_{sd} that is governed by the size dependence of the free energy cost of the fivefold grain boundary.

3. Growth of ice 0

In this section, we consider the growth of Ice 0 nuclei with the seeding technique at $T = 208$ K. We run simulations with an Ice 0 seed and determine the critical size to be around $n_c = 400$ molecules. We then run 100 independent simulations and focus on the trajectories where the size of the nucleus grows. We observe two different scenarios, occurring with similar probability (number of trajectories that follow one growth mechanism or the other): (i) growth of a defect free ice 0 crystal and (ii) cross-nucleation of ice I_c on the ice 0 seed. A snapshot from a configuration following scenario (ii) is shown in Fig. 11. While the first scenario is the expected growth mode for a metastable nucleus, the observation of cross nucleation of cubic ice poses some interesting questions. First, cross nucleation at such small sizes implies that the surface penalty between ice 0 and cubic ice should be considerably low, which is unexpected given the differences between the two different crystalline structures. Second, the question remains why only the cubic form is cross-nucleated, and not the hexagonal phase.

To answer these questions, we run growth simulations with the direct coexistence method, where a slab of Ice 0 is prepared in contact with the supercooled liquid phase. In addition, in this case, we observe the cross-nucleation of defect free cubic ice from the (100)

FIG. 11. Seeding of an ice 0 nucleus at $T = 208$ K. Ice 0 particles are colored in green, and the cubic ice that cross nucleates on the surface of ice 0 is colored in blue.

plane of Ice 0. The (100) plane is the same that minimizes the cluster energy, as we described in Fig. 4.

In Fig. 12, we plot a snapshot from such configurations. (d) shows that a coherent grain boundary is formed between the face (100) of Ice 0 (b) and the (310) face of cubic ice (a). The two faces join coherently (without breaking of hydrogen bonds), as shown, for example, in (c). From these equilibrium simulations, we measure a very small strain (around 2%) of the cubic unit cell compared to its equilibrium value close to the interface. This signals a very low surface free energy cost of the defect. The cost of coherent grain boundaries between two solid surfaces of the same material (mW) originates almost entirely from the strain energy. We can estimate this cost as

$$\Delta E_{c0} = e_I - \frac{e_0 + e_c}{2},$$

where e_I is the energy of an interface atom, while e_0 and e_c are the energy of atoms in the bulk Ice 0 and I_c phases, respectively. We use our simulations to estimate these energies as $e_I = 94.71$ kJ/mol, $e_0 = 96.23$ kJ/mol, and $e_c = 97.18$ kJ/mol for $\Delta E_{c0} = 1.99$ kJ/mol. e_I is obtained by measuring the average energy of the molecules at the interface between ice 0 and cubic ice from the direct-coexistence simulations. Since the interface is coherent, it is very sharp (one particle width), and there is no ambiguity in the selection of the correct particles. γ_{c0} is obtained by assuming (as usually done for coherent interfaces) that the dominant contribution to the free energy is energetic (given by ΔE_{c0}) and then dividing the total surface energy for the area of the surface. This leads to a surface energy of stacking between the (100) plane of Ice 0 and the (310) plane of cubic ice of $\gamma_{c0} = 29.67$ mJ/m². This energy is much larger than the stacking surface energy of cubic and hexagonal layers in Ice I (that we estimated earlier as $\gamma_{sf} \sim 0.16$ mJ/m² at similar thermodynamic conditions), but is well below the limit of 200 mJ/m² that is conventionally used to separate coherent from semicoherent grains.

FIG. 12. Ice0/cubic interface. Surface projections of slabs of thickness 4.3 Å for the (310) plane of cubic ice (a) and the (100) plane of Ice 0 (b). (c) shows the coherent interface between the two crystals and (d) the direct coexistence box. The slab thickness corresponds to the range of the mW potential.

We observe that the low cost of the grain boundary also makes it difficult to obtain free energy barriers of Ice 0 (as shown for the other phases in Fig. 5), as we always observe the appearance of cubic ice in the umbrella sampling simulations.

One important consequence of this grain boundary is the fact that the (100) face of Ice 0 appears to be an ideal template for the cross nucleation of Ice I_c . Our simulations always show the absence of hexagonal stacking, and the formation of a crystal with cubicity of $c = 1$. The reason for this is that the grain boundary involves a high-order face of cubic ice, the (310) face, and so the new cubic phase grows along a direction incompatible with stacking faults.

IV. CONCLUSIONS

In this article, we have focused on crystalline clusters in the mW model of water. In the spirit of Frank's original argument, we have shown that there is a competition between different crystalline clusters and that the most energetic clusters at small sizes include five-membered rings. Here, we have demonstrated this with the Ice 0 structure, which is a metastable crystal for the mW potential. Five-membered rings allow small clusters to increase the ratio between number of bonds and number of particles in the cluster, without significant penalties to the bond energy. As the clusters grow in size, six-membered rings (as found in the stable Ice I forms) become favorable due to their optimal bond energy. Frustration effects between local structures can suppress the nucleation of crystals and are thus linked to glass-forming ability.^{2,8,99,100} The

competition between different local structures also plays an important role in polymorph selection, i.e., which metastable phase is nucleated.

To study the growth of crystalline clusters in supercooled conditions, we have used both direct simulations, seeding, and umbrella sampling biasing techniques. In particular, we have highlighted the role of grain boundaries in the growth process of the clusters. The lowest energy grain boundary is the stacking fault between hexagonal and cubic layers that can appear both on the basal plane of Ice I_h and the (111) plane of Ice I_c , which we estimate as $\gamma_{sf} = 0.16 \pm 0.05 \text{ mJ m}^{-2}$ at $T = 218 \text{ K}$. Given the negligible bulk free energy difference between I_c and I_h , this predicts, at low temperatures, the formation of Ice I_{sd} as the first stage of nucleation, favored by the configurational entropy associated with its disordered stacking sequence. However, our results also show that cubic ice still plays an important role in ice nucleation, thanks to another coherent grain boundary. Differently from Ice I_h and I_{sd} , cubic ices can generate stacking faults in four different directions. The associated cost of intersecting these stacking faults, which would otherwise suppress the nucleation of small cubic crystallites, is instead accommodated by the formation of a coherent fivefold symmetric grain boundary. At deep supercooled conditions ($T = 208 \text{ K}$), we estimate the free energy difference between nuclei with and without the fivefold grain boundary to be of the order of $0.37 k_B T$ in favor of the first ones. The abundance of configurations with the fivefold grain boundary is an indirect measure of the importance of the nucleation of cubic ice. Due to its fivefold symmetric nature, the grain boundary cost will increase rapidly with nucleus size, due to the bond strain far

from the core of the boundary. We can thus expect that the grain boundary will also be a contributing factor in selecting one growth direction for cubic ice, which would then gradually transform into Ice I_{sd}, in accordance with Ostwald's step rule.

We have also considered seeding simulations of Ice 0 nuclei and found that its growth is also controlled by a coherent grain boundary. The new grain boundary is formed between the (100) face of Ice 0 and the (310) face of cubic ice. We estimated the surface free energy of the grain boundary to be $\gamma_{c0} = 29.67 \text{ mJ/m}^2$. The fact that the grain boundary involves a high order face of cubic ice makes the (100) face of Ice 0 an ideal template for the cross-nucleation of pure cubic ice. Indeed, our simulations show growth of cubic ice without stacking faults.

ACKNOWLEDGMENTS

J.R. and F.L. acknowledge support from the European Research Council (Grant No. DLV-759187). J.R. acknowledges support from the Royal Society University Research Fellowship and travel support from the Institute of Industrial Science (IIS), University of Tokyo for a short stay at IIS. This study was partially supported by Grants-in-Aid for Specially Promoted Research (Grant No. JP25000002) and Scientific Research (A) (Grant No. 18H03675) from the Japan Society for the Promotion of Science (JSPS).

REFERENCES

- 1 F. C. Frank, *Proc. R. Soc. London, Ser. A* **215**, 43 (1952).
- 2 H. Tanaka, R. Shi, H. Tong, and J. Russo, *Nat. Rev. Phys.* **1**, 333 (2019).
- 3 H. Tanaka, *Eur. Phys. J. E* **35**, 23 (2012).
- 4 H. Shintani and H. Tanaka, *Nat. Phys.* **2**, 200 (2006).
- 5 M. Leocmach and H. Tanaka, *Nat. Commun.* **3**, 974 (2012).
- 6 C. P. Royall and S. R. Williams, *Phys. Rep.* **560**, 1 (2015).
- 7 C. P. Royall, F. Turci, S. Tatsumi, J. Russo, and J. Robinson, *J. Phys.: Condens. Matter* **30**, 363001 (2018).
- 8 J. Russo, F. Romano, and H. Tanaka, *Phys. Rev. X* **8**, 021040 (2018).
- 9 J. Russo and H. Tanaka, *J. Chem. Phys.* **145**, 211801 (2016).
- 10 H. Tanaka, *J. Chem. Phys.* **112**, 799 (2000).
- 11 J. Russo and H. Tanaka, *Nat. Commun.* **5**, 3556 (2014).
- 12 R. Shi, J. Russo, and H. Tanaka, *J. Chem. Phys.* **149**, 224502 (2018).
- 13 R. Shi, J. Russo, and H. Tanaka, *Proc. Natl. Acad. Sci. U. S. A.* **115**, 9444 (2018).
- 14 P. H. Poole, F. Sciortino, U. Essmann, and H. E. Stanley, *Nature* **360**, 324 (1992).
- 15 O. Mishima and H. E. Stanley, *Nature* **396**, 329 (1998).
- 16 P. Gallo and F. Sciortino, *Phys. Rev. Lett.* **109**, 177801 (2012).
- 17 P. Gallo, K. Amann-Winkel, C. A. Angell, M. A. Anisimov, F. Caupin, C. Chakravarty, E. Lascaris, T. Loerting, A. Z. Panagiotopoulos, J. Russo *et al.*, *Chem. Rev.* **116**, 7463 (2016).
- 18 P. H. Handle, T. Loerting, and F. Sciortino, *Proc. Natl. Acad. Sci. U. S. A.* **114**, 13336 (2017).
- 19 F. Martelli, *J. Chem. Phys.* **150**, 094506 (2019).
- 20 J. Russo, F. Romano, and H. Tanaka, *Nat. Mater.* **13**, 733 (2014).
- 21 A. V. Brukhno, J. Anwar, R. Davidchack, and R. Handel, *J. Phys.: Condens. Matter* **20**, 494243 (2008).
- 22 E. B. Moore and V. Molinero, *Nature* **479**, 506 (2011).
- 23 T. L. Malkin, B. J. Murray, C. G. Salzmann, V. Molinero, S. J. Pickering, and T. F. Whale, *Phys. Chem. Chem. Phys.* **17**, 60 (2015).
- 24 J. Shilling, M. Tolbert, O. Toon, E. Jensen, B. J. Murray, and A. K. Bertram, *Geophys. Res. Lett.* **33**, L17801, <https://doi.org/10.1029/2006gl026671> (2006).
- 25 L. Dowell and A. Rinfret, *Nature* **188**, 1144 (1960).
- 26 I. Kohl, E. Mayer, and A. Hallbrucker, *Phys. Chem. Chem. Phys.* **2**, 1579 (2000).
- 27 T. Loerting, I. Kohl, W. Schustereder, K. Winkel, and E. Mayer, *ChemPhysChem* **7**, 1203 (2006).
- 28 P. Geiger, C. Dellago, M. Macher, C. Franchini, G. Kresse, J. Bernard, J. N. Stern, and T. Loerting, *J. Phys. Chem. C* **118**, 10989 (2014).
- 29 G. Arnold, E. Finch, S. Rabideau, and R. Wenzel, *J. Chem. Phys.* **49**, 4365 (1968).
- 30 W. Kuhs, D. Bliss, and J. Finney, *J. Phys. Colloq.* **48**, C1 (1987).
- 31 T. Hansen, M. Koza, and W. Kuhs, *J. Phys.: Condens. Matter* **20**, 285104 (2008).
- 32 T. Hansen, M. Koza, P. Lindner, and W. Kuhs, *J. Phys.: Condens. Matter* **20**, 285105 (2008).
- 33 B. J. Murray and A. K. Bertram, *Geophys. Res. Lett.* **34**, L16810, <https://doi.org/10.1029/2007gl030471> (2007).
- 34 M. A. Carignano, *J. Phys. Chem. C* **111**, 501 (2007).
- 35 P. Pirzadeh and P. G. Kusalik, *J. Am. Chem. Soc.* **133**, 704 (2010).
- 36 D. Rozmanov and P. G. Kusalik, *J. Chem. Phys.* **137**, 094702 (2012).
- 37 M. Seo, E. Jang, K. Kim, S. Choi, and J. S. Kim, *J. Chem. Phys.* **137**, 154503 (2012).
- 38 T. L. Malkin, B. J. Murray, A. V. Brukhno, J. Anwar, and C. G. Salzmann, *Proc. Natl. Acad. Sci. U. S. A.* **109**, 1041 (2012).
- 39 S. Choi, E. Jang, and J. S. Kim, *J. Chem. Phys.* **140**, 014701 (2014).
- 40 D. Quigley, *J. Chem. Phys.* **141**, 121101 (2014).
- 41 W. F. Kuhs, C. Sippel, A. Falenty, and T. C. Hansen, *Proc. Natl. Acad. Sci. U. S. A.* **109**, 21259 (2012).
- 42 C. Desgranges and J. Delhommelle, *J. Am. Chem. Soc.* **128**, 10368 (2006).
- 43 A. H. Nguyen and V. Molinero, *J. Chem. Phys.* **140**, 084506 (2014).
- 44 L. Lupi, A. Hudait, B. Peters, M. Grünwald, R. Gotchy Mullen, A. H. Nguyen, and V. Molinero, *Nature* **551**, 218–222 (2017).
- 45 E. A. Engel, A. Anelli, M. Ceriotti, C. J. Pickard, and R. J. Needs, *Nat. Commun.* **9**, 2173 (2018).
- 46 M. Matsumoto, S. Saito, and I. Ohmine, *Nature* **416**, 409 (2002).
- 47 R. Radhakrishnan and B. L. Trout, *Phys. Rev. Lett.* **90**, 158301 (2003).
- 48 V. Molinero and E. B. Moore, *J. Phys. Chem. B* **113**, 4008 (2008).
- 49 E. B. Moore and V. Molinero, *J. Chem. Phys.* **132**, 244504 (2010).
- 50 E. B. Moore and V. Molinero, *Phys. Chem. Chem. Phys.* **13**, 20008 (2011).
- 51 D. Quigley and P. M. Rodger, *J. Chem. Phys.* **128**, 154518 (2008).
- 52 T. Li, D. Donadio, G. Russo, and G. Galli, *Phys. Chem. Chem. Phys.* **13**, 19807 (2011).
- 53 A. Reinhardt, J. P. Doye, E. G. Noya, and C. Vega, *J. Chem. Phys.* **137**, 194504 (2012).
- 54 A. Reinhardt and J. P. Doye, *J. Chem. Phys.* **139**, 096102 (2013).
- 55 J. Benet, L. G. MacDowell, and E. Sanz, *Phys. Chem. Chem. Phys.* **16**, 22159 (2014).
- 56 J. Benet, L. G. MacDowell, and E. Sanz, *J. Chem. Phys.* **141**, 034701 (2014).
- 57 B. Cheng, G. A. Tribello, and M. Ceriotti, *Phys. Rev. B* **92**, 180102 (2015).
- 58 J. R. Espinosa, C. Vega, and E. Sanz, *J. Phys. Chem. C* **120**, 8068 (2016).
- 59 J. R. Espinosa, A. Zaragoza, P. Rosales-Pelaez, C. Navarro, C. Valeriani, C. Vega, and E. Sanz, *Phys. Rev. Lett.* **117**, 135702 (2016).
- 60 M. Ambler, B. Vorselaars, M. P. Allen, and D. Quigley, *J. Chem. Phys.* **146**, 074701 (2017).
- 61 T. Koop and B. J. Murray, *J. Chem. Phys.* **145**, 211915 (2016).
- 62 B. Cheng and M. Ceriotti, *J. Chem. Phys.* **146**, 034106 (2017).
- 63 B. Cheng, G. A. Tribello, and M. Ceriotti, *J. Chem. Phys.* **147**, 104707 (2017).
- 64 L. Lupi, A. Hudait, and V. Molinero, *J. Am. Chem. Soc.* **136**, 3156 (2014).
- 65 A. Reinhardt and J. P. Doye, *J. Chem. Phys.* **141**, 084501 (2014).
- 66 S. J. Cox, S. M. Kathmann, B. Slater, and A. Michaelides, *J. Chem. Phys.* **142**, 184704 (2015).
- 67 G. C. Sosso, G. A. Tribello, A. Zen, P. Pedevilla, and A. Michaelides, *J. Chem. Phys.* **145**, 211927 (2016).
- 68 L. Lupi, B. Peters, and V. Molinero, *J. Chem. Phys.* **145**, 211910 (2016).
- 69 J. C. Johnston and V. Molinero, *J. Am. Chem. Soc.* **134**, 6650 (2012).
- 70 T. Li, D. Donadio, and G. Galli, *Nat. Commun.* **4**, 1887 (2013).
- 71 A. Hudait and V. Molinero, *J. Am. Chem. Soc.* **136**, 8081 (2014).

- ⁷²A. Haji-Akbari, R. S. DeFever, S. Sarupria, and P. G. Debenedetti, *Phys. Chem. Chem. Phys.* **16**, 25916 (2014).
- ⁷³S. M. Malek, G. P. Morrow, and I. Saika-Voivod, *J. Chem. Phys.* **142**, 124506 (2015).
- ⁷⁴A. Haji-Akbari and P. G. Debenedetti, *J. Chem. Phys.* **147**, 060901 (2017).
- ⁷⁵E. B. Moore, E. de la Llave, K. Welke, D. A. Scherlis, and V. Molinero, *Phys. Chem. Chem. Phys.* **12**, 4124 (2010).
- ⁷⁶E. Gonzalez Solveyra, E. de la Llave, D. A. Scherlis, and V. Molinero, *J. Phys. Chem. B* **115**, 14196 (2011).
- ⁷⁷G. Bullock and V. Molinero, *Faraday Discuss.* **167**, 371 (2013).
- ⁷⁸G. D. Soria, J. R. Espinosa, J. Ramirez, C. Valeriani, C. Vega, and E. Sanz, *J. Chem. Phys.* **148**, 222811 (2018).
- ⁷⁹M. Conde, M. Rovere, and P. Gallo, *Phys. Chem. Chem. Phys.* **19**, 9566 (2017).
- ⁸⁰L. Rovigatti, J. Russo, and F. Romano, *Eur. Phys. J. E* **41**, 98 (2018).
- ⁸¹B. C. Knott, V. Molinero, M. F. Doherty, and B. Peters, *J. Am. Chem. Soc.* **134**, 19544 (2012).
- ⁸²N. E. Zimmermann, B. Vorselaars, D. Quigley, and B. Peters, *J. Am. Chem. Soc.* **137**, 13352 (2015).
- ⁸³J. R. Espinosa, C. Vega, C. Valeriani, and E. Sanz, *J. Chem. Phys.* **144**, 034501 (2016).
- ⁸⁴J. Espinosa, C. Navarro, E. Sanz, C. Valeriani, and C. Vega, *J. Chem. Phys.* **145**, 211922 (2016).
- ⁸⁵Y. Lifanov, B. Vorselaars, and D. Quigley, *J. Chem. Phys.* **145**, 211912 (2016).
- ⁸⁶J. Russo, K. Akahane, and H. Tanaka, *Proc. Natl. Acad. Sci. U. S. A.* **115**, E3333 (2018).
- ⁸⁷S. Mossa and G. Tarjus, *J. Phys. Chem.* **119**, 8069 (2003).
- ⁸⁸P. J. Steinhardt, D. R. Nelson, and M. Ronchetti, *Phys. Rev. B* **28**, 784 (1983).
- ⁸⁹G. Tarjus, S. A. Kivelson, Z. Nussinov, and P. Viot, *J. Phys.: Condens. Matter* **17**, R1143 (2005).
- ⁹⁰D. Quigley, D. Alfè, and B. Slater, "Communication: On the stability of ice 0, ice i, and I_h," *J. Chem. Phys.* **141**, 161102 (2014).
- ⁹¹M. Matsumoto, A. Baba, and I. Ohmine, *J. Chem. Phys.* **127**, 134504 (2007).
- ⁹²A. Reinhardt and J. P. Doye, *J. Chem. Phys.* **136**, 054501 (2012).
- ⁹³A. Hudait, S. Qiu, L. Lupi, and V. Molinero, *Phys. Chem. Chem. Phys.* **18**, 9544 (2016).
- ⁹⁴B. Cheng, C. Dellago, and M. Ceriotti, *Phys. Chem. Chem. Phys.* **20**, 28732 (2018).
- ⁹⁵F. Romano, J. Russo, and H. Tanaka, *Phys. Rev. B* **90**, 014204 (2014).
- ⁹⁶J. Wedekind and D. Reguera, *J. Phys. Chem. B* **112**, 11060 (2008).
- ⁹⁷J. Russo, A. C. Maggs, D. Bonn, and H. Tanaka, *Soft Matter* **9**, 7369 (2013).
- ⁹⁸W. Lechner and C. Dellago, *J. Chem. Phys.* **129**, 114707 (2008).
- ⁹⁹P. Ronceray and P. Harrowell, *Phys. Rev. E* **96**, 042602 (2017).
- ¹⁰⁰D. Wei, J. Yang, M.-Q. Jiang, L.-H. Dai, Y.-J. Wang, J. Dyre, I. Douglass, and P. Harrowell, *J. Chem. Phys.* **150**, 114502 (2019).