Fry, C. (2020). The Lower Urinary Tract Science Group: ICI-RS. 5th June 2019. *Neurourology and Urodynamics*, 39(S3).
<https://doi.org/10.1002/nau.24209>

Peer reviewed version

Link to published version (if available):
[10.1002/nau.24209](https://doi.org/10.1002/nau.24209)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/abs/10.1002/nau.24209?af=R> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Record of Meeting at Bristol University, UK

Co-ordinator CH Fry. School of Physiology, Pharmacology and Neuroscience, University of Bristol, UK (chris.fry@bristol.ac.uk)

High frequency stimulation of the pelvic nerve or S1 sacral nerve root to inhibit urinary voiding - which is best?

Thelma Lovick

School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol BS8 1TD, UK

Introduction and Aims

‘On-demand’ high frequency (1-3kHz) stimulation of the pelvic nerve (PN) in rats inhibits imminent voids^{1,2} and could be an attractive novel therapeutic strategy for controlling urinary urge incontinence in humans. Whilst PN stimulation is effective, translation to the clinic is limited by the invasive surgery required. We therefore investigated whether stimulation of the sacral nerve root containing pelvic nerve afferents, might be a preferable target.

Methods

Urethane-anaesthetised female Wistar rats were instrumented for continuous cystometry and stimulation of the pelvic preganglionic nerve bundle (PN) and S1 spinal nerve root; the level at which PN afferents enter the cord in rats. Saline was infused into the bladder (6ml hr⁻¹) to evoke repeated voids.

Results

When initiated at the onset of the steep rise in bladder pressure signalling an imminent void, stimulation of PN or S1 (1-3kHz, 1mA for 60s) suppressed the void. Whilst stimulation at both PN and S1 was effective, PN stimulation was accompanied by transient rises in blood pressure and abdominal pressure and contraction of abdominal muscles, which were not evoked by S1 stimulation.

Conclusions

High frequency stimulation at PN and S1 are both effective in suppressing imminent voids. However, ease of surgical access to S1 and minimal side-effects produced by stimulation compared to PN, combined with extensive clinical experience with sacral neuromodulation techniques is preferable. This indicates that high frequency S1 stimulation may be a better target for developing suppression of unwanted voids ‘on-demand’ to manage urinary urge incontinence.

Supported by MRC. Grant G1002251 and EPSRC-NIHR HTC Partnership award (IMPRESS) EP/M000109/1

References

1. Crook JJ, Lovick TA. High frequency stimulation of the pelvic nerve inhibits urinary voiding in anaesthetized rats. *Front Physiol* 2017; 8: 623.
2. Brouillard CBJ, Crook JJ, Irazoqui PP et al. Suppression of urinary voiding by conditional high frequency stimulation of the pelvic nerve in conscious rats. *Front Physiol* 2018; 9: 437.

Neural network analysis to assist differentiation of male outlet obstruction from underactivity

Li Rui¹, Gammie Andrew², Zhu Quanmin¹, Nibouche Mokhtar¹, Chen Chen¹

¹Faculty of Environment and Technology, University of the West of England. Bristol, BS16 1QY, UK.

²Bristol Urological Institute, Southmead Hospital, Bristol BS10 5 NB, UK

Introduction and Aims

Detrusor underactivity (DU) and bladder outlet obstruction (BOO) are two prevalent urinary dysfunctions in males. Currently the only standardised methods for diagnosis is pressure-flow studies, but the similarity of low a flow rate in each means that non-invasive methods cannot differentiate between the two conditions. Several studies have indicated that uroflowmetry parameters could serve as additional indicators for differentiation, however poor accuracy in individual assessments hampers their possible clinical use. In this analysis, an artificial neural network is assessed by employing non-invasive parameters to differentiate between DU from BOO.

Methods

The parameters employed are same as in the previous statistical model¹. The neural network was designed with one hidden layer to avoid overfitting, and the number of hidden neurons was chosen by five cross-fold validations. The data cohort contained free-flow data in 293 males, 135 with DU and 158 with BOO. These were divided randomly into 70% for training, 15% for validation and 15% for testing.

Results

The designed neural network yielded 79.5% overall diagnosing accuracy with 75.6% sensitivity and 82.9% specificity to differentiate between DU with BOO. The diagnosing accuracies in training, validation and testing were 79%, 77.3% and 84.1% respectively.

Conclusions

The results show neural network analysis is more robust compared to other linear models, for instance multivariate analysis. This initial trial employed a feed-forward neural network to assess its ability to assist non-invasive differentiation between DU from BOO by employing free-flow derived parameters. Further analysis will increase the data number, assess convolutional or feed-forward neural networks, and include data from normal individuals for non-invasive diagnosing.

Reference

1. Li R, Gammie A, Zhu Q, et al. Multivariate analysis of variance for maximising the diagnosing accuracy in differentiating DU with BOO in male. *Neurourol Urodyn* 2018; 21(S5): 327-328.

Urofacial syndrome and innervation of the bladder outflow.

Imerjit Manak¹, Alison Gurney², Adrian Woolf^{1,3}, Neil Roberts¹.

¹Division of Cell Matrix Biology and Regenerative Medicine, School of Biological Sciences, University of Manchester, M13 9PL, UK. ²Division of Pharmacy and Optometry, School of Medical Sciences, University of Manchester, M13 9PL, UK. ³Royal Manchester Children's Hospital, Manchester University NHS Foundation Trust, Manchester, M13 9WL, UK.

Introduction and hypothesis

Urofacial syndrome is an autosomal recessive disease characterised by bladder dysfunction with a high-pressure bladder and outflow dyssynergia. If untreated it leads to kidney failure due to urine reflux from the bladder to the kidney. We found that individuals with urofacial syndrome carry homozygous null mutations in either *heparanase 2 (HPSE2)* or *leucine-rich repeats and immunoglobulin-like domains 2 (LRIG2)*. Until recently, nothing was known about the pathophysiology underlying urofacial syndrome. The aim of this study was to define the bladder defect in mouse models of the disease.

Methods

Analysis of *Hpse2* and *Lrig2* homozygous mutant mice. Voiding stain on filter paper analysis of mouse voiding patterns. Whole tissue bladder immunostaining and confocal imaging; bladder body and outflow physiological analyses using isolated tissue myography with chemical and electrical field stimulation.

Results

The genetic mouse models of urofacial syndrome had urination defects resembling those in human patients. Immunostaining whole bladders revealed a patterning defect of the nerves that control urinary voiding.¹ Physiological analysis of the bladder body revealed a hypercontractile response to the muscarinic receptor agonist carbachol, with no change to relaxation profiles no relaxation of contraction defect was observed in the outflow.

Conclusion

Collectively, these results provide compelling evidence that a peripheral neuropathy underlies bladder dysfunction in urofacial syndrome. Further investigation is required to reveal whether the hypercontractile muscarinic response in the body is a primary or secondary phenomenon, and whether the neuropathy affects the coordination of bladder body contraction and outflow relaxation.

Supported by Kidney Research UK Grant PDF_005_20151126

Reference

1. Roberts NA, Hilton EN, Lopes FM et al. *Lrig2* and *Hpse2*, mutated in urofacial syndrome, pattern nerves in the urinary bladder. *Kidney Int.* 2019; 95: 1138-1152

A proposal for non-invasive abdominal straining measurement combined with uroflowmetry

Andrew Gammie

Clinical Engineer, Bristol Urological Institute, Southmead Hospital, Bristol BS10 5NB, UK

Introduction and Hypothesis

Differentiating bladder outlet obstruction (BOO) from detrusor underactivity (DU) in symptomatic men requires invasive urodynamics. An analysis of a large urodynamic database [1] showed a significant proportion of underactive patients describe straining during voiding, whereas an earlier study [2] showed obstructed men gain little benefit from straining. We therefore aim to examine whether the effect of abdominal straining on urine flow patterns can diagnose each group noninvasively. We hypothesise that an obstructed man will not by straining vary the urine flow greatly, due to the strain increasing the obstructive effect of an enlarged prostate.

Methods

Thirty-two male patients (16 BOO, 16 DU) will be recruited and asked to void twice, once while not straining and once while straining. During each void, abdominal pressure will be measured to identify the straining episodes, using a rectally placed water-filled catheter. In addition, simultaneous measurement with a tocodynamometer and abdominal EMG surface electrodes will be made, to identify whether these methods will reliably and acceptably quantify the strains, so future studies can be done without the rectal catheter. Comparison will be made of the ability of each method to detect straining and the variations in urine flow caused by straining. Ethical approval has been granted for the study.

Conclusions

This study will test the hypothesis that straining during voiding affects urine flow differently, depending upon whether the male patient is obstructed or underactive. The prospect of less invasive, quicker and cheaper diagnosis will benefit both patients and healthcare providers.

References

1. Gammie A, Kaper M, Dorrepaal C, Kos T, Abrams P. Signs and symptoms of detrusor underactivity: an analysis of clinical presentation and urodynamic tests from a large group of patients undergoing pressure flow studies. *Eur Urol* 2016; 69:361-9.
2. Reynard JM, Peters TJ, Lamond E, Abrams P. The significance of abdominal straining in men with lower urinary tract symptoms. *Br J Urol* 1995; 75:148-53.

Expression of a transcription factor protein in human prostate cancer.

Callum Arthurs¹, Rui Henrique², Michael Millar³, Fiona Inglis³, Aamir Ahmed¹

¹Prostate Cancer Research Centre, Centre for Stem Cells and Regenerative Medicine, Guy's Hospital, London SE1 9RT, UK. ²Department of Pathology and Molecular Immunology, Abel Salazar Institute of Biomedical Sciences, University of Porto, Portugal. ³Queen's Medical Research Institute, University of Edinburgh, Edinburgh, EH16 4TJ, UK.

Introduction and Aims

Prostate cancer (PCa) is the most commonly diagnosed cancer in UK men and Early detection is key to reducing mortality rates. Therefore, there is a need to develop quantitative and reliable clinical biomarkers for its early detection and prognostication. We have previously shown that the Wnt signalling pathway is dysregulated in PCa.¹ In this study we investigated the expression of the transcription factor, Pygopus-1, chosen for its association with the Wnt pathway. Transcription factor activation is a key event in carcinogenesis. Here we evaluate the performance of Pygopus-1 as a biomarker of PCa.

Methods

Tissue arrays (TA) were constructed from human archival formalin-fixed paraffin embedded samples, containing areas identified as PCa and PCa-adjacent by an expert pathologist. Using automated immunohistochemical (IHC) techniques,² 6µm sections were stained for Pygopus-1 by labelling with 3,3'-Diaminobenzidine (DAB). Slides were scanned using a bright field slide scanner (Hammamatsu) and tissue core images were exported from the scanned slide for quantification.

Results

DAB labelling of Pygopus-1 revealed expression of the transcription factor in both PCa and PCa-adjacent tissue cores. In PCa tissue cores, staining appeared to localise to the acinar cells. Quantification of DAB staining revealed an increase in protein expression of Pygopus-1 in prostate cancer tissue cores ($P < 0.0001$, Mann Whitney U).

Conclusions

Initial findings indicate that Pygopus-1 could play a role as a putative biomarker of PCa. Furthermore, a biomarker analysis pipeline is being developed based on the principle of measuring protein expression on stained TA.

References

- 1 Symes AJ, Eilertsen M, Millar M, Nariculam J, Freeman A, Notara M, Feneley MR, Patel HR, Masters JR, Ahmed A. Quantitative analysis of BTF3, HINT1, NDRG1 and ODC1 protein over-expression in human prostate cancer tissue. PLoS One 2013; 8: e84295.
- 2 Arthurs C, Murtaza BN, Thomson C, Dickens K, Henrique R, Patel HRH, Beltran M, Millar M, Thrasivoulou C, Ahmed A. Expression of ribosomal proteins in normal and cancerous human prostate tissue. PLoS One 2017; 12: e0186047.

Connexin 43 (Cx43) and Wnt signaling

Mohammad Khan¹, Xiaoming Hou^{2¶}, Mark Turmaine³, Christopher Thrasivoulou⁴, David Becker⁵, Aamir Ahmed^{1,2}

¹Prostate Cancer Research Centre, Centre for Stem Cells and Regenerative Medicine, Guy's Hospital, London SE1 9RT, UK; ² Prostate Cancer Research Centre, University College London, London W1W 7EJ, UK; ³Division of Biosciences, University College London, London, WC1E 6BT, UK; ⁴The Centre for Cell and Molecular Dynamics, University College London, London WC1E 6JJ, UK; ⁵Nanyang Technological University, Mandalay Road, Singapore, 308232.

Introduction and Hypothesis

Wnt signalling plays a key role during development and disease. The transcription factor β -catenin, intracellular Ca^{2+} are major transducers of Wnt signalling. Wnt signalling activation releases β -catenin which translocates to the nucleus. Connexin 43 (Cx43), a member of the gap junction family is a target of Wnt-induced transcription and regulation.¹ However, the inter-relationship between Wnt signaling and Cx43 protein remains unknown. We hypothesised that Wnt signalling activation modulates Cx43 mobilisation in PC3, a prostate cancer cell line.

Methods

Cell fractionation of PC3 cells with or without Wnt5A was performed using a Qproteome cell compartment kit (Qiagen). Cell fractions were loaded onto an SDS PAGE gel and probed with Cx43 antibody using Western blotting;² signals were quantified with ImageJ.

Results

Figure 1 shows an increase of Cx43 signal in the nuclear and membrane fractions after Wnt treatment. Similar results were obtained for another Wnt ligand, Wnt 9B.

Figure 1. Quantification of the intensity of Cx43 expression signal from Western blots in PC3 cells with Wnt5A treatment (black bars) and in control untreated cells (white bars); A.U., arbitrary units.

Conclusion

In response to Wnt signalling activation triggers intracellular mobilisation of Cx43, in a similar manner to translocation of β -catenin and may suggest a cross-talk between Wnt signalling and gap junction proteins.

References

- 1 van der Heyden MA, Rook MB, Hermans MM, Rijksen G, Boonstra J, Defize LH, Destree OH. Identification of connexin43 as a functional target for Wnt signalling. J Cell Sci 1998; 111: 1741-1749.
- 2 Hou X, Khan MRA, Turmaine M, Thrasivoulou C, Becker DL, Ahmed A. Wnt signalling regulates cytosolic translocation of connexin 43. Am J Phys Regul Integr Comp Phys 2019; doi: 10.1152/ajpregu.00268.2018

Psychological stress and bladder function

Russ Chess-Williams, Eliza West, Donna Sellers, Catherine McDermott

Centre for Urology Research, Bond University, Gold Coast, QLD 4226, Australia

Introduction and Hypothesis: Stress greatly influences the development of bladder symptoms,¹ but the mechanisms involved are little understood. This study investigates the hypothesis that psychological stress in mice induces bladder dysfunction via altered contractile mechanisms.

Methods: Female mice were placed on a pedestal surrounded by water for 1hr/day for 10 days, to induce water avoidance stress (WAS). Controls were not exposed to stress. 24-hours after the final stress exposure, or following 10 days recovery, animals were euthanised, a blood sample taken for a corticosterone assay and whole bladders were isolated, catheterised and intravesical pressure recorded.

Results: Plasma corticosterone levels and voiding frequency were increased in the WAS group compared to controls (Figure 1). Bladders from stressed mice showed greater contractility in response to carbachol ($p < 0.05$) and to the purinergic agonist ATP (1mM, $p < 0.05$) compared to controls. Voiding frequency was reduced following 10-days stress-free recovery (Figure 1B). This was accompanied by an increase in bladder compliance (Figure 1C), which had been unchanged during the stress protocol.

Figure 1: A) Plasma corticosterone levels in control and WAS mice; B) number of voiding events during a 4-hour observation and C) bladder compliance in control, WAS and Recovery mice.

Conclusions: Repeated exposure to environmental stress produces a hormonal stress response and an overactive bladder phenotype. A stress-free recovery period reduced voiding frequency, but this was due to compensation via increased bladder compliance, rather than a reversal of the stress-induced changes.

Reference

1. Rothrock NE, Lutgendorf SK. Kreder KJ, Rafliff T, Zimmerman B. Stress and symptoms in patients with interstitial cystitis: a life stress model. *Urology* 2001; 57: 422– 427.

Heterogeneity of PDGFR α (+) cells in the urogenital system

Hikaru Hashitani¹, Retsu Mitsui¹, Richard Lang²

¹Department of Cell Physiology, Nagoya City University, Nagoya 4678601, Japan., ²Department of Physiology, Monash University, Clayton 3800, Australia

Introduction and Aims

PDGFR α (+) cells in the gastrointestinal tract are considered to suppress muscle excitability by generating hyperpolarising signals¹. PDGFR α (+) cells are also distributed in the bladder and generate SK3 outward currents in response to P2Y1 stimulation². However, a functional coupling of PDGFR α (+) cells with detrusor smooth muscle (DSM) cells has not been demonstrated. We have examined whether PDGFR α (+) cells in other regions of the urogenital system have similar properties.

Methods

PDGFR α -GFP mice were used and the distribution of PDGFR α (+) cells and their SK3 expression were investigated. Ca²⁺ dynamics in PDGFR α (+) cells were also visualized using Cal-590 fluorescence.

Results

In the bladder, PDGFR α (+) cells in DSM but not mucosal layer expressed SK3 immunoreactivity. PDGFR α (+) cells in the both layers develop P2Y1-mediated Ca²⁺ transients. However, these Ca²⁺ transients were not associated with inhibition of Ca²⁺ transients in DSM cells. In the renal pelvis, atypical smooth muscle cells (ATSMCs) expressed weak PDGFR α -GFP fluorescence compared with two other distinct populations of PDGFR α (+) cells, but appear not to be directly activated upon P2Y1-stimulation. In the seminal vesicle, PDGFR α (+) cells in the musculature but not mucosa expressed SK3 immunoreactivity, while PDGFR α (+) cells in both layers responded to P2Y1-stimulation.

Conclusions

PDGFR α (+) cells in the urogenital system display a heterogeneity in terms of their SK3 expression and P2Y1 response. There was no evidence that PDGFR α (+) cells suppress the excitability of DSM cells, while PDGFR α (+) ATSMCs generate depolarising rather than hyperpolarising signals.

Supported by MRC. Grant-in-Aid for Scientific Research (C) (No. 17K11187) from JSPS.

References

1. Kurahashi M, Zheng H, Dwyer L, Ward SM, Koh SD, Sanders KM. A functional role for the 'fibroblast-like cells' in gastrointestinal smooth muscles. *J Physiol.* 2011; 589: 697-710.
2. Lee H, Koh BH, Peri LE, Sanders KM, Koh SD. Purinergic inhibitory regulation of murine detrusor muscles mediated by PDGFR α + interstitial cells. *J Physiol.* 2014; 592: 1283-93.

The distribution of aquaporins in the pig bladder wall and its relevance to water movements

Marian Manso¹, Christopher Fry², Marcus Drake³, Bahareh Vahabi¹

¹Faculty of Health and Applied Sciences, University of the West of England. Bristol, BS16 1QY, UK.

²School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol BS8 1TD, UK.

³School of Medicine, University of Bristol, BS8 2PL, UK.

Introduction and Aim

Bladder urothelium expresses aquaporins (AQPs) with AQP1-4, AQP7, AQP9 and AQP11 found in various species suggesting they could regulate urothelial cell volume and water transport.^{1,2} The aim was to investigate AQP expression and function in porcine bladder, an alternative to human bladder.

Methods

Mucosa (urothelium/suburothelium) was dissected from female pig (*Sus scrofa*) bladders (≈6 months) obtained from the local abattoir. AQP1-11 transcription and protein expression were tested by qPCR and Western blot. Localisation was by immunohistochemistry, using an avidin-biotin-peroxidase system. Water (deuterium oxide, D₂O) apical-to-basolateral flux was measured across a mucosa sheet in an Ussing chamber, ±the non-selective AQP inhibitor, HgCl₂ (300 μM), starting apical concentration 40% v/v. Data are mean±SEM, significant (p<0.05) differences between sets were determined by Student's *t*-tests.

Results

AQP1,3,9,11 mRNA and protein expression were found in the mucosa, with AQP3,9,11 localised to urothelium and AQP1 to blood vessels. There was a constant apical-basolateral D₂O flux when the apical side was exposed to hypertonic or hypotonic solutions. The average diffusion rate for D₂O in the absence of HgCl₂ (0.41±0.06 D₂O%.hr⁻¹.cm⁻², *n*=9) was significantly (p<0.01) greater than with HgCl₂, 0.20±0.02 D₂O%.hr⁻¹.cm⁻²), with flux rates greater in hypotonic compared to hypertonic media.

Conclusions

The demonstration of AQPs in pig bladder urothelium and a significant HgCl₂-sensitive water flux, dependent on the osmotic gradient, shows a role in regulating net water flux from urine to blood. The magnitude of water flux and if this can significantly alter bladder urine osmolality remains to be determined.

Supported by a fellowship from Ferring Pharmaceuticals

References

- 1 Rubenwolf PC, Georgopoulos NT, Kirkwood LA et al. Aquaporin expression contributes to human transurothelial permeability in vitro and is modulated by NaCl. PLoS One 2012; 24: 7: e45339
- 2 Spector DA, Wade JB, Dillow R et al. Expression, localization, and regulation of aquaporin-1 to -3 in rat urothelia. Am J Physiol Renal Physiol 2002; 282: F1034-1042.

Bladder contractile function in children with congenital bladder anomalies

Navroop Johal¹, Peter Cuckow¹, Christopher Fry²

¹Dept Urology, Great Ormond St Hospital for Sick Children, London, WC1N 3JH, UK ²School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol BS8 1TD, UK.

Introduction.

Several congenital anomalies are associated with poor bladder function, even after successful surgical correction of the defect, with life-long consequences. The causes of diminished contractile function are unknown and we investigated using bladder tissue biopsies retrieved during corrective surgery.

Methods.

Biopsies were obtained from children with: obstructed bladders from posterior urethral valves ($n=7$); exstrophy ($n=11$) or myelomeningocele ($n=12$). A control group were children with normal bladder function. All tissue was obtained with informed consent and ethical committee approval. Isolated detrusor strips were used to record isometric tension and for histology to measure smooth muscle and connective tissue (van Gieson stain).¹ Intracellular Ca^{2+} was measured in isolated myocytes with the fluorochrome Fura-2

Results.

Contractile function was reduced to a similar extent in all pathology groups compared to control, whether force was generated by EFS or 1 μM carbachol. This was mirrored by an increase of the connective tissue to smooth muscle ratio. With detrusor myocytes intracellular Ca^{2+} responses to carbachol or high-K solutions were similar in all pathology and control groups. Atropine-resistance of EFS contractions was similar in all groups.

Discussion and Conclusions. Reduced contractile function in all three congenital anomalies was mirrored by a reduced smooth muscle content in detrusor samples. However, similar intracellular Ca^{2+} responses to contractile agonists implies no loss of intrinsic detrusor contractility. Data are consistent with increased passive mechanical stiffness in exstrophy tissue.¹ The ubiquitous presence of atropine resistance suggests it is normal in neonatal/ paediatric bladders, as this declines in normal bladders toward adolescence.²

Supported by the Royal College of Surgeons of England and the Children's Research Fund, Liverpool.

References.

- 1 Johal NS, Arthurs C, Cuckow P, Cao K, Wood DN, Ahmed A, Fry CH. Functional, histological and molecular characteristics of human exstrophy detrusor. *J Pediatr Urol* 2019; 15: 154.
- 2 Johal N, Wood DN, Wagg AS, Cuckow P, Fry CH. Functional properties and connective tissue content of pediatric human detrusor muscle. *Am J Physiol Renal Physiol* 2014; 307: F1072-1079.

Effects of soluble guanylyl cyclase activators on contractile performance

Basu Chakrabarty¹, Marcus Drake², Anthony Kanai³, Christopher Fry¹

¹School of Physiology, Pharmacology, and Neuroscience, University of Bristol, Bristol BS8 1TD UK.

²Bristol Medical School, University of Bristol, UK ³Departments of Medicine & Pharmacology and Chemical Biology, University of Pittsburgh, USA

Introduction and Aims:

Phosphodiesterase type 5 (PDE5) inhibitors like sildenafil alleviate lower urinary tract symptoms¹. PDE5 inhibitors block cGMP hydrolysis and recently sildenafil was shown to inhibit the purinergic components of nerve-mediated contractions². Soluble guanylyl cyclase (sGC) catalyses cGMP synthesis. The aim of this study was to assess the effects of an sGC activator BAY 58-2667 on nerve-mediated contractions and ATP release.

Methods: Isolated intact bladder strips from 12-week old C57BL/6 male mice were tied to isometric force transducers. Nerve-mediated contractions were generated by electrical field stimulation (EFS: 0.1 ms pulses, 1-40 Hz, 3-s train every 90-s), and force-frequency relationships were generated. Nerve-mediated ATP release was measured using a luciferin-luciferase assay². Data are means±SD.

Results: The addition of BAY 58-2667 (10μM, n=6) resulted in the reduction of nerve-mediated contractions (two-way ANOVA, $p<0.05$), from 0.64 ± 0.18 mN.mg⁻¹ to 0.40 ± 0.14 mN.mg⁻¹ at 2Hz, and from 1.57 ± 0.57 mN.mg⁻¹ to 1.61 ± 0.63 mN.mg⁻¹ at 20Hz, with an increase in $f_{1/2}$ value from 2.97 ± 1.98 Hz to 5.16 ± 1.48 Hz (Student's paired t-test, $p<0.01$). BAY 58-2667 reduced nerve-mediated ATP release (two-way ANOVA, $p<0.001$), from 102.3 ± 14.5 pmoles to 59.2 ± 12.0 pmoles at 8Hz stimulation.

Conclusions: The data are consistent with the hypothesis that an increase in cGMP levels in detrusor preparations reduces nerve-mediated contractions and ATP release. Nerve-mediated ATP release is a feature of detrusor contractility from overactive human bladder. The attenuation of the purinergic components of nerve-mediated contractions by sGC activators suggests that modulating cGMP in efferent nerve terminals may be a target to reduce overactive bladder contractions.

Supported by the United States National Institutes of Health Grant NIH R01 DK098361.

References

1. McVary KT, Monnig W, Camps JL, Young JM, Tseng LJ, van den Enge G. Sildenafil citrate improves erectile function and urinary symptoms in men with erectile dysfunction and lower urinary tract symptoms associated with benign prostatic hyperplasia: A randomized, double-blind trial. *J Urol* 2007; 177: 1071-1077.
2. Chakrabarty B, Ito H, Ximenes M, Nishikawa N, Vahabi B, Kanai AJ, et al. Influence of sildenafil on the purinergic components of nerve-mediated and urothelial ATP release from the bladder of normal and spinal cord injured mice. *Br J Pharm* 2019; 176: 2227-2237.

Investigation of P2X7R physiology in urothelial cells

Breen C, Srivastava K, McCloskey KD.

Centre for Cancer Research and Cell Biology, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, 97 Lisburn Road, Belfast, BT9 7AE, Northern Ireland, UK.

Introduction and Hypothesis

The bladder expresses a number of purinergic receptors including P2X7R (1,2). P2X7R functions as a non-selective ion channel for influx of Na⁺, Ca²⁺ and K⁺. P2X7R activation is also associated with formation of large pores, enabling transport of larger molecules. Urothelial cells express P2X7R however, little is known of their physiology. Here, we hypothesised that P2X7R were functional ion channels in urothelial cells.

Methods

Immortalised human urothelial cells, SVHUC, were used in physiological experiments including patch-clamp and fluorescence Ca²⁺-imaging. P2X7R knockdown was conducted via siRNA molecules and confirmed by Western blotting. YO-PRO-1 dye-uptake experiments were performed to determine whether ATP could evoke pore formation.

Results

SVHUC were patch-clamped with Cs⁺-filled electrodes, in Hank's physiological solution at room temperature. Application of ATP (1 mM, to activate P2X7R), evoked inward currents in cells held at -60mV. Inward currents were also evoked by a selective P2X7R agonist, BzATP (0.5mM). SVHUC exhibited Ca²⁺-transients when exposed to ATP (1 mM) that were reduced by the P2X7R antagonist, A740003 (10 µM). siRNA knockdown markedly reduced P2X7R protein expression; these cells had smaller ATP-evoked Ca²⁺-responses compared to scrambled controls. Dye uptake experiments indicated that ATP did not evoke pore formation in SVHUC in normal Hank's solution.

Conclusions

The findings support the hypothesis that urothelial cells express P2X7R which function as ion channels when activated with ATP or BzATP. There was no evidence indicating that under these conditions, P2X7R formed large pores in SVHUC.

Funding from the Medical Research Council (MR/M012425/1) is acknowledged.

References

- Dunton CL, Purves JT, Hughes FM, Jr., Jin H, Nagatomi J. Elevated hydrostatic pressure stimulates ATP release which mediates activation of the NLRP3 inflammasome via P2X4 in rat urothelial cells". *Int Urol Nephrol* 2018; 50: 1607-1617.
- Negoro H, Urban-Maldonado M, Liou LS, Spray DC, Thi MM, Suadicani SO. Pannexin 1 channels play essential roles in urothelial mechanotransduction and intercellular signaling". *PloSOne* 2014; 9: e106269.

Spontaneous ATP transients in guinea-pig detrusor muscle

Christopher Fry¹, Carly McCarthy²

¹School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol BS8 1TD, UK.

²Facultad de Ciencias Biomédicas, Austral University, Argentina

Introduction and Hypothesis or Aims

Spontaneous contractions are a feature of the urinary bladder and are greater in pathological conditions, but their origins are unknown. We tested the hypothesis that a contributor is spontaneous release of ATP from within the bladder wall

Methods

Isolated guinea-pig detrusor muscle strips, with the mucosa removed, were superfused with Tyrode's solution at 36°C and tied to an isometric force transducer. An amperometric enzyme-based ATP electrode was placed on the surface of the strip, with a null (enzyme-free) electrode as reference (Sarissa, UK). Electrical field-stimulation (EFS) was used to elicit nerve-mediated contractions. Interventions were added to the superfusate.

Results

EFS generated frequency-dependent contractions and ATP transients that were abolished by 1 μ M TTX and greatly attenuated by nifedipine (1 μ M). Atropine (1 μ M) reduced force by about 50% but had no effect on ATP transients. With no EFS stimulation spontaneous contractions developed that were mirrored by ATP transients with a close correlation in respective amplitudes. Tension and ATP transients persisted in the presence of TTX, atropine and nifedipine. Immediately after EFS spontaneous contractions and ATP transients were temporally suppressed and recovered after about 40 s.

Conclusions

We propose that spontaneous contractions arise from ATP release from nerve-terminals, due to their temporary suppression by previous stimulation of motor nerves that releases ATP and generates contractions. The pharmacology of the transient events is different from nerve-mediated contractions suggesting ATP release is not due to excitation of motor nerves.

Supported by St Peter's Trust for Kidney, Bladder and Prostate Research

The action of calcineurin inhibitors on bladder contractile function and ATP release

Rita Jabr

Faculty of Health & Biomedical Sciences, University of Surrey, GU2 7HX, UK.

Introduction.

Overactive bladder is characterised by spontaneous contractions. Several hypotheses include: increased functionality between mucosa and detrusor; increased mucosal ATP release and augmented detrusor contractions. Increase of the Ca^{2+} /calmodulin-dependent phosphatase calcineurin (Cn) activity is implicated in the aetiology of defective intercellular signalling in myocardium,¹ but its role in signalling in bladder wall tissues is unknown.

Methods.

Detrusor strips with intact mucosa from *ex vivo* guinea-pig bladders euthanised by cervical stimulation and superfused with Tyrode's solution at 37°C. Carbachol (0.1 μM) contractures or spontaneous contractions were recorded in unstimulated preparations. Stretch-activated mucosal ATP release was measured during a stretch of 20% resting length in superfusate samples with a luciferin-luciferase technique. Human detrusor biopsies from normal and OAB human bladders were analysed for calpain (-5,-13) mRNA levels.

Results.

The Cn-inhibitor cyclosporin-A (CsA, 5 and 10 μM) decreased carbachol contractures to $74.7 \pm 4.8\%$ and $75.9 \pm 4.8\%$ control, respectively; 1 μM CsA had no significant effect ($n=6$). Spontaneous contractions were also reduced by CsA (1,10 μM ; $36.2 \pm 2.5\%$, $39.6 \pm 7.3\%$ control respectively, $n=5$). CsA (1 μM) reduced ATP release from 145 ± 15 to 75 ± 6 fmol/mg wet weight. mRNA quantification showed gene expression of calpain-5 and calpain-13 increased 1.77 ± 0.32 and 1.84 ± 0.41 -fold respectively ($n=4$).

Discussion and Conclusions. CsA diminished agonist-induced and spontaneous contractions, the latter more sensitive to CsA, and stretch-activated mucosal ATP release. mRNA analyses showed that calpain, which renders Cn constitutively active, was significantly increased in OAB human bladder samples. These demonstrate that Cn can upregulate contractile and ATP-signalling functions and may be augmented in OAB.

Supported by EU FP7 InComb

Reference

1. Jabr RI, Hatch FS, Salvage SC, Orlowski A, Lampe PD, Fry CH. Regulation of gap junction conductance by calcineurin through Cx43 phosphorylation: implications for action potential conduction. *Pflügers Arch* 2016; 468: 1945-1955.