Allen, K. J., Fitzharris, L. E., & McConnell, A. K. (2019). Inspiratory Muscle Training and Testing: Rationale, Development and Feasibility. *Equine Veterinary Journal*. <https://doi.org/10.1111/evj.13217>

Peer reviewed version

Link to published version (if available):
[10.1111/evj.13217](https://doi.org/10.1111/evj.13217)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://beva.onlinelibrary.wiley.com/doi/abs/10.1111/evj.13217> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Inspiratory Muscle Training and Testing: Rationale, Development and Feasibility

Formatted

Summary

Background

Inspiratory muscle training (IMT) applies a training stimulus directly to the inspiratory muscles and is distinct from whole-body training. The potential benefits of IMT have yet to be explored in horses.

Objectives

The objectives were: 1) to develop an equine-specific method of testing and training inspiratory muscles; 2) to assess tolerance and feasibility in a pilot study in a commercial Thoroughbred training establishment.

Study Design

Field study

Methods

A mask was used to interface commercial human IMT equipment. Ten horses undertook IMT once daily while stood in the stable approximately 5 days/week over a 9-week period. Inspiratory muscle strength testing (IMST) employed a continuous incremental inspiratory loading protocol alternating two loaded and two minimally loaded breaths until failure to tolerate the load occurred or the maximum 60 breaths were completed. The IMST was undertaken twice; firstly in 10 horses with minimal acclimatisation and secondly in 8 horses experienced with the IMT programme.

Results

The ten horses undertook IMT for a median of 42 days, reaching a median peak training load of 32.5cmH₂O. One horse did not tolerate the mask with repeated snorting and was replaced. All horses completed the IMST. The median peak value in IMST 1 was 27cmH₂O and in IMST 2 was 41cmH₂O. Two of 10 horses reached the maximum possible value in IMST 1; therefore, the test was adapted to permit a higher maximum value, despite this 3/8 horses reached the maximum possible value in IMST 2.

Main limitations

A small number of horses were assessed. The IMST was refined during the study and requires additional refinement.

Conclusion

Inspiratory muscle testing and training were feasible and tolerated in horses. Further research is required to understand whether the IMST values obtained correlate with other physiological/performance outcomes. The potential benefits and/or adverse effects of IMT warrant further investigation.

1 **Introduction**

2 Respiratory muscle training (RMT) applies a training stimulus directly to the respiratory muscles and
3 is distinct from whole body training. The respiratory muscles (upper airway and respiratory pump
4 muscles, including the diaphragm) are skeletal muscles, which are morphologically and functionally
5 like locomotor muscles. Evidence from human and rodent studies indicates that upper airway and
6 respiratory pump muscles can adapt to exercise training^{1, 2}. Thus the respiratory muscles appear to be
7 the primary component of the equine respiratory system that may respond to training³. The most
8 common form of RMT in humans is inspiratory pressure threshold loading, which involves inhaling
9 against a resistance, which induces adaptations that improve the strength, power and endurance of the
10 inspiratory muscles. This training method is commonly known as inspiratory muscle training (IMT).

11
12 Inspiratory muscle training has been used in humans for the management/treatment of a range of
13 medical conditions, including exercise-induced laryngeal obstruction (EILO) and in the treatment of
14 breathlessness for patients with chronic lung and/or heart disease⁴. It has also gained popularity
15 amongst athletes as an ergogenic aid, by improving respiratory muscle performance.

16
17 The respiratory musculature of humans has been shown to fatigue during exercise, resulting in complex
18 mechanisms of performance limitation⁶. One of these mechanisms is a respiratory muscle fatigue
19 (RMF) -induced metaboreflex that results in a decrease in blood flow to the exercising limb muscles⁷.
20 The use of IMT in human athletes has been shown to improve athletic performance^{4,7,8}, by delaying the
21 onset of the RMF and postponing activation of the respiratory muscle metaboreflex. Existence of the
22 respiratory muscle metaboreflex has been confirmed in dogs⁹, but it is unknown whether RMF occurs
23 in horses nor whether the respiratory muscle metaboreflex is active during race/competition conditions.
24 As an athletic species the horse is limited by the respiratory system to a greater extent than in most
25 human athletes¹⁰. However, ventilation in the horse may be facilitated by back flexion and locomotor
26 respiratory coupling thereby reducing the work of the diaphragm in comparison to humans. Despite
27 these differences the respiratory musculature in both maximally exercising humans and ponies
28 commands a similar proportion (~14-20%) of cardiac output¹¹.

29

30 Inspiratory muscle training has been shown to activate the upper airway muscles and has been used to
31 treat EILO in human athletes^{4,12}. The high prevalence of upper airway obstructions in racehorses poses
32 significant performance, health and welfare concerns to the Thoroughbred industry. Exercise-induced
33 laryngeal/pharyngeal obstructions can occur when the stabilising muscles of the upper airway are
34 unable to withstand the dramatic increases in airflows and pressures that occur during exercise. Most
35 surgical treatments try to provide a mechanical solution and there is no treatment currently available to
36 improve the strength of the upper airway musculature, thereby providing a functional treatment. If the
37 upper airway muscles in the horse do respond to a training stimulus, this could not only be applied as a
38 non-surgical treatment option, but also as a prophylactic strategy.

39

40 In order to set training loads and to monitor improvement, it would be useful to develop methods to
41 measure respiratory muscle performance. Several indices of respiratory muscle function are available
42 in humans, including maximal inspiratory mouth pressures, sniff nasal pressure, oesophageal, gastric
43 and trans-diaphragmatic pressures¹³. Functional outcomes, such as maximal inspiratory muscle
44 shortening velocity, power and endurance, have also been developed. The most common method of
45 assessing respiratory muscle strength is maximal inspiratory pressure (MIP) measured at the mouth
46 during a maximal, volitional inspiratory effort against an occluded airway, which provides a non-
47 invasive index of global inspiratory muscle strength⁴. However, functional testing of the respiratory
48 system in horses is limited by the inability to undertake maximal volitional efforts. Surrogate measures
49 of inspiratory pressure have been described in unconscious or ventilated human patients, including
50 incremental threshold loading and occlusion tests^{14,15}. A non-invasive index of inspiratory muscle
51 strength (IMS), not requiring maximal volitional effort, and suitable for routine clinical use in the horse,
52 would benefit the evaluation of the responses of the equine inspiratory muscles to IMT.

53

54 Thus, the purpose of this study is to describe an equine-specific method of testing and training
55 inspiratory muscles and the feasibility of implementing this in a pilot study of 10 Thoroughbreds within
56 a commercial training establishment.

Materials and Methods

Equipment:

Equipment for the application of IMT and IMST had previously been developed and [undergone preliminary evaluation](#) for use in the horse (unpublished data). Briefly, an airtight mask (Figure 1) made from PETG plastic, covering the entire muzzle, was developed. A tight-fitting latex rubber and Velcro®-fastened head piece secured the mask. An opening at the level of the nares allowed the attachment of IMT pressure threshold valves^{a,b} or electronic equipment^c. The valves^{a,b} [had been](#) tested in conjunction with the mask using a laboratory based flow/volume simulator pump at flows and volumes commensurate with those [measured](#) in resting horses¹⁵. These tests confirmed the valves created the prescribed inspiratory pressures and did not restrict airflow through flow resistance.

Population:

A convenience sample of 10 healthy Thoroughbred National Hunt racehorses in active training with a single trainer were recruited. All horses were race fit and free from clinical signs of respiratory disease at the start of the study. The study was conducted towards the end of the National Hunt season, from January to March, when the horse's exercise regimens would remain constant during the study period.

Inspiratory Muscle Training (IMT):

Horses were initially introduced to wearing a loose-fitting mask with large air holes for the first session. Gradually, the IMT mask and low load valves^a (5cmH₂O) were applied by an experienced user. The yard staff were taught how to undertake the daily IMT and given a user guide document; safety aspects were explicitly emphasised and communicated. Subsequently, all IMT sessions were performed by yard staff whilst the horse was standing in its stable. [In humans, a wide range of IMT programmes have been used clinically and within research, depending on the patient and outcome of interest. The literature indicates training loads should exceed 30% MIP and typical guidelines in healthy people are to](#)

undertake IMT at values of 50-70% MIP or at the maximum that can be sustained for 30 consecutive breaths⁴. In young adults typical MIP values are between 97-128cmH₂O¹⁶.

As the ideal intensity and duration of an IMT programme in horses is unknown, a preliminary training programme was developed in conjunction with an expert in human IMT. The programme was further adapted to promote tolerance, as this was considered key for equine IMT and hence the load is at a lower level with a more gradual increase than programmes used in humans. All horses undertook the IMT programme over a 9-week period, which involved two sessions of 30 'loaded' breaths performed back-to-back with a short break (~2 mins) in-between, completed 5 days/week. The inspiratory load was increased gradually every 4 days (5, 10, 12.5, 15, 20 cmH₂O using one type of valve^a then in increments of 2.5cmH₂O using the second valve^b) over the 9-week period, with minor adjustments according to the horse's tolerance to increases in load. The guideline IMT programme was provided for all horses with the ability for users to increase more quickly if the horse was finding the training easy (readily completed both sets of 30 breaths maintaining normal breath duration throughout) and to remain at a level for longer if the user felt the horse was not ready to increase to the next level (unable to complete the 30 breaths, taking short, clipped breaths, or displaying avoidance behaviour such as crossing the jaw). Inspiratory muscle training was not performed for 48 hours prior to racing. The horse's normal exercise and racing schedule continued throughout.

The yard staff were given a daily training diary to complete to log the IMT undertaken and asked to complete a paper questionnaire (anonymously) at the end of the study. Training diaries and yard staff questionnaires were analysed to determine number of training days, acceptance and training level. The trainer was also questioned about subjective interpretation of health and performance of each horse at the start and end of the study.

Inspiratory Muscle Strength Test (IMST):

An electronic inspiratory loading device (POWERbreathe® K5^c) designed for IMT in human beings was used to implement a continuous incremental pressure loading protocol (Figure 2). The device has

112 been validated in human patients¹⁷ and used to assess changes following IMT¹⁸. Briefly, the device can
113 measure average load [=pressure] (cmH₂O), breath volume (l), peak flow (l/s), peak power [product of
114 inspiratory load and flow] (W) and work [product of load and volume] (J) continuously at 500Hz,
115 enabling a comprehensive real time analysis of the physical task being undertaken by the inspiratory
116 muscles during loading. Since it is impossible to elicit maximal voluntary efforts from horses, an
117 incremental loading protocol was developed and designed to progressively increase the inspiratory load
118 to the limit of tolerance; the test is analogous to an incremental exercise test. Briefly, two identical
119 loaded breaths were interspersed with two minimally loaded breaths, with the loaded breaths increasing
120 incrementally in magnitude by 2cmH₂O each step, to a maximum of 60 breaths in total^c. By starting the
121 protocol with a low-level loading, the horses had a period of habituation, during which they became
122 accustomed to the unusual sensations of the load, before it became too challenging. The novel
123 introduction of the minimally loaded breaths mitigated the effects of accumulated fatigue and, by
124 permitting full tidal excursions, prevented the development of hypercapnia and lung hyperinflation. The
125 test continued until the horse made an inspiratory effort but was unable to open the valve or all 60
126 breaths were completed. The index of IMS (IMSi) was the highest load (cmH₂O) at which the horse
127 was able to open the valve. Prior to embarking on this feasibility trial, the IMST had been assessed in
128 10 horses (7 Thoroughbred, 3 mixed breed) and ages (median age 5 years, range 3-16 years), in which
129 IMS was measured on two separate occasions, in order to assess repeatability. A median value of
130 21cmH₂O and a peak value of 30cmH₂O were recorded, with no significant differences in the test re-
131 test values (unpublished data).

132 The horses underwent 3 days of acclimatisation to the mask before undertaking IMST 1, which had a
133 maximum IMSi of 33cmH₂O if the horse completed all 60 breaths^c (Figure 2). Horses then undertook
134 the IMT programme over a 9-week period, followed by IMST 2, which had a maximum IMSi of
135 45cmH₂O (figure 2). All IMST were undertaken by the same persons who were not blinded to horse
136 identity. The author that lead the IMST was blinded to the progress and the IMT level reached by each
137 horse.

Data Analysis:

The IMSi (cmH₂O), defined as the highest load within the IMST protocol that the horse opened the valve, was considered to be the primary outcome measure of interest for assessing inspiratory strength. However, average load (cmH₂O), breath volume (l), peak flow (l/s), peak power (W) and work (J) were extracted for the paired loaded breaths for IMST 1. Data were assessed graphically and with the use of a Shapiro-Wilk test to determine normality. SPSS (version 24)^c was used for statistical analysis. Wilcoxon Signed Rank tests were used to compare average load (cmH₂O), breath volume (l), peak flow (l/s), peak power (W) and work (J) for the paired loaded breaths for IMST 1. Significance was set at P<0.05.

Results

Ten Thoroughbred horses were recruited initially. One horse was immediately replaced as repeated snorting occurred when the mask was fitted; this horse was diagnosed subsequently with lower airway inflammation. Ten horses completed the IMST 1. A further horse was removed from the yard for unrelated reasons after completing the first test, but before starting IMT, and was replaced with another horse that did not undertake IMST 1. The ten horses that underwent the IMT were all geldings with a median age of 8 years (range 5-10). The number of days of IMT undertaken, the final training level and results of the two IMST are shown in Table 1. The exact number of days of IMT completed by each horse varied slightly according to racing schedule and staff availability. One horse (horse 3) was restricted to box rest for the final three weeks of the IMT period. Two horses were unavailable for the second IMST due to racing commitments. Five of the horses followed the guideline IMT programme exactly, 2 horses were a level (2.5cmH₂O) ahead of the suggested programme and three horses a level (2.5cmH₂O) behind.

It was necessary to adapt the IMST protocol after IMST 1 because two horses unexpectedly completed the maximum number of breaths possible with the equipment at IMST 1, creating a ceiling effect for IMSi. The IMST 2 had the same number of loaded breaths, same maximum number of breaths, and same increase in load but started at a higher load to allow for a higher end point (figure 2). Despite this,

three horses also reached the maximum load of 45cmH₂O in IMST 2. One horse (11) repeatedly crossed his jaw during IMST 2 and failed to take normal breath volumes, thus the validity of this IMSi was questioned. The median peak IMSi value for IMST 1 was 27cmH₂O and for IMST 2 was 41cmH₂O.

The median peak training load reached was 32.5cmH₂O thus, at the end of the programme IMT was undertaken at approximately 79% of the IMSi from IMST 2.

The analysis of the loaded breaths in IMST 1 showed that there were no significant differences in load (p=0.849), volume (p=0.135), flow (p=0.745), power (p=0.170) or work (p=0.355) between first and second of each pair of breaths.

Tolerance for the IMST under the conditions specified was good. Of the 18 tests performed on one occasion a horse repeatedly crossed his jaw which resulted in loss of the air-tight seal; thus, the test result was likely lower than the true value. During IMT, nostril dilation is observed, and thorax movement was clear. Subjectively, horses also appeared to make an obvious inspiratory effort in response to loading. Horse acceptance of IMT was very good, on no occasion were yard staff unable to undertake training due to horse behaviour. The trainer questionnaire did not identify any change in health or performance and no adverse effects were reported by trainers or yard staff.

Discussion

The purpose of this pilot study was to assess the feasibility and tolerance of inspiratory muscle training and testing in the horse. The results show that IMT can be undertaken in a commercial Thoroughbred training establishment. Further research is now required to evaluate whether IMT results in any potential benefits or adverse effects to health and performance.

In both IMST 1 and 2, horses outperformed our expectations from our initial research. As a result, the IMST protocol was changed between the two tests; despite increasing the demands of the IMST 2, horses still reached the test's upper limit, thus a true IMSi may not have been obtained as horses may have been capable of higher values. The change in protocol, the ceiling effect and lack of control group

prevent robust comparisons of the IMST results from tests 1 and 2 but highlight the importance of undertaking a small feasibility trial prior to a larger case control study. Further research will help demonstrate the effect of IMT on IMS and will elucidate whether any increases result from respiratory muscle hypertrophy, improved neuromuscular activation as a result of IMT, or simply an improved ability to tolerate the test.

As horses reached the upper limit of both IMSTs, further adaptation will be required to permit an even higher endpoint. This could be developed through having a higher start point, larger increments between or a decrease in the number of loaded or minimally loaded breaths. As no significant differences were found between breath variables in the first and second loaded breath, we suggest the protocol be amended to one loaded breath. This will permit a higher maximum value, whilst maintaining only small incremental increases and retaining two minimally loaded breaths to maintain tolerance.

During development of the IMST the primary objective was to determine the highest inspiratory load (cmH₂O), the horse was able to open. This was termed the index of inspiratory muscle strength. The equipment used also records additional breath data, such as load (cmH₂O), volume (l), peak flow (l/s), peak power (W) and work (J). In humans, when undertaking a strength test at maximal volitional effort, all these variables typically increase following IMT. For equine testing, further analysis of these variables on an identical test pre and post-IMT will aid understanding of their value. However, it is anticipated that they are of less value in equine testing as it is likely that the horse will do the minimum work required to open the valve, rather than giving a maximum effort, therefore the primary parameter to improve should be the ability to open load i.e. the index of inspiratory muscle strength.

Interestingly the pressures/loads generated by the horse during training and testing were substantially lower than those typically achieved by humans. Although this can be partly explained by inspiratory testing in humans involving a maximal volitional effort, which is not possible in the horse, it is also likely that the scale and anatomy of the equine diaphragm and rib cage alters the relationship between muscle tension and the resultant static pressure. It remains unclear how representative the IMSi is of

true maximal inspiratory strength. Research in humans has shown that the values achieved during incremental threshold loading are on average 69% of MIP¹⁴. The equine training programme was developed to prioritise tolerance; nevertheless, the final IMT level was approximately 79% of our index of IMS. Further research is required to understand whether the IMSi value obtained correlates with other physiological/performance variables.

From an ethical and welfare perspective the application of IMT in horses requires considerable care. The horses were introduced to the mask gradually by the authors, who oversaw the programme. At any point if the horse failed to open the valve during IMT or IMST it was removed. Safety measures were included when the equipment was developed so both the valve and the mask can be rapidly removed in any adverse situation.

In conclusion, this study has shown the novel application of IMT and the development of an innovative method of IMS testing in the horse. Undertaking IMT in the horse is feasible but further research is required to understand any potential benefits or adverse effects.

250 **Manufacturer Addresses**

251 a CPEEP valves, Intersurgical, Wokingham, Berkshire UK

252 b POWERbreathe® Medic Classic, POWERbreathe International Ltd, Southam, Warwickshire UK

253 c POWERbreathe K5®, POWERbreathe International Ltd, Southam, Warwickshire UK

254 d IBM SPSS Statistics for Windows, version 24.0 Armonk, NY:IBM Corp

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

Figure 1: Equipment for inspiratory muscle training

Figure 2: Inspiratory muscle strength test protocols (a) IMST 1 (b) IMST 2. For both tests the maximum number of breaths was 60, of which the maximum number of loaded breaths was 30. The loaded breaths increase by an increment of 2cmH₂O. IMST 2 had a higher start point enabling a higher end point of 45cmH₂O compared to 33cmH₂O in IMST 1.

Horse	<u>IMSi from IMST</u> <u>1</u> (cmH ₂ O)	<u>IMSi from</u> <u>IMST 2</u> (cmH ₂ O)	No. Days IMT	Peak training load reached (cmH ₂ O)
1.	21	35	40	32.5
2.	33*	43	42	35
3.	27	33	44	32.5
4.	21	-	-	-
5.	31	45^	43	32.5
6.	23	-	38	32.5
7.	21	45^	42	32.5
8.	27	-	37	27.5
9.	33*	45^	40	32.5
10.	31	39	44	35
11.	-	27	42	32.5
Median	27	41	42	32.5

Table 1 shows the number of days of IMT completed by each horse, the peak training load (cmH₂O), and the IMSi (cmH₂O) from tests 1 and 2. *= horse completed all 60 breaths, 33cmH₂O was the maximum inspiratory load tested. ^= horse completed all 60 breaths, 45cmH₂O was the maximum inspiratory load tested.

References

1. Powers S.K., Coombes J. and Demirel H. (1997) Exercise Training-Induced changes in respiratory muscles. *Sports Med.* **24**, 120-131.
2. Vincent H.K., Shanely R.A., Stewart D.J., Demirel H.A., Hamilton K.L., Ray A.D., Michlin C., Farkas G.A. and Powers S.K. (2002) Adaptation of upper airway muscles to chronic endurance exercise. *Am. J. Crit. Care Med.* **166**, 287-293.
3. Roberts C.A., Marlin D.J. and Lekeux P. (1999) The effects of training on ventilation and gas exchange in exercising Thoroughbreds. *Equine Vet. J. Suppl.* **30**, 57-61.
4. McConnell A.K. (2013) *Respiratory Muscle Training: Theory and Practice*. Reed Elsevier, Oxford.
5. Romer L.M. and Polkey M.I. (2008) Exercise-induced respiratory muscle fatigue: implications for performance. *J. Appl. Physiol.* **104**, 879-888.
6. Harms C.A., Babcock M.A., McClaran S.R., Pegelow D.F., Nিকে G.A., Nelson W.B. and Dempsey J.A. (1997) Respiratory muscle work compromises leg blood flow during maximal exercise. *J. Appl. Physiol.* **82**, 1573-1583.
7. HajGhanbari B., Yamabayashi C., Buna T.R., Coelho J.D., Freedman K.D., Morton T.A., Palmer S.A., Toy M.A., Walsh C., Sheel A.W. and Reid W.D. (2013) Effects of respiratory muscle training on performance in athletes: a systematic review with meta-analyses. *J. Strength Cond. Res.* **27**, 1643-63.
8. Illi S.K., Held U., Frank I. and Spengler C.M. (2012) Effect of respiratory muscle training on exercise performance in healthy individuals: a systematic review and meta-analysis. *Sports Med.* **42**, 707-24.
9. Rodman J.R., Henderson K.S., Smith C.A. and Dempsey J.A. (2003) Cardiovascular effects of the respiratory muscle metaboreflexes in dogs: rest and exercise. *J. Appl. Physiol.* **95**, 1159-1169.
10. Franklin S.H., Van Erck-Westergren E. and Bayly W.M. (2012) Respiratory responses to exercise in the horse. *Equine Vet. J.* **44**, 726-732.

11. Sheel A.W., Bousel R. and Dempsey J.A. (2018) Competition for blood flow distribution between respiratory and locomotor muscles: implications for muscle fatigue. *J. Appl. Physiol.* **125**, 820-831.
12. How S.C., McConnell A.K., Taylor B.J. and Romer L.M. (2007). Acute and chronic responses of the upper airway to inspiratory loading in healthy awake humans: an MRI study. *Respir. Physiol. Neurobiol.* **157**, 270-280.
13. ATS/ERS Statement on Respiratory Muscle Testing (2002) *Am. J. Respir. Crit. Care Med.* **166**, 518-624.
14. Johnson P.H., Cowley A.J. and Kinnear W.J. (1997) Incremental threshold loading: a standard protocol and establishment of a reference range in naïve normal subjects. *Eur. Respir. J.* **10**, 2686-2671.
15. The respiratory system: anatomy, physiology and adaptations to exercise and training (2014) In: *The Athletic Horse: principles and practice of equine sports medicine*. Eds D.R. Hodgson, C.M. McGowan and K. McKeever. Elsevier Saunders. pp125-154.
16. Schlauser Pessoa I.M., Franco Parreira V., Fregonezi G.A., Sheel A.W., Chung F. and Reid W.D. (2014) Reference values for maximal inspiratory pressure: a systematic review. *Can. Respir. J.* **21**, 43-50.
17. Langer D., Jacombe C., Charusisin N., Scheers H., McConnell A., Decramer M. And Gosselink R. (2013) Measurement validity of an electronic inspiratory loading device during a loaded breathing task in patients with COPD. *Respir. Med.* **107**, 633-635.
18. Charusisin N., Gosselink R., Decramer M., Demeyer H., McConnell A., Saey D., Maltais F., Derom E., Vermeersch S., Heijdra Y.F., van Helvoort H., Garms L., Schneeberger T., Kenn K., Gloeckl R. and Langer D. (2018) Randomised controlled trial of adjunctive inspiratory muscle training for patients with COPD. *Thorax.* **73**, 942-950.