Manzanares, E., Escudero-Mozo, M. J., Ferrón, H., Martínez-Pérez, C., & Botella, H. (2020). Middle Triassic sharks from the Catalan Coastal ranges (NE Spain) and faunal colonization patterns during the westward transgression of Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 539, [109489].
<https://doi.org/10.1016/j.palaeo.2019.109489>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.palaeo.2019.109489](https://doi.org/10.1016/j.palaeo.2019.109489)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S0031018219305693> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Manuscript Details

Manuscript number	PALAEO_2019_457_R2
Title	Middle Triassic sharks from the Catalan Coastal Ranges (NE Spain) and faunal colonization patterns of East Iberia during the Triassic westward transgression of Tethys
Article type	Research Paper

Abstract

Palaeogeographic changes that occurred during the Middle Triassic in the westernmost Tethyan domain were governed by a westward marine transgression of the Tethys Ocean. It flooded for the first time during the Mesozoic, covering wide areas of the eastern part of Iberia with the subsequent evolution of new epicontinental shallow-marine environments and their colonization by diverse faunas, including chondrichthyans. In East Iberia, this general transgression is recorded by two successive transgressive–regressive cycles: (1) middle–late Anisian and (2) late Anisian–early Carnian. Here, we describe the chondrichthyan fauna recovered from several Middle Triassic stratigraphic sections (Pelsonian-Longobardian) located at the Catalan Coastal Basin (western-most Tethys). The assemblage consists of isolated teeth of the species *Hybodus plicatilis*, *Omanoselache bucheri*, *O. contrarius* and *Pseudodolatias henarejensis*. Our data complement a series of recent studies on chondrichthyan faunas from Middle-Late Triassic marine basins of the Iberian Peninsula, allowing to evaluate the patterns of faunal colonization of the new shallow epicontinental seas of the western Tethys. Thus, sharks that reach East Iberia during the first transgressive pulse (Anisian) show strong affinities with the Sephardic bioprovince (related with the Neotethys Ocean) whereas a posterior increase in chondrichthyan diversity (during the Ladinian transgressive pulse) is due to the arrival of taxa with clear affinities to European faunas (i.e. from the Palaeotethys). This contrasts with recently evaluated colonization patterns of invertebrates and conodonts, where Middle–Upper Anisian assemblages are composed of taxa with clear affinities to the Alpine/Germanic bioprovince whereas faunas from the Sephardic bioprovince only reached the shallow waters of the Iberian peninsula during the second (Ladinian) transgressive pulse. The differences between chondrichthyan and invertebrate colonisation patterns are explained here on the basis of their different dispersal strategies. This requires some adjustment of previous palaeogeographic reconstructions of the westernmost Tethyan realm for the Middle Triassic including the possible existence, during the Anisian, of connections between Palaeo- and NeoTethys through the western Cimmerian microcontinent, which served as pathways for the entrance of chondrichthyans from the Sephardic bioprovince.

Keywords	dispersal strategies; palaeocurrents; Anisian; Ladinian; coastal chondrichthyans.
Manuscript region of origin	Europe
Corresponding Author	Hector Botella
Corresponding Author's Institution	University of Valencia
Order of Authors	Esther Manaznares Ubeda, Maria José Escudero-Mozo, Humberto G. Ferrón, Carlos Martínez Pérez, Hector Botella
Suggested reviewers	Christian Klug, Charlie Underwood, Ana Marquez-Aliaga, Detlev Thies, Jürgen Kriwet

1 **Middle Triassic sharks from the Catalan Coastal Ranges (NE Spain)**
2 **and faunal colonization patterns during the westward transgression of**
3 **Tethys**

4 Manzanares, E.¹, Escudero-Mozo, M.J.², Ferrón, H.^{3,4}, Martínez-Pérez, C.^{3,4}, and
5 Botella, H.^{3*}

6
7 ¹*Botany and Geology Department, University of Valencia, Avda. Dr. Moliner, 50, 46100 Burjassot*
8 *(Valencia), Spain; Esther.Manzanares@uv.es*

9 ²*Instituto de Geociencias (UCM, CSIC), Calle del Dr. Severo Ochoa, 7, 28040 Madrid, Spain;*
10 *mjescudero@geo.ucm.es*

11 ³*Institut Cavanilles de Biodiversitat i Biologia Evolutiva, Universitat de Valencia, Paterna, 46980,*
12 *Valencia, Spain; Carlos.Martinez-Perez@bristol.ac.uk; Hector.Botella@uv.es.*

13 ⁴*School of Earth Sciences, University of Bristol, Bristol, BS8 1TQ, UK; Humberto.Ferron@uv.es*

14 **Corresponding author*
15
16

17 **Abstract.** Palaeogeographic changes that occurred during the Middle Triassic in the
18 westernmost Tethyan domain were governed by a westward marine transgression of the
19 Tethys Ocean. The transgression flooded wide areas of the eastern part of Iberia,
20 forming new epicontinental shallow-marine environments, which were subsequently
21 colonized by diverse faunas, including chondrichthyans. The transgression is recorded
22 by two successive transgressive–regressive cycles: (1) middle–late Anisian and (2) late
23 Anisian–early Carnian. Here, we describe the chondrichthyan fauna recovered from
24 several Middle Triassic stratigraphic sections (Pelsonian-Longobardian) located at the
25 Catalan Coastal Basin (western-most Tethys). The assemblage consists of isolated teeth
26 of the species *Hybodus plicatilis*, *Omanoselache bucheri*, *O. contrarius* and

Pseudodalatias henarejensis. Our data complement a series of recent studies on chondrichthyan faunas from Middle-Late Triassic marine basins of the Iberian Peninsula, allowing us to evaluate patterns of faunal colonization. Sharks that reached East Iberia during the first transgressive pulse (Anisian) show strong affinities with the Sephardic bioprovince (related with the Neotethys Ocean) whereas a later increase in chondrichthyan diversity (during the Ladinian transgressive pulse) is due to the arrival of taxa with affinities to European faunas (Palaeotethys). This contrasts with recently evaluated colonization patterns of invertebrates and conodonts, where Middle–Upper Anisian assemblages are composed of taxa with affinities to the Alpine/Germanic bioprovince whereas faunas from the Sephardic bioprovince only reached the shallow waters of the Iberian peninsula during the second (Ladinian) transgressive pulse. The differences between chondrichthyans vs. invertebrate and conodonts colonisation patterns are explained here on the basis of differing dispersal strategies. Our finding suggests the existence, during the Anisian, of connections between Palaeo- and NeoTethys through the western Cimmerian microcontinent, which served as pathways for the entrance of chondrichthyans from the Sephardic bioprovince.

Keywords: dispersal strategies; palaeocurrents; Anisian; Ladinian; coastal chondrichthyans.

1. Introduction

The Triassic is a key period to understand the evolutionary history of chondrichthyans, with the decline or extinction of several major Palaeozoic groups (such as xenacanthiforms, symmoriiforms, ctenacanthiformes and non-chimaeroid holocephalans), the increasing dominance of hybodontiforms, and the emergence of the

modern sharks (elasmobranchii). The Triassic was also an important period for the Iberian Peninsula in terms of palaeogeography. During these times, the western Tethys domain was characterised by important palaeogeographic changes and the subsequent advance of the broad westward marine transgression of the Tethys Sea that reached East Iberia during the Middle Triassic (Anisian) (Escudero-Mozo et al., 2015). This transgression, for the first time during the Mesozoic, flooded wide areas of the eastern part of Iberia, connecting some internal basins (i.e. Iberian and Catalan), which until this moment had been independent (e.g. Arche et al., 2004; Galán-Abellán, et al., 2013; Escudero-Mozo et al., 2015), causing the demise of continental sedimentation and the consequent deposition of marine carbonates and mixed coastal facies (Muschelkalk).

In East Iberia, this general transgression is recorded by two transgressive-regressive cycles that show important palaeogeographic and fossil content changes (Escudero-Mozo et al., 2015). The first one (middle–late Anisian) took place from the NE to the Catalan Coastal Basin reaching the Iberian Basin through a narrow corridor bounded by palaeohighs. The second transgressive-regressive cycle (late Anisian–early Carnian) was characterised by an extensive transgression that covered a wide area of E Iberia including some of the previous topographic highs. After that, during the early Carnian, this area is characterised by the development of an extensive siliciclastic-evaporitic setting of the Keuper. These transgressive events resulted in the evolution of a diversity of epicontinental shallow coastal environments and associated colonization by marine faunas, including chondrichthyans. Accordingly, Middle Triassic faunas from the Iberian Peninsula reflect the transformation from continental settings to a wide variety of coastal and marine depositional environments owing to the Muschelkalk transgression, which occurred in the middle Anisian (Escudero Mozo et al., 2015).

In this paper, we document for the first time the presence of chondrichthyans in the Middle Triassic rocks of the Catalan Coastal ranges (CCR). This paper is complementary to Pla et al.'s (2013) and Manzanares et al.'s (2018) studies on Middle–Late Triassic chondrichthyans from other areas of the Iberian Peninsula, i.e. the Iberian (IR) and Betic ranges (BR), respectively. Together, these three papers give the first general overview of the chondrichthyan communities that inhabited the shallow epicontinental seas of the westernmost Tethys during the Middle to Late Triassic. This allows us to evaluate here the patterns of colonization of the new shallow epicontinental seas of the western Tethys followed by coastal sharks and to compare it with invertebrate faunas. Findings require some adjustments of previous palaeogeographic reconstructions of the westernmost Tethyan realm for the Middle Triassic.

2. Geographical and geological settings

The Catalanian Coastal range is located in the north-eastern Iberian Peninsula (Fig. 1A) and includes two mountain chains that run parallel to the Mediterranean Coast in a NE–SW direction, from the Gerona region towards the Ebro delta, separated by the Prelitoral Depression (Sopeña and De Vicente, 2004; Escudero-Mozo et al., 2015).

The Catalanian Basin constituted an intracratonic basin that was mainly developed during Mesozoic times when it experienced some extensional periods with several syn-rift and post-rift phases (Vargas et al., 2009; De Vicente et al., 2009). It is dominated by longitudinal NE–SW faults and to a lesser extent by a conjugated NW–SE fault system (Vegas, 1975); the latter divided this basin into three domains with different subsidence (from SW to NE): Priorat-Baix Ebre, Prades, and Gaià-Montseny (Marzo, 1980; Calvet and Ramón, 1987; Ramón and Calvet, 1987; Gaspar-Escribano et al., 2004). During the Cenozoic, Alpine compressional tectonics gave rise to the present-day Iberian range and

Catalan Coastal range.

In the Catalanian Basin, the succession shows the three typical subdivisions of the Germanic Triassic, namely Buntsandstein, Muschelkalk and Keuper facies. In the Catalan Coastal range, the Muschelkalk facies is represented by two carbonate units (“Lower” and “Upper” Muschelkalk) that record the deposition of shallow to outer carbonate ramps, separated by a mixed evaporite-siliciclastic unit (“Middle Muschelkalk”). This facies, combined with the uppermost part of the Buntsandstein (or Röt) facies and the lower part of the Keuper facies, records two major transgressive–regressive cycles, respectively of middle–late Anisian and late Anisian–early Carnian age (Escudero-Mozo et al., 2014, 2015), each one related to the development of a wide carbonate ramp. The studied sections are constituted from materials of the “Lower Muschelkalk” (L’Ametlla section) from the Gaia–Montseny domain, and “Upper Muschelkalk” (Rasquera-Benifallet and Tivissa sections) from the Priorat-Baix-Ebre (Fig. 1B).

In the Gaia–Montseny domain, the Lower Muschelkalk shows an average thickness of 120 m, showing a sharp lower contact with lutite, carbonate and evaporite units (Röt facies) and a transitional upper contact with the Middle Muschelkalk. The unit was divided by Calvet and Ramón (1987) and Calvet et al. (1990) into four members, which correspond to El Brul (shallow subtidal to supratidal deposits), Olesa (lagoonal bioclastic carbonates), Vilella Baixa (open marine burrowed deposits) and Colldejou (white peritidal dolomite facies). Based on ammonoids, foraminifera, pollen and brachiopods, the Lower Muschelkalk unit is dated as middle–upper Anisian (Calvet and Marzo, 1994; Goy, 1995; López-Gómez et al., 1998; Márquez-Aliaga et al., 2000). Chondrichthyan remains are concentrated in different levels of the Olesa unit.

The L'Ametlla section is located close to the village of L'Ametlla del Vallès in the province of Barcelona, Catalonia, 35 km northeast of Barcelona City (coordinates at the base; 41° 43' 18" N and 2° 15' 26" W; Fig. 1A). The section and its fossil record have been studied previously by Márquez-Aliaga et al. (2000), Plasencia (2009, and references therein).

In the Baix-Ebre Priorat domain, the Upper Muschelkak carbonates show a gradual to sharp lower contact with the "Middle" Muschelkalk and a gradual upper contact with the Keuper, and show an average thickness of 140 m. In this area, the carbonate unit was divided by Calvet et al. (1987) into several members from base to top that are: 1) Rojals Member (shallow subtidal to supratidal oolitic limestone and dolostone); 2) Benifallet Member (low energy subtidal bioturbated limestone and dolostone); 3) Rasquera Member (outer to middle ramp limestone, dolostone and shales with *Daonella*); 4) Tivissa Member (outer to inner ramp carbonates); and 5) Capafons Member (peritidal dolomites, marls and shales). Based on the ammonoids yielded by the Rasquera and Tivissa members and the sequence stratigraphic analysis, the Upper Muschelkalk has been dated as late Illyrian–late Longobardian (Goy, 1995; Escudero-Mozo et al., 2015). Conodonts from the same samples confirm a Ladinian age (Plasencia, 2009).

Both the Rasquera–Benifallet and Tivissa sections expose sediments of the five members as defined by Calvet et al. (1987). These sections are located in the province of Tarragona, Catalonia (Fig. 1A). The Rasquera–Benifallet section is along the road of Rasquera to Benifallet villages (coordinates at the base 40° 59' 38" N and 0° 32' 22" W), and the Tivissa section is close to Tivissa village (coordinates at the base 41° 02' 33" N and 0° 44' 48" W). Samples that have yielded chondrichthyan remains came from

both the Rasquera and Tivissa members (Fig. 1B). Conodonts from the same samples confirm a Ladinian age (Plasencia, 2009).

3. Methodology

The specimens studied here were obtained by acid etching in 5–10% acetic acid, of more than 60 Kg of limestones (in samples of around 1.5 kg each) from the Rasquera–Benifatell, Tivissa and Ametlla sections. After dilution residues were sieved with sieves meshes of 2, 0.125 and 0.063 mm respectively, which created three fractions refer as the “total sample”. Figure 1B shows the stratigraphic position of all samples that have yielded chondrichthyan remains. Fossils were picked under a binocular microscope and photographed with an HITACHII S-4800 Electronic Microscope hosted at the Microscopy Services of the University of Valencia. The archived specimens are mainly isolated teeth and scales, housed in the Museum of Natural History of the University of Valencia (former Geological Museum of Valencia University), Spain (MGUV-36111 to MGUV-36113; and MGUV-36117 to MGUV-36131).

4. Results

The chondrichthyan assemblage found in the Middle Triassic of the Catalan Basin includes four species belonging to three different genera; i.e. *Pseudodalatias henarejensis*, *Omanoselache bucheri*, *O. contrarius* and *Hybodus plicatilis* (Fig. 2). A detailed Systematic Palaeontology section is provided as Supplementary Material.

5. Discussion

175 5.1. Middle Triassic sharks from the Catalan Basin in the context of the Iberian
176 Peninsula

177 This association is similar to, but less diverse than, the chondrichthyan faunas
178 recently described by Pla et al. (2013) and Manzanares et al. (2018) from other Middle-
179 Late Triassic basins of the Iberian Peninsula (see also Botella et al., 2009; Ferrón et al.,
180 2014; Manzanares et al., 2017). In fact, *P. henarejensis*, *O. bucheri*, *O. contrarius* and
181 *H. plicatilis* are a typical selection of Middle Triassic forms from Iberia being also
182 present in the Ladinian of both the Iberian and Betic Ranges (Fig. 3 B).

183 In general, the Middle Triassic shark communities from the epicontinental seas
184 of the westernmost margin of the Tethys ocean (i.e. east of the actual Iberian Peninsula)
185 were largely dominated, in terms of diversity of species and abundance, by durophagous
186 sharks with crushing dentitions (i.e. *O. bucheri*, *O. contrarius*, *Lissodus* aff. *L. lepagei*,
187 *Lonchidion derenzii*), well adapted for grabbing and crushing prey with a hardened
188 exoskeleton (e.g. crustaceans and mollusks). The teeth in general are small, suggesting
189 that these sharks were also of small size, possibly in turn subsidiary prey of some of the
190 larger aquatic reptiles, e.g. pachypleurosaurs, nothosaurs or ichthyosaurians, that are
191 common in the same areas (Fortuny et al., 2011, Márquez-Aliaga et al., 2019), being
192 these (most probably) the apex predator of these shallow marine environments.

193 The stratigraphic distribution of taxa in East Iberia (i.e. Catalanian, Iberian and
194 Betic basins) is summarized in Figure 3 A. Figure 3B provided synthetic information on
195 the specific abundance. Teeth of *O. bucheri*, *O. contrarius* and *H. plicatilis* from
196 samples of the Olesa Unit in section L’Ametlla represent the first report of Anisian
197 (Pelsonian–lower Illyrian) sharks of the Iberian Peninsula showing that chondrichthyans
198 reached the epicontinental shallow waters of NE Iberia during the first (middle–late
199 Anisian) Triassic transgression. The stratigraphic record of all these three species

extends up into the Carnian (Late Triassic) in the Iberian Peninsula, being also the most representative and abundant taxa.

On the other hand, an important increase of the chondrichthyan diversity in the Triassic Iberian basins occurs in the Longobardian, during the second transgressive–regressive cycle, with the appearance of several taxa, such as *Palaeobates angustissimus*, *Lissodus* aff. *L. lepagei*, *Rhomaleodus*, or *Pseudodalatias henarejensis* (Fig. 3A).

5.2 Biogeographical affinities of chondrichthyan vs. invertebrate faunas from East Iberia during the Triassic westward transgressions of the Tethys Sea

Recent evaluations on the evolution of invertebrate faunas from the Middle Triassic carbonate platforms in eastern Iberia reflect a drastic palaeobiogeographic change during the Upper Anisian. Thus, fossil associations (mainly bivalves, foraminifers, brachiopods and conodonts) from the Landete Formation (Iberian Range) and lower Muschelkalk (Catalan Coastal Range), both dated as Middle–Upper Anisian (Pelsonian–early Illyrian), are almost exclusively comprised by taxa with affinities to the Alpine/Germanic bioprovince, related with the Palaeotethys Ocean. This has been interpreted as the Iberian and Catalanian Basins acting as a palaeogeographic gulf opening northwards, only connected with the Palaeotethys. The presence of geographical barriers should have prevented the entrance of faunas from the Sephardic bioprovince, related with the Neotethys (Escudero-Mozo et al. 2015; see fig. 19). In contrast, fossil associations of invertebrates and conodonts from the upper Anisian to upper Ladinian (late Illyrian–Longobardian; i.e. Cañete Fm., Iberian Range and upper Muschelkalk, Catalan Coastal Range) show a mixed origin, where, along with endemic species, typical species of both the Alpine/Germanic and Sephardic bioprovinces

appear. This palaeobiogeographical heterogeneity of faunas has been linked to the second and most severe, transgressive–regressive cycle (late Anisian–early Carnian), which allowed for the colonization of shallow waters of the Iberian Peninsula by faunas coming from both Tethys provinces: the Germanic/Alpine to the north and the Sephardic to the south.

Noticeably, the stratigraphic distribution of Middle–Upper Triassic chondrichthyans in the Iberian Peninsula differs from this pattern of colonization. Thus, taxa that reached the new shallow-water environments of NE Iberia during the middle Anisian transgression (*i.e.* *Hybodus plicatilis*, *O. bucheri* and *O. contrarius*) do not show the clear Alpine/Germanic affinities reflected in invertebrate and conodont faunas. Only *Hybodus plicatilis* has been reported from other localities in Europe (see supplementary information.) Poland (Niedzwiedzki, 2008); Luxembourg (Delsate and Duffin, 1999); Bulgaria (Stefanov, 1966); Germany and UK (Korneisel et al., 2015). However, this taxon might be a cosmopolitan species from the Tethys realm as teeth assigned to *Hybodus* sp. aff. *Hybodus plicatilis* have been reported from the Middle Triassic of Saudi Arabia (Vickers-Rich et al., 1999). In European localities, *H. plicatilis* occurs in association with other “*Hybodus*”, *Acrodus*, *Polyacrodus*, *Lissodus* and some neoselachian species (Niedzwiedzki, 2008; Delsate and Duffin, 1999; Korneisel et al., 2015), which are absent in the Anisian of the Iberian Peninsula. In the same sense, the genus *Omanoselache* has never been reported from the other Triassic localities in Europe apart from the Iberian Peninsula occurrences. *O. contrarius* occurs in the Middle Triassic of Canada (Johns et al., 1997) and in the Middle–Late Triassic of Gualing, China (Chen et al., 2007), whereas *O. bucheri* is known from the Middle Triassic of Nevada, U.S.A. (Rieppel et al., 1996; Cuny et al., 2001). Pla et al. (2013) suggested that this species might have reached the Iberian Peninsula through the

Sephardic province (i.e., the southern margin of the Tethys Ocean). The recent discoveries of several new *Omanoselache* species from the Middle Permian (Koot et al., 2013) and Lower Triassic (Koot et al., 2015) of Oman confirms the Sephardic affinities of the genus. Then again, the increase of the chondrichthyan diversity occurred during the second transgressive–regressive cycle (Longobardian, Fig 3A) is due to the appearance of several taxa with clear affinities with Alpine/Germanic faunas, such as *Palaeobates angustissimus*, *Lissodus*, *Rhomaleodus*, or *Pseudodolatias* (Fig. 3).

5.3 Dispersal strategies, oceanic palaeocurrents and readjustment of palaeogeographic reconstructions; an overview.

Oceanic currents therefore act as distinctive unidirectional corridors for passive dispersion of organisms with a microscopic pelagic life-stage (Zinsmeister and Emerson, 1979; Jablonski and Lutz, 1983). Consequently, palaeontologists have for a long time recognized that distributions of some marine fossils are useful in reconstructing ancient marine current patterns. As Jablonski and Lutz (1983) pointed out, low (or high) faunal similarity between regions does not necessarily imply large (or small) geographic disjunctions, but rather it can reflect the prevailing current system.

In this sense, dispersion of bivalves, ammonites and foraminifers was largely determined by the dominant systems of marine currents. Most marine invertebrates include a microscopic, free-living, pelagic life history stage—the larval form. Larvae feed in the plankton (except for lecithotrophs) exhibiting long pelagic durations and high dispersal capacity. Distribution of these taxa is largely determined by ocean currents transporting larvae between distant patches of suitable habitat for adults, often over great distances (Scheltema, 1986; Jablonski and Lutz, 1983; Pechenik, 1999; Kinlan et al., 2005; Trakhtenbrot et al., 2005; Trembl et al., 2008; Wangensteen et al.,

2017). On the other hand, ammonites underwent a direct development lacking a distinct larval stage (as other, extant cephalopods: nautilus, cuttlefishes, squids and octopuses) consisting of four ontogenetic stages (embryonic/egg, neanic, juvenile, and adult; see e.g. Westermann, 1996; Klug et al 2015; Zell and Stinnesbeck, 2016; Tajika *et al*, 2018). Nevertheless, most authors consider that ammonite hatchlings – the neanic stage – were planktic, with ocean currents enhancing their dispersion. In fact, the patchy yet widespread palaeogeographic distribution of numerous ammonite species suggests that extensive planktic dispersal with ocean currents played an important role in their dispersion patterns (Zell and Stinnesbeck, 2016). The same is valid for foraminifers. Thus, for benthic attached species, such as those fossils found in NE Iberia (L. Marquez pers. comm. (2018), but see also the list of species in Escudero-Mozo et al., 2015), the most common ways of dispersion were, based on living forms (1) release of gametes, zygotes, or of embryonic agamonts/gamonts to the water column, and (2) adaption to a meroplanktonic juvenile stage with passive spread by currents. Moreover, shallow-water taxa can also be transported long distances by floating objects, larger animals and/or by suspended particles (Myers, 1936, Murray, 2006, Nomura et al, 2010, Alve 1999).

In contrast, chondrichthyans lack any planktic stage but undergo a direct development after internal fertilization. They possess large eggs with a high amount of yolk and produce ovoviviparous or oviparous juveniles (there are some viviparous species). In any case, the distribution of the chondrichthyan species largely depends on the dispersal abilities of adults and much less on marine currents. In this sense, it is well known that the vagility of sharks is directly proportional to body size (e.g. Musick et al., 2004). As a whole, small-bodied coastal sharks present a very low vagility and short forage distances, with oceanic distances, deep channels or strong regional currents

300 acting as dispersal barriers (Musick et al., 2004; Munroe et al., 2015; Whitney et al.,
301 2012). Thus, dispersion of coastal species usually requires continuous appropriate
302 habitats across regionally small geographical scales.

304 *5.4. Palaeogeography and dominant marine currents at the westernmost Tethys*

305 To consider the different dispersal capability of organisms is important in order
306 to evaluate the timing and nature of palaeobiogeographical barriers. Taxa with planktic
307 stages can provide considerable information for reconstructing marine palaeocurrent
308 systems while taxa lacking these stages (or with low-dispersal larvae) can be used to
309 infer continuity among continental shelves, both groups being affected differentially by
310 palaeobiogeographic barriers (see e.g. Shuto, 1974; Jablonski and Lutz, 1983).

311 Following these general rules, the existing differences in the colonizing patterns of
312 invertebrate vs chondrichthyan faunas in the Triassic shallow waters of E Iberia can be
313 reasonably well explained. However, it requires some adjustments of previous
314 palaeogeographic reconstructions of the westernmost Tethyan realm (see Escudero-
315 Mozo et al. 2015; their figure 19 and references therein).

316 We suggest that the colonization by bivalves, ammonites and foraminifers, during both
317 the late Pelsonian–early Illyrian and the late Illyrian–Longobardian marine
318 transgressions, was largely determined (or enhanced) by the prevailing marine currents
319 in the westernmost Tethys and by the transgression of the sea itself (Fig. 4).

320 Thus, in late Pelsonian–early Illyrian times, marine currents from the
321 Palaeotethys enhanced the arrival of planktic organisms to the new Catalan and Iberian
322 marine basins. At this time, the Cimmerian microcontinent, which controlled the
323 regional palaeogeography in the westernmost Tethys (Yin and Song, 2013; Escudero-
324 Mozo et al., 2015), would have acted as a barrier for marine currents (if any) of the

incipient Neotethys. However, the occurrence of chondrichthyans from the Sephardic domain indicates the formation of narrow, probably temporary connections (acting as alternative routes for the entrance of sharks) between Palaeo- and Neotethys during the northward displacement of the Cimmerian microcontinent (Fig. 4). During the next greater and more generalized sea level rise (Illyrian–Longobardian transgression), the Tethys sea flooded vast areas of the westernmost Cimmerian microcontinent, such as Alboran, Majorca, Minorca, Sardinia, Corsica, Tuscany, Apulia, Mani (Escudero-Mozo, 2015: figure 19). Subsequently, the Palaeotethys and Neotethys became broadly connected, allowing for the entrance of planktonic organisms (i.e. bivalve larvae, naetic ammonites and foraminifers, among others) into E Iberia, which were transported by marine currents from the Neotethys (Fig. 4).

Notably, during both colonization events, conodonts follow the same dispersion patterns as invertebrates and not those of chondrichthyans. Conodonts are considered by some authors a basal group of vertebrates (e.g. Donoghue et al., 2000) although their placement in this group is seriously questioned by other authors (e.g. Turner et al., 2010). Little is known about their reproductive strategies (with the finding of putative conodont eggs, von Bitter and Pohl, 2009), although it has been suggested, based on their crown growth and function (see Martínez-Pérez et al, 2014), that conodonts could had developed a larval stage (Shirley et al., 2018). Data here discussed are compatible with conodonts having a passive dispersive planktic stage. Planktic larval stages (or eggs) are present in several osteichthyan (bony fishes; common in teleosteans) but unknown in other fishes. A larval stage is present in agnathan lampreys (but not in mixines) and consists of a large sedentary burrowed larva in freshwater streams (e.g. Kelly and King, 2001), and could to be present in other extinct Palaeozoic agnathan fishes (Ferrón et al. 2018). The cosmopolitanism and high dispersal of Palaeozoic

conodonts might also agree with a planktic dispersion drifted by ocean currents; however, further and more extensive studies on this topic are necessary to support this idea.

It should further be noted that, despite that our interpretation are well supported by data from Pla et al.'s (2013), Manzanares et al. (2018) and here, the number of specimens and taxa in the Iberian Peninsula is relatively low. Moreover, triassic shark faunas are still poorly known in general. Therefore, new findings could change (or refine) this interpretation in the future.

6. Conclusions

This work completes a series of studies on the chondrichthyan faunas from Middle–Late Triassic of Iberia which has changed the previously inaccurate perception that chondrichthyans were rare, or completely absent, in the Iberian Peninsula. These studies provide now a detailed overview of the evolution of chondrichthyan communities in the new shallow-water environments developed in E Iberia (westernmost Tethys) during the Triassic westward transgression of the Tethys Sea. Chondrichthyan communities were largely dominated by small durophagous coastal sharks with crushing dentitions well adapted for preying on crustaceans and mollusks. Shark that reach NE Iberia during the first transgressive pulse (middle–late Amirian) of the Tethys Sea show clear affinities with taxa from the Sephardic bioprovince, related to the Neotethys Ocean (i.e. *Omanoselache*). The notable increase in chondrichthyan diversity that occurs in E Iberia during the second transgressive pulse (Ladinian) is due to the arrival of new taxa with clear affinities with European faunas from the Palaeotethys Ocean (i.e. *Paleobates angustissimus*, *Lissodus*, *Hybodus*, *Rhomaleodus*, or *Pseudodalatias*). This colonization pattern is somewhat inverse to those recently

evaluated for other fossil groups. Thus, Anisian bivalves, foraminifers, brachiopods and conodont faunas from E Iberia are almost exclusively composed of taxa from the Alpine/Germanic bioprovince. In contrast invertebrate and conodonts from the Sephardic bioprovince reach the shallow waters of the Iberian Peninsula for the first time during the second (Ladinian) transgressive pulse.

The existing differences between the colonization patterns of sharks and other groups are explained on the basis of their dissimilar dispersion strategies: dispersion of bivalves, ammonites and foraminifers, is largely determined by oceanic currents whereas dispersion of coastal sharks depends of adult vagility (very low) and requires continuous appropriate habitats. Based on this, we propose some adjustments of previous palaeogeographic reconstructions including: (1) the existence of narrow, probably short-term, shallow-water corridors between Palaeo- and Neo-Tethys during the Anisian; (2) the Cimmerian microcontinent would have acted as a barrier for the oceanic currents in the incipient Neotethys, while currents from the Palaeotethys transported planktic organisms to the new Catalan and Iberian Marine Basins, and (3) during the next sea-level rise (Ladinian), the Tethys sea overlapped vast areas of the westernmost Cimmerian microcontinent allowing marine currents from the Neotethys to extend toward the Palaeotethys reaching E Iberian epicontinental seas.

6. Acknowledgements

This work has been partially funded by the Ministry of Science, Innovation and Universities, Research Project CGL2014-52662, the Formación de Personal de Investigación (FPI) Grant BES-2015-072618 and the European Regional Development Fund (ERDF) “A Way to Make Europe.” We also want to thank Dr. Susan Turner (Brisbane) for the English review. We thank Dr. Christian Klug, Dr. Jürgen Kriwet and

Prof. Howard Falcon-Lang whose comments helped to improve and clarify this manuscript.

7. References

Arche, A., López-Gómez, J., Marzo, M., and Vargas, H. 2004. The siliciclastic Permian-Triassic deposits in central and northeastern Iberian Peninsula (Iberian, Ebro and Catalan basins): a proposal for correlation. *Geol. Acta*, 2(4):305–320.
<https://doi.org/10.1344/105.000001422>

Alve, E. 1999. Colonization of new habitats by benthic foraminifera: a review. *Earth-Sci. Rev.*, 46(1-4):167-185.

Botella, H., Plasencia, P., Márquez-Aliaga, A., Cuny, G., and Dorka, M. 2009. *Pseudodalatias henarejensis* nov. sp. A new pseudodalatiid (Elasmobranchii) from the Middle Triassic of Spain. *J. Vertebr. Paleontol.*, 29:1–20.
<https://doi.org/10.1671/039.029.0425>

Calvet, F., and Marzo, M. 1994. El Triásico de las cordilleras costero catalanas: estratigrafía, sedimentología, y análisis secuencial. *Alfredo Arche*. 53 p.

Calvet, F., and Ramon, X., 1987. Estratigrafía, sedimentología y diagénesis del Muschelkalk inferior de los Catalánides. *Cuadernos de Geología Ibérica*, 11: 141-169.

Calvet, F., March, M., and Pedrosa, A. 1987. Estratigrafía, sedimentología y diagénesis del Muschelkalk superior de los Catalánides. *Cuadernos Geología Ibérica*, 11:171–197.

Calvet, F., Tucker, M. E., and Henton, J. M. 1990. Middle Triassic carbonate ramp systems in the Catalan Basin, northest Spain: facies, systems tracks, sequences

and controls. Institut of Association of Sedimentology, Special Publications,
9:79– 08.

Chen, L., Cuny, G., and Wang, X. 2007. The chondrichthyan fauna from the Middle–
Late Triassic of Guanling (Guizhou province, SW China). *Hist. Biol.*, 19:291–
300. <https://doi.org/10.1080/08912960701248234>

Crasquin, S., Forel, M. B., Feng, Q., Yuan, A., Baudin, F., and Collin, P. Y. 2010.
Ostracods (Crustacea) through the Permian-Triassic boundary in South China:
the Meishan stratotype (Zhejiang Province). *J. Syst. Palaeontol.*, 8:331-370.
<https://doi.org/10.1080/14772011003784992>

Cuny, G., Rieppel, O., and Sander, P. M. 2001. The shark fauna from the Middle
Triassic (Anisian) of North–Western Nevada. *Zool. J. Linn. Soc-Lond.*, 13:285–
30. <https://doi.org/10.1006/zjla.2000.0273>

Delsate, D., and Duffin, C. J. 1999. A new fish fauna from the Middle Triassic (Upper
Muschelkalk) of Moersdorf (Grand Duchy of Luxembourg). *Travaux
Scientifiques du Musée National d'Histoire Naturelle de Luxembourg*, 32 :5-53.

De Vicente, G., Vegas, R., Muñoz-Martín, A., Van Wees, J. D., Casas-Sáinz, A.,
Sopeña, A., Sánchez-Moya, Y., Arche, A., López-Gómez, J., Olaiz, A., and
Fernández-Lozano, J. 2009. Oblique strain partitioning and transpression on an
inverted rift: The Castilian Branch of the Iberian Chain. *Tectonophysics*,
470(3) :224–242. <https://doi.org/10.1016/j.tecto.2008.11.003>

Donoghue, P. C. J., Forey, P. L., and Aldrige, R. J. 2000. Conodont affinity and
chordate phylogeny. *Biol. Rev.* 75(02):191 – 251.
<https://doi.org/10.1111/j.1469-185X.1999.tb00045.x>

- Escudero-Mozo, M. J. 2015. The Triassic carbonate platforms (Muschelkalk) of eastern Iberia and Minorca: implications for the paleogeographic evolution of Western Tethys. PhD dissertation. Universidad Complutense de Madrid, Madrid, 370 p.
- Escudero-Mozo, M. J., Márquez-Aliaga, A., Goy, A., Martín-Chivelet, J., López-Gómez, J., Márquez, L., Plasencia, P., Pla C., Marzo, M., and Sánchez-Fernández, D. 2015. Middle Triassic carbonate platforms in eastern Iberia: Evolution of their fauna and palaeogeographic significance in the western Tethys. *Palaeogeogr., Palaeoclimatol.*, 417:236–260.
<https://doi.org/10.1016/j.palaeo.2014.10.041>
- Escudero-Mozo, M.J., Martín-Chivelet, J., Goy, A., and López-Gómez, J. 2014. Middle-Upper Triassic carbonate platforms in Minorca (Balearic islands): implications for Western Tethys correlations. *Sediment. Geol.*, 310:41–58.
<https://doi.org/10.1016/j.sedgeo.2014.06.002>
- Ferrón, H., Pla, C., Martínez-Pérez, C., Escudero-Mozo, M. J., and Botella, H. 2014. Morphometric discriminant analysis of isolated chondrichthyan scales for palaeoecological inferences: the Middle Triassic of the Iberian Chain (Spain) as a case of study. *J. Iber. Geol.*, 40(1):87–97.
https://doi.org/10.5209/rev_JIGE.2014.v40.n1.44089
- Ferrón, H. G., Martínez-Pérez, C., Turner, S., Manzanares, E., and Botella, H. 2018. Patterns of ecological diversification in thelodonts. *Palaeontology*, 61(2):303–315. <https://doi.org/10.1111/pala.12347>
- Fortuny, J., Bolet, A., Sellés, A. G., Cartanyà, J., and Galobart, À. 2011. New insights on the Permian and Triassic vertebrates from the Iberian Peninsula with emphasis on the Pyrenean and Catalanian basins. *J. Iber. Geol.*, 37(1):65–86.
https://doi.org/10.5209/rev_JIGE.2011.v37.n1.5

- Galán-Abellán, B., López-Gómez, J., Barrenechea, J. F., Marzo, M., De la Horra, R.,
and Arche, A. 2013. The beginning of the Buntsandstein cycle (Early–Middle
Triassic) in the Catalan Ranges, NE Spain: sedimentary and palaeogeographic
implications. *Sediment. Geol.*, 296:86–102.
<https://doi.org/10.1016/j.sedgeo.2013.08.006>
- Gaspar-Escribano, J. M., Garcia-Castellanos, D., Roca, E., and Cloetingh, S. A. P. L.
(2004). Cenozoic vertical motions of the Catalan Coastal Ranges (NE Spain):
The role of tectonics, isostasy, and surface transport. *Tectonics*, 23(1):1–18.
<https://doi.org/10.1029/2003TC001511>
- Goy, A. 1995. Ammonoideos del Triasico Medio de España: bioestratigrafía y
correlaciones. *Cuadernos de Geología Ibérica*, 19:21–60.
- Jablonski, D., and Lutz, R. A. 1983. Larval ecology of marine benthic invertebrates:
paleobiological implications. *Biol. Rev.*, 58(1):21-89.
- Johns, M. J., Barnes, C. R., and Orchard M. J. 1997. Taxonomy and Biostratigraphy of
Middle and Late Triassic elasmobranch ichthyoliths from northeastern British
Columbia. *Geol. Surv. Canada*, 502:1–235.
- Kelly, F. L., and King, J. J. 2001. A review of the ecology and distribution of three
lamprey species, *Lampetra fluviatilis* (L.), *Lampetra planeri* (Bloch) and
Petromyzon marinus (L.): a context for conservation and biodiversity
considerations in Ireland. In *Biology and Environment: Proceedings of the Royal
Irish Academy* Proc. Royal Ir. Acad. (ed.).165-185
- Kinlan, B. P., Gaines, S. D., and Lester, S. E. 2005. Propagule dispersal and the scales
of marine community process. *Diver. Distrib.*, 11(2):139-148.
<https://doi.org/10.1111/j.1366-9516.2005.00158.x>

497 Klug, C., Korn, D., De Baets, K., Kruta, I., and Mapes, R. H. 2015. Ammonoid
 498 paleobiology: from anatomy to ecology (Vol. 43). Springer.

499 Koot, M. B., Cuny, G., Tintori, A., and Twitchett, R. J. 2013. A new diverse shark
 500 fauna from the Wordian (Middle Permian) Khuff Formation in the interior
 501 haushi-huqf area, Sultanate of Oman. *Palaeontology*, 56(2):303–343.
 502 <https://doi.org/10.1111/j.1475-4983.2012.01199.x>

503 Koot, M. B., Cuny, G., Orchard, M. J., Richoz, S., Hart, M. B., and Twitchett, R. J.
 504 2015. New hybodontiform and neoselachian sharks from the Lower Triassic of
 505 Oman. *J. Syst. Palaeontol.*, 13(10):891-917.
 506 <https://doi.org/10.1080/14772019.2014.963179>

507 Korneisel, D., Gallois, R. W., Duffin, C. J., and Benton, M. J. 2015. Latest Triassic
 508 marine sharks and bony fishes from a bone bed preserved in a burrow system,
 509 from Devon, UK. *Proc. Geol. Ass.*, 126(1):130– 42.
 510 <https://doi.org/10.1016/j.pgeola.2014.11.004>

511 López-Gómez, J., Arche, A. Calvet, F., and Goy, A. 1998. Epicontinental marine
 512 carbonate sediments of the Middle and Upper Triassic in the westernmost part of
 513 the Tethys Sea, Iberian Peninsula. In: *Epicontinental Triassic*. Bachmann, G.H.
 514 and Lerche, I. (eds). E. Schweizerbart'sche Verlagsbuchhandlung (Nägele u.
 515 Obermiller) Stuttgart. *Zbl. Geo. Pal.*, 2(9-10):1033–1084.

516 Manzanares, E., Pla, C., Ferrón, H., and Botella, H. 2018. Middle-Late Triassic
 517 chondrichthyan remains from the Betic Range (Spain). *J. Iber. Geol.*, 44(1):129-
 518 138. <https://doi.org/10.1007/s41513-017-0027-1>

519 Manzanares, E., Pla, C., Martínez-Pérez, C., Ferrón, H., and Botella, H. 2017.
 520 *Lonchidion derenzii* sp. nov., a new lonchidiid shark (Chondrichthyes,
 521 Hybodontiforms) from the Upper Triassic of Spain, with remarks on lonchidiid

522 enameloid. J. Verteb. Paleontol., 37(1):e1253585.

523 <https://doi.org/10.1080/02724634.2017.1253585>

524 Márquez-Aliaga, A., Valenzuela-Ríos, J. I., Calvet, F., and Budurov, K. 2000. Middle
525 Triassic conodonts from northeastern Spain: biostratigraphic implications. Terra
526 Nova, 12:77–83. <https://doi.org/10.1111/j.1365-3121.2000.00273.x>

527 Martindale, R. C., Foster, W., and Velledits, F. 2019. The survival, recovery, and
528 diversification of metazoan reef ecosystems following the end-Permian mass
529 extinction event. Palaeogeogr. Palaeocl. 513:100–115.
530 <https://doi.org/10.1016/j.palaeo.2017.08.014>

531 Martínez-Pérez, C., Plasencia, P., Jones, D.O., Kolar-Jurkovsek, T., Sha, J., Botella, H.,
532 and Donoghue, P.C.J. 2014b. There is no general model for occlusal kinematics
533 in conodonts. Lethaia.47:547–555. <https://doi.org/10.1130/G37547.1>

534 Marzo, M. 1980. El Bundstandstein de los Catalánides: Estratigrafía y procesos
535 sedimentarios. PhD. dissertation, Universitat de Barcelona, Barcelona, p. 192.

536 Munroe, S. E., Heupel, M. R., Fisk, A. T., Logan, M., and Simpfendorfer, C. A. 2015.
537 Regional movement patterns of a small-bodied shark revealed by stable-isotope
538 analysis. J. Fish Biol., 86(5):1567-1586. <https://doi.org/10.1111/jfb.12660>

539 Murray, J. W. 2006. Ecology and applications of benthic foraminifera. Cambridge
540 University Press (ed.), 318 p.

541 Musick, J. A., Harbin, M. M., and Compagno, L. J. 2004. Historical zoogeography of
542 the Selachii. In *Biology of sharks and their relatives*, 2004:3354.

543 Myers, E. H. 1936. The life-cycle of *Spirillina vivipara* Ehrenberg, with notes on
544 morphogenesis, systematics and distribution of the foraminifera. J. Roy.
545 Microsc. Soc., 56(2):120-146.

546 Niedzwiedzki, R. 2008. Triassic selachians in Poland, a review. *Ichthyolith Issues*,
547 Special Publication, 11:38-42.

548 Nomura, R., Seto, K., and Tsujimoto, A. 2010. High tide dispersion of marine benthic
549 foraminifera into brackish waters: implications for dispersion processes during
550 sea-level rise. *Laguna*, 17:15-21

551 Pla, C., Márquez-Aliaga, A., and Botella, H. 2013. The chondrichthyan fauna from the
552 Middle Triassic (Ladinian) of the Iberian Range (Spain). *J. Verteb. Palaeontol.*,
553 33(4): 770-785. <https://doi.org/10.1080/02724634.2013.748668>

554 Plasencia, P. 2009. Bioestratigrafía y paleobiología de conodontos del Triásico Medio
555 del Sector Oriental de la Península Ibérica. PhD. dissertation, Universitat de
556 València, Valencia, 408 p.

557 Pechenik, J. A. 1999. On the advantages and disadvantages of larval stages in benthic
558 marine invertebrate life cycles. *Mar. Ecol. Prog. Ser.*, 177:269-297. doi:
559 10.3354/meps177269

560 Ramon, X., and Calvet, F., 1987. Estratgrafia y sedimentología del Muschelkalk inferior
561 del dominio Montseny-Llobregat (Catalánides). *Estudios geológicos*, 43(5-
562 6):471-487.

563 Rieppel, O., Kindlimann, R., and Bucher, H. 1996. A new fossil fish fauna from the
564 Middle Triassic (Anisian) of North-Western Nevada. In *Mesozoic Fishes*,
565 *Systematics and Paleoecology*. Proceedings of the 1st International Meeting
566 Eichstätt 1993, G. Arratia and G. Viohl (eds.), 501–512.

567 Scheltema, R. S. 1986. On dispersal and planktonic larvae of benthic invertebrates: an
568 eclectic overview and summary of problems. *Bull. Mar. Sci.*, 39(2):290-322.

- Scotese, C. R. 1994. Early Triassic paleogeographic map. In *Pangea: Paleoclimate, Tectonics, and Sedimentation During Accretion, Zenith, and Breakup of a Supercontinent: Boulder, Colorado*. G. D. Klein (ed.). Geol. S. Am. S. 288 p.
- Shirley, B., Grohgan, M., Bestmann, M., and Jarochovska, E. 2018. Wear, tear and systematic repair: testing models of growth dynamics in conodonts with high resolution imaging. *Proc. R. Soc. B* 285: 20181614.
<http://dx.doi.org/10.1098/rspb.2018.1614>
- Shuto, T. 1974. Larval ecology of prosobranch gastropods and its bearing on biogeography and paleontology. *Lethaia*, 7(3):239-256.
- Sopeña, A., and de Vicente, G. 2004. Cordilleras Ibérica y Costero-Catalana. Rasgos generales. In: *Geología de España*. J. A. Vera, (ed). SGE-IGME, 467-470.
- Stampfli, G. M., and Borel, G. D. 2002. A plate tectonic model for the Palaeozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons. *Earth Planet. Sc. Lett.*, 196:17-33
[https://doi.org/10.1016/S0012-821X\(01\)00588-X](https://doi.org/10.1016/S0012-821X(01)00588-X)
- Stefanov, S. 1966. Fischreste aus der Trias Bulgariens. *Academie Bulgare des Sciences, Travaux sur la Geologie de Bulgarie, Série Paléontologie*, 8:123–129.
- Tajika, A., Nützel, A., and Klug, C. 2018. The old and the new plankton: ecological replacement of associations of mollusc plankton and giant filter feeders after the Cretaceous? *PeerJ*, 6:e4219- <https://doi.org/10.7717/peerj.4219>
- Trakhtenbrot, A., Nathan, R., Perry, G., and Richardson, D. M. 2005. The importance of long-distance dispersal in biodiversity conservation. *Diver. Distrib.*, 11(2):173-181. <https://doi.org/10.1111/j.1366-9516.2005.00156.x>
- Treml, E. A., Halpin, P. N., Urban, D. L., and Pratson, L. F. 2008. Modeling population connectivity by ocean currents, a graph-theoretic approach for marine

conservation. *Landscape Ecol.*, 23(1):19-36. <https://doi.org/10.1007/s10980-007-9138-y>

Turner, S., Burrow, C. J., Schultze, H. P., Blieck, A., Reif, W. E., Rexroad, C. B., Bultynck, P., and Nowlan, G. S. 2010. False teeth: conodont-vertebrate phylogenetic relationships revisited. *Geodiversitas*. 32(4) :545-595. <https://doi.org/10.5252/g2010n4a1>

Vargas, H., Gaspar-Escribano, J. M., López-Gómez, J., Van Wees, J. D., Cloetingh, S., de La Horra, R., and Arche, A. 2009. A comparison of the Iberian and Ebro Basins during the Permian and Triassic, eastern Spain: A quantitative subsidence modelling approach. *Tectonophysics*, 474(1):160–183. doi:10.1016/j.tecto.2008.06.005

Vegas, R. 1975. Wrench (transcurrent) fault system of the south-western Iberian Peninsula, palaeontogeographic and morphostructural implications. *Geol. Res.*, 64:266–278.

Vickers-Rich, P., Rich, T. H., and Coffa, F. 1999. Wildlife of Gondwana: dinosaurs and other vertebrates from the ancient supercontinent (p. 304). Bloomington and Indianapolis: Indiana University Press.

von Bitter P. H., and Pohl, J. A. 2009. Possible Conodont Eggs from the Pennsylvanian (Middle Desmoinesian) of NW Illinois, U.S.A. *Permophiles*. 53(1):8-9.

Wangenstein, O. S., Turon, X., and Palacín, C. 2017. Reproductive strategies in marine invertebrates and the structuring of marine animal forests. In *Marine Animal Forests*. Rossi, S. (ed.) 1-24. doi: 10.1007/978-3-319-17001-5_52-1

Westermann, G. E. 1996. Ammonoid life and habitat. In *Ammonoid paleobiology*. Springer (ed.). 607-707.

- Whitney, N. M., Robbins, W. D., Schultz, J. K., Bowen, B. W., and Holland, K. N. 2012. Oceanic dispersal in a sedentary reef shark (*Triaenodon obesus*): genetic evidence for extensive connectivity without a pelagic larval stage. *J. Biogeogr.*, 39(6):1144-1156. <https://doi.org/10.1111/j.1365-2699.2011.02660.x>
- Yan, J. X., and Zhao, K. 2001. Permo-Triassic paleogeographic, paleoclimatic and paleoceanographic evolutions in eastern Tethys and their coupling. *Sci. China Ser. D Earth Sci.*, 44:968–978. <https://doi.org/10.1007/BF02875390>
- Yin, H., and Song, H. 2013. Mass extinction and Pangea integration during the Paleozoic-Mesozoic transition. *Sci. China Earth Sci.*, 56(11):1791-1803. <https://doi.org/10.1007/s11430-013-4624-3>
- Zell, P., and Stinnesbeck, W. 2016. Paleobiology of the latest Tithonian (Late Jurassic) ammonite *Salinites grossicostatum* inferred from internal and external shell parameters. *PloS one*, 11(1), e0145865.
- Zinsmeister, W. J., and Emerson, W. K. 1979. Role of passive dispersal in the distribution of hemipelagic invertebrates, with examples from the tropical Pacific Ocean. *Veliger*, 22(1):32-40.

Figure captions

Fig. 1. Geographical and geological settings A) Geographical setting of Catalanian Coastal Range (CCR) with location of sections studied here, modified from Escudero-Mozo (2015); B) Stratigraphic columns of the sections studied. Only levels where chondrichthyans remains were recovered are marked in the columns; for complete columns see Plasencia (2009).

Fig. 2. Chondrichthyan teeth from the Middle Triassic of the Catalan Coastal

Ranges, Spain. A-D) *Hybodus plicatilis*; A-B) labial and lingual view, MGUV-36111; C-D) basal and occlusal view, MGUV-36112; E-J) *Omanoselache bucheri* E) labial view, MGUV-36121; F) labial view, MGUV-36122; G-H) labial and lingual view, MGUV-36123; I-J) labial and lingual view, MGUV-36127; K-P) *Omanoselache contrarius*; K-L) labial and lingual view, MGUV-36127; M-N) labial and lingual view, MGUV-36120; O-P) MGUV-36118. Q-R) *Pseudodalatias henarejensis* in labial and lingual view, respectively, MGUV-36113. All scale bars equal 100 μ m.

Fig. 3. Stratigraphic distribution and abundance of chondrichthyans in Middle-Late

Triassic of the Iberian Peninsula A) Stratigraphic distribution of taxa. *H. plicatilis*, *O. bucheri*, *O. contrarius* and *P. henarejensis* are the typical fauna that can be found in any Middle Triassic basin of the Iberian Peninsula. *Lissodus* aff. *L. lepagei* appears only in the Iberian Ranges and Betic Ranges, whereas *Hybodus bugarensis* and *Palaeobates angustissimus* have been only found in the Iberian Ranges. *Lonchidion derenzii* and cf. *Rhomaleodus budurovi* are only present at the Betic Ranges. B) Histograms showing the number of chondrichthyan teeth recovered (for each species) from in the three basins of the Iberian Peninsula considered in this study (the Iberian Range, the Betic Range and the Catalan-Coastal Range).

Fig. 4. Faunal colonization patterns of East Iberia during the Anisian and

Ladinian westward transgression of the Tethys ocean. A) Global palaeogeographic map, Early Triassic (Induan), with schematic surface-current circulation. B-C) detailed palaeogeographic reconstructions of the westernmost Tethyan realm for the Anisian (B) and Ladinian (C) with interpreted colonization routes for chondrichthyans and

667 invertebrates. Palaeogeographic maps modified from Scotese (1994) (A), Stampfli and
668 Borel (2002) and Escudero-Mozo et al. (2015) (B); oceanic currents interpreted
669 according with Yan and Zhao (2001); Crasquin et al. (2010) and Martindale et al.
670 (2019). The black star marks the position of the Catalan Coastal Range, the black
671 circle marks the Betic Range, the black diamond represents the Iberian Range.

B

A

B

A Early Triassic (250 Ma)

B Anisian (240 Ma)

C Ladinian (230 Ma)

We confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome. The manuscript has been read and approved by all named authors and that there are no other people who satisfied the criteria for authorship who are not listed.

We further confirm that the order of authors listed in the manuscript has been approved by all of us. We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have follow the regulations of our institutions concerning intellectual property.

Signed by all authors as follows:

Esther Manzanares

María José Escudero Mozo

Humberto Ferrón

Carlos Martínez Pérez

Héctor Botella Sevilla

Supplementary material

Systematic Palaeontology

Class CHONDRICHTHYES Huxley, 1880

Subclass ELASMOBRANCHII Bonaparte, 1838

Order EUSELACHII Hay, 1902

Superfamily HYBODONTOIDEA Owen, 1846

Family HYBODONTIDAE Owen, 1846

Genus *HYBODUS* Agassiz, 1837

Type species—*Hybodus reticulatus* Agassiz, 1837

Occurrence— Lower Jurassic, Lias of Lyme Regis, England (Woodward, 1916).

HYBODUS PLICATILIS Agassiz, 1843

Figure 2, A–D

Material—Two incomplete teeth from the Rasquera–Benifallet section (CAME-14 level: MGUV-36111/-36112).

Description—Most of our material is broken, with only almost complete two multicuspid teeth. They show a central cusp flanked by up to two pairs of cusplets, all of them lingually inclined (Fig. 2 A–D). Ornamentation ridges run from the tooth shoulder towards the tip of the cusp and cusplets. In the main cusp, the ridges do not reach the apical part; they cannot be seen in the lateral cusplets due to their damage (Fig. 2A–B). The vascularization is of the anaulacorhize type (Fig. 2 A–B).

Remarks —Several recent authors agree that the genus *Hybodus* is a polyphyletic group and that it does not correspond to a natural group (Rees, 1999; Underwood and Rees, 2002; Rees and Underwood, 2008; Ginter et al., 2010; Cappetta, 2012; Leuzinger et al., 2017), a statement with which we fully agree. A large number of isolated teeth described in

the literature have been assigned to *Hybodus*, although a revision of this material is required to verify, which of them fit completely the original diagnosis of the genus. However, this is beyond the scope of the present work; nevertheless, our material shows characteristics of the type species *Hybodus reticulatus*: well-defined and well-ornamented conical cusps; lateral cusplets decreasing in height, frequently inclined lingually; and a fairly shallow base (see Maisey, 1987; Duffin, 1993; Ginter et al., 2010). Thus, we choose to maintain the name of the genus *Hybodus* until a revision of the genus is done.

Occurrence—Muschelkalk, Middle Triassic of Schwenningen, Germany (Agassiz, 1843); Muschelkalk, Middle Triassic of Monte Giorgio, Switzerland (von Meyer, 1849; Rieppel, 1981); Muschelkalk, Middle Triassic of Luxemburg (Delsate, 1992, 1993); Middle Triassic, Iberian Ranges, Spain (Pla et al., 2013), Middle–Late Triassic of the Betic Ranges, Spain (Manzanares et al., 2018); Middle Triassic, Catalan Coastal Ranges, Spain.

Order *INCERTAE SEDIS*

Family HOMALODONTIDAE Mutter, De Blanger and Neuman, 2008

Genus *OMANOSELACHE* Koot, Cuny, Tintori and Twitchett, 2013

Type species—*Omanoselache hendersoni* Koot, Cuny, Tintori and Twitchett, 2013

Occurrence— Middle Permian, Wordian from the Khuff Formation, Oman (Koot et al., 2013)

OMANOSELACHE BUCHERI (Cuny, Rieppel and Sander, 2001)

Figure 2, E-J

Material—Eight complete teeth and more than 20 fragments from the Rasquera–Benifallet section (CRB-1 level: MGUV-36121/-36122/-36126/-361627/-36128; and CRB-3 levels MGUV-MGUV-361216/36128; MGUV/-36123/-36130/-36131), l’Ametlla section

(CAME-12: MGUV-36124; and CAME-13 levels; MGUV-36125) and the Tivissa section (CT-8 level: MGUV-36129).

Description—Teeth are labiolingually compressed and elongated mesiodistally. All display a well-developed occlusal crest and a pyramidal main cusp (Fig. 2 E-H). Undulated vertical ornamentation ridges originate on the occlusal crest and appear on the labial and lingual sides (Fig. 2 E-F, I). A rounded lingual peg, characteristic of this species, is located under the main cusp (Fig. 2 E-G, I). Vascularisation is anaulacorhize with a row of well-developed foramina opening on the labial and lingual sides (Fig. 2 E, G).

Remarks—*Omanoselache bucheri* teeth are the most widespread, abundant and well-preserved chondrichthyan remains from the Iberian Peninsula (Pla et al., 2013; Manzanares et al., 2017). Pla et al. (2013), described three different morphotypes related to the different position they occupied in the jaws: symphyseal, mesial, or latero-distal teeth. Most of our material can be classified as their mesial teeth. These teeth exhibit one main cusp and are arched, showing a ‘boomerang’ appearance (Fig. 2 G-H). A few show a blunt and reduced main cusp and an overall more rectilinear appearance (Fig. 2 I-J), which are the characteristics of latero-distal teeth. The presence of a lingual peg, but no labial one, and the rounded edges (Fig. 2E-F, G, I) permit us to differentiate them from *Omanoselache contrarius* (Johns, Barnes and Orchard, 1997).

Occurrence—Middle Triassic, North-West of Nevada (Rieppel et al., 1996; Cuny et al., 2001); Middle Triassic, Iberian Ranges, Spain (Pla et al., 2013); Middle–Late Triassic from the Betic Ranges of Spain (Manzanares et al., 2018); Middle Triassic, Catalan Coastal Ranges, Spain.

OMANOSELACHE CONTRARIUS (Johns, Barnes and Orchard, 1997)

Figure 2 K-P

Material—Four complete teeth and several fragments from the Rasquera–Benifallet section (CRB-1 level: MGUV-36117; and CRB-3 levels: MGUV-3619/-36120) and l’Ametlla section (CAME-12 level: MGUV-36118).

Description—Teeth elongated mesiodistally and compressed labiolingually. All of them show one main cusp flanked by up to two lateral cusplets (Fig. 2 K-P). Ornamentation consists of vertical ridges that originate from the apex of the cusps and extend to the tooth shoulder. The teeth present two pegs (labial and lingual) under the main cusp (Fig. 2 M-N). As in *O. bucheri*, the vascularization is anaulacorhize (Fig. 2 K-P).

Remarks—All the teeth recovered of this species present evident wear signs on the cusps.

Occurrence—Middle Triassic of British Columbia, Canada (Johns et al., 1997); Middle–Late Triassic of Yang Liu Jing, China (Chen et al., 2007); Middle Triassic, Iberian Ranges, Spain (Pla et al., 2013); Middle–Late Triassic, Betic Ranges, Spain (Manzanares et al., 2018); Middle Triassic, Catalan Coastal Ranges, Spain.

Order *INCERTAE SEDIS*

Family PSEUDODALATIIDAE Reif, 1978

Genus *PSEUDODALATIAS* Reif, 1978

Type species—*Pseudodalatias bartonensis* (Sykes, 1971)

Occurrence — Upper Triassic, Rhaetian of Barnstone, England (Skyes, 1971)

PSEUDODALATIAS HENAREJENSIS Botella, Plasencia, Márquez-Aliaga, Cuny and Dorka, 2009

Figure 2 Q-R

Material—One tooth from the Tivissa section (CT-8 level: MGUV-36113).

Description—The tooth is almost complete, but poorly preserved with the base missing (Fig. 2Q-R). It shows the characteristic features of *Pseudodalatias henarejensis*: a serrated-triangular morphology, with three (damaged) denticles on the edges and often lacking the base (Botella et al., 2009). The cusp is also damaged and the whole tooth surface is eroded.

Remarks—*Pseudodalatias henarejensis* is an endemic species from the Iberian Peninsula, which shows a dignathic heterodonty, as described in Pla et al. (2013). The tooth found in Tivissa belongs to the lower jaw.

Occurrence —Middle Triassic from the Iberian Chain of Spain (Botella et al., 2009; Pla et al., 2013); Middle–Late Triassic of the Betic Ranges (Manzanares et al., 2018); Middle Triassic of the Catalan Coastal Ranges, Spain.

REFERENCES

- Agassiz, L. 1833–44. Recherches sur les poissons fossiles, 3. Imprimerie de Petitpierre, Neuchâtel:32–390.
- Bonaparte, C. L. 1838. Synopsis Vertebratorum Systematis. Nuovi Annali delle Scienze Naturali, Bologna:105–133.
- Botella, H., Plasencia, P., Márquez-Aliaga, A., Cuny, G., and Dorka, M. 2009. *Pseudodalatias henarejensis* nov. sp. A new pseudodalatiid (Elasmobranchii) from the Middle Triassic of Spain. J. Vertebr. Paleontol., 29:1–20.
<https://doi.org/10.1671/039.029.0425>

- Cappetta, H. 2012. Handbook of Paleoichthyology, Vol. 3E: Chondrichthyes – Mesozoic and Cenozoic Elasmobranchii: teeth. Verlag Dr. Friedrich Pfeil, 512 p.
- Chen, L., Cuny, G., and Wang, X. 2007. The chondrichthyan fauna from the Middle–Late Triassic of Guanling (Guizhou province, SW China). *Hist. Biol.*, 19:291–300.
<https://doi.org/10.1080/08912960701248234>
- Cuny, G., Rieppel, O., and Sander, P. M. 2001. The shark fauna from the Middle Triassic (Anisian) of North–Western Nevada. *Zool. J. Linn. Soc-Lond.*, 13:285–30.
<https://doi.org/10.1006/zjla.2000.0273>
- Delsate, D. 1992. Chondrichthyens mésozoïques du Luxembourg. Note préliminaire. *Bull. Soc. Nat. Luxemb.*, 93 :181–193.
- Delsate, D. 1993. Synthèse des faunes d'Elasmobranches du Trias et du Jurassique de Lorraine. *Cossmanniana*, Hors-série, 2:52–55.
- Duffin, C. J. 1993. Mesozoic Chondrichthyan faunas 1. Middle Norian (Upper Triassic) of Luxembourg. *Palaeontogr. Abt. A*, 229:15–36.
- Ginter M., Hampe, O., and Duffin, C. J. 2010. Chondrichthyes. Paleozoic Elasmobranchii: Teeth. In: *Handbook of Paleoichthyology* 3D. H.-P. Schultze (ed.), 168 p.
- Hay, O. P. 1902. Bibliography and Catalogue of the fossil Vertebrata of North America. *B U.S Geol. Surv.*, 179:1–168.
- Huxley, T. H. 1880. On the application of the laws of evolution to the arrangement of the Vertebrata and more particularly of the Mammalia. *Proc. Zool. Soc. Lond.*, 1880:649–662.
- Johns, M. J., Barnes, C. R., and Orchard M. J. 1997. Taxonomy and Biostratigraphy of Middle and Late Triassic elasmobranch ichthyoliths from northeastern British Columbia. *Geol. Surv. Canada*, 502:1–235.

- Koot, M. B., Cuny, G., Tintori, A., and Twitchett, R. J. 2013. A new diverse shark fauna from the Wordian (Middle Permian) Khuff Formation in the interior haushi-huqf area, Sultanate of Oman. *Palaeontology*, 56(2):303–343.
<https://doi.org/10.1111/j.1475-4983.2012.01199.x>
- Koot, M. B., Cuny, G., Orchard, M. J., Richoz, S., Hart, M. B., and Twitchett, R. J. 2015. New hybodontiform and neoselachian sharks from the Lower Triassic of Oman. *J. Syst. Palaeontol.*, 13(10):891-917. <https://doi.org/10.1080/14772019.2014.963179>
- Leuzinger, L., Cuny, G., Popov, E., and Billon-Bruyat, J.-P. 2017. A new chondrichthyan fauna from the Late Jurassic of Swiss Jura (Kimmeridgian) dominated by hybodonts, chimaeroids and guitarfishes. *Pap. Palaentol.*, 3(4):471-511.
<https://doi.org/10.1002/spp2.1085>
- Maisey, J. G. 1987. Cranial anatomy of the Lower Jurassic shark *Hybodus reticulatus* (Chondrichthyes: Elasmobranchii), with comments on hybodontid systematics. *Am. Mus. Novit.*, 2878:1–39.
- Manzanares, E., Pla, C., Ferrón, H., and Botella, H. 2018. Middle-Late Triassic chondrichthyan remains from the Betic Range (Spain). *J. Iber. Geol.*, 44(1):129-138. <https://doi.org/10.1007/s41513-017-0027-1>
- Manzanares, E., Pla, C., Martínez-Pérez, C., Ferrón, H., and Botella, H. 2017. *Lonchidion derenzii* sp. nov., a new lonchidiid shark (Chondrichthyes, Hybodontiforms) from the Upper Triassic of Spain, with remarks on lonchidiid enameloid. *J. Verteb. Paleontol.*, 37(1):e1253585. <https://doi.org/10.1080/02724634.2017.1253585>
- Mutter, R., De Blanger, K., and Neuman, A. G. 2008. *Homalodontus* nom. nov., a replacement name for *Watipiodus* Mutter, de Blanger and Neuman, 2007

- (Homalodontidae nom. nov.,? Hybodontoides), preoccupied by *Watipiodus* Orchard, 2005. Zool. J. Linn. Soc-Lond. 154:419–420.
- Owen, R. 1846. Lectures on the comparative anatomy and physiology of the vertebrate animals, delivered at the Royal College of Surgeons of England in 1844 and 1846. Part 1. *Fishes*. Longman, Brown, Green and Longmans, London: xi + 304 p.
- Pla, C., Márquez-Aliaga, A., and Botella, H. 2013. The chondrichthyan fauna from the Middle Triassic (Ladinian) of the Iberian Range (Spain). J. Verteb. Palaeontol., 33(4): 770-785. <https://doi.org/10.1080/02724634.2013.748668>
- Reif, W. E. 1978. Types of morphogenesis of the dermal skeleton in fossil sharks. Palaeont. Z., 52:110–128.
- Rees, J. 1999. Late Cretaceous hybodont sharks from the Hasle Kristianstad Basin, southern Sweden. Neu. Jb. Geol. und Palaöntol, Mh., 1999(2):257–270.
- Rees, J., and Underwood, C. J. 2002. The status of the shark genus *Lissodus* Brough, 1935, and the position of nominal species of *Lissodus* within the Hybodontoides (Selachii). J. Verteb. Paleontol., 22:471–479. [https://doi.org/10.1671/0272-4634\(2002\)022\[0471:TSOTSG\]2.0.CO;2](https://doi.org/10.1671/0272-4634(2002)022[0471:TSOTSG]2.0.CO;2)
- Rees, J., and Underwood, C. J. 2008. Hybodont sharks of the English Bathonian and Callovian (Middle Jurassic). Palaeontology, 51(1):117-147. <https://doi.org/10.1111/j.1475-4983.2007.00737.x>
- Rieppel, O. 1981. The Hybodont sharks from the Middle Triassic of Monte Giorgio, Switzerland. Neues Jahrb. Geol. P-A., 161:324–353.
- Rieppel, O., Kindlimann, R., and Bucher, H. 1996. A new fossil fish fauna from the Middle Triassic (Anisian) of North-Western Nevada. In *Mesozoic Fishes, Systematics and*

- Paleoecology*. Proceedings of the 1st International Meeting Eichstätt 1993, G. Arratia and G. Viohl (eds.), 501–512.
- Sykes, J. H. 1971. A new dalatiid fish from the Rhaetic Bond Bed at Barnstone, Nottinghamshire. *Mercian Geologist*, 4:13–22.
- Underwood, C., and Rees, J. 2002. Selachian faunas from the earliest Cretaceous Puberck Group of Dorset, southern England. *Spec. Pap. Palaeontol.*, 68:83– 01.
- von Meyer, H. 1849. Fossile Fische aus dem Muschelkalk von Jena, Querfurt und Esperstädt. *Palaeontographica*, 1:195–208.
- Woodward, A. S. 1916. The Fossil Fishes of the English Wealden and Purbeck Formations. Part I. Pages 1–48, Plates I–X. *Monographs of the Palaeontographical Society*, 69(334) :1-48.