Xie, W., Yang, F., Meng, S., Scarpa, F., & Wang, L. (2020). Perforation of needle-punched carbon-carbon composites during high-temperature and high-velocity impacts. *Composite Structures*, 245, [112224]. <https://doi.org/10.1016/j.compstruct.2020.112224>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.compstruct.2020.112224](https://doi.org/10.1016/j.compstruct.2020.112224)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.compstruct.2020.112224> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

glucosamines.^{10a} Upon successful ligation to $[\text{Rh}(\text{cod})\text{Cl}]_2$, the carbohydrate-based NHC complexes (e.g. **1**, Scheme 1A, eqn. (1)) were evaluated in the asymmetric hydrosilylation of ketones (e.g. **2**, Scheme 1A, eqn. (2)).^{11,2} Ultimately, catalyst **1**, featuring permethylated β -glucoside residues attached to the NHC moiety, was most efficient and provided phenyl-1-ethanol (**3**) in modest er (81:19 *R:S*).¹⁰

One of the advantages of carbohydrate chiral synthons is the ability to chemoselectively incorporate functional groups at each of the different OH groups. We thus sought to achieve greater asymmetric induction from **1** by sterically differentiating the C1 and C3 positions which flank the connecting position (C2) between the carbohydrate and the imidazolylidene scaffold (**4a-b**, Scheme 1B). Practically, this can be realised in two different ways: (1) by employing a carbohydrate unit with a large group (R_L) at C1 and a small group (R_S) at C3, as depicted in **4a**, or (2) vice versa as in **4b**. We surmised that an aryl group would serve as an effective R_L candidate as this motif is readily available. From a synthetic viewpoint, installation of an aryl residue (e.g. phenol or thiophenol) at C1 by stereoselective glycosylation is well known.¹² Accessing a related C3-OH arylated carbohydrate derivative is conceptually more challenging, however, this could be achieved using $S_N\text{Ar}$ -based methodology recently developed in our laboratory.¹³ Therefore, our aims were to synthesize sterically differentiated carbohydrate amines, access the corresponding C2-symmetric imidazolium salts, ligate to a Rh source and subsequently evaluate the complexes in the hydrosilylation of acetophenone (**2**).

Scheme 2. Synthesis of a carbohydrate-based imidazolium salt with steric bulk at C1

Thioglycoside **5**¹⁴, synthesized in two steps from D-glucosamine-HCl in 70% yield (see ESI for details), was chosen as it was a commonly used building block in carbohydrate synthesis and thus easily accessible, was globally deprotected

by stirring with NaOMe in methanol to remove the ester groups. This was followed by addition of aq. NaOH at reflux to give amine intermediate **6**¹⁵ (Scheme 2A). Cu-catalyzed azide formation from **6** with TfN_3 gave triol **7** in 78% yield over the three steps. Methylation of the remaining hydroxyl positions using MeI and NaH and subsequent azide hydrogenation (H_2 , Pd/C) gave C1 sterically differentiated amine **8a** in an 81% yield. It is pertinent to note that a methyl ether (OMe) was selected as the R_S group based on our previous observations.^{10a} Elaboration of **8a** into imidazolium salt **9a** was achieved in 41%

yield by condensation with aq. glyoxal and then cyclization with MOMCl (Scheme 2B).

Scheme 3. Synthesis of a carbohydrate-based imidazolium salt with steric bulk at C3

With a C1 sterically differentiated carbohydrate-based NHC-HCl salt in hand, our attention turned to the C3 analogue (Scheme 3A-B).¹⁶ The synthesis of carbohydrate-aryl ether **10** was accomplished by an $S_N\text{Ar}$ reaction between the appropriate fluoro-arene and the corresponding OH-bearing carbohydrate.¹³ Next, under sonication, TsOH acid catalyzed deprotection of the benzylidene acetal revealed the O4 and O6 hydroxyls, which were methylated as before to give **11** in 95% overall yield. Heterogeneous hydrogenation gave amine **8b** and the corresponding imidazolium salt (**9b**) was accessed in 44% yield utilizing the same procedure outlined before.

Scheme 4. Ligation of **9a-b** to [Rh(cod)Cl]₂

Imidazolium salt **9a** was deprotonated by ^tBuONa at rt and ligated to [Rh(cod)Cl]₂ which afforded complex **12a** in usable yield (Scheme 4, (1)). NHC-HCl **9b** required the action of ^tBuOK at 100 °C to effect ligation and this allowed isolation of **12b** in 72% yield (Scheme 4, (2)). Unfortunately, neither complex was crystalline but HRMS confirmed product formation (see the Experimental Section) and the carbenic ¹³C NMR resonance was visible for **12b** (δ_c : 189.2 ppm). The ¹H NMR data of **12a-b** is similar to other β -glucoside-based NHC-Rh compounds and the spectra also featured significant peak broadening, which further supports metal conjugation.^{10a} In **12b** the backbone imidazolylidene protons were not obscured and appeared at δ_H : 6.95 and 6.80 ppm respectively, indicating that the complex does not possess C₂-symmetry and, therefore, it is likely the pyranoside rings are each in unique environments. From this information we infer that our NHC ligands are dynamic in solution. Indeed, Grubbs and co-workers have observed rotation around C_{Carbohydrate}-N_{Imidazolylidene} bonds in metathesis catalysts featuring carbohydrate-based NHC ligands.^{10c}

Table 1. Evaluation of **12a-b** in the hydrosilylation of acetophenone

Entry	12	Yield (%) ^a	Er (R:S) ^b
1	a	75	30:70
2	b	95	75:25

^a Isolated yield. ^b Determined by chiral HPLC.

Next, complexes **12a-b** were evaluated in the hydrosilylation of acetophenone as a model system and both were found to be catalytically competent (Table 1). Interestingly, although the enantiomer ratios (er) of the product were modest, **12a** resulted in the isolation of **3** with a distinguishable preference for the *S*-enantiomer (30:70 *R*:*S*, entry 1), whereas **12b** gave preference for the *R*-enantiomer (75:25 *R*:*S*, entry 2). Carbohydrates largely exist in nature in the *D*-configuration and this causes complications when employing them as ligands because sufficient quantities of the antipode are not normally available. Therefore, an asymmetric process must often rely on pseudo-enantiomeric ligands to give access to each enantiomer of the product.¹⁷ While pseudo-enantiomerism has been documented in carbohydrate-based ligands, such as phosphinites¹⁸ and phosphites,¹⁹ and carbohydrate-based organocatalysts,²⁰ to the best of our knowledge this is the first example concerning a carbohydrate-based NHC ligand.

Encouraged by the results, we envisaged optimizing the asymmetric induction potential of the carbohydrate-based NHC ligands by altering the size and substitution pattern of the C3 aryl group.²¹ Commercially available fluoro-aryls bearing a range of structural motifs (*ortho*-, *meta*- and *para*- substitution

patterns; biaryls and bicyclics) were introduced using S_NAr, expediently delivering a small library of carbohydrate-aryl ethers (**13a-e**) as previously described.¹³ The ethers were then transformed into the required amines (**15a-e**) by benzylidene acetal cleavage, methylation of the OHs (**14a-e**) and reduction of the azide functionality as before (Scheme 5).

Scheme 5. Synthesis of S_NAr modified amines **15a-e** **Scheme 6.** Synthesis of complexes **17a-e**

The S_NAr modified amines were swiftly elaborated into the corresponding imidazolium salts (**16a-e**) by treatment with aq. glyoxal and then MOMCl (Scheme 6). Deprotonation with ^tBuOK and ligation to Rh was performed in THF at reflux to give complexes **17a-e** in 40-99% yields. While the yields for both of these steps were predominantly good, the example featuring Ar₁ (bearing *ortho-ortho* aryl substitution, e.g. **13a-17a**) was consistently low yielding despite our best efforts and this was attributed to the level of steric bulk present in the system.

Table 2. Evaluation of **17a-e** in the hydrosilylation of acetophenone.

Entry	17	Yield (%) ^a	Er (R:S) ^b
1	a	72	51:49
2	b	15	50:50
3	c	89	86:14
4	d	70	61:39
5	e	73	78:22

^a Isolated yield. ^b Determined by chiral HPLC.

Complexes **17a-e** were then evaluated in the hydrosilylation of **2** and the results are displayed in Table 2. Catalysts **17a-b**, featuring an *ortho*-CF₃ substituted aryl group (Ar₁₋₂), gave negligible stereoselection in **3** (entries 1-2). However, **17c** (entry 3) resulted in the isolation of **3** in high yield and with the greatest *er* to date (86:14). Both **17d** and **17e** (entries 4-5) afforded **3** with detectable *er*'s but were ultimately less efficient systems.

Conclusions

In summary, a series of pseudo-enantiomeric carbohydrate-based NHC-ligands featuring sterically differentiated groups at positions C1 or C3 of the carbohydrate were synthesized by simple glycosylation or S_NAr strategies and evaluated in the Rh-catalyzed hydrosilylation of acetophenone. We show that ligands bearing a bulky group at C1 lead to a preference for (*S*)-phenyl-1-ethanol, while the presence of large groups at C3 afforded preferentially the (*R*)-enantiomer. Our results suggest this effect is achieved by the steric differentiation of the carbohydrate's C1 and C3 positions which flank the attachment to the imidazolylidene scaffold. This "pseudoenantiomeric" design feature is shown in Figure 1.²²

Figure 1. Pseudo-stereochemical ligands **12a** and **17c**.

While our efforts to achieve a high level of asymmetric induction in the benchmark hydrosilylation reaction¹¹⁻¹² have been less successful, we have surpassed our previous result⁹ and shown, for the first time to the best of our knowledge, that the judicious functionalization of carbohydrate synthons can lead to pseudo-enantiomeric NHC-ligands for asymmetric catalysis.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

ASH thanks EPSRC Bristol Chemical Synthesis CDT EP/G036764/1, JFB thanks Royal Society for a University Research Fellowship and MCG thanks the ERC (COG: 648239) and EPSRC CAF EP/L001926/1 and EPSRC EP/J002542/1 for funding.

Notes and references

- (a) W. A. Herrmann, C. Köcher, *Angew. Chem., Int. Ed. Engl.* 1997, **36**, 2162. (b) *N-Heterocyclic Carbenes*; S. P. Nolan, Ed.; Wiley-VCH: Weinheim, 2014.
- A. Doddi, M. Peters, M. Tamm, *Chem. Rev.* 2019, **119**, 6994.3. (a) S. Kumar, *J. Heterocyclic Chem.* 2019, **56**, 1168. (b) S. Díez-González, N. Marion, S. P. Nolan, *Chem. Rev.* 2009, **109**, 3612.
- (a) V. César, S. Bellemin-Lapponnaz, L. H. Gade, *Chem. Soc. Rev.* 2004, **33**, 619. (b) F. Wang, L.J. Liu, W. Wang, S. Li, M. Shi, *Coord. Chem. Rev.* 2012, **256**, 804. (c) D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.* 2007, **107**, 5606.
- Selected examples: (a) D. Enders, H. Gielen, G. Raabe, J. Runsink, J. H. Teles, *Chem. Ber.* 1996, **129**, 1483. (b) W. A. Herrmann, L. J. Goossen, C. Köcher, G. R. J. Artus, *Angew. Chem., Int. Ed.* 1996, **35**, 2805. (c) ST. J. eiders, D. W. Ward, R. H. Grubbs, *Org. Lett.* 2001, **3**, 3225.
- Selected examples: (a) E. P. Kündig, T. M. Seidel, Y. X. Jia, G. Bernardinelli, *Angew. Chem. Int. Ed.* 2007, **46**, 8484. (b) C. T. Check, K. P. Jang, C. B. Schwamb, A. S. Wong, M. H. Wang, K. A. Scheidt, *Angew. Chem. Int. Ed.* 2015, **54**, 4264.
- Selected examples: (a) F. Glorius, G. Altenhoff, R. Goddard, C. Lehmann, *Chem. Commun.* 2002, 2704. (b) S. Würzt, C. Lohre, R. Fröhlich, K. Bergander, F. Glorius, *J. Am. Chem. Soc.* 2009, **131**, 8344. (c) P. J. Czerwinski, M. Michalak, *Synthesis-Stuttgart* 2019, **51**, 1689.
- L. Benhamou, E. Chardon, G. Lavigne, S. Bellemin-Lapponnaz, V. César, *Chem. Rev.* 2011, **111**, 2705.
- Selected reviews: (a) M. Diéguez, O. Pàmies, C. Claver, *Chem. Rev.* 2004, **104**, 3189. (b) Carbohydrates: Tools for Stereoselective Synthesis; M. M. K. Boysen, Ed.; Wiley-VCH: Weinheim, 2013. (c) A. S. Henderson, J. F. Bower, M. C. Galan, *Org. Biomol. Chem.* 2016, **14**, 4008. (d) R. Pretorius, J. Olguin, M. Albrecht, *Inorg. Chem.* 2017, **56**, 12410. (e) M. E. Cucciolito, A. D'Amora, G. De Feo, G. Ferraro, A. Giorgio, G. Petruk, D. M. Monti, A. Merlino, F. Ruffo, *Inorg. Chem.* 2018, **57**, 3133. (f) J. C. Shi, N. Lei, Q. S. Tong, Y. R. Peng, J. F. Wei, L. Jia, *Eur. J. Inorg. Chem.* 2007, 2221. (g) J. P. Byrne, P. Musembi, M. Albrecht, *Dalton Trans.* 2019, **48**, 11838. (h) B. K. Keitz, R. H. Grubbs, *Organometallics* 2010, **29**, 403. (i) W. H. Zhao, V. Ferro, M. V. Baker, *Coordinat. Chem. Rev.* 2017, **339**, 1. (j) Zhou, Z. G.; Yuan, Y. Y.; Xu, G. H.; Chen, Z. W.; Li, M. *Prog. Chem.* 2019, **31**, 351; (k) T. Shibata, H. Hashimoto, I. Kinoshita, S. Yano, T. Nishioka, *Dalton Trans.* 2011, **40**, 4826.
- (a) A. S. Henderson, J. F. Bower, M. C. Galan, *Org. Biomol. Chem.* 2014, **12**, 9180. (b) F. Tewes, A. Schlecker, K. Harms, F. Glorius, *J. Organomet. Chem.* 2007, **692**, 4593. (c) B. K. Keitz, R. H. Grubbs, *Organometallics* 2009, **29**, 403. (d) M. Guitet, P. Zhang, F. Marcelo, C. Tugny, J. Jiménez-Barbero, O. Buriez, C. Amatore, V. Mouries-Mansuy, J.-P. Goddard, L. Fensterbank, Y. Zhang, S. Roland, M. Ménand, M. Sollogoub, *Angew. Chem. Int. Ed.* 2013, **52**, 7213.
- For selected examples on asymmetric Rh-catalysed hydrosilylation of ketones see: (a) K. Riener, M. P. Högerl, P. Gigler, F. E. Kühn, *ACS Catalysis* 2012, **2**, 613. (b) W.-L. Duan, M. Shi, G.-B. Rong, *Chem. Commun.* 2003, 2916. (c) L. H. Gade, V. César, S. Bellemin-Lapponnaz, *Angew. Chem. Int. Ed.* 2004, **43**, 1014. (d) N. Schneider, M. Finger, C. Haferkemper, S. Bellemin-Lapponnaz, P. Hofmann, L. H. Gade, *Angew. Chem. Int. Ed.* 2009, **48**, 1609. (e) J. Lee, H. Hahm, J. Kwak, M. Kim, *Adv. Synth. Catal.* 2019, **361**, 1479.
- Selected reviews: (a) P. J. Garegg, In *Adv. Carbohydr. Chem. Biochem.*; H. Derek, Ed.; Academic Press: 1997; Vol. 52, 179. (b) M. Jacobsson, J.; Malmberg, U. Ellervik, *Carbohydr. Res.* 2006, **341**, 1266.
- A. S. Henderson, S. Medina, J. F. Bower, M. C. Galan, *Org. Lett.* 2015, **17**, 4846.
- T. Nishiyama, Y. Kusumoto, K. Okumura, K. Hara, S. Kusaba, K. Hirata, Y. Kamiya, M. Isobe, K. Nakano, H. Kotsuki, Y. Ichikawa, *Chemistry* 2010, **16**, 600.
- T. Buskas, P. J. Garegg, P. Konradsson, J. L. Maloisel, *Tetrahedron-Asymm.* 1994, **5**, 2187.

16. The use of an identical aryl group for the C1 and C3 sterically differentiated NHC ligands would have been ideal, however, for proof of principle aryls compatible with the respective glycosylation and SNAr methodologies were initially used.
17. For a review see: V. Benessere, M. Lega, F. Ruffo, *J. Organomet. Chem.* 2014, **771**, 105.
18. T. V. RajanBabu, T. A. Ayers, G. A. Halliday, K. K. You, Calabrese, J. C. *J. Org. Chem.* 1997, **62**, 6012.
19. M. T. Reetz, T. Neugebauer, *Angew. Chem. Int. Ed.* 1999, **38**, 179.
20. L. Fang, L. Yan, F. Haeffner, J. P. Morken, *J. Am. Chem. Soc.* 2016, **138**, 2508.
21. It is pertinent to note that given the pseudo-enantiomeric relationship between the C1 and C3 positions, a promising aryl moiety at C3 could also be mirrored at C1 by glycosylation with the appropriate phenol or thiol.
22. Sulfur-containing molecules can act as ligands outright, however, we have no evidence for a Rh-S interaction in complex **12a**. For a review on sulfur-containing ligands in asymmetric catalysis see: H. Pellissier, *Tetrahedron* 2007, **63**, 1297