Carnevale Neto, F., Guaratini, T., Colepicolo, P., Gates, P. J., & Lopes, N. P. (2020). Characteristic product ions of acetylene carotenoids by electrospray and nanospray ionization tandem mass spectrometry. *Rapid Communications in Mass Spectrometry*, 34(16), [e8811]. <https://doi.org/10.1002/rcm.8811>

Peer reviewed version

Link to published version (if available):
[10.1002/rcm.8811](https://doi.org/10.1002/rcm.8811)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/epdf/10.1002/rcm.8811> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Characteristic product ions of acetylene carotenoids by electrospray and nanospray ionization tandem mass spectrometry

Fausto Carnevale Neto^{1,2}, Thais Guaratini^{1,3}, Pio Colepicolo⁴, Paul J. Gates⁵, Norberto Peporine Lopes^{1†}

¹ Núcleo de Pesquisa em Produtos Naturais e Sintéticos (NPPNS), Departamento de Física e Química, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, 14040-903, São Paulo, Brazil.

² Northwest Metabolomics Research Center, Department of Anesthesiology and Pain Medicine, University of Washington, 850 Republican St., Seattle, WA 98109, United States.

³ Lychnoflora Pesquisa e Desenvolvimento em Produtos Naturais LTDA, Incubadora SUPERA, Campus da Universidade de São Paulo, Ribeirão Preto, 14040-903, São Paulo, Brazil.

⁴ Departamento de Bioquímica, Instituto de Química de São Paulo, Universidade de São Paulo, São Paulo, 05508-000, São Paulo, Brazil.

⁵ School of Chemistry, University of Bristol, Cantock's Close, BS8 1TS, Bristol, UK.

† Corresponding Author:

Norberto P. Lopes, Universidade de São Paulo
Faculdade de Ciências Farmacêuticas de Ribeirão Preto,
Av. do Café S/N, CEP 14040-903, Ribeirão Preto-SP, Brazil.
E-mail: npelopes@fcfrp.usp.br
Contract/grant sponsor: FAPESP.
Contract/grant sponsor: CAPES.
Contract/grant sponsor: CNPq.

Keywords: high-resolution tandem mass spectrometry, electrospray ionization, acetylenic carotenoids, alloxanthin, diatoxanthin, diadinoxanthin.

Carotenoids constitute a large group of natural pigments produced by photosynthetic organisms (plants and algae) and non-photosynthetic bacteria and fungi.¹ They are responsible for several biochemical events that regulate the adaptation and survival of different organisms, including light-harvesting, protection against UV radiation, and quenching of reactive oxygen species.²⁻⁵

In animals, they play important roles in visual communication and sex behavior⁶, modulation of the immune system⁷, growth factors and intracellular signaling pathways, regulation of cell differentiation⁸, and as precursors for the pigment vitamin A.⁹

Carotenoids normally contain a C40 linear and symmetrical skeleton with an extensive polyene chain, formed by head-to-tail condensation of eight isoprene units.¹⁰ These pigments possess unique chemical reactivity and light-absorbing properties due to the long central conjugated polyene chain, a chromophore system with λ_{max} . 400–500 nm. Modifications in the basic structure, such as cyclization, hydrogenation, dehydrogenation, hydroxylation, epoxidation, rearrangement, and addition of lateral groups, give rise to a complex variety of similar compounds with widespread isomerism.¹¹

About 750 carotenoids have been isolated from natural sources. More than 250 are of marine origin and show an interesting structural diversity, such as allenic (e.g., fucoxanthin) and all the acetylenic carotenoids (alloxanthin).¹² Albeit most carotenogenesis have been proposed, still little is known about the final biosynthetic steps leading to the formation of both allenic and acetylenic bonds.^{13,14}

The economic relevance of carotenoids has driven the development of reliable analytical methodologies to analyze these pigments in food and other biological matrices.¹⁵ Mass spectrometry (MS) is currently the preferred technique for determining the identity of carotenoids in complex samples. It allows one to distinguish between co-eluting compounds, when coupled with chromatographic systems, and it provides structural information on the basis of the molecular masses, using high-resolution instruments (HRMS), and key fragment ions through application of tandem mass spectrometry (MS/MS or MS2).¹⁶⁻¹⁹

Various approaches have explored carotenoid behavior through different ionization techniques, including electron impact (EI), fast atom bombardment (FAB), chemical ionization (CI), matrix-assisted laser desorption/ionization (MALDI), electrospray (ESI) and atmospheric pressure chemical ionization (APCI).²⁰⁻²³ EI has been applied to evaluate the fragmentation pathways of carotenoids through radical ion formation and redox

reactions.^{24–27} Most recently, APCI combined with collision-induced dissociation (CID) experiments have attempted to identify characteristic fragments that can be associated with specific functional groups in the structures of these compounds.^{20,22}

Previously, we reported the fragmentation of protonated carotenoids using ESI- and nanospray-CID-MS/MS.²⁸ The results were compared with the characteristic carotenoid ions proposed by APCI-MS/MS.²⁰ It was demonstrated that product ions of protonated carotenoids derived from charge-remote electrocyclic and retro-ene eliminations. These mechanisms rely on the correct conformation and distribution of hydrogens, achieved during protonation of the polyene chain, which form a carbocation intermediate and induce cis-trans isomerization.²⁸ In addition to the characteristic elimination of the aromatic ring from the polyene chain by $8\pi/6\pi$ electron electrocyclization,²⁹ we also provided diagnostic product-ions for keto-carotenoids, generated by vinyl-allyl cleavage,³⁰ and epoxy-carotenes, with formation of oxonium ions.^{24,28}

Herein we investigate the gas-phase reactions of acetylenic carotenoids by ESI- and nanospray-CID-MS/MS, and provided key product ions for their rapid detection.

The MS/MS analysis of the carotenoid pigments alloxanthin, diadinoxanthin and diatoxanthin (Figure S-1) by two tandem mass spectral techniques* exhibited the product ions m/z 217 and 199. Accurate mass measurements by ultra-high-resolution ESI-FTICR MS/MS analysis (see Figures 1A-C) allow unequivocal determination of the molecular formulae of the two product ions, as shown in Table 1. These product ions are only observed in carotenoids that contain acetylenic bonds in the polyene chain. Similar fragments could also be detected on MS/MS spectra deposited in the public MASSBANK database (see Supporting Information).³¹

The fragmentation of acetylenic carotenoids initiated with selection of the protonated molecule $[M+H]^+$ as the precursor ion. Protonation is favored when using protic solvents mixed with 0.5% formic acid prior the ionization. According to previous studies, protonation of carotenoids preferably occurs between C7-C11, even in the presence of oxygen atoms in the xanthophyll.^{28,32,33} During the ionization and subsequent CID fragmentation, the internal energy of the analytes increases sufficiently to induce isomerization of the polyene chain.

We observed dehydration of hydroxyl substituents, similar to those previously detected for xanthophylls,²⁰ and sequential cleavages of the polyene chain, which includes the diagnostic loss of toluene and/or xylene.²⁹ The formation of the key product ion m/z 217 arises by the transfer of the γ -hydrogen atom to an unsaturated center via a

six-electron cyclic transition state, resulting in the formation of a conjugated diene and an allyl ion (Figure 1D). Several cleavages of the polyene chain by sequential pericyclic charge-remote reactions are characteristic of carotenoids. Due to the sp orbitals at C7-C8, this retro-ene reaction is chosen to attain the proper structural conformation and the hydrogen requirements. The product ion m/z 199 occurs through a further dehydration of m/z 217, possibly via a 1,2 elimination.

As a carotenoid epoxide, diadinoxanthin also showed the fragment m/z 221, formed by epoxide-furanoxide rearrangement with formation of cyclic oxonium ion, as summarized in Figure 2.²⁸

The ultra-high-resolution accurate-mass analysis supports the evidence of $C_{15}H_{21}O^+$ as m/z 217 and $C_{15}H_{19}^+$ as m/z 199, and corresponds to key fragments for the rapid detection of acetylenic carotenoids. These results expand the set of diagnostic product ions for carotenoids analyzed by ESI-MS/MS, and provide structural information to assist their characterization via collision induced dissociation mass spectrometry.

* Ultra-high resolution accurate-mass ESI-MS/MS analyses were performed on a 7 Tesla Apex IV Fourier-transform ion cyclotron resonance instrument (Bruker Daltonics, Billerica, MA, USA) using an Apollo off-axis ESI source. Samples were directly infused from a syringe pump at 100 $\mu\text{L min}^{-1}$. The acquisition parameters were: capillary voltage, 4600 V; end plate voltage, 3500 V; capillary exit potential, 200 V and drying gas temperature, 200°C. Spectra were obtained by summing 80 0.5 s scans. CID-MS/MS fragmentation analyses were performed on the isolated parent ions using N_2 as the collision gas. Nanospray ionization analyses were performed on a QStar-XL quadrupole time-of-flight hybrid instrument (Applied Biosystems, Warrington, UK) using a Nanomate 100 automatic chip based nanospray system (Advion Biosciences, Norwich, UK). The QStar acquisition parameters were: ion source gas flow rate, 50 arbitrary units; curtain gas flow rate, 20 arbitrary units; ionspray voltage, 2700 V; declustering potential, 75 V; focusing potential, 280 V (see ESI-MS/MS spectra on Supporting Information).

Acknowledgment

This work was funded by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for fellowship support (2014/12343-2 to FCN, and 2014/50265-3 to

NPL), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)

References

1. Alcaíno J, Baeza M, Cifuentes V. Carotenoid distribution in nature. In: Stange C, ed. *Carotenoids in Nature. Biosynthesis, Regulation and Function*. Cham: Springer International Publishing; 2016:3-33. doi:10.1007/978-3-319-39126-7_1
2. Mathews MM, Sistrom WR. Function of carotenoid pigments in non-photosynthetic bacteria. *Nature*. 1959;184(4702):1892-1893. doi:10.1038/1841892a0
3. Leverenz RL, Sutter M, Wilson A, et al. A 12 Å carotenoid translocation in a photoswitch associated with cyanobacterial photoprotection. *Science*. 2015;348(6242):1463 LP - 1466. doi:10.1126/science.aaa7234
4. Palozza P, Parrone N, Simone R, Catalano A. Role of lycopene in the control of ROS-mediated cell growth: implications in cancer prevention. *Curr Med Chem*. 2011;18(12):1846-1860. doi:10.2174/092986711795496845
5. Havaux M. Carotenoids as membrane stabilizers in chloroplasts. *Trends Plant Sci*. 1998;3(4):147-151. doi:10.1016/S1360-1385(98)01200-X
6. Svensson PA, Wong BBM. Carotenoid-based signals in behavioural ecology: a review. *Behaviour*. 2011;148(2):131-189. doi:10.1163/000579510X548673
7. Chew BP, Park JS. Carotenoid action on the immune response. *J Nutr*. 2004;134(1):257S-261S. doi:10.1093/jn/134.1.257S
8. Palozza P, Serini S, Di Nicuolo F, Calviello G. Modulation of apoptotic signalling by carotenoids in cancer cells. *Arch Biochem Biophys*. 2004;430(1):104-109. doi:10.1016/j.abb.2004.02.038
9. Dingle JT, Lucy JA. Vitamin A, carotenoids and cell function. *Biol Rev*. 1965;40(3):422-458. doi:10.1111/j.1469-185X.1965.tb00809.x
10. Moise AR, Al-Babili S, Wurtzel ET. Mechanistic aspects of carotenoid biosynthesis. *Chem Rev*. 2014;114(1):164-193. doi:10.1021/cr400106y
11. Amorim-Carrilho KT, Cepeda A, Fente C, Regal P. Review of methods for analysis of carotenoids. *TrAC Trends Anal Chem*. 2014;56:49-73. doi:https://doi.org/10.1016/j.trac.2013.12.011
12. Galasso C, Corinaldesi C, Sansone C. Carotenoids from marine organisms: biological functions and industrial applications. *Antioxidants*. 2017;6(4):96. doi:10.3390/antiox6040096
13. Maresca JA, Graham JE, Bryant DA. The biochemical basis for structural diversity in the carotenoids of chlorophototrophic bacteria. *Photosynth Res*. 2008;97(2):121-140. doi:10.1007/s11120-008-9312-3
14. Dambek M, Eilers U, Breitenbach J, Steiger S, Buchel C, Sandmann G. Biosynthesis of fucoxanthin and diadinoxanthin and function of initial pathway genes in *Phaeodactylum tricornutum*. *J Exp Bot*. 2012;63(15):5607-5612. doi:10.1093/jxb/ers211
15. Rivera SM, Canela-Garayoa R. Analytical tools for the analysis of carotenoids in diverse materials. *J Chromatogr A*. 2012;1224:1-10. doi:10.1016/j.chroma.2011.12.025
16. Carnevale Neto F, Andréo MA, Raftery D, et al. Characterization of aporphine alkaloids by electrospray ionization tandem mass spectrometry and density

- functional theory calculations. *Rapid Commun Mass Spectrom.* 2019;n/a(n/a):rcm.8533. doi:10.1002/rcm.8533
17. Pilon AC, Gu H, Raftery D, et al. Mass spectral similarity networking and gas-phase fragmentation reactions in the structural analysis of flavonoid glycoconjugates. *Anal Chem.* 2019;91(16):10413-10423. doi:10.1021/acs.analchem.8b05479
 18. Demarque DP, Crotti AEM, Vessecchi R, Lopes JLC, Lopes NP. Fragmentation reactions using electrospray ionization mass spectrometry: an important tool for the structural elucidation and characterization of synthetic and natural products. *Nat Prod Rep.* 2016;33:432-455. doi:10.1039/c5np00073d
 19. de Oliveira G, Carnevale Neto F, Demarque D, et al. Dereplication of flavonoid glycoconjugates from *Adenocalymma imperatoris-maximilianii* by untargeted tandem mass spectrometry-based molecular networking. *Planta Med.* 2016;83(07):636-646. doi:10.1055/s-0042-118712
 20. Rivera SM, Christou P, Canela-Garayoa R. Identification of carotenoids using mass spectrometry. *Mass Spectrom Rev.* 2014;33(5):353-372. doi:10.1002/mas.21390
 21. Crupi P, Milella RA, Antonacci D. Simultaneous HPLC-DAD-MS (ESI+) determination of structural and geometrical isomers of carotenoids in mature grapes. *J Mass Spectrom.* 2010;45(9):971-980. doi:10.1002/jms.1794
 22. van Breemen RB, Dong L, Pajkovic ND. Atmospheric pressure chemical ionization tandem mass spectrometry of carotenoids. *Int J Mass Spectrom.* 2012;312:163-172. doi:10.1016/j.ijms.2011.07.030
 23. Carnevale J, Cole ER, Nelson D, Shannon JS. Chemical ionization mass spectrometry of carotenoids. *Biol Mass Spectrom.* 1978;5(11):641-646. doi:10.1002/bms.1200051109
 24. Baldas J, Porter QN, Cholnoky L, Szabolcs J, Weedon BCL. Mass spectrometry of carotenoid epoxides and furanoid oxides. *Chem Commun.* 1966;(23):852. doi:10.1039/c19660000852
 25. Baldas J, Porter QN, Leftwick AP, Holzel R, Weedon BCL, Szabolcs J. Mass spectrometry of carotenoid ketones. *J Chem Soc D Chem Commun.* 1969;(8):415. doi:10.1039/c29690000415
 26. Enzell CR, Francis GW, Liaaen-Jensen S, et al. Mass spectrometric studies of carotenoids. 2. A survey of fragmentation reactions. *Acta Chem Scand.* 1969;23(8):727-750. doi:10.3891/acta.chem.scand.23-0727
 27. Uarrota VG, Moresco R, Coelho B, et al. Metabolomics combined with chemometric tools (PCA, HCA, PLS-DA and SVM) for screening cassava (*Manihot esculenta* Crantz) roots during postharvest physiological deterioration. *Food Chem.* 2014;161:67-78. doi:10.1016/j.foodchem.2014.03.110
 28. Carnevale Neto F, Guaratini T, Costa-Lotufo L, Colepicolo P, Gates PJ, Lopes NP. Re-investigation of the fragmentation of protonated carotenoids by electrospray ionization and nanospray tandem mass spectrometry. *Rapid Commun Mass Spectrom.* 2016;30(13):1540-1548. doi:10.1002/rcm.7589
 29. Guaratini T, Lopes NP, Pinto E, Colepicolo P, Gates PJ. Mechanism for the elimination of aromatic molecules from polyenes in tandem mass spectrometry. *Chem Commun.* 2006;(39):4110-4112. doi:10.1039/b609672g
 30. Hsu F-F, Turk J. Elucidation of the double-bond position of long-chain unsaturated fatty acids by multiple-stage linear ion-trap mass spectrometry with electrospray ionization. *J Am Soc Mass Spectrom.* 2008;19(11):1673-1680. doi:10.1021/jasms.8b03093

31. Horai H, Arita M, Kanaya S, et al. MassBank: a public repository for sharing mass spectral data for life sciences. *J Mass Spectrom.* 2010;45(7):703-714. doi:10.1002/jms.1777
32. Konovalov V, Kispert L. AM1, INDO/S and optical studies of carbocations of carotenoid molecules. Acid induced isomerization. *J Chem Soc Perkin Trans 2.* 1999;(4):901-910. doi:10.1039/A800551F
33. Kildahl-Andersen G, Lutnaes BF, Krane J, Liaaen-Jensen S. Structure elucidation of polyene systems with extensive charge delocalization carbocations from allylic carotenols. *Org Lett.* 2003;5(15):2675-2678. doi:10.1021/ol034987w

Table 1. Accurate masses and calculated molecular formulae of the main product ions observed in the FTICR MS/MS analysis of the $[M+H]^+$ acetylenic carotenoids **1-3** (see Fig. 1A-C).

compound	precursor ion	precursor m/z	product ion m/z	molecular formula
Alloxanthin (1)	$[M+H]^+$	565.4036 (-0.9) ^a	217.1582 (-2.3)	$C_{15}H_{21}O^+$
			199.1482 (0.0)	$C_{15}H_{19}^+$
Diatoxanthin (2)	$[M+H]^+$	567.4187 (-1.8)	217.1588 (0.5)	$C_{15}H_{21}O^+$
			199.1482 (0.0)	$C_{15}H_{19}^+$
Diadinoxanthin (3)	$[M+H]^+$	583.4145 (-0.2)	217.1592 (2.3)	$C_{15}H_{21}O^+$
			199.1488 (3.0)	$C_{15}H_{19}^+$

^a The mass measurement error (ppm) is given in parentheses.

Figure caption

Figure 1. Selected ESI-FTICR MS/MS spectrum of (A) alloxanthin (**1**), precursor ion $[M+H]^+$ m/z 565.4036, (B) diatoxanthin (**2**), precursor ion $[M+H]^+$ m/z 567.4187 and (C) diadinoxanthin (**3**), precursor ion $[M+H]^+$ m/z 583.4145. (D) Proposed fragmentation mechanism for the formation of product ions $C_{15}H_{21}O^+$ (m/z 217.1587) and $C_{15}H_{19}^+$ (m/z 199.1482).

Figure 2. Main product ions of acetylenic carotenoids by ESI-MS/MS. The bond colored in red represents the site of C-C cleavage.