Tarusikirwa, V., Mutamiswa, R., English, S., Chidawanyika, F., & Nyamukondiwa, C. (2020). Thermal plasticity in the invasive South American tomato pinworm *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Journal of Thermal Biology*, [102598].
<https://doi.org/10.1016/j.jtherbio.2020.102598>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.jtherbio.2020.102598](https://doi.org/10.1016/j.jtherbio.2020.102598)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/abs/pii/S0306456519306047> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Thermal plasticity in the invasive South American tomato pinworm *Tuta absoluta*
(Meyrick) (Lepidoptera: Gelechiidae)

Vimbai Tarusikirwa¹, Reyard Mutamiswa², Sinead English³, Frank Chidawanyika² and
Casper Nyamukondiwa^{1*}

¹Department of Biological Sciences and Biotechnology, Botswana International University of
Science and Technology, P. Bag 16, Palapye, Botswana.

²Department of Zoology and Entomology, University of the Free State, P.O. Box 339,
Bloemfontein 9300, South Africa

³School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, UK

*Author for correspondence: nyamukondiwa@biust.ac.bw

Abstract

South American tomato pinworm, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) is a devastating invasive global insect pest of tomato, *Solanum lycopersicum* (Solanaceae). In nature, pests face multiple overlapping environmental stressors, which may significantly influence survival. To cope with rapidly changing environments, insects often employ a suite of mechanisms at both acute and chronic time-scales, thereby improving fitness at sub-optimal thermal environments. For *T. absoluta*, physiological responses to transient thermal variability remain under explored. Moreso, environmental effects and physiological responses may differ across insect life stages and this can have implications for population dynamics. Against this background, we investigated short and long term plastic responses to temperature of *T. absoluta* larvae (4th instar) and adults (24–48 h old) field populations. We measured traits of temperature tolerance *vis* critical thermal limits [critical thermal minima (CT_{min}) and maxima (CT_{max})], heat knockdown time (HKDT), chill coma recovery time (CCRT) and supercooling points (SCP). Our results showed that at the larval stage, Rapid Cold Hardening (RCH) significantly improved CT_{min} and HKDT but impaired SCP and CCRT. Heat hardening in larvae impaired CT_{min}, CCRT, SCP, CT_{max} but not HKDT. In adults, both heat and cold hardening generally impaired CT_{min} and CT_{max}, but had no effects on HKDT, SCP and CCRT. Low temperature acclimation significantly improved CT_{min} and HKDT while marginally compromising CCRT and CT_{max}, whereas high temperature acclimation had no significant effects on any traits except for HKDT in larvae. Similarly, low and high temperature acclimation had no effects on CT_{min}, SCPs and CT_{max}, while high temperature acclimation significantly compromised adult CCRT. Our results show that larvae are more thermally plastic than adults and can shift their thermal tolerance in short and long timescales. Larval plasticity advantage over adults reported here suggest asymmetrical ecological role of the larva relative to adults in facilitating *T. absoluta* invasion.

Keywords: Acclimation, environmental stress; hardening; invasive species; Pinworm, thermal tolerance

1. Introduction

Global climate change characterised by increased magnitude and frequency of extreme climatic conditions such as heat waves, cold snaps, severe droughts and floods poses a great threat to biodiversity (Bozinovic et al., 2013; IPCC, 2014; Colinet et al., 2015). Such changes to the biophysical environments may result in changes in the abundance and geographic distribution of invasive species (Dukes and Mooney, 1999; Walther et al., 2009; Hulme, 2017). Owing to this is the notion that invasive alien species are more eurythermal, or able to maintain physiological functionality across variable temperatures. This is in contrast to stenothermy in non-invasive or indigenous species, which tolerate a narrower thermal range, and thus experience compromised fitness under changing thermal environments (Boher et al., 2016). Furthermore, climate change may also alter the spatio-temporal availability of biotic resources, creating bottlenecks in survival of organisms (Huey and Kingsolver, 2019). Overcoming these environmental challenges is the first of several potential barriers determining whether a species may become established, naturalised and, ultimately, invasive (Richardson and Pysek, 2006; Nyamukondiwa et al., 2010; Mbande et al., 2019). Given that most insects are ectotherms and their body temperature closely matches environmental temperature, dramatic changes in climate and more specifically weather, may affect their survival (Gray, 2013). Indeed, temperature is one of the key abiotic factors known to directly influence reproduction, development, activity, fitness, survival and spatio-temporal distribution in insects (Bowler and Terblanche, 2008; Klepsatel et al., 2019). Thus, the potential to tolerate environmental temperature stress is important for fitness, survival and adaptation under global change (Karl et al., 2014; Boher et al., 2016). For example, insects have been reported to employ short and long term morphological, behavioural, physiological and molecular mechanisms to withstand extreme environmental conditions (Lachenicht et al., 2010). Physiologically, insects compensate through genetic adaptation which involves allele frequency changes (Karl et al.,

2014) as well as phenotypic plasticity, the ability of a single genotype to change morphological, biochemical and physiological characteristics of an organism under heterogeneous environments (Whitman and Ananthakrishnan, 2009; Sgrò et al., 2016; Griffith et al., 2019). Phenotypic plasticity regarding thermal tolerance encompasses a suite of mechanisms such as rapid cold- and heat-hardening (RCH and RHH respectively) (in the short term) (Chidawanyika and Terblanche, 2011a), acclimation (under managed laboratory conditions) (Mutamiswa et al., 2018a) or acclimatisation (in the long term under field conditions) (Sgrò et al., 2016). The underling basis of these mechanisms is the pre-exposure to sub-lethal temperatures (Chidawanyika and Terblanche, 2011a; Chidawanyika et al., 2017) which may lead to molecular responses such as the expression of heat shock proteins that act as chaperones against protein denaturation, in the case of high temperature exposure (Hoffmann et al., 2003; Nyamukondiwa et al., 2010). For cold tolerance, exposure to sub-lethal temperatures has been associated with increased polyols and lipid content, which protects the insects against extracellular freezing (see review Overgaard and MacMillan, 2017).

Phenotypic plasticity can be adaptive and has been reported to improve survival in Lepidoptera (Stotter and Terblanche, 2009; Chidawanyika and Terblanche, 2011a; Fischer et al., 2010; Mutamiswa et al., 2018a), Diptera (Overgaard and Sørensen, 2008; Kalosaka et al., 2009; Nyamukondiwa et al., 2010), Coleoptera (Chidawanyika et al., 2017; Nyamukondiwa et al., 2018) and related taxon e.g. Collembola (see Chown et al., 2007; Sengupta et al., 2017). Studies have also shown that phenotypic plasticity varies across ontogeny (Marais and Chown, 2008) and that immobile stages generally have inherent higher plasticity to compensate for their inability to behaviorally adapt through seeking benign microhabitats. During insect development, different life stages respond differently to abiotic stress due to the variable microenvironments they experience during their development. This creates differential

adaptations across ontogeny, including differences in behavioral compensation abilities (Mutamiswa et al., 2019).

Different hypotheses have been postulated to explain how phenotypic plasticity facilitates adaptation under changing environments. The beneficial acclimation hypothesis has received considerable interest among insect physiologists and ecologists (Klepsatel et al., 2019) although it is debatable (Woods and Harrison, 2002b). This hypothesis posits that acclimation to a particular environment confers performance and fitness advantage to an organism in that environment over another organism that has not been exposed to that particular environment (Leori et al., 1994; Wilson and Franklin, 2002; Woods and Harrison, 2002a). The ‘hotter-is-better’ hypothesis states that organisms thermally adapted to warmer environments may have a better performance and fitness advantage than those adapted to colder or benign environments (Klepsatel et al., 2019). Similarly, the optimal developmental temperature hypothesis postulates that organisms raised at optimal temperature have greater survival advantage across many thermal environments (Woods and Harrison 2002b; Klepsatel et al., 2019). In particular, beneficial acclimation and thermal plasticity may play an important role in facilitating the success of invasive pests, although the generality of these effects are not known.

The South American tomato pinworm, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) native to South America is one of the key global destructive insect pests of tomatoes causing yield losses of 80–100% (Desneux et al., 2010; Brévault et al., 2014). Following its invasion into North Africa around 2008, the past decade has been marked by its rapid spread and establishment in East, West and Southern Africa threatening food security and livelihoods (Desneux et al., 2010; Biondi et al., 2018; Mansour et al. 2018). The successful spread and establishment of *T. absoluta* is a result of a number of reasons. These include a wide host range (cultivated and wild) (Machekano et al., 2018), high fecundity (Uchôa-Fernandes et al., 1995; Tropea Garzia et al., 2012; Biondi et al., 2018), trade globalisation and increased human travel

(Williamson, 1996; Pimentel et al., 2001; Simberloff and Rejmanek, 2001) and tolerance to diverse environmental conditions including temperature (Tropea Garzia et al., 2012; Ponti et al., 2015; Santana et al., 2019).

Projected thermal variability under climate change may have contrasting effects on species adaptation, compromising plastic responses in some species thereby limiting their potential to spread and establish, while enhancing such responses in other species (Sgrò et al., 2016). One of the major points to emerge from recent work on thermal tolerance is that there may be a link between susceptibility to climate change and the magnitude of phenotypic plasticity (e.g. Kellett et al., 2005; Balanya et al., 2006; Calosi et al., 2008; Nyamukondiwa et al., 2011; though see Gunderson and Stillman, 2015; Gunderson et al., 2017). Given the dramatic increase in the risk and invasion rate globally in the past decades, an understanding of plastic responses of invasive species to thermal stress is of fundamental importance in predicting climate change effects on their population dynamics, spatio-temporal distribution (Pimentel et al., 2001; Hance et al., 2007; Ma et al., 2014) and potential establishment in new environments (Nyamukondiwa et al., 2010; Buckley and Huey, 2016).

While several reports have documented significant ecological effects of RCH (Stotter and Terblanche, 2009; Basson et al., 2011; Mutamiswa et al., 2018b), RHH (Chidawanyika and Terblanche, 2011a; 2011b; Mutamiswa et al., 2018b) and acclimation (Lagerspetz, 2006; Fischer et al., 2010; Chidawanyika and Terblanche, 2011a; Sgrò et al., 2016), little is known on phenotypic plasticity of *T. absoluta* and its thermal tolerance (Han et al. 2018). Previous studies on *T. absoluta* have only focused on thermal requirements for development (Krechmer and Foerster, 2015; but see van Damme et al., 2015). To the best of our knowledge, no study has been done to date with focus on its plastic responses to temperature, and likely contributory effects on invasion success. In addition, studies on acclimation responses to temperature treatment, a major mechanism used by insects to cope with temperature variation both in

medium-term timescales (Overgaard and Sørensen, 2008; Nyamukondiwa and Terblanche, 2010; Nyamukondiwa et al., 2010) and over the longer term (e.g. seasonal timescales, Nyamukondiwa et al., 2013) also remain underexplored. In the present study, we investigated the thermal tolerance of *T. absoluta* larvae (4th instar) and adults (24–48 h old) through determining their short- (RCH and RHH) and long -term (acclimation) plastic responses of fitness and survival. We hypothesize that *T. absoluta* is thermally plastic at variable thermal regimes. Understanding such factors associated with successful invasion provides a basis for a predictive framework important for mitigating the introduction, establishment or spread of potentially invasive pest species, and informing suitable control measures (e.g. when and where to focus efforts).

2. Materials and Methods

2.1 Insect culture

Wild populations of *T. absoluta* larvae were collected from infested tomatoes plants at Genesis Farm (S21°8'54.239"; E027°38'48.42"), Matshelagabedi village in North East District of Botswana. All specimens were collected in austral summer (December-February), so that all test organisms have the same environmental history. Larvae were reared in the laboratory on organically produced and pesticide-free excised fresh tomato (“Rodade”) leaves inside Bugdorm rearing cages (240mm³; Bugdorm-BD43030F, Megaview Science Co., Ltd, Taiwan) under optimum conditions, i.e., 28°C; 65±10% relative humidity (RH) (Guimapi et al., 2016) and 12L:12D photocycle in climate chambers (HPP 260; Memmert GmbH + Co.KG, Germany) until adult emergence (for adult experiments). For all the experiments, 4th instar wild larvae (field collected) and 24–48h old adults (eclosed from field collected larvae) were used.

2.2 Low temperature plasticity

Tuta absoluta hardening experiments were performed using established experimental protocols for Diptera and Lepidoptera (e.g., Nyamukondiwa et al., 2010; Mutamiswa et al., 2018b). To assess the effects of RCH and RHH on *T. absoluta* physiological traits, hardening temperatures for the two developmental treatment stages (larvae and adults) were established through preliminary assays. Cold hardening temperature for larvae was derived from critical thermal minima (CT_{min}) (Hoffmann et al., 2003; Lee and Denlinger, 2010) and was defined as 6°C below CT_{min} (Mutamiswa et al., 2018b) (Table 1). For adults, low hardening temperature was defined as 10°C above the lower discriminating temperature, consistent with other studies (see Nyamukondiwa et al., 2010) (Table 1). Lower discriminating temperature (-10°C), defined as the temperature that causes 80–100% mortality upon 2 h exposure to a stressful low temperature was derived from Machekano et al. (2018). These hardening temperatures suffice to elicit RCH responses in similar organisms under laboratory experimental conditions (Hoffmann et al., 2003; Lee and Denlinger, 2010). Larval and adult *T. absoluta* specimens were first placed in a climate chamber at 28°C and 65±10% RH for 30 min to allow for equilibration. Thereafter, the insects were placed in a zip lock bag and submerged in a programmed water bath (Systronix Scientific, Industria, South Africa) with 1:1 water: propylene glycol and hardened for 2 h at low hardening temperature (-2°C (larvae) and 1°C (adults)). Following treatment, insects were allowed to recover under benign conditions (28°C; 65±10% RH) in a climate chamber for 30 min before measuring physiological traits. Control insects were kept at benign conditions during the same treatment before measuring thermal fitness traits.

As for RHH effects, hardening high temperature for larvae was derived from critical thermal maxima (CT_{max}) (Hoffmann et al., 2003; Lee and Denlinger, 2010) and defined as 7°C below CT_{max} , (see e.g. Mutamiswa et al., 2018b). For the adults, hardening high temperature was defined as 5°C below the discriminating high temperature (42°C), derived from Machekano et al. (2018). This hardening temperature is generally adequate to elicit an RHH response in

similar insects (Hoffmann et al., 2003). Organisms were first temperature equilibrated in a climate chamber at 28°C and 65±10% RH for 30 min before being hardened for 2 h at high temperature (see table 1). Following hardening, organisms were given 30 min recovery time to allow the *de novo* synthesis of heat shock proteins (HSPs) (Nyamukondiwa et al., 2010; Mutamiswa et al., 2018a) before measuring physiological traits. Control organisms were maintained at optimum conditions (28°C; 65% RH) before measuring physiological traits. Traits measured include critical thermal minima (CT_{min}), chill coma recovery time (CCRT) and supercooling points (SCPs).

Long term acclimation effects on low thermal tolerance were tested using standardised protocols (see Nyamukondiwa and Terblanche, 2010; Mutamiswa et al. 2018b). Acclimation temperatures were defined as 5°C above and below the optimal temperature (28°C), for low (23°C) and high (33°C) temperature pre-conditioning respectively. The control organisms were acclimated to an optimum temperature of 28°C and 65% RH. Following three-day acclimation, physiological traits were measured. Such temperature duration is also sufficient to elicit acclimation responses in similar insect taxa (Nyamukondiwa and Terblanche, 2010; Weldon et al., 2011; Mutamiswa et al. 2018a).

2.2.1 Critical thermal minima (CT_{min})

Critical thermal minima were measured on *T. absoluta* larvae and adults using standardized protocols as outlined by Nyamukondiwa and Terblanche (2009). Ten replicate organisms were individually placed in a series of numbered borosilicate glass tubes ('organ pipes') connected to an insulated double-jacketed chamber which was linked to a programmable water bath before decreasing the temperature at a rate of 0.25°C/min until their CT_{min} were recorded. This was repeated twice to yield sample sizes of $n = 20$ per treatment. CT_{min} was defined as the

temperature at which each individual insect lost coordinated muscle function and ability to respond to mild stimuli using a thermally inert object.

2.2.2 Chill coma recovery time (CCRT)

Chill coma recovery time experiments were conducted following established protocols (e.g. Weldon et al., 2011; Nyamukondiwa et al., 2018). Ten organisms from each temperature treatment i.e., larvae and adults were individually placed in 2ml Eppendorf tubes and then loaded into a zip-lock bag which was then submerged in a water bath (1:1 water: propylene glycol) set at 0°C (larvae) and -2 °C (adults) for 1 h. After 1 h under chill coma temperature, the vials were removed from the water bath and placed in a climate chamber set at 28°C, 65±10% RH for recovery. The chamber was connected to a video recording camera (HD Covert Network Camera, DS-2CD6412FWD-20; Hikvision Digital Technology Co. Ltd, Hangzhou, China) which was linked to a computer for recording observations. This was repeated twice to yield a sample size of n = 20. Chill coma recovery time was defined as the time (in min) required for the insects to regain consciousness, e.g., movement for larvae and standing upright on legs (for adults) following recovery from chill-coma.

2.2.3 Supercooling points (SCPs)

Supercooling points were assayed as outlined by Nyamukondiwa et al., (2013). Twenty organisms from each developmental stage (larvae and adults) and temperature treatment were individually placed into 2 ml Eppendorf tubes. These organisms were ‘fasted’ for 24 hours to clear gut contents and circumvent confounding effects the food gut particles may have on SCPs. Each insect was placed in contact with the tip of a type-T copper-constantan thermocouple, inserted through the gauzed lid of the vial and both the insect and thermocouple were secured in contact by a cotton wool. Thermocouples were connected to one of two 8-channel Picotech

TC-08 thermocouple interfaces and temperatures were recorded at 1s intervals using PicoLog software for Windows. In all treatments the insects were first exposed to a temperature of 15°C for 10 min before decreasing the temperature at a rate of 0.5°C/ min until SCPs were recorded. SCP for each organism was determined as the lowest temperature recorded before a spike in temperature due to latent heat of crystallization (Nyamukondiwa et al., 2013).

2.3 High temperature plasticity

Hardening and acclimation experiments were conducted using standardised protocols, consistent with low temperature plasticity experiments (see section 2.2) (e.g. Nyamukondiwa et al., 2010; Nyamukondiwa and Terblanche, 2010; Mutamiswa et al., 2018b). Physiological traits were measured following hardening (RCH and RHH) and three-day acclimation at low (23°C), optimum (28°C) and high (33°C) temperature for both larvae and adults. Control organisms were also maintained at optimum temperature (28°C; 65% RH) before measuring physiological traits. Fitness traits measured following acclimation treatments include CT_{max} and HKDT.

2.3.1 Critical thermal maxima (CT_{max})

Critical thermal maxima were measured using the same protocol as that for CT_{min} except that temperature was ramped up at 0.25°C/min until the endpoint. Critical thermal maximum was defined as the temperature at which each individual insect lost coordinated muscle function and ability to respond to mild stimuli using a thermally inert object.

2.3.2 Heat knock down time (HKDT)

Heat knockdown experiments were assayed as outlined by Mutamiswa et al. (2018a). Ten replicate individuals in both larva and adult stage were individually placed in 0.65ml

microcentrifuge tubes and placed in a climate chamber connected to a camera linked to a computer. The tubes carrying the insects were then exposed to an acute knockdown temperature (50°C) in the climate chamber. This knockdown temperature was selected basing on preliminary investigations of CT_{max} for larvae and adults ranging 49±0.13°C and 48±0.2°C respectively. This was repeated twice to yield sample sizes of $n = 20$. All observations were made from the climate chamber video recording system. Heat knockdown time was defined as the time (in minutes) at which insects lost activity following exposure to knockdown temperature in the climate chamber.

2.4 Statistical Analyses

Data analyses were done using STATISTICA, version 13.0 (Statsoft Inc., Tulsa, Oklahoma) and R version 3.3.0 (R Development Core Team, 2019). Critical thermal limits (CT_{max}, CT_{min}), SCP, CCRT and HKDT data were first checked for normality and equality of variances using the Shapiro-Wilk and Hartley-Bartlett tests and met assumptions of ANOVA in all traits except for HKDT. Therefore CT_{max}, CT_{min}, SCP and CCRT results were analysed using factorial ANOVA in STATISTICA. Tukey-Kramer's *post-hoc* tests were used to separate statistically heterogeneous groups. Heat knockdown time results were analysed using generalized linear models (GLZ) assuming a Gaussian distribution and an identity link function in R. Kruskal–Wallis *post hoc* tests were used to separate statistically homogeneous groups for HKDT data.

3. Results

3.1 Low temperature plasticity

3.1.1 Critical thermal minima

Hardening (RCH and RHH) significantly affected CT_{min} in both larvae and adults, but the direction of effect differed between the life stages (treatment x life stage interaction, $p < 0.001$,

Table 2). Rapid cold hardening significantly improved CT_{min} in larvae while impairing it in adults (Fig. 1A). However, RHH significantly compromised CT_{min} in adults (Fig. 1A), and marginally improved in larvae. Acclimation also significantly influenced cold tolerance (CT_{min}) ($P < 0.001$) such that low temperature acclimation (23°C) significantly improved CT_{min} in larvae while it had no effects on adults. Similarly, high temperature acclimation (33°C) had no significant effects for CT_{min} for both larval and adult *T. absoluta* (Table 2; Fig. 1B). As such, life stage $\times$ acclimation interaction effect was not significant ($P > 0.05$) (Table 2).

3.1.2 Chill coma recovery time

Hardening significantly influenced CCRT in larvae and adults ($P < 0.001$) (Table 3; Fig. 2A). Rapid heat hardening significantly compromised CCRT in adults, and marginally so in larvae, albeit the latter was non-significant (Fig. 2A). There were no significant differences in CCRT between larvae and adults following RCH, between the controls and RHH larval treatment ($P > 0.05$) (Fig. 2A). Nevertheless, larval CCRT did not significantly differ from the controls. There was a higher magnitude of difference in cold tolerance (CCRT) between larvae and adults following RHH (14.63 min) than RCH (0.23 min) (Fig. 2A). In addition, life stage $\times$ treatment interaction effect was significant ($P < 0.001$) (Table 3). Acclimation also significantly affected CCRT of *T. absoluta* larvae and adults ($P < 0.001$) (Table 3; Fig. 2B). Low temperature acclimation (23°C) marginally improved CCRT in larvae and adults (recovered faster from chill coma) whereas high temperature acclimation (33°C) compromised it (took more time to recover) in both developmental stages (Fig. 2B). Nevertheless, life stage $\times$ acclimation interaction effect was not significant ($P > 0.05$) (Table 3).

3.1.3 Supercooling points

Supercooling points for larvae and adults were significantly affected by hardening ($P<0.001$; Fig. 3A). For larval *T. absoluta*, both RCH and RHH significantly impaired supercooling ability (more positive SCPs), while for adults, both hardening treatments had no significant effects (Table 4; Fig. 3A). There was a significant life stage $\times$ treatment interaction effect ($P<0.001$) following hardening (Table 4). There were no significant acclimation effects on larval and adult SCPs (Table 4; Fig. 3B). For all developmental stages, there was no significant difference between the controls and that of the two treatments (23 and 33°C) (Fig. 3B). However, there was developmental stage related difference in SCPs following acclimation to high temperature (33°C), with larvae having more negative SCPs than the adults (Fig. 3B). In addition, life stage $\times$ acclimation interaction effect was significant ($P<0.05$) (Table 4).

3.2 High temperature plasticity

3.2.1 Critical thermal maxima

Heat tolerance (CT_{max}) in larvae and adults was significantly affected by hardening ($P<0.001$) (Table 5; Fig. 4A). Hardening significantly reduced CT_{max} in adults, and more significantly so for the RHH treatment (Fig. 4A). However, there was no significant difference in CT_{max} for larvae following both RCH and RHH (Fig. 4A). Similarly, life stage $\times$ treatment interaction effect was significant ($P<0.001$, Table 5). In addition, acclimation effects were not significant for larvae and adults CT_{max} (Table 5; Fig. 4B). For both developmental stages, low (23°C) and high (33°C) temperature treatments did not significantly affect CT_{max} from the control level (Fig. 4B). Nevertheless, it appeared larvae had higher CT_{max} than adults across all the treatments (Fig. 4B). The interaction between life stage and acclimation was also not significant (Table 5).

3.2.2 Heat knockdown time

As in CT_{max} , hardening significantly affected HKDT of larvae and adults ($P<0.001$) (Table 6; Fig 5A). Rapid cold hardening significantly compromised HKDT for larvae, while it showed no significant effects for the adults (Fig. 5A). Similarly, no significant differences in HKDT were noted in adults following both hardening treatments, and between controls and RHH treatments for larvae (Fig. 5A). In addition, there was a significant life stage $\times$ treatment interaction effect ($P<0.001$) (Table 6). Generally, larvae had higher HKDT (took longer to be knocked down following acute heat stress) than adults across all treatments (Fig. 5A). Acclimation also significantly influenced larvae and adults HKDT ($P<0.001$) (Table 6; Fig 5B). Low temperature acclimation ($23^{\circ}C$) significantly improved HKDT in both larvae and adults whereas high temperature acclimation ($33^{\circ}C$) impaired it in both life stages (Fig. 5B). Similarly, there was a significant life stage $\times$ acclimation interaction effect (Table 6). Consistent with other traits, it also appeared larvae had significantly higher heat tolerance (higher HKDT) across all treatments (Fig. 5B).

4. Discussion

The success of invasive species may stem from their ability to withstand stressful ambient novel environments. As such, basal and phenotypic plasticity traits play a pivotal role in invasive species' successful spread and establishment in new environments. Indeed, fitness and survival may be dependent on adjusting to novel environmental conditions through physiological acclimatization/acclimation and genetic adaptation (Webster et al., 2017; Castañeda et al., 2019; Griffith et al., 2019). It is increasingly being documented that invasive species may particularly have flexible life history traits (Agosta et al., 2018), and this potentially explains their success under heterogeneous environments.

The present study showed that rapid hardening and acclimation affects plasticity of thermal tolerance in *T. absoluta*, albeit asymmetrically for larvae and adults. Larvae appeared to be

more thermally plastic than adults, suggesting that *T. absoluta* invasion propagules may be carried in the form of larvae under thermally extreme environments. Hence, this enhanced larval plasticity coupled with the limited one in adults may aid each other to enhance invasion success (see discussions in Nyamukondiwa et al., 2010).

It appeared there was higher plasticity for low temperature compared to high temperature following both short- and long-term hardening. Changes in thermal limits as a result of acclimation are thought to be more pronounced at the lower- than upper -thermal thresholds (Klok and Chown, 2003; Weldon et al., 2011; Hoffmann et al., 2013). This is in agreement with our current results where both RCH and low temperature acclimation had a positive effect on CT_{min} of *T. absoluta* larvae. Coello Alvarado et al. (2015) reported improved CT_{min} in *Gryllus pennsylvanicus* adults following long- and -short term acclimation. Similar results have also been reported for *C. partellus* larvae, adult parasitoid, *Cotesia flavipes* (Mutamiswa et al. (2018a) and *Busseola fusca* larvae (Mutamiswa et al., 2018b).

In terms of low temperature tolerance (CCRT and CT_{min}), we found that adults had higher basal tolerance than larvae, corroborating a recent study by Machekano et al. (2018) who reported superior basal cold tolerance in adults. Interestingly, larvae showed strikingly higher plasticity of low temperature tolerance than adults. Insects often trade off basal low temperature tolerance for phenotypic plasticity (Nyamukondiwa et al., 2011), consistent with the current results which showed compromised basal low temperature tolerance and improved plasticity of cold tolerance (CCRT and CT_{min}) for larvae. In addition, this also affirms the notion that insects with high basal thermal tolerance may be constrained in phenotypic plasticity of thermal traits (Stillman, 2003; Coello Alvarado et al., 2015). Studies have also shown that less or non-mobile life-stages are generally considered to have evolved inherent high phenotypic plasticity to abiotic stress, to compensate for limited ability of behavioural compensation (Vernon and

388 Vannier, 1996; Klok and Chown, 2001; Jensen et al., 2007; Marais et al., 2009; Mutamiswa et
389 al., 2019).

390 Although Coello Alvarado et al. (2015) reported improved CCRT in *G. pennsylvanicus*
391 following cold acclimation and RCH, our results report no effects for both *T. absoluta*
392 developmental stages for both RCH and low temperature acclimation. The reason for this
393 absence is unknown but may, to a lesser extent be explained by the developmental stages we
394 tested, and choice of acclimation treatments (see discussions in Mutamiswa et al., 2019).
395 Nevertheless, this result indicates that plasticity of thermal tolerance in insects may be species
396 or trait dependent (Nyamukondiwa et al., 2018). It also supports the notion that differences in
397 responses to hardening and acclimation depend on duration of acclimation exposure hence this
398 may consequently determine the direction of induced plasticity and any associated costs and/or
399 benefits (Bowler and Terblanche, 2008). In addition, RHH and high temperature acclimation
400 (33°C) generally impaired CT_{min} (more positive) in both developmental stages. This result is
401 in keeping with Mutamiswa et al. (2018a; 2018b) who reported compromised cold tolerance
402 (CT_{min}) in larvae and adults of *C. partellus* and *B. fusca* larvae following high temperature
403 acclimation and RHH respectively. Nevertheless, impaired CCRT in adults following high
404 temperature acclimation corroborates a previous study on *C. flavipes* adults (Mutamiswa et al.,
405 2018a). A comparison of larvae and adults showed that the former recovered faster than latter
406 from chill coma following RHH. This likely means that mechanisms associated with tolerance
407 to low and high temperature for *T. absoluta* are decoupled (Hoffmann et al., 2003;
408 Nyamukondiwa et al., 2011). Rapid cold hardening has also been shown to improve thermal
409 fitness in other insect species e.g., *Drosophila melanogaster* (Overgaard and Sørensen, 2008);
410 *Ceratitis rosa* (Nyamukondiwa et al., 2010) and *Cydia pomonella* (Chidawanyika and
411 Terblanche, 2011a).

The thermal physiology of any organism is related to the range of environmental temperature where it evolved (Bryant et al., 2002; Hoffmann et al., 2003; Weldon et al., 2011; Ismail and Brookes, 2016; Griffith et al., 2019). *Tuta absoluta* larvae dwell in tomato leaves and feed by mining the leaf mesophyll (Biondi et al., 2018), where it is somewhat buffered from environmental heterogeneity. Thus, the higher larval plasticity to low temperature reported here may be an evolutionary adaptation to cope with changing ambient environments. Faced with stressful environments, adults on the other hand, can adjust their behaviour through e.g. flight (Bowler and Terblanche, 2008). This may help explain the limited plasticity reported here for the adults.

Both short- and long-term acclimation did not have any significant effects on SCPs in adults and larvae of *T. absoluta*. The results are in keeping with Mutamiswa et al. (2018a) who reported compromised SCPs for *C. partellus* pupae and adults following low and high temperature acclimation. Interesting, a comparison of larvae and adults showed an enhanced SCPs (more negative temperature) for the larvae relative to adults across all treatments, suggesting higher cold hardiness for the larvae. Boardman et al., 2012 reported that 24 h fasting removes all gut contents in *Thaumatotibia leucotreta* larvae resulting in a decrease in SCPs. While clearance of gut contents was not tested after 24 h treatment in this study, we hypothesise this may likely not have affected our SCPs results here. Some studies have reported that *T. absoluta* may struggle to survive in low temperatures (e.g. Cuthbertson et al., 2013; Biondi et al., 2018), and that it may not be able to overwinter under sub-zero environmental conditions. Nevertheless, few empirical studies have tested its cold hardiness under laboratory or field conditions (but see van Damme et al., 2015; Han et al., 2018). This therefore calls for imminent studies looking at *T. absoluta* cold hardiness and overwintering physiology, including ontogenic freeze strategy to help explain how this may affect its current and future geographic

distribution as well as population dynamics with a view to developing sustainable pest management programs (Tonnang et al., 2015; Santana et al., 2019).

Our study demonstrates effects of short- and long-term acclimation on *T. absoluta* high temperature tolerance. While RHH, RCH, low and high temperature acclimation did not have a significant effect on *T. absoluta* larvae, it appeared to come at a significant cost for adults (see Fig. 4A). Our results are in keeping with Mutamiswa et al. (2018b) who reported impaired heat tolerance (CT_{max}) in *B. fusca* and *Sesamia calamistis* larvae following RCH as well as low and high temperature acclimation. Some insects have been reported to trade off plasticity for other physiological or fitness traits (Liefting and Ellers, 2008; Angilletta, 2009; Basson et al., 2011; Murren et al., 2015).

Thermal conditioning to a single environmental stress often has deleterious consequences upon exposure to a different or same stress (reduced fitness). This may partially be accounted for by the cumulative effects of the stress combinations (see discussions in Gotcha et al., 2017). Fitness maladaptations owing to cumulative abiotic stresses reported here somewhat vary and depend on methodology employed e.g. short versus long term acclimation and the magnitude of stress severity (see Mittler, 2006). Similarly, RCH also significantly reduced HKDT, suggesting similar heat tolerance costs of short term low temperature acclimation (Fig. 5A). Conversely, acclimation to low temperature improved HKDT for both adult and larval *T. absoluta* larvae suggesting significant cross-tolerance effects. Phenotypic plasticity to one stress factor often confers fitness advantages to a different stressor, a phenomenon called ‘cross-tolerance’ (Sinclair et al., 2015). Such a phenomenon shows that shared co-evolutionary physiological stress resistance mechanisms exist across seemingly heterogeneous stress factors and may help species survive divergent environments.

In conclusion, we document larval and adult *T. absoluta* plasticity of thermal tolerance, and their likely contributions to invasion success. To our knowledge, this is the first study to

describe short- and long -term plastic responses to temperature in this species, using field collected populations. Determining the invasion pathway and pest risk assessment is significant for management of invasive pest species. As such, determining risk factors and mechanisms facilitating invasive species upon introduction to novel environments is critical in focusing control efforts. Although our results show the effects of acclimation on *T. absoluta* are complex and may vary considerably depending on the traits and life stage examined, it appears larvae are generally more plastic under short- and long- term acclimation than adults. Therefore, we suggest that larval propagules may be contributing more to the ongoing invasion of *T. absoluta*, owing to their better ability to shift their thermal phenotypes under heterogeneous environments. More studies could explore the role of parental history including testing invasive versus non-invasive species or endemic versus invasive environments to tease apart the exact role of plasticity for invasion success (as in e.g. Jarošík et al., 2015). Furthermore, the limited plasticity in adults ought to be further explored, including the role of other temperature, time and RH combinations in eliciting acclimation (see Mutamiswa et al., 2019). These results are important for pest risk analysis and for informing decision making in management this invasive species.

Declaration of interest

The authors declare no conflict of interest

Funding

This study received no specific grant from any funding agency in the public, commercial or non-profit organisations.

Acknowledgements

Authors would like to acknowledge the generous financial support from Botswana International University of Science and Technology (BIUST) to VT, RM and CN; University of the Free State (UFS) funding to RM and FN and a University of Bristol Global Challenges Research Fund Partnerships Award to SE and CN. SE was supported by a Royal Society Dorothy Hodgkin Fellowship. We also want to thank Mmabaledi Buxton and Virgil Joseph for assistance with insects filed collection, and to several anonymous referees for the valuable inputs in a previous version of this manuscript.

References

- Agosta, S.J., Joshi, K.A., Kester, K.M., 2018. Upper thermal limits differ among and within component species in a tritrophic host-parasitoid hyperparasitoid system. *PLoS One*. 13: e0198803.
- Angilletta, M.J., 2009. *Thermal Adaptation*. A Theoretical and Empirical Synthesis. Oxford University Press, Oxford, New York.
- Balanya, J., Oller, J., Huey, R., Gilchrist, G., Serra, L., 2006. Global Genetic Change Tracks Global Climate Warming in *Drosophila subobscura*. *Science*. 313: 1773-1775.
- Basson, C.H., Nyamukondiwa, C., Terblanche, J.S., 2011. Fitness costs of rapid cold hardening in *Ceratitis capitata*. *Evolution*. 66: 296-304.
- Biondi, A., Guedes, R.N.C., Wan, F-H., Desneux, N., 2018. Ecology, Worldwide Spread, and Management of the Invasive South American Tomato Pinworm, *Tuta absoluta*: Past, Present, and Future. *Annu. Rev. Entomol.* 63: 239–58.
- Boardman, L., Grout, T. G., Terblanche, J. S., 2012. ‘False codling moth *Thaumatotibia leucotreta* (Lepidoptera, Tortricidae) larvae are chill-susceptible’, *Insect Sci.* 19: 315–328.
- Boher, F., Trefault, N., Estay, S.E., Bozinovic, F., 2016. Ectotherms in variable thermal landscapes: A physiological evaluation of the invasive potential of fruit flies species. *Front. Physiol.* 7: 302.

512 Bowler, K. and Terblanche, J.S., 2008. Insect thermal tolerance: what is the role of ontogeny,
513 ageing and senescence. *Biol. Rev.* 83: 339–355.

514 Bozinovic, F., Catalan, T.P., Estay, S.A., Sabat, P., 2013. Acclimation to daily thermal
515 variability drives the metabolic performance curve. *Evol. Ecol. Res.* 15: 579-587.

516 Brévault, T., Sylla, S., Diatte, M., Bernadas, G., Diarra, K., 2014. *Tuta absoluta* Meyrick
517 (Lepidoptera: Gelechiidae): A new threat to tomato production in Sub-Saharan Africa. *Afr.*
518 *Entomol.* 22: 441-444.

519 Bryant, S.R., Thomas, C.D., Bale J.S., 2002. The influence of thermal ecology on the
520 distribution of three nymphalid butterflies. *J. Appl. Ecol.* 39: 43–55.

521 Buckley, L.B. and Huey, R.B., 2016. How extreme temperatures impact organisms and the
522 evolution of their thermal tolerance. *Integr. Comp. Biol.* 56: 98–109.

523 Calosi, P., Bilton, D.T., Spicer, J.I., 2008. Thermal tolerance, acclimatory capacity and
524 vulnerability to global climate change. *Biol. Lett.* 4: 99-102.

525 Castañeda, L.E., Romero-Soriano, V., Mesas, A., Roff, D.A., Santos, M., 2019. Evolutionary
526 potential of thermal preference and heat tolerance in *Drosophila subobscura*. *J. Evol. Biol.* 32:
527 818–824.

528 Chidawanyika, F., and Terblanche, J.S., 2011a. Rapid thermal responses and thermal tolerance
529 in adult codling moth *Cydia pomonella* (Lepidoptera: Tortricidae). *J. Insect Physiol.* 57: 108-
530 117.

531 Chidawanyika, F., and Terblanche, J.S., 2011b. Costs and benefits of thermal acclimation for
532 codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae): implications for pest control and
533 the sterile insect release programme. *Evol. Appl.* 4: 534– 544.

534 Chidawanyika, F., Nyamukondiwa, C., Strathie, L., Fischer, K., 2017. Effects of thermal
535 regimes, starvation and age on heat tolerance of the Parthenium Beetle *Zygogramma bicolorata*
536 (Coleoptera: Chrysomelidae) following dynamic and static protocols. *PLoS One.* 12:
537 e0169371.

538 Chown, S.L., Slabber, S., McGeouch, M., Janion, C., Leinaas, H.P., 2007. Phenotypic plasticity
539 mediates climate change responses among invasive and indigenous arthropods. *Proc. Biol. Sci.*
540 274: 2531–2537.

541 Coello Alvarado, L.E., MacMillan, H.A., Sinclair, B.J., 2015. Chill-tolerant *Gryllus* crickets
542 maintain ion balance at low temperatures. *J. Insect Physiol.* 77: 15–25.

543 Colinet, H., Sinclair, B. J., Vernon, P., Renault, D., 2015. Insects in fluctuating thermal
544 environments. *Annu. Rev. Entomol.* 60: 123-140.

545 Cuthbertson, A.G.S., Mathers, J.J., Blackburn, L.F., *et al.*, 2013. Population development of
546 *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under simulated UK glasshouse
547 conditions. *Insects*, 4: 185-197.

548 Desneux, N., Wajnberg, E., Wyckhuys, K.A.G., Burgio, G., Arpaia, S., Narva´ez-Vasquez,
549 C.A., et al., 2010. Biological invasion of European tomato crops by *Tuta absoluta*: ecology,
550 geographic expansion and prospects for biological control. *J. Pest Sci.* 83: 197-215.

551 Dukes, J. S. and Mooney, H. A., 1999. Does global change increase the success of biological
552 invaders? *Trends Ecol. Evol.* 14: 135–139.

553 Fischer, K., Dierks, A., Franke, K., Geister, T.L., Liszka, M., Winter, S., Pflicke, C., 2010.
554 Environmental effects on temperature stress resistance in the tropical butterfly *Bicyclus*
555 *anyana*. *PLoS One.* 5: e15284.

556 Gotcha, N., Terblanche, J.S., Nyamukondiwa, C., 2018. Plasticity and cross-tolerance to
557 heterogeneous environments: divergent stress responses co-evolved in an African fruit fly. *J.*
558 *Evol. Biol.* 31: 98–110.

559 Gray, E.M., 2013. Thermal acclimation in a complex life cycle: The effects of larval and adult
560 thermal conditions on metabolic rate and heat resistance in *Culex pipiens* (Diptera: Culicidae).
561 *J. Insect Physiol.* 59: 1001–1007.

562 Griffith, T. C., Paterson, I.D., Owen, C.A., Coetzee, J.A., 2019. Thermal plasticity and
563 microevolution enhance establishment success and persistence of a water hyacinth biological
564 control agent. *Entomol. Exp. Appl.* 167: 616-625.

565 Guimapi, R.Y.A., Mohamed, S.A., Okeyo, G.O., et al. 2016. Modeling the risk of invasion and
566 spread of *Tuta absoluta* in Africa. Ecol. Complex., 28: 77–93.

567 Gunderson, A. R. and Stillman, J. H., 2015. Plasticity in thermal tolerance has limited potential
568 to buffer ectotherms from global warming. Proc. Royal Soc. B. 282: 20150401.

569 Gunderson, A. R., Dillon, M.E., Stillman, J. H., 2017. Estimating the benefits of plasticity in
570 ectotherm heat tolerance under natural thermal variability. Funct. Ecol. 31: 529-1539.

571 Han, P., Zhang, Y., Lu, Z., Wang, S., Ma, D., Biondi, A., Desneux, N., 2018. Are we ready for
572 the invasion of *Tuta absoluta*? Unanswered key questions for elaborating an Integrated Pest
573 Management package in Xinjiang, China. Entomol. Gen. 38: 113-125.

574 Hance, T., Van Baaren, J., Vernon, P., Boivin, G., 2007. Impact of extreme temperatures on
575 parasitoids in a climate change perspective. Annu. Rev. Entomol. 52:107–126.

576 Hoffmann, A.A., Chown, S.L., Clusella-Trullas, S., 2013. Upper thermal limits in terrestrial
577 ectotherms: how constrained are they? Funct. Ecol. 27: 934–949.

578 Hoffmann, A.A., Sørensen, J.G., Loeschcke, V., 2003. Adaptation of *Drosophila* to
579 temperature extremes: bringing together quantitative and molecular approaches. J. Therm.
580 Biol. 28: 175–216.

581 Huey, R.B. and Kingsolver, J.G., 2019. "Climate Warming, Resource Availability, and the
582 Metabolic Meltdown of Ectotherms," Am. Nat. 194: E140-E150.

583 Hulme, P. E., 2017. Climate change and biological invasions: evidence, expectations, and
584 response options. Biol. Rev. 92: 1297–1313.

585 IPCC., 2014. Climate change 2014: Synthesis report. Contribution of working groups I, II and
586 III to the fifth assessment report of the intergovernmental panel on climate change (eds.Core
587 Writing Team, R.K. Pachauri & L.A. Meyer), pp. 151. IPCC, Geneva, Switzerland.

588 Ismail, M. and Brookes, M., 2016. The impact of geographical origin of two strains of the
589 herbivore, *Eccritotarsus catarinensis*, on several fitness traits in response to temperature. J.
590 Therm. Biol. 60: 222–230.

591 Jarošík, V., Kenis, M., Honěk, A., Skuhrovec, J., Pyšek, P., 2015. Invasive insects differ from
 592 non-invasive in their thermal requirements. PloS One, 10. [https://doi.org/10.1371/journal.](https://doi.org/10.1371/journal.pone.0131072)
 593 pone.0131072.

594 Jensen, D., Overgaard, J., Sørensen, J.G., 2007. The influence of developmental stage on cold
 595 shock resistance and ability to cold-harden in *Drosophila melanogaster*. J. Insect Physiol. 53:
 596 179 – 186.

597 Kalosaka, K., Soumaka, E., Politis, N., Mintzas, A.C., 2009. Thermotolerance and Hsp70
 598 expression in Mediterranean fruit fly *Ceratitis capitata*. J. Insect Physiol. 55: 568-573.

599 Karl, I., Becker, M., Hinzke, T., Mielke, M., Schiffler, M., Fischer, K., 2014. Interactive effects
 600 of acclimation temperature and short-term stress exposure on resistance traits in the butterfly
 601 *Bicyclus anynana*. Physiol. Entomol. 39: 222-228.

602 Kellett, M., Hoffmann, A.A., Mckechnie, S.W., 2005. Hardening capacity in the *Drosophila*
 603 *melanogaster* species group is constrained by basal thermotolerance. Funct. Ecol. 19: 853–858.

604 Klepsatel, P., Girish, T.N., Dirksen, H., Gáliková, M., 2019. Reproductive fitness of
 605 *Drosophila* is maximised by optimal developmental temperature. J. Exp. Biol. 222: jeb202184.

606 Klok, C.J. and Chown, S.L., 2001. Critical thermal limits, temperature tolerance and water
 607 balance of a sub-Antarctic kelp fly, *Paractora druexi* (Diptera: helcomyzidae). J. Insect
 608 Physiol. 47: 95 – 109.

609 Klok, J.C. and Chown, S.L., 2003. Resistance to temperature extremes in sub-Antarctic
 610 weevils: interspecific variation, population differentiation and acclimation. Biol. J. Linn. Soc.
 611 78: 401–414.

612 Krechemer, F.S. and Foerster, L.A., 2015. *Tuta absoluta* (Lepidoptera: Gelechiidae): Thermal
 613 requirements and effect of temperature on development, survival, reproduction and longevity.
 614 Eur. J. Entomol. 112: 658–663.

615 Lachenicht, M.W., Clusella-Trullas, S., Boardman, L., Le Roux, C., Terblanche, J.S., 2010.
 616 Effects of acclimation temperature on thermal tolerance, locomotion performance and
 617 respiratory metabolism in *Acheta domestica* L. (Orthoptera: Gryllidae). J. Insect Physiol. 56:
 618 822–830.

619 Lagerspetz, K.Y.H., 2006. What is thermal acclimation? J. Therm. Biol. 31: 332-336.

620 Lee, R.E. and Denlinger, D.L., 2010. Rapid cold-hardening: ecological significance and
621 underpinning mechanisms, in *Low Temperature Biology of Insects*, ed. by Denlinger DL and
622 Lee RE. Cambridge University Press, UK, 35–58.

623 Leori, A.M., Bennett, A.F., Lenski, R.E., 1994. Temperature acclimation and competitive
624 fitness: an experimental test of the beneficial acclimation assumption. Proc. Natl. Acad. Sci.
625 USA. 91: 1917-1921.

626 Liefting, M. and Ellers, J., 2008. Habitat-specific differences in thermal plasticity in natural
627 populations of a soil arthropod. Biol. J. Linn. Soc. 94: 265– 271.

628 Ma, F-Z., Lü, Z-C., Wang, R., Wan, F-H., 2014. Heritability and evolutionary potential in
629 thermal tolerance traits in the invasive Mediterranean cryptic species of *Bemisia tabaci*
630 (Hemiptera: Aleyrodidae). PLoS One 9: e103279.

631 Machekano, H., Mutamiswa, R., Nyamukondiwa, C., 2018. Evidence of rapid spread and
632 establishment of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in semi-arid Botswana.
633 Agriculture & Food Security, 7: 48.

634 Mansour, R., Brévault, T., Chailleux, A., Cherif, A., Grissa-Lebdi, K., Haddi, K., et al., 2018.
635 Occurrence, biology, natural enemies and management of *Tuta absoluta* in Africa. Entomol.
636 Gen., 38: 83–112.

637 Marais, E. and Chown, S.L., 2008. Beneficial acclimation and the Bogert effect. Ecol. Lett. 11:
638 1027 – 1036.

639 Marais, E., Terblanche, J.S., Chown, S.L., 2009. Life-stage related differences in hardening
640 and acclimation of thermal tolerance traits in the kelp fly, *Paractora dreuxi* (Diptera:
641 helcomyzidae). J. Insect Physiol. 55: 336-343.

642 Mbande, A., Tedder, M., and Chidawanyika, F., 2019. Offspring diet supersedes the
643 transgenerational effects of parental diet in a specialist herbivore *Neolema abbreviata* under
644 manipulated foliar nitrogen variability. Insect Sci. 2: 361-374.

645 Mittler, R., 2006. Abiotic stress, the field environment and stress combination. Trends Plant
646 Sci. 11: 15-19.

647 Murren, C.J., Auld, J.R., Callahan, H., Ghalambor, C.K., et al., 2015. Constraints on the
648 evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. *Heredity*. 115:
649 293–301.

650 Mutamiswa, R., Chidawanyika, F., Nyamukondiwa, C., 2018a. Thermal plasticity potentially
651 mediates the interaction between host *Chilo partellus* Swinhoe (Lepidoptera: crambidae) and
652 endoparasitoid *Cotesia flavipes* Cameron (Hymenoptera: braconidae) under rapidly changing
653 environments. *Pest Manag. Sci.* 74: 1335-1345.

654 Mutamiswa, R., Chidawanyika, F., Nyamukondiwa, C., 2018b. Superior basal and plastic
655 thermal responses to environmental heterogeneity in invasive exotic stemborer *Chilo partellus*
656 Swinhoe over indigenous *Busseola fusca* (Fuller) and *Sesamia calamistis* Hampson. *Physiol.*
657 *Entomol.* 43: 108–119.

658 Mutamiswa, R., Machekano, H., Chidawanyika, F., Nyamukondiwa, C., 2019. Life-stage
659 related responses to combined effects of acclimation temperature and humidity on the thermal
660 tolerance of *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). *J. Therm. Biol.* 79: 85-94.

661 Nyamukondiwa, C. and Terblanche, J.S., 2009. Thermal tolerance in adult Mediterranean and
662 Natal fruit flies (*Ceratitis capitata* and *Ceratitis rosa*): effects of age, gender and feeding status.
663 *J. Therm. Biol.* 34: 406-414.

664 Nyamukondiwa, C. and Terblanche, J.S., 2010. Within-generation variation of critical thermal
665 limits in adult Mediterranean and Natal fruit flies *Ceratitis capitata* and *Ceratitis rosa*: thermal
666 history affects short-term responses to temperature. *Physiol. Entomol.* 35: 255–264.

667 Nyamukondiwa, C., Chidawanyika, F., Machekano, H., et al., 2018. Climate variability
668 differentially impacts thermal fitness traits in three coprophagic beetle species. *PLoS One*. 13:
669 e0198610.

670 Nyamukondiwa, C., Kleynhans, E., Terblanche, J.S., 2010. Phenotypic plasticity of thermal
671 tolerance contributes to the invasion potential of Mediterranean fruit flies (*Ceratitis capitata*).
672 *Ecol. Entomol.* 35: 565–575.

673 Nyamukondiwa, C., Terblanche, J.S., Marshall, K.E., Sinclair, B.J., 2011. Basal cold but not
674 heat tolerance constraints plasticity among *Drosophila* species (Diptera: Drosophilidae). *J.*
675 *Evol. Biol.* 24: 1927–1938.

676 Nyamukondiwa, C., Weldon, C.W., Chown, S.L., le Roux, P.C., Terblanche, J.S., 2013.
 677 Thermal biology, population fluctuations and implications of temperature extremes for the
 678 management of two globally significant insect pests. *J. Insect Physiol.* 59: 1199–1211.

679 Overgaard, J. and MacMillan, H. A., 2017. The Integrative Physiology of Insect Chill
 680 Tolerance. *Annu. Rev. Physiol.* 79: 187–208.

681 Overgaard, J. and Sørensen, J.G., 2008. Rapid thermal adaptation during field temperature
 682 variations in *Drosophila melanogaster*. *Cryobiology.* 56: 159–162.

683 Pimentel, D., McNair, S., Janecka, J., et al., 2001. Economic and environmental threats of alien
 684 plant, animal, and microbe invasions. *Agric. Ecosy. Environ.* 84: 1–20.

685 Ponti, L., Gilioli, G., Biondi, A., Desneux, N., Gutierrez, A.P., 2015. Physiologically based
 686 demographic models streamline identification and collection of data in evidence-based pest
 687 risk assessment EPPO Bulletin 45: 317–322.

688 Richardson, D. and Pyšek, P., 2006. Plant invasions: merging the concepts of species
 689 invasiveness and community invasibility. *Prog. Phys. Geog.* 30: 409–431.

690 Santana, P.A., Kumar, L., Da Silva, R.S., Picanço, M.C., 2019. Global geographic distribution
 691 of *Tuta absoluta* as affected by climate change. *J. Pest Sci.* 92: 1373–1385.

692 Sengupta, S., Ergon, T., Leinaas, H. P., 2017. Thermal plasticity in postembryonic life history
 693 traits of a widely distributed Collembola: Effects of macroclimate and microhabitat on
 694 genotypic differences. *Ecol. Evol.* 7: 8100–8112.

695 Sgrò, C.M., Terblanche, J.S., Hoffmann, A.A., 2016. What can plasticity contribute to insect
 696 responses to climate change? *Annu. Rev. Entomol.* 61: 433–451.

697 Simberloff, D. and Rejmanek, M., 2001. *Encyclopedia of biological invasions*. Berkeley and
 698 Los Angeles, USA: University of California Press.

699 Sinclair, B.J., Coello Alvarado, L.E., Ferguson, L.V., 2015. An invitation to measure insect
 700 cold tolerance: methods, approaches, and workflow. *J. Therm. Biol.* 53: 180–197.

701 Stillman, J.H., 2003. Acclimation capacity underlies susceptibility to climate change. *Science.*
 702 301, 65.

703 Stotter, R.L. and Terblanche, J.S., 2009. Low temperature tolerance of false codling moth
704 *Thaumotibia leucotreta* (Meyrick) (Lepidoptera: Tortricidae) in South Africa. J. Therm. Biol.
705 34: 320–325.

706 Tonnang, H.E.Z., Mohamed, S.A., Khamis, F., Ekesi, S. 2015. Correction: Identification and
707 Risk Assessment for Worldwide Invasion and Spread of *Tuta absoluta* with a Focus on Sub-
708 Saharan Africa: Implications for Phytosanitary Measures and Management. PLoS One 10:
709 e0138319.

710 Tropea Garzia, G., Siscaro, G., Biondi, A., et al., 2012. *Tuta absoluta*, a South American pest
711 of tomato now in the EPPO region: biology, distribution and damage. OEPP. EPPO. Bull. 42:
712 205–210.

713 Uchôa-Fernandes, M.A., Della Lucia, T.M.C., Vilela, E.F., 1995. Mating, oviposition and
714 pupation of *Scrobipalpula absoluta* (Meyrick) (Lepidoptera: Gelechiidae). An. Soc. Entomol.
715 Bras. 24: 159–164.

716 Van Damme, V., Berkvens, N., Moerkens, R., Berckmoes, E., Wittemans, L., De Vis, R.,
717 Casteels, H., Tirry, L., De Clercq, P., 2015. Overwintering potential of the invasive leafminer
718 *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) as a pest in greenhouse tomato production
719 in Western Europe. J. Pest Sci. 88: 533–541.

720 Vernon, P. and Vannier, G., 1996. Developmental patterns of supercooling capacity in a
721 subantarctic wingless fly. Experientia. 52: 155 – 158.

722 Walther, G.R., Roques, A., Hulme, P.E., Sykes, M.T., Pyšek, P., Kühn, I., Zobel, M., Bacher,
723 S., Botta-Dukat, Z., Bugmann, H., Czucz, B., 2009. Alien species in a warmer world: risks and
724 opportunities. Trends Ecol. Evolut. 24: 686–693.

725 Webster, M.S., Colton, M.A., Darling, E.S., Armstrong, J., Pinsky, M.L., Knowlton, N.,
726 Schindler, D.E., 2017. Who Should Pick the Winners of Climate Change? Trends Ecol. Evolut.
727 32:167-173.

728 Weldon, C.W., Terblanche, J.S., Chown, S.L., 2011. Time-course for attainment and reversal
729 of acclimation to constant temperature in two *Ceratitis* species. J. Therm. Biol. 36: 479-485.

730 Whitman, D.W. and Ananthakrishnan, T.N., 2009. *Phenotypic Plasticity of Insects.*
731 *Mechanisms and Consequences.* Science Publishers, Enfield, NH, USA.

732 Williamson, M., 1996. *Biological Invasions.* New York, USA: Chapman & Hall.

733 Wilson, R.S. and Franklin, C.E., 2002. Testing the beneficial acclimation hypothesis. Trends
734 Ecol. Evol. 17: 66-70.

735 Woods, H. A. and Harrison, J. F., 2002b. Interpreting rejections of the beneficial acclimation
736 hypothesis: when is physiological plasticity adaptive? Evolution. 56: 1863-1866.

737 Woods, H.A. and Harrison, J.F., 2002a. The beneficial acclimation hypothesis versus
738 acclimation of specific traits: physiological change in water stressed *Manduca sexta*. Physiol.
739 Biochem. Zool. 74: 32-44.

740

Figure 1: Effects of short-term acclimation- (A) 2-h pre-treatments/hardening of *Tuta absoluta* larvae (42°C for RHH; -2°C for RCH) and adults (37°C for RHH; 1°C for RCH) and long-term acclimation (B)- three-day acclimation (23°C; 28°C and 33°C) on critical thermal minima (CT_{min}). Error bars represent 95% confidence limits (CLs) ($n=20$ per group). Means with the same letter are not significantly different from each other. RHH and RCH represents rapid heat- and -cold hardening respectively.

Figure 2: Effects of short-term acclimation (A)- 2-h pre-treatments/hardening of *Tuta absoluta* larvae (42°C for RHH; -2°C for RCH) and adults (37°C for RHH; 1°C for RCH) and long-term acclimation (B)- three-day acclimation (23 °C; 28 °C and 33 °C) on chill coma recovery time (CCRT). Error bars represent 95% confidence limits (CLs) ($n=20$ per group). Means with the same letter are not significantly different from each other. RHH and RCH represents rapid heat- and -cold hardening respectively.

Figure 3: Effects of long-term acclimation (A)- 2-h pre-treatments/hardening of *Tuta absoluta* larvae (42°C for RHH; -2°C for RCH) and adults (37°C for RHH; 1°C for RCH) and long-term acclimation (B)- three-day acclimation (23°C; 28°C and 33°C) on supercooling points (SCPs). Error bars represent 95% confidence limits (CLs) ($n=20$ per group). Means with the same letter are not significantly different from each other. RHH and RCH represents rapid heat- and -cold hardening respectively.

Figure 4: Effects of long-term acclimation (A)- 2-h pre-treatments/hardening of *Tuta absoluta* larvae (42°C for RHH; -2°C for RCH) and adults (37°C for RHH; 1°C for RCH) and long-term acclimation (B)- three-day acclimation (23°C; 28°C and 33°C) on critical thermal maxima (CT_{max.}) Error bars represent 95% confidence limits (CLs) ($n=20$ per group). Means with the same letter are not significantly different from each other. RHH and RCH represents rapid heat- and -cold hardening respectively.

Figure 5: Effects of long-term acclimation (A)- 2-h pre-treatments/hardening of *Tuta absoluta* larvae (42°C for RHH; -2°C for RCH) and adults (37°C for RHH; 1°C for RCH) and long-term acclimation (B)- three-day acclimation (23°C; 28°C and 33°C) on heat knockdown time (HKDT). Error bars represent 95% confidence limits (CLs) ($n=20$ per group). Means with the same letter are not significantly different from each other. RHH and RCH represents rapid heat- and -cold hardening respectively.

Table 1. Summary table showing hardening temperatures and duration for *T. absoluta* larvae and adults

Developmental stage	Hardening high temperature (°C)	Hardening low temperature (°C)	Control temperature(°C)
Larvae	42 (2h)	-2 (2h)	28
Adults	37 (2h)	1 (2h)	28

Table 2. Summary table on short term responses to high temperature after 2-h pre-treatments/hardening of *T. absoluta* larvae (42°C; -2°C) and adults (37°C; 1°C) and three-day acclimation (23°C; 28°C and 33°C) on CT_{min}. SS = sum of squares, df = degrees of freedom, MS = means of squares.

Trait	Effect	SS	df	MS	F	p
CT _{min}	<i>Hardening</i>					
	Intercept	191.77	1	191.77	345.80	< 0.001
	Life stage	117.81	1	117.81	212.43	< 0.001
	Treatment	177.68	2	88.84	160.19	< 0.001
	Life stage×Treatment	209.38	2	104.69	188.78	< 0.001
	Error	63.22	114	0.55		
	<i>Acclimation</i>					
	Intercept	34.03	1	34.03	68.47	< 0.001
	Life stage	503.07	1	503.07	1012.24	< 0.001
	Acclimation	8.66	2	4.33	8.72	< 0.001
	Life stage×Acclimation	1.03	2	0.52	1.04	0.36
	Error	56.66	114	0.5		

Table 3. Summary table on short term responses to high temperature after 2-h pre-treatments/hardening of *T. absoluta* larvae (42°C; -2°C) and adults (37°C; 1°C) and three-day acclimation (23°C; 28°C and 33°C) on CCRT. SS = sum of squares, df = degrees of freedom, MS = means of squares.

Trait	Effect	SS	df	MS	F	p
CCRT	<i>Hardening</i>					
	Intercept	3714.64	1	3714.64	88.82	< 0.001
	Life stage	668.59	1	668.59	15.99	< 0.001
	Treatment	2037.07	2	1018.54	24.35	< 0.001
	Life stage×Treatment	1473.12	2	736.56	17.61	< 0.001
	Error	4767.79	114	41.82		
	<i>Acclimation</i>					
	Intercept	931.3969	1	931.4	932.89	< 0.001
	Life stage	0.8031	1	0.80	0.80	0.37
	Acclimation	44.4514	2	22.22	22.26	< 0.001
	Life stage×Acclimation	1.3589	2	0.68	0.68	0.51
	Error	113.82	114	0.99		

Table 4. Summary table on short term responses to high temperature after 2-h pre-treatments/hardening of *T. absoluta* larvae (42°C; -2°C) and adults (37°C; 1°C) and three-day acclimation (23°C; 28°C and 33°C) on SCPs. SS = sum of squares, df = degrees of freedom, MS = means of squares.

Trait	Effect	SS	df	MS	F	p
SCP	<i>Hardening</i>					
	Intercept	35942.14	1	35942.14	50592.06	< 0.001
	Life stage	184.09	1	184.09	259.13	< 0.001
	Treatment	10.62	2	5.31	7.48	< 0.001
	Life stage×Treatment	10.45	2	5.22	7.35	< 0.001
	Error	80.99	114	0.71		
	<i>Acclimation</i>					
	Intercept	37655.36	1	37655.36	3564.70	< 0.001
	Life stage	139.04	1	139.04	13.16	< 0.001
	Acclimation	13.75	2	6.87	0.65	0.52
	Life stage×Acclimation	73.76	2	36.88	3.49	0.03
	Error	1204.23	114	10.56		

Table 5. Summary table on short term responses to high temperature after 2-h pre-treatments/hardening of *T. absoluta* larvae (42°C; -2°C) and adults (37°C; 1°C) and three-day acclimation (23°C; 28°C and 33°C) on CT_{max}. SS = sum of squares, df = degrees of freedom, MS = means of squares.

Trait	Effect	SS	df	MS	F	p
CT _{max}	<i>Hardening</i>					
	Intercept	273435.6	1	273435.6	352409.3	< 0.001
	Life stage	168	1	168	216.6	< 0.001
	Treatment	33.9	2	17	21.9	< 0.001
	Life stage×Treatment	35.5	2	17.7	22.9	< 0.001
	Error	88.5	114	0.8		
	<i>Acclimation</i>					
	Intercept	274075.7	1	274075.7	204793.5	< 0.001
	Life stage	104.3	1	104.3	78	< 0.001
	Acclimation	3.2	2	1.6	1.2	0.31
	Life stage×Acclimation	1.6	2	0.8	0.6	0.56
	Error	152.6	114	1.3		

Table 6. Summary table on short term responses to high temperature after 2-h pre-treatments/hardening of *T. absoluta* larvae (42°C; -2°C) and adults (37°C; 1°C) and three-day acclimation (23°C; 28°C and 33°C) on HKDT. SS = sum of squares, df = degrees of freedom, MS = means of squares.

Trait	Effect	SS	df	MS	F	p
HKDT	<i>Hardening</i>					
	Treatment	380	2	190	11.44	< 0.001
	Life stage	7635	1	7635	459.94	< 0.001
	Treatment×Life stage	523	2	261	15.75	< 0.001
	Residuals	1892	114	17		
	<i>Acclimation</i>					
	Acclimation	3775	2	1888	28.1	< 0.001
	Life stage	5754	1	5754	85.67	< 0.001
	Acclimation×Life stage	2268	2	1134	16.88	< 0.001
	Residuals	7657	114	67		