Wang, G-W., Boyd, O., Young, T. A., Bertrand, S. M., & Bower, J. F. (2020). Rhodacyclopentanones as Linchpins for the Atom Economical Assembly of Diverse Polyheterocycles. *Journal of the American Chemical Society*, 142(4), 1740-1745.
<https://doi.org/10.1021/jacs.9b12421>

Peer reviewed version

Link to published version (if available):
[10.1021/jacs.9b12421](https://doi.org/10.1021/jacs.9b12421)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via American Chemical Society at <https://pubs.acs.org/doi/10.1021/jacs.9b12421> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Rhodacyclopentanones as Linchpins for the Atom Economical Assembly of Diverse Polyheterocycles

Gang-Wei Wang,[†] Olivia Boyd,[†] Tom A. Young,[†] Sophie M. Bertrand[‡] and John F. Bower^{*,†}

[†] School of Chemistry, University of Bristol, Bristol, BS8 1TS, United Kingdom

[‡] GlaxoSmithKline R&D, Medicines Research Centre, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, United Kingdom

Supporting Information Placeholder

ABSTRACT: We outline a conceptual blueprint that provides direct and atom economical access to a wide range of complex polyheterocycles. Our method capitalizes on the ambiphilic reactivity of rhodacyclopentanones that arise upon exposure of cyclopropanes to Rh(I)-catalysts and CO. Using this approach, a wide array of polycyclizations are achieved, including variants that involve powerful dearomatizations and medium ring formations.

Expeditious strategies for the assembly of complex heterocycles are of significant value to synthetic chemists. In particular, there is high demand for protocols that generate challenging scaffolds, such as medium rings, sp^3 -rich systems and complex polycycles.¹ Within this context, we have developed a suite of by-product free processes where cyclopropane-derived metallacycles engage tethered π -unsaturates or nucleophiles in cycloadditions² or “capture-collapse” heterocyclizations,³ respectively.⁴ The latter exploits the electrophilicity of rhodacyclopentanones **I** that arise upon directing group controlled carbonylative C-C bond activation of aminocyclopropanes **1** (Scheme 1A).⁵ These can be accessed with a variety of substitution patterns and, where appropriate, in enantiopure form. Our heterocyclizations harness these features to provide high levels of regioselectivity and stereospecificity in the formation of challenging 7- and 8-membered N-heterocycles **2**.³ These are generated from key metallacyclic intermediate **II** by a sequence of C-Nu reductive elimination (to **III**) and protodemetalation.

The C-C bond activation triggered heterocyclizations we have developed so far generate one new ring, where C-H bond formation is the terminating step (**III** to **2**). If protodemetalation could be averted then the tantalizing opportunity of using C(sp^3)-Rh(I)-intermediates related to **III** in subsequent C-C bond formations would emerge (Scheme 1B). This design offers tremendous flexibility because a wide range of *endo*- or *exo*-polycyclizations can be envisaged by varying the nature and position of the electrophile. Significantly, this approach uses the rhodacyclopentanone as a linchpin for the assembly of two rings rather than one – its electrophilicity enables the first annulation, whereas its latent nucleophilicity facilitates the second (via **IV/V**). Accordingly, the rhodacyclopentanone functions as a synthetic equivalent to ambiphilic synthon **VI** (Scheme 1C), with this reactivity mode unveiled simply by carbonylative C-C bond activation of the easily installed aminocyclopropane moiety. In this report, we outline extensive studies concerning this broad concept. These results show how C-C bond activation of simple functionality can enable by-product free access to powerful and unusual reactivity patterns.⁶

The development of the polycyclizations in Scheme 1B was driven by the observation that cyclopropane **3a** is converted to

Scheme 1. Polyheterocycles via rhodacyclopentanones.

complex polycycle **4a** in 10% yield when exposed to [Rh(cod)₂]OTf (7.5 mol%), P(3,5-(CF₃)₂C₆H₃)₃ (15 mol%), PhOCH₂CO₂H (30 mol%) and CO (1 atm) at 140 °C (Table 1A). This process can be rationalized by invoking *endo*-polycyclization of key alkyl-Rh(I) intermediate **VII** (cf. **IV**) onto the C2-C3 π -system of the indole (vide infra). Extensive studies were undertaken to improve efficiency; ultimately, we found that the combination of [Rh(cod)₂]OTf (7.5 mol%) and 4-NMe₂C₆H₄CO₂H (30 mol%) delivers **4a** in 81% yield. The inclusion of phosphine ligands resulted in reduced yields, and neutral Rh-sources were ineffective. The acid additive had the most profound effect, with 4-Me₂NC₆H₄CO₂H (30 mol%) emerging as optimal from a broad

screen (see the SI). Higher or lower CO pressures offered no benefits and lower reaction temperatures were ineffective. The inclusion of Na₂SO₄ as a desiccant provided a small but reproducible benefit.⁷

Table 1. Dearomatizing *endo*-polycyclizations of indole systems.

^a Run at 140 °C; ^b Run in DCB (0.2 M); ^c [Rh(cod)₂]OTf (10 mol%) was used; ^d Run at 150 °C; ^e Run at 125 °C.

The process has broad scope with respect to the indole unit (Table 1A). Systems **3c-j** possessing electronically diverse substituents generated polycycles **4c-j** in good to excellent yield. Substitution at C7 results in diminished efficiencies such that **4k** was formed in only 21% yield. The process can be transferred to substituted aminocyclopropanes (Table 1B); for example, polycyclization of *trans*-1,2-disubstituted systems **3l-n** generated **4l-n** with excellent levels of diastereo- and regiocontrol. These features arise from selective cleavage of the less hindered proximal C-C bond *a* of **3l-n**, followed by transfer of cyclopropane stereochemistry to the product.⁸ The starting materials are easily accessed in enantioenriched form, and this allowed enantiospecific conversion of **3m** to **4m** (98.5:1.5 e.r.). Polycyclizations of trisubstituted cyclopropanes **3o** and **3p** resulted in efficient desymmetrization to provide **4o** and **4p**

with exquisite levels of diastereocontrol (Table 1C). For non-symmetrical system **3q**, C-C bond activation was selective for benzylic C-C bond *a* leading to regioselective generation of **4q**, again with complete diastereocontrol.

The processes in Table 1 validate the conceptual blueprint in Scheme 1B, and also provide a notable contribution to indole dearomatization chemistry.⁹ Uniquely, the method enables the concurrent formation of two C-C bonds at C-2 (i.e. “dual C-H functionalization”)¹⁰ such that this position formally functions as a carbene equivalent (Table 1D). Prior methods for accessing similar structures require preinstallation of a substituent at C-2.¹¹ Other catalytic dual functionalizing indole dearomatizations usually generate new C-C/C-X bonds at both C-2 and C-3.¹²

Polycyclization of **3a** in the presence of D₂O (300 mol%) delivered *deuterio-3aa* and *deuterio-4aa* in 32% and 42% yield, respectively (Equation 1). For *deuterio-3aa*, deuterium incorporation was observed at C2-H (17%) and C3-H (43%). Exchange at these positions is dependent on the presence of the Rh-catalyst but not the cyclopropane (Equation 4) or the acid additive (see the SI); these data support exchange by reversible C-H activation of **3a**. For *deuterio-4aa*, deuterium incorporation was observed at C2-H (27%), C11-H_a (20%) and C11-H_b (49%). When the same reaction was run for 72 hours, a similar pattern of deuterium incorporation was observed (Equation 2). Deuterium incorporation at C11-H_b of *deuterio-4aa/4ab* likely occurs at the stage of **3a** because similar levels

Scheme 2. Mechanistic Experiments.

of exchange are observed at C3-H of this system. When **4a** was re-subjected to the reaction conditions, but in the presence of D₂O, deuterium incorporation was observed at C2-H, likely due to enolization (Equation 3). Deuterium incorporation at C11-H_a of **deuterio-4aa/4ab** is consistent with *syn*-carbometallation of the C2-C3 π -system (from **VII**) prior to protodemetalation. The necessary proton originates from either the acid additive or C2-H of **3a**. When **deuterio-3ab** was exposed to standard conditions, the deuterium label was transferred predominantly to C11-H_b (Equation 5). Finally, exposure of equimolar quantities of **3a** and **deuterio-3ac** to standard conditions revealed a small KIE ($k_H/k_D = 1.21$), suggesting that C-H cleavage is not turnover limiting (Equation 6).¹³

Scheme 3. DFT analysis of C-C bond activation and C-C bond forming pathways.^a

^a Calculations performed at the B3PW91-D3BJ/6-311+G(2d,p),Rh(LANTZ(f))/SMD(DCB)/B3PW91-D3BJ/6-311G(d),Rh(MWB28) level of theory, with thermochemical corrections calculated at $T = 403$ K and a 1 M standard state.

The experiments above are consistent with a pathway involving stereoretentive protodemetalation of a C11-alkyl-Rh(I) species. To provide insight into (a) the mode of C-C bond activation and (b) the C-C bond forming sequence from the rhodabicycle (cf. **II**), we undertook DFT studies (Scheme 3). The detail of the ligand envi-

ronment around rhodium cannot be confirmed, and so the calculations are informative purely from a comparative viewpoint. It is also challenging to compare the relative stabilities of intermediates with different overall charge (neutral vs cationic);¹⁴ accordingly, we rely on the observation that C-H metalation is reversible, such that **A** and **B** are not separated by more than a few kcal mol⁻¹ and zero points in energy can be set for these species (Scheme 3A). Directed C-C bond activation of **A** to form **E** is endergonic ($\Delta G = 7.1$ kcal mol⁻¹) with a barrier of 22.5 kcal mol⁻¹. Migratory insertion of CO proceeds with a lower barrier (12.7 kcal mol⁻¹ vs 15.4 kcal mol⁻¹ for **E** to **A**) to provide rhodacyclopentanone **H** ($\Delta G = -22.1$ kcal mol⁻¹).¹⁵ Equation 4 suggests that a C2-H metallation complex akin to **B** is accessible; however, C-C bond activation from this (via **D**) is discounted due to the higher energy barrier ($\Delta G = 32$ kcal mol⁻¹). Accordingly, we favor a carbonyl directed C-C bond activation pathway for the generation of **H**, from which C2-H metallation (cf. Equation 6) provides **I** (Scheme 3B). Two different sequences could lead to alkyl-Rh(I) species **P**. In one scenario, 5-ring C(sp²)-C(sp³) reductive elimination to **M** is followed by carbometallation of the indole C2-C3 π -system via **O**. Both steps have high barriers (32.4 and 31.9 kcal mol⁻¹). The second option is in accord with Table 1A, and is supported by Scheme 2, Equation 5. Here, 8-ring C(sp²)-C(sp²) reductive elimination to **L** is followed by carbometallation of the indole π -system via **N**. Both steps are accessible with barriers of 17.8 and 23.5 kcal mol⁻¹, respectively. Thus, the analysis supports a sequence involving C(sp²)-C(sp²) reductive elimination from **I**. Additional evidence discounting the alternate C(sp²)-C(sp³) reductive elimination pathway lies in the observation that products arising from decarbonylation of intermediates of type **M** have not been observed.¹⁶ The relative facility of C(sp²)-C(sp²) vs C(sp²)-C(sp³) reductive elimination from Rh(III) has been demonstrated in other studies.¹⁷ It is striking that this reactivity facet overrides the energetic penalty associated with formation of a strained 8-membered ring (**L**).¹⁸ The role of the acid additive (4-NMe₂C₆H₄CO₂H) is likely to facilitate protodemetalation of **P**; alternate or additional roles are possible (e.g. as a carboxylate ligand facilitating CMD-type conversion of **H** to **I**).¹⁹

The mechanistic pathway proposed above suggested that further polycyclizations should be achievable by carbometallation of other π -unsaturated units at the stage of **VII** (see Table 1A). In particular, we envisaged accessing distinct ring systems via *exo*-selective trapping of this intermediate (cf. Scheme 1B, **5** to **6**). To this end alkenyl and alkynyl systems **3r** and **5a** were exposed to carbonylative polycyclization conditions, but in the presence of P(4-FC₆H₄)₃ (15 mol%), an additive that suppresses the dearomatization processes in Table 1 (Table 2A). **3r** afforded solely dearomatization product **4r**, and the product of alkene carbometallation was not observed. Conversely, cyclization of **5a**, which contains a strongly coordinating alkyne, led to **6a** in 26% yield and the corresponding dearomatization product was not observed. **6a** formed as a single geometric isomer, which is consistent with *syn*-stereospecific alkyne carbometallation from **VII**.²⁰ When these processes were conducted without P(4-FC₆H₄)₃ (cf. Table 1) **4r** formed in 42% yield whereas **6a** was not observed. Optimization of **5a** to **6a** was achieved primarily by switching the P-ligand to P(3,5-(CH₃)₂C₆H₃)₃; using this modification **6a** was generated in 55% yield (Table 2B). This protocol extended to a variety of related polycyclizations, which provided indole and pyrrole systems **6b-e** in 39-82% yield. When **5d** to **6d** was run in the presence of D₂O, deuterium incorporation (10%) was observed only at the alkenyl C-H (see the SI).²¹ This is consistent with the second ring forming via a *syn*-carbometallation-protodemetalation sequence from an intermediate akin to **VII**.

The processes above provided the impetus for exploring the scope of the nucleophilic component (Table 2B). 2-Carbamoyl indole, pyrrole and benzofurans are also effective and **6f-i** were generated in 43-67% yield. *Exo*-selective polycyclizations via the C-2

position of heteroarenes with C3 directing groups led to **6j-n**; here, competing cyclization via C4 was not observed. By blocking C-2, pyrrole **5o** could be induced to cyclize via C-4 to provide **6o** in 40% yield.

Table 2. Exo-polycyclizations.

yield. Use of less nucleophilic arenes (e.g. *N*-benzoyl) for the first ring formation was unsuccessful. Rh-catalyzed 8-membered ring formations have been reported previously, but they are not transferable to the polycyclizations described here.²²

Other types of bond formation can be incorporated into these cascades. Directed rhodacyclopentanone formation from **5p** was followed by N-H metallation to generate **VIII** (Table 2C). At this stage, 7-ring C-N reductive elimination occurs prior to alkyne hydrometallation and protodemetalation, which afforded **6p** in 73% yield. This example is unique because reductive elimination (a) generates a C-N rather than a C-C bond, and (b) provides a 7- rather than 8-membered ring. To examine scope further, we subjected 1,3-diene **5q** to polycyclization conditions (Table 2D). Remarkably, this led to 8,7-fused ring system **6q** as a single diastereomer. Here, formation of the first ring likely follows a pathway akin to that outlined earlier. However, the second ring formation is most easily rationalized via a mechanistically distinct hydrometallation pathway, wherein protonation of the alkyl-Rh(I) intermediate provides Rh(III)-hydride **IX**.²³ Hydrometallation of the 1,3-diene generates a Rh- π -allyl which undergoes C-C reductive elimination to **6q**. The intermediacy of Rh- π -allyl species is favored because $\pi \rightarrow \sigma \rightarrow \pi$ isomerization can account for the switch from the *trans*-C3-C4 linkage in **5q** to the *cis*-geometry in **6q**.²⁴

In summary, the latent ambiphilic reactivity of rhodacyclopentanones can be harnessed for the design of polycyclization cascades. The key intermediates are unveiled by carbonylative C-C bond activation of cyclopropanes, an initiating motif that is otherwise stable and can be accessed easily (if required) in a stereocontrolled manner. The method offers high flexibility for the construction of diverse polyheterocycles that are challenging or inaccessible using other methods. We anticipate that reactivity platforms of the type outlined here will be of increasing importance for accessing molecular scaffolds that lie in underexplored regions of chemical space.

ASSOCIATED CONTENT

Supporting Information

Experimental details, characterization data and crystallographic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

john.bower@bris.ac.uk

Notes

The authors declare no competing financial interest

ACKNOWLEDGMENT

We thank the University of Bristol X-ray crystallography service and The Centre for Computational Chemistry, EPSRC and GlaxoSmithKline (studentship to O.B.), the European Research Council (Grant 639594 CatHet), the Royal Society (K. C. Wong Fellowship to G.-W.W. and URF to J.F.B.) and the Leverhulme Trust (Philip Leverhulme Prize to J.F.B.).

REFERENCES

- (1) (a) Taylor, R. D.; MacGoss, M.; Lawson, A. D. G. *Rings in Drugs*. *J. Med. Chem.* **2014**, *57*, 5845. (b) Karawajczyk, A.; Giordanetto, F.; Benninghof, J.; Hamza, D.; Kalliokoski, T.; Pouwer, K.; Morgentin, R.; Nelson, A.; Müller, G.; Piechotod, A.; Tzalis, D. Expansion of Chemical Space for Collaborative Lead Generation and Drug Discovery: the European Lead Factory Perspective. *Drug Discov. Today* **2015**, *20*, 1310. (c) Lovering, F.; Bikker, J.; Humblet, C. Escape

from Flatland: Increasing Saturation as an Approach to Improving Clinical Success. *J. Med. Chem.* **2009**, *52*, 6752.

(2) (a) Shaw, M. H.; Melikhova, E. Y.; Kloer, D. P.; Whittingham, W. G.; Bower, J. F. Directing Group Enhanced Carbonylative Ring Expansions of Amino-Substituted Cyclopropanes: Rhodium-Catalyzed Multicomponent Synthesis of N-Heterobicyclic Enones. *J. Am. Chem. Soc.* **2013**, *135*, 4992. (b) Shaw, M. H.; McCreanor, N. G.; Whittingham, W. G.; Bower, J. F. Reversible C–C Bond Activation Enables Stereocontrol in Rh-Catalyzed Carbonylative Cycloadditions of Aminocyclopropanes. *J. Am. Chem. Soc.* **2015**, *137*, 463. (c) Shaw, M. H.; Croft, R. A.; Whittingham, W. G.; Bower, J. F. Modular Access to Substituted Azocanes via a Rhodium-Catalyzed Cycloaddition–Fragmentation Strategy. *J. Am. Chem. Soc.* **2015**, *137*, 8054. (d) Shaw, M. H.; Whittingham, W. G.; Bower, J. F. Directed Carbonylative (3+1+2) Cycloadditions of Amino-Substituted Cyclopropanes and Alkynes: Reaction Development and Increased Efficiencies Using a Cationic Rhodium System. *Tetrahedron* **2016**, *72*, 2731. (e) Wang, G.-W.; McCreanor, N. G.; Shaw, M. H.; Whittingham, W. G.; Bower, J. F. New Initiation Modes for Directed Carbonylative C–C Bond Activation: Rhodium-Catalyzed (3 + 1 + 2) Cycloadditions of Aminomethylcyclopropanes. *J. Am. Chem. Soc.* **2016**, *138*, 13501. (f) Dalling, A. G.; Yamauchi, T.; McCreanor, N. G.; Cox, L.; Bower, J. F. Carbonylative C–C Bond Activation of Electron-Poor Cyclopropanes: Rhodium-Catalyzed (3+1+2) Cycloadditions of Cyclopropylamides. *Angew. Chem. Int. Ed.* **2019**, *58*, 221.

(3) (a) McCreanor, N. G.; Stanton, S.; Bower, J. F. Capture–Collapse Heterocyclization: 1,3-Diazepanes by C–N Reductive Elimination from Rhodacyclopentanones. *J. Am. Chem. Soc.* **2016**, *138*, 11465. (b) Wang, G.-W.; Bower, J. F. Modular Access to Azepines by Directed Carbonylative C–C Bond Activation of Aminocyclopropanes. *J. Am. Chem. Soc.* **2018**, *140*, 2743. (c) Boyd, O.; Wang, G.-W.; Sokolova, O. O.; Calow, A. D. J.; Bertrand, S. M.; Bower, J. F. Modular Access to Eight-Membered N-Heterocycles by Directed Carbonylative C–C Bond Activation of Aminocyclopropanes. *Angew. Chem. Int. Ed.* **2019**, *58*, 18844.

(4) Review: Dalling, A. G.; Bower, J. F. Synthesis of Nitrogen Heterocycles via Directed Carbonylative C–C Bond Activation of Cyclopropanes. *Chimia* **2018**, *72*, 595. At the present level of development these processes do not extend to cyclobutanes.

(5) See also: (a) Murakami, M.; Tsuruta, T.; Ito, Y. Lactone Formation by Rhodium-Catalyzed C–C Bond Cleavage of Cyclobutanone. *Angew. Chem. Int. Ed.* **2000**, *39*, 2484. (b) Zhang, Y.-L.; Guo, R.-T.; He, J.-H.; Wang, X.-C. Catalytic Intermolecular Coupling of Rhodacyclopentanones with Alcohols Enabled by Dual Directing Strategy. *Org. Lett.* **2019**, *21*, 4239. (c) Shaw, M. H.; Bower, J. F. Synthesis and Applications of Rhodacyclopentanones Derived from C–C Bond Activation. *Chem. Commun.* **2016**, *52*, 10817.

(6) Selected reviews on C–C bond activation methodologies: (a) Soullart, L.; Cramer, N. Catalytic C–C Bond Activations via Oxidative Addition to Transition Metals. *Chem. Rev.* **2015**, *115*, 9410. (b) Fumagalli, G.; Stanton, S.; Bower, J. F. Recent Methodologies that Exploit C–C Single Bond Cleavage of Strained Ring Systems by Transition Metal Complexes. *Chem. Rev.* **2017**, *117*, 9404. (c) Murakami, M.; Ishida, N. Potential of Metal-Catalyzed C–C Single Bond Cleavage for Organic Synthesis. *J. Am. Chem. Soc.* **2016**, *138*, 13759. (d) Kim, D.-S.; Park, W.-J.; Jun, C.-H. Metal–Organic Cooperative Catalysis in C–H and C–C Bond Activation. *Chem. Rev.* **2017**, *117*, 8977. (e) Chen, P.-H.; Billett, B. A.; Tsukamoto, T.; Dong, G. “Cut and Sew” Transformations via Transition-Metal-Catalyzed Carbon–Carbon Bond Activation. *ACS Catal.* **2017**, *7*, 1340.

(7) The major side product for the processes described throughout was protodecyclopropanated starting material ($\text{R}_2\text{Nc-Pr} \rightarrow \text{R}_2\text{NH}$). This likely arises from hydrolysis of the corresponding enamide, which is generated by rhodacyclobutane decomposition (see reference 2a).

(8) (a) Powell, K. G.; McQuillin, F. J. Reactions of Cyclopropanes with Dicarbonylchlororhodium: Carbonyl Insertion and Isomerisation. *J. Chem. Soc. D* **1971**, 931. (b) McQuillin, F. J.; Powell, K. G. Stereoselectivity in the Carbonyl Insertion Reaction between Tetracarbonyldichlororhodium and Substituted Cyclopropanes. *J. Chem. Soc., Dalton Trans.* **1972**, 2129.

(9) (a) Zhuo, C.-X.; Zhang, W.; You, S.-L. Catalytic Asymmetric Dearomatization Reactions. *Angew. Chem. Int. Ed.* **2012**, *51*, 12662. (b) Ding, Q.; Zhou, X.; Fan, R. Recent Advances in Dearomatization of Heteroaromatic Compounds. *Org. Biomol. Chem.* **2014**, *12*, 4807. (c) Roche, S. P.; Tendoung, J.-Y.; Tréguier, B. Advances in Dearomatization Strategies of Indoles. *Tetrahedron* **2015**, *71*, 3549.

(10) Seregin, V.; Gevorgyan, V. Direct Transition Metal-Catalyzed Functionalization of Heteroaromatic Compounds. *Chem. Soc. Rev.* **2007**, *36*, 1173.

(11) (a) Qin, X.; Lee, M. W. Y.; Zhou, J. S. Nickel-Catalyzed Asymmetric Reductive Heck Cyclization of Aryl Halides to Afford Indolines. *Angew. Chem. Int. Ed.* **2017**, *56*, 12723. (b) Shen, C.; Liu, R.-R.; Fan, R.-J.; Li, Y.-L.; Xu, T.-F.; Gao, J.-R.; Jia, Y.-X. Enantioselective Arylative Dearomatization of Indoles via Pd-Catalyzed Intramolecular Reductive Heck Reactions. *J. Am. Chem. Soc.* **2015**, *137*, 4936.

(12) (a) Kimura, M.; Futamata, M.; Mukai, R.; Tamaru, Y. Pd-Catalyzed C3-Selective Allylation of Indoles with Allyl Alcohols Promoted by Triethylborane. *J. Am. Chem. Soc.* **2005**, *127*, 4592. (b) Lian, Y.; Davies, H. M. L. Rhodium-Catalyzed [3 + 2] Annulation of Indoles. *J. Am. Chem. Soc.* **2010**, *132*, 440.

(13) Simmons, E. M.; Hartwig, J. F. On the Interpretation of Deuterium Kinetic Isotope Effects in C–H Bond Functionalizations by Transition-Metal Complexes. *Angew. Chem. Int. Ed.* **2012**, *51*, 3066.

(14) (a) Takano, Y.; Houk, K. N. Benchmarking the Conductor-like Polarizable Continuum Model (CPCM) for Aqueous Solvation Free Energies of Neutral and Ionic Organic Molecules. *J. Chem. Theory Comput.* **2005**, *1*, 70. (b) Mirzaei, S.; Ivanov, M. V.; Timerghazin, Q. K. Improving Performance of the SMD Solvation Model: Bondi Radii Improve Predicted Aqueous Solvation Free Energies of Ions and pKa Values of Thiols. *J. Phys. Chem. A* **2019**, *123*, 9498.

(15) Roundhill, D. M.; Lawson, D. N.; Wilkinson, G. New Complexes Derived from the Interaction of Dicarbonylchlororhodium (I) and Tris(triphenylphosphine) chlororhodium (I) with Cyclopropane, Butadiene, and Perfluorobutadiene. *J. Chem. Soc. A* **1968**, 845.

(16) Doughty, D. H.; Pignolet, L. H. Catalytic Decarbonylation of Aldehydes. *J. Am. Chem. Soc.* **1978**, *100*, 7083.

(17) Gatard, S.; Çlenligil-Çetin, R.; Guo, C.; Foxman, B. M.; Ozerov, O. V. Carbon-Halide Oxidative Addition and Carbon–Carbon Reductive Elimination at a (PNP)Rh Center. *J. Am. Chem. Soc.* **2006**, *128*, 2808.

(18) Galli, C.; Mandolini, L. The Role of Ring Strain on the Ease of Ring Closure of Bifunctional Chain Molecules. *Eur. J. Org. Chem.* **2000**, 3117.

(19) Ackermann, L. Carboxylate-Assisted Transition-Metal-Catalyzed C–H Bond Functionalizations: Mechanism and Scope. *Chem. Rev.* **2011**, *111*, 1315.

(20) Hayashi, T.; Inoue, K.; Taniguchi, N.; Ogasawara, M. Rhodium-Catalyzed Hydroarylation of Alkynes with Arylboronic Acids: 1,4-Shift of Rhodium from 2-Aryl-1-alkenylrhodium to 2-Alkenylaryl rhodium Intermediate. *J. Am. Chem. Soc.* **2001**, *123*, 9918.

(21) Low deuterium incorporation can be explained by 4-NMe₂C₆H₄CO₂H, C2-H of **5d** and adventitious water serving as competing proton sources.

(22) Selected examples: (a) Wender, P. A.; Correa, A. G.; Sato, Y.; Sun, R. Transition Metal-Catalyzed [6+2] Cycloadditions of 2-Vinylcyclobutanones and Alkenes: A New Reaction for the Synthesis of Eight-Membered Rings. *J. Am. Chem. Soc.* **2000**, *122*, 7815. (b) Evans, P. A.; Robinson, J. E.; Baum, E. W.; Fazal, A. N. Intermolecular Transition Metal-Catalyzed [4 + 2 + 2] Cycloaddition Reactions: A New Approach to the Construction of Eight-Membered Rings. *J. Am. Chem. Soc.* **2002**, *124*, 8782. (c) Wender, P. A.; Christy, J. P. Rhodium(I)-Catalyzed [4+2+2] Cycloadditions of 1,3-Dienes, Alkenes, and Alkynes for the Synthesis of Cyclooctadienes. *J. Am. Chem. Soc.* **2006**, *128*, 5354. (d) Yu, R. T.; Friedman, R. K.; Rovis, T. Enantioselective Rhodium-Catalyzed [4+2+2] Cycloaddition of Dienyl Isocyanates for the Synthesis of Bicyclic Azocine Rings. *J. Am. Chem. Soc.* **2009**, *131*, 13250. (e) Lee, D. J.; Ko, D.; Yoo, E. J. Rhodium(II)-Catalyzed Cycloaddition Reactions of Non-Classical 1,5-Dipoles for the Formation of Eight-Membered Heterocycles. *Angew. Chem. Int. Ed.* **2015**, *54*, 13715. (f) Miede, F.; Meyer, C.; Cossy, J. Rhodium-Catalyzed Cycloisomerization Involving Cyclopropanes: Efficient Stereoselective Synthesis of Medium-Sized Heterocyclic Scaffolds. *Angew. Chem. Int. Ed.* **2011**, *50*, 5932.

(23) (a) Yang, X.-H.; Lu, A.; Dong, V. M. Rhodium-Catalyzed Hydrofunctionalization: Enantioselective Coupling of Indolines and 1,3-Dienes. *J. Am. Chem. Soc.* **2017**, *139*, 1774. (b) Yang, X.-H.; Lu, A.; Dong, V. M. Intermolecular Hydroamination of 1,3-Dienes to Generate Homoallylic Amines. *J. Am. Chem. Soc.* **2017**, *139*, 14049.

(24) (a) Lawson, D. N.; Osborn, J. A.; Wilkinson, G. Interaction of Tris(triphenylphosphine)chlororhodium(I) with Iodomethane, Methylallyl, and Allyl Chloride. *J. Chem. Soc. A* **1966**, 1733. (b) Ariafard, A.; Lin, Z. Structure and Bonding of d⁸ Allyl Complexes M(η^3 -allyl)L₃ (M = Co, Rh, Ir; L = Phosphine or Carbonyl). *Organometallics* **2005**, *24*, 3800.