Williams, M. G. L., Liang, K., & Bucciarelli-Ducci, C. (2020). Transient recurrent takotsubo cardiomyopathy mimicking apical hypertrophic cardiomyopathy. *European Heart Journal - Cardiovascular Imaging*, [jeaa274]. <https://doi.org/10.1093/ehjci/jeaa274>

Peer reviewed version

Link to published version (if available):
[10.1093/ehjci/jeaa274](https://doi.org/10.1093/ehjci/jeaa274)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/ehjcmimaging/advance-article/doi/10.1093/ehjci/jeaa274/5957229> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1 Main text

2 A 50-year-old woman presented in 2010 with chest pain with lateral ST-elevation on
3 the electrocardiogram. Invasive coronary angiography revealed non-obstructive coronary
4 disease and she had a mildly raised troponin (342 ng/L). Echocardiography suggested severe
5 LV systolic dysfunction (ejection fraction 35%) with hyperdynamic basal segments and
6 akinesia of the mid to apical segments. Cardiac magnetic resonance (CMR) imaging
7 demonstrated hypokinesia of the mid to apical segments associated with apical hypertrophy
8 (14mm, normal range < 6mm) and diffuse mid to apical ventricular oedema on T2-STIR
9 images (Figure 1 A-B, supplementary video 1). The patient underwent repeat imaging 6
10 months later which revealed complete resolution of the changes in keeping with a diagnosis
11 of resolved takotsubo cardiomyopathy (TCM) (Figure 1 C-D, supplementary video 2). She
12 was reassured and discharged with no medical therapy.

13 The patient represented in 2019 with breathlessness, chest pain and elevated
14 troponin (95 ng/L). Non-obstructive coronary artery disease was again demonstrated on
15 angiography (supplemental video 3-5). The electrocardiogram was normal, and the
16 echocardiogram demonstrated mild apical septal hypokinesia (supplemental video 6). A
17 further CMR demonstrated recurrence of the apical oedema and hypertrophy (11 mm)
18 (Figure 1 E-F, supplemental video 7). Repeat imaging 3 months later again showed LV
19 functional recovery with near complete resolution of the changes (Figure 1 G-H,
20 supplemental video 8) in keeping with recurrent TCM.

21 This case illustrates few important points: 1) recurrent episodes of TCM can occur
22 10 years apart with similar clinical presentation and identical CMR appearances; 2) TCM can
23 mimic apical hypertrophic cardiomyopathy (previously unreported) as the acute myocardial
24 oedema swelling the myocardial wall; however, this is a transient process and, upon

1 resolution of myocardial oedema, the wall thickness reduces returning to normal 3) caution
2 should be used when interpreting thickened myocardial walls in the presence of extensive
3 myocardial oedema, avoiding to overcall hypertrophic cardiomyopathy where macroscopic
4 myocardial oedema is normally not present.

5

6

7

8