Ha, K. P., Clarke, R. S., Kim, G-L., Brittan, J. L., Rowley, J. E., Mavridou, D. A. I., Parker, D., Clarke, T. B., Nobbs, A. H., & Edwards, A. M. (2020). Staphylococcal DNA Repair Is Required for Infection. *mBio*, 11, [e02288-20]. <https://doi.org/10.1128/mBio.02288-20>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1128/mBio.02288-20](https://doi.org/10.1128/mBio.02288-20)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via American Society for Microbiology at <https://mbio.asm.org/content/11/6/e02288-20> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Staphylococcal DNA Repair Is Required for Infection

Kam Pou Ha,^a Rebecca S. Clarke,^a Gyu-Lee Kim,^b Jane L. Brittan,^c Jessica E. Rowley,^a Despoina A. I. Mavridou,^{a,d} Dane Parker,^b Thomas B. Clarke,^a Angela H. Nobbs,^c Andrew M. Edwards^a

^aMRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, United Kingdom

^bDepartment of Pathology, Immunology and Laboratory Medicine, Center for Immunity and Inflammation, Rutgers New Jersey Medical School, Newark, New Jersey, USA

^cBristol Dental School, University of Bristol, Bristol, United Kingdom

^dDepartment of Molecular Biosciences, University of Texas at Austin, Austin, Texas, USA

ABSTRACT To cause infection, *Staphylococcus aureus* must withstand damage caused by host immune defenses. However, the mechanisms by which staphylococcal DNA is damaged and repaired during infection are poorly understood. Using a panel of transposon mutants, we identified the *rexBA* operon as being important for the survival of *Staphylococcus aureus* in whole human blood. Mutants lacking *rexB* were also attenuated for virulence in murine models of both systemic and skin infections. We then demonstrated that RexAB is a member of the AddAB family of helicase/nuclease complexes responsible for initiating the repair of DNA double-strand breaks. Using a fluorescent reporter system, we were able to show that neutrophils cause staphylococcal DNA double-strand breaks through reactive oxygen species (ROS) generated by the respiratory burst, which are repaired by RexAB, leading to the induction of the mutagenic SOS response. We found that RexAB homologues in *Enterococcus faecalis* and *Streptococcus gordonii* also promoted the survival of these pathogens in human blood, suggesting that DNA double-strand break repair is required for Gram-positive bacteria to survive in host tissues. Together, these data demonstrate that DNA is a target of host immune cells, leading to double-strand breaks, and that the repair of this damage by an AddAB-family enzyme enables the survival of Gram-positive pathogens during infection.

IMPORTANCE To cause infection, bacteria must survive attack by the host immune system. For many bacteria, including the major human pathogen *Staphylococcus aureus*, the greatest threat is posed by neutrophils. These immune cells ingest the invading organisms and try to kill them with a cocktail of chemicals that includes reactive oxygen species (ROS). The ability of *S. aureus* to survive this attack is crucial for the progression of infection. However, it was not clear how the ROS damaged *S. aureus* and how the bacterium repaired this damage. In this work, we show that ROS cause breaks in the staphylococcal DNA, which must be repaired by a two-protein complex known as RexAB; otherwise, the bacterium is killed, and it cannot sustain infection. This provides information on the type of damage that neutrophils cause *S. aureus* and the mechanism by which this damage is repaired, enabling infection.

KEYWORDS respiratory burst, oxidative burst, DNA damage, *Enterococcus*, SOS system, *Staphylococcus*, *Streptococcus*, neutrophils

The ability of *Staphylococcus aureus* to maintain the integrity of its DNA in the face of reactive oxygen and nitrogen species produced by host immune defenses is crucial for the establishment of infection. However, despite the importance of DNA repair for staphylococcal survival in the host, little is known about the processes responsible, with most functions based on inferences from work done with the model organisms *Bacillus subtilis* or *Escherichia coli* (1, 2).

Citation Ha KP, Clarke RS, Kim G-L, Brittan JL, Rowley JE, Mavridou DA, Parker D, Clarke TB, Nobbs AH, Edwards AM. 2020. Staphylococcal DNA repair is required for infection. mBio 11:e02288-20. <https://doi.org/10.1128/mBio.02288-20>.

Editor Tarek Msadek, Institut Pasteur

Copyright © 2020 Ha et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Andrew M. Edwards, a.edwards@imperial.ac.uk.

Received 13 August 2020

Accepted 14 October 2020

Published 17 November 2020

This is important because *S. aureus* is responsible for a raft of serious invasive infections, including bacteremia, infective endocarditis, and osteomyelitis (3). Despite a potent immune response, many infections become chronic or recurrent (4), implying either that *S. aureus* does not experience DNA damage during infection or that it has efficient mechanisms for damage repair.

The entry of *S. aureus* into normally sterile tissues triggers the infiltration of neutrophils to control infection (5–7). Neutrophils phagocytose *S. aureus* and expose the bacterium to a cocktail of antimicrobial peptides and proteases (8–10), reactive nitrogen species, and reactive oxygen species (ROS) that are generated by the respiratory burst (also known as the oxidative burst) (11–16). While the contribution of each ROS to bactericidal activity is the subject of investigation, there is compelling evidence that the respiratory burst is crucial for the control of *S. aureus* infection (13, 16, 17). For example, individuals with chronic granulomatous disease (CGD) are particularly prone to staphylococcal infections because their neutrophils are defective for the respiratory burst (6, 18). In keeping with this, *S. aureus* survives better in mice defective for the respiratory burst than in wild-type animals, while the treatment of human neutrophils with an inhibitor of the respiratory burst increased staphylococcal survival relative to untreated immune cells (7–9, 13, 19). However, even when the respiratory burst is functional, there is evidence that some *S. aureus* cells can survive in neutrophils, which contributes to the progression of infection (8, 9).

Despite the importance of the respiratory burst in combating staphylococcal infection, relatively little is known about how it kills the pathogen. Studies with single oxidants such as H_2O_2 indicate that the molecular targets of the respiratory burst are broad and include proteins, lipids, and DNA (20). To survive this damage, *S. aureus* employs several stress response regulators and repair systems. For example, previous work has shown that DNA damage caused by *S. aureus* exposure to H_2O_2 leads to the initiation of the DNA repair SOS response (21), which facilitates the excision of damaged bases or the repair of double-strand breaks (DSBs) by homologous recombination (22). However, H_2O_2 is a suboptimal model for the ROS produced by the respiratory burst because it is typically used at concentrations that exceed those generated by the respiratory burst (23). Therefore, it is unclear whether staphylococcal DNA is damaged by neutrophil-generated ROS, what the nature of this damage is, how it is repaired by *S. aureus*, and the impact of this damage on infection.

To address this gap in our knowledge, we examined mutants defective for DNA repair and found that a member of the AddAB helicase/nuclease family of enzymes was required for staphylococcal survival in blood and murine models of systemic and skin infections. We also demonstrated that this complex is required for the repair of DNA double-strand breaks caused by ROS produced by the respiratory burst of neutrophils, which leads to the induction of the mutagenic SOS response. Similar complexes were required for the survival of the infective endocarditis pathogens *Streptococcus gordonii* and *Enterococcus faecalis* in human blood, demonstrating that DNA damage repair is an important mechanism by which Gram-positive pathogens withstand host defenses.

RESULTS

RexAB is required for staphylococcal survival in host tissues. To determine whether DNA damage occurs under infection conditions, we assembled a panel of *S. aureus* mutants from the ordered Nebraska transposon library produced by the Network on Antimicrobial Resistance in *S. aureus* (NARSA) (24). Each mutant was defective for a different protein associated with DNA repair, which would enable the nature of any damage to be identified. We then assessed the contribution of each repair protein to bacterial survival in the host by measuring CFU counts of mutants in an *ex vivo* whole human blood model of infection. Previous characterizations of this model system by our group and others have shown that *S. aureus* is rapidly phagocytosed by neutrophils and exposed to ROS and other killing mechanisms (25–27).

Wild-type (WT) *S. aureus* JE2 survived relatively well in human blood, with >60% of the inoculum being viable after 2 h (see Fig. S1 in the supplemental material). In

FIG 1 Survival of the *rexB* mutant is significantly reduced in human blood and *in vivo* murine infection models. (A and B) Survival of the *S. aureus* wild type (WT), the *rexB* mutant, the empty vector (pEmpty), and the complemented mutant (*prexBA*) in the SH1000 (A) and JE2 (B) backgrounds after 6 h of incubation in whole human blood. Empty vectors and complemented mutants were supplemented with 100 ng/ml AHT (anhydrotetracycline) to induce *rexBA* expression ($n \geq 3$). (C and D) Numbers of CFU recovered from the peritoneal cavities of mice at 6 h postinfection (each circle represents a single mouse; $n = 4$ to 5). (E) CFU per milliliter recovered from mouse skin biopsy specimens at 5 days postinfection (each circle represents a single mouse; $n = 12$). (F) Sizes of skin lesions observed on mice at up to 5 days postinfection ($n = 12$). Data in panels A and B were analyzed by one-way ANOVA with Tukey's *post hoc* test relative to the WT (*, $P < 0.05$). Data in panels C to F were analyzed by a Mann-Whitney test (*, $P < 0.05$). NS, not significant.

contrast, the survival of mutants defective for *rexA* or *rexB* was $<5\%$ of the inoculum (Fig. S1). Since the transposon insertion in *rexB* contains a terminator sequence that prevents the transcription of *rexA*, the second gene in the operon, *rexB* mutants are defective for both *rexB* and *rexA* and are effectively *rexBA* mutants (24). Therefore, we confirmed the importance of RexAB for the survival of two distinct strains (JE2 and SH1000) in blood by complementation of the *rexB* mutants with a plasmid containing the *rexBA* operon (*prexBA*), which restored bacterial survival to wild-type levels (Fig. 1A and B). In contrast, the survival of *rexB* mutants transformed with the vector alone (pEmpty) was not changed from that of the mutant (Fig. 1A and B).

Having shown that RexAB contributed to staphylococcal tolerance of neutrophil-mediated killing in human blood, we then tested its role in staphylococcal survival *in vivo* using a murine model of systemic infection. Mice were infected via the peritoneal cavity, which results in the recruitment of neutrophils within 2 h, with wild-type or *rexB* mutant strains of *S. aureus* SH1000 or JE2 (28). After 6 h, the mice were sacrificed, and the peritoneal cavity was washed with phosphate-buffered saline (PBS) to recover bacteria, which were quantified by CFU counts. This revealed that *rexB* mutants in both genetic backgrounds were significantly attenuated for survival *in vivo*, with ~5-fold-lower CFU counts than those of the respective wild-type bacteria ($P \leq 0.05$), confirming that RexAB contributes to staphylococcal resistance to host immune defenses and the progression of systemic infection (Fig. 1C and D).

Because *S. aureus* causes many different types of infection, and the associated immune responses might vary, we next assessed the survival of wild-type strain JE2 and the *rexB* mutant in a murine skin infection model. Mice were infected via subcutaneous injection, and infection was allowed to progress for 5 days before CFU counts at inoculation sites were determined. This revealed that wild-type bacteria were present at 2- to 3-fold-higher levels than the *rexB* mutant ($P \leq 0.05$) (Fig. 1E). We also measured the sizes of the skin lesions generated by injected *S. aureus*. Wild-type JE2 caused a lesion that progressively increased in size over time (Fig. 1F). In contrast, the lesion caused by the *rexB* mutant did not increase after day 2 and was significantly smaller than that caused by the wild type for days 2, 3, 4, and 5 (Fig. 1F; Fig. S2). This indicated that RexAB was also required for skin infection progression.

To understand whether the reduced CFU counts of the *rexB* mutants relative to the wild type in animal models were due to differences in growth rates or reduced virulence factor production, we measured bacterial growth, hemolysin production, staphyloxanthin levels, and catalase levels across JE2 and SH1000 wild-type and mutant strains. For both JE2 and SH1000, the *rexB* mutant replicated at a slightly lower rate than the wild type, but there were no significant differences in the production of hemolysin, staphyloxanthin, or catalase (Fig. S3). Taken together, these findings demonstrate that RexAB significantly promotes staphylococcal survival in host tissues and is required for infection progression.

RexAB is a member of the AddAB family of ATP-dependent helicase/nucleases.

The *rexA* and *rexB* genes form a two-gene operon (*rexBA*) on the staphylococcal chromosome and are proposed to encode an AddAB helicase/nuclease enzyme on the basis of sequence homology (29–32). However, this had not been demonstrated experimentally.

Our initial *in silico* structural analysis of the predicted *rexBA* gene products supported predictions that this operon encodes an AddAB-type ATP-dependent helicase/nuclease that contributes to the processing and repair of DNA DSBs (Fig. S4). AddAB enzymes process DSBs to generate a 3' single-stranded DNA (ssDNA) overhang that is necessary for RecA-mediated homologous recombination (33). In support of the structural predictions, phenotypic testing of *rexBA* mutants showed that they were >8-fold more susceptible than wild-type strains to the DNA-damaging antibiotics ciprofloxacin and mitomycin C, both of which cause DNA DSBs (34, 35) (Fig. S5).

To confirm the ATP-dependent helicase/nuclease activity of the *S. aureus* RexAB complex, recombinant RexAB protein was generated, and the helicase and nuclease activities were measured over 1 h. Nuclease activity assays were performed under conditions of high free Mg^{2+} , which has been previously shown to activate nuclease activity in AddAB enzymes (36). We found that DNA was degraded over time by the recombinant complex in the presence of ATP, whereas this degradation was minimal in its absence, demonstrating that RexAB has ATP-dependent nuclease activity (Fig. 2A).

We measured helicase activity under conditions of low free Mg^{2+} levels and observed a loss of double-stranded DNA (dsDNA) concomitant with increased ssDNA formation over time (Fig. 2B and C). These experiments were repeated in the absence of single-stranded DNA binding protein (SSB), which is required to prevent

FIG 2 RexAB possesses ATP-dependent nuclease and helicase activities. (A) Nuclease activity of recombinant *S. aureus* RexAB protein in the presence or absence of ATP ($n = 4$). (B to E) Helicase activity of recombinant *S. aureus* RexAB protein in the presence or absence of ATP, determined by measuring the unwinding of dsDNA and the formation of ssDNA. (B and C) Single-stranded DNA binding protein (SSB) was added to prevent the reannealing of DNA ($n = 4$). (D and E) Equivalent assays were performed without SSB to confirm helicase activity ($n = 4$). Data in panels A to C were analyzed by two-way ANOVA with Sidak's *post hoc* test comparing activities without versus those with ATP (*, $P < 0.05$). Error bars represent standard deviations of the means.

the reannealing of DNA. In the absence of SSB, no DNA unwinding or ssDNA formation was observed, demonstrating that RexAB has ATP-dependent helicase activity (Fig. 2D and E).

Combined, these results confirm that RexAB is a functional member of the AddAB family of DNA repair complexes. In turn, this demonstrates that DNA DSBs occur during staphylococcal infection and must be repaired for bacterial survival.

RexAB enables staphylococcal tolerance of ROS produced by the respiratory burst. Having confirmed that DNA DSBs occur in *S. aureus* during infection, we wanted

FIG 3 RexAB protects *S. aureus* from ROS produced by the respiratory burst of neutrophils and when exposed to H₂O₂. (A and B) Survival of the *S. aureus* WT, the *rexB* mutant, the empty vector (pEmpty), and the complemented mutant (prexBA) in the SH1000 (A) and JE2 (B) backgrounds in whole human blood after 6 h of incubation in the presence of the respiratory burst inhibitor DPI or an identical volume of the DMSO solvent alone. Empty vectors and complemented mutants were supplemented with 100 ng/ml AHT to control *rexB* expression ($n = 3$). (C and D) Survival of *S. aureus* WT and *rexB* mutant strains incubated with purified human neutrophils for 3 h ($n = 4$). (E and F) Survival of the *S. aureus* WT, the *rexB* mutant, the empty vector (pEmpty), and the complemented mutant (prexBA) after 1 h of incubation with 10 mM H₂O₂. Empty vectors and complemented mutants were supplemented with 100 ng/ μ l AHT to induce *rexB* expression ($n = 3$). Data in panels A and B were analyzed by one-way ANOVA with Tukey's *post hoc* test relative to the WT (*, $P < 0.05$). Data in panels C and D were analyzed by Student's *t* test relative to the WT (*, $P < 0.05$). Error bars represent standard deviations of the means. Data in panels E and F were analyzed by one-way ANOVA with Dunnett's *post hoc* test relative to the WT (*, $P < 0.05$).

to determine whether this was due to ROS produced by the respiratory burst of neutrophils. Therefore, we incubated wild-type and *rexB* mutant strains in whole human blood in the presence of diphenyleneiodonium chloride (DPI) to block the NADPH oxidase-generated respiratory burst or dimethyl sulfoxide (DMSO) alone as a solvent control.

As shown in Fig. 1A and B, the survival of *rexB* mutants in whole human blood was significantly reduced relative to that of wild-type bacteria (Fig. 3A and B). However, the presence of DPI promoted the survival of *rexB* mutants to wild-type levels, indicating that the survival deficit observed for bacteria lacking RexAB was due to increased

sensitivity to ROS produced by the respiratory burst (Fig. 3A and B). To ensure that the killing of *rexB* mutants in blood was due to neutrophils, *S. aureus* strains were incubated with purified human neutrophils, and survival was measured via CFU counts. Similar to whole blood, *rexB* mutants were more susceptible to neutrophil-mediated killing than the wild type, but the presence of DPI restored the survival of the *rexB* mutants to wild-type levels (Fig. 3C and D). These data strongly indicated that RexAB contributed to staphylococcal survival of DNA damage caused by ROS produced by the respiratory burst.

To confirm that *rexB* mutants were more susceptible to oxidative damage, we measured the survival of *S. aureus* strains in H_2O_2 , which is one of several different ROS produced in the respiratory burst (6). As observed for whole human blood and purified neutrophils, *rexB* mutants were more susceptible to H_2O_2 than the wild-type or complemented strains (Fig. 3E and F).

Together, these data demonstrate that neutrophils cause DNA DSBs in *S. aureus* via ROS produced by the respiratory burst. This damage must be repaired by RexAB to enable staphylococcal survival in the host.

RexAB is required for induction of the SOS response during exposure to ROS produced by the respiratory burst. The processing of DNA DSBs by AddAB proteins leads to the generation of a 3' overhang. This results in the formation of a RecA filament, which triggers the SOS response, a multicomponent DNA repair mechanism that mediates the repair of the DNA DSB (1, 22). However, the induction of the SOS response also leads to a transient increase in the mutation rate, which promotes the emergence of mutants with resistance to antibiotics or host-adapted phenotypes such as small-colony variants (2, 37).

Therefore, we tested whether ROS produced by the respiratory burst triggers the SOS response and whether this was dependent upon RexAB. To do this, we used a *PrecA-gfp* reporter construct and validated it by showing dose-dependent activity with mitomycin C, a well-established trigger of the SOS response (38) (Fig. S6). We then incubated tetramethyl rhodamine isothiocyanate (TRITC)-labeled *S. aureus* JE2 wild type and *rexB* mutant strains containing the reporter with neutrophils for 30 min and used flow cytometry to measure phagocytosis and reporter activity (green fluorescent protein [GFP] fluorescence) (Fig. S7) (39). As reported previously, >95% of *S. aureus* cells were associated with neutrophils within 30 min (Fig. 4A) (26). Also, by 30 min, there was an increase in the GFP signal from wild-type *S. aureus* relative to the start of the assay (Fig. 4B). In contrast, there was no increase in the GFP signal from the *rexB* mutant (Fig. 4B), indicating that neutrophils trigger the SOS response in *S. aureus* via DNA processing by RexAB.

To further explore the requirement for RexAB for the induction of the SOS response during exposure to oxidative stress, wild-type and *rexB* mutant bacteria were incubated with various subinhibitory doses of paraquat, which results in the generation of endogenous superoxide, which dismutates to H_2O_2 . For wild-type bacteria, there was a clear dose-dependent increase in GFP-mediated fluorescence, indicative of SOS induction (Fig. 4C). For the *rexB* mutant, while there also appeared to be a dose-dependent induction of the SOS response, it was at considerably lower levels than those seen for the wild type (Fig. 4C). Therefore, the induction of SOS in response to oxidative stress is almost entirely dependent upon RexAB-mediated processing of DNA DSBs.

RexAB is required for survival of streptococci and enterococci in human blood.

Since homologues of RexAB are present in most Gram-positive bacteria (31–33), we next tested whether this repair complex contributes to the survival of other bacteria exposed to neutrophils. Like *S. aureus*, *Enterococcus faecalis* and *Streptococcus gordonii* are frequent causes of infective endocarditis, which brings these species into close contact with neutrophils (40, 41). Therefore, *rexBA* was deleted in representative strains of each species, and their sensitivity to the microbicidal activity of neutrophils was determined using the *ex vivo* whole human blood model. To confirm that RexAB mediates the repair of DNA DSBs in both bacteria, we also assessed their susceptibility to the antibiotics ciprofloxacin, which causes DNA DSBs (42), and gentamicin, which

FIG 4 RexAB is required for induction of the SOS response during exposure to ROS produced by the respiratory burst of neutrophils and when exposed to the ROS generator paraquat. (A and B) Phagocytosis (both ingested and surface-bound bacteria [39]) of JE2 wild-type and *rexB* mutant *recA-gfp* reporter strains by neutrophils (A) and the resulting induction of the SOS response measured by GFP expression (RFU) after 30 min of exposure ($n = 4$) (B). (C) Induction of the SOS response of JE2 wild-type and *rexB* mutant strains as measured by GFP expression upon exposure to sublethal concentrations of paraquat. GFP fluorescence was normalized by the OD₆₀₀ to determine the induction of SOS relative to the cell density ($n = 4$). OD₆₀₀ measurements alone are shown in Fig. S5 in the supplemental material. Data in panels A and B were analyzed by a Mann-Whitney test (*, $P < 0.05$). Where shown, error bars represent standard deviations of the means. Error bars were omitted from panels C and D for clarity.

targets protein synthesis and thus acted as a negative control (43). For both *S. gordonii* and *E. faecalis*, the Δ *rexBA* mutants were significantly more susceptible to ciprofloxacin than the wild type, confirming that RexAB in these bacteria contributes to DNA DSB repair (Table S1). In contrast, both the wild type and the Δ *rexBA* mutants were equally susceptible to the antibiotic gentamicin (Table S1).

Wild-type *E. faecalis* survived at high levels in human blood, with ~100% of the inoculum remaining viable during the full 6-h duration of the assay, but the loss of *rexB* reduced enterococcal survival by ~50% (Fig. 5A). This indicated that *E. faecalis* suffers DNA damage that results in DSBs while in blood, but it can be tolerated via DNA repair (Fig. 5A). However, in contrast to *S. aureus*, DNA damage in *E. faecalis* was not due to ROS produced by the respiratory burst since the survival of both wild-type and *rexB* mutant bacteria in blood was unaffected by the presence of DPI (Fig. 5A).

S. gordonii was much more susceptible to host defenses in blood than *E. faecalis*, with ~5% of wild-type bacteria remaining viable after 6 h (Fig. 5A). However, the survival of the *rexB* mutant was still reduced relative to the wild type, with ~1% of streptococci surviving after 6 h, indicating that DNA DSB repair also contributes to the survival of *S. gordonii* in blood (Fig. 5A). As for *E. faecalis*, the presence of DPI did not increase the survival of *S. gordonii* in blood, indicating that DNA damage was not due to ROS produced by the respiratory burst in either bacterium. In support of these findings, the *rexB* mutants of both *S. gordonii* and *E. faecalis* were no more susceptible to H₂O₂ than wild-type bacteria (Fig. 5B).

FIG 5 RexAB protects *Enterococcus faecalis* and *Streptococcus gordonii* from host defenses in blood. (A) Survival of *E. faecalis* and *S. gordonii* wild-type and $\Delta rexBA$ mutant strains in whole human blood after 6 h in the presence of the respiratory burst inhibitor DPI or an identical volume of the DMSO solvent alone ($n = 3$). (B) Survival of *E. faecalis* and *S. gordonii* wild-type and $\Delta rexBA$ mutant strains exposed to H_2O_2 for 1 h ($n = 3$). Data in panels A and B were analyzed by Student's *t* test comparing the $\Delta rexBA$ mutant versus the WT (*, $P < 0.05$). Error bars represent standard deviations of the means.

Therefore, as for *S. aureus*, neutrophils damage DNA of both *S. gordonii* and *E. faecalis*, the repair of which by RexAB promotes bacterial survival. However, in contrast to *S. aureus*, ROS generated by the neutrophil respiratory burst do not appear to contribute to DNA damage in these bacteria.

DISCUSSION

Neutrophils are an essential host defense against *S. aureus* and many other bacterial pathogens. However, our understanding of the mechanisms by which these immune cells kill staphylococci is limited. The data presented here demonstrate that neutrophils cause DNA damage in *S. aureus* via ROS produced by the respiratory burst, the repair of which requires the RexAB complex and leads to the induction of the SOS response. We also confirmed that RexAB is a member of the AddAB helicase/nuclease family of enzymes involved in the processing of DNA DSBs for repair via homologous recombination. Thus, ROS generated by neutrophils cause DNA DSBs in *S. aureus*, which are lethal if not repaired. This provides new information on both the target of the ROS generated by the respiratory burst and the mechanisms by which *S. aureus* repairs and survives this damage.

The importance of RexAB for staphylococcal survival during infection was demonstrated in murine models of both systemic and skin infections as well as an *ex vivo* whole human blood model of bacteremia. Since *S. aureus* is a frequent cause of both superficial and systemic infections (3), these findings confirm the importance of DNA repair for staphylococcal survival in relevant but distinct host tissues.

Previous work demonstrated the requirement of DNA DSB repair for the survival of Gram-negative pathogens *in vivo*. For example, AddAB was shown to be required

for the infection of chickens and mice by *Campylobacter jejuni* and *Helicobacter pylori*, respectively (44, 45). Furthermore, the virulence of *Salmonella enterica* in a murine model of bacteremia was dependent upon the RecBCD DNA DSB repair complex (46). However, the importance of such systems for Gram-positive bacteria was unclear.

DNA damage occurred within 30 min of phagocytosis of *S. aureus* by neutrophils, which corresponds to the time at which ROS are maximally generated in these immune cells (47, 48). The finding that the respiratory burst leads to DNA DSBs in *S. aureus* is in keeping with previous reports that ROS damage the DNA of Gram-negative pathogens such as *Escherichia coli*, *Salmonella enterica*, and *Coxiella burnetii*, with the survival of bacteria phagocytosed by macrophages being dependent upon RecBCD (46, 49, 50). Therefore, DNA is a common target of ROS produced by phagocytic immune cells for several different human pathogens.

The findings from this work build on previous studies of oxidative DNA damage and repair and enable us to understand the sequence of events that occur during the exposure of *S. aureus* to ROS of the respiratory burst. Of the ROS produced by neutrophils, only H_2O_2 can cross the membrane due to its lack of charge (14, 15, 23). In the cytoplasm, H_2O_2 reacts with iron in a process known as the Fenton reaction, which leads to the generation of highly reactive hydroxyl radicals (14, 15, 20, 23). These can damage DNA as well as the pool of nucleotides, leading to various types of lesions (14, 15, 20, 23). Based on our findings, physiological concentrations of ROS produced by neutrophils lead to DNA DSBs in *S. aureus* despite the numerous antioxidant defenses of this pathogen (21, 25, 27). While some bacteria can engage in nonhomologous end joining, most DSBs are repaired via homologous recombination (31, 33, 36). RexAB processes the broken ends to produce single-stranded DNA to which RecA binds (1, 31, 33, 36). The resulting RecA nucleoprotein filament triggers the SOS response by initiating the autocleavage of the LexA transcriptional repressor (1, 22, 51). As shown here, DNA damage processed by RexAB results in the expression of *recA*, leading to homologous recombination and survival of the bacterium. The SOS response also leads to the expression of the low-fidelity DNA polymerase UmuC, which leads to a transient increase in the mutation rate (21, 22). Previous work from our group has shown that mutagenesis due to SOS induction as a consequence of oxidative stress leads to the acquisition of mutations conferring antibiotic resistance and the small-colony phenotype associated with chronic infection and resistance to neutrophil-mediated killing (26, 37). Therefore, the processing of DNA DSBs by RexAB not only promotes the survival of bacteria exposed to ROS produced by the respiratory burst but also may promote the emergence of mutants that are more resistant to neutrophil-mediated killing by triggering the SOS response. However, this remains to be tested.

In addition to *S. aureus*, the RexAB system was also demonstrated to be important for the survival of the infective endocarditis pathogens *E. faecalis* and *S. gordonii*. However, while neutrophils in blood caused DNA DSBs in these pathogens, this did not appear to be due to ROS because the inhibition of NADPH oxidase with DPI had no effect on the survival of the wild type or the *rexBA* mutants. Neutrophils employ several different antibacterial elements to kill invading pathogens, including reactive oxygen and nitrogen species, proteases, and antimicrobial peptides. Several studies have indicated that bacteria vary in their susceptibility to each of these microbicides. In agreement with our findings, Standish and Weiser showed that *S. aureus* but not *Streptococcus pneumoniae* was killed by ROS (52). However, while *S. aureus* can grow in the presence of nitric oxide, the replication of several other pathogens, including *Pseudomonas aeruginosa* and *Streptococcus pyogenes*, is inhibited (53). Since reactive nitrogen species can cause DNA damage, this may provide an explanation for the finding that mutants of *E. faecalis* and *S. gordonii* lacking RexAB are more susceptible to killing by neutrophils in which the respiratory burst is blocked.

While the consequences of RexAB-mediated processing of DNA DSBs are relatively predictable for *S. aureus*, they are less so for *E. faecalis* and *S. gordonii*. This is particularly the case for *Streptococcus* since this genus lacks the LexA repressor that is central to the

control of the SOS response (54–56). However, *S. gordonii* encodes RecA, which promotes survival during exposure to UV light, and there is evidence that the role of LexA is fulfilled by HdiR in at least some streptococci and *Lactococcus lactis* (56–59). For example, *Streptococcus uberis* has been found to encode a UmuC error-prone DNA polymerase that appears to be regulated by HdiR and induced in response to UV-mediated DNA damage, so it is possible that RexAB-mediated processing of DNA DSBs leads to the induction of an SOS-like response in *S. gordonii*, including homologous recombination and mutagenic DNA repair (56, 59).

In contrast to *S. gordonii*, *E. faecalis* encodes both RecA and LexA and appears to have a DNA damage-inducible UmuC polymerase, suggesting a SOS response similar to that described above for *S. aureus* (60–63). However, as for *S. aureus*, it remains to be seen whether neutrophil-mediated DNA damage leads to an increase in the mutation rate in either *S. gordonii* or *E. faecalis*.

In addition to providing protection from oxidative damage caused by neutrophils, we have previously shown that RexAB provides *S. aureus* with tolerance to the combination antibiotic co-trimoxazole (38). That work showed that DNA damage was partly due to endogenous oxidative stress that occurred during exposure to the combination antibiotic, in addition to direct damage to DNA caused by thymidine limitation (38). The data presented here show that the loss of RexAB also sensitizes *S. aureus* to ciprofloxacin, even though the JE2 strain is resistant. In addition, there is growing evidence that multiple classes of antibiotics cause endogenous ROS production in *S. aureus*, suggesting that RexAB may provide an important defense against damage caused by both host defenses and multiple classes of antibiotics, the two key threats to staphylococcal survival in the host. Further studies are needed to test this, but the identification of RexAB as being important for staphylococcal survival during exposure to both neutrophils and at least two antibiotics makes this complex a potential target for novel therapeutics, particularly as the lack of RexAB homologues in eukaryotes reduces the likelihood of host toxicity (64).

Inhibitors of AddAB and RecBCD have been previously reported in the literature, but problems include limited *in vivo* stability, poor oral bioavailability, and a suboptimal mechanism of action (65–68). However, Amundsen et al. identified several small-molecule inhibitors of *Helicobacter pylori* AddAB and *E. coli* RecBCD, in particular ML328 (69), indicating that the development of stable, potent inhibitors is possible. More recently, a derivative of ML328 (IMP-1700) was found to be capable of potentiating antibiotic activity such that a resistant *S. aureus* strain was sensitized to ciprofloxacin (70). However, the ability of IMP-1700 to inhibit AddAB/RecBCD activity under *in vivo* conditions remains to be determined. Further work in this area may lead to broad-spectrum therapeutics that promote bacterial susceptibility to both host defenses and antibiotics as well as inhibiting the induction of the mutagenic SOS response associated with the acquisition of drug resistance and host adaptation (37).

In summary, staphylococcal, streptococcal, and enterococcal DNA is damaged by the host immune system, leading to DNA DSBs that are lethal if not repaired by RexAB. These findings suggest that the RexAB complex is a potentially viable target for novel therapeutics, capable of sensitizing Gram-positive pathogens to neutrophil-mediated killing and blocking the SOS response associated with the emergence of drug resistance.

MATERIALS AND METHODS

Bacterial strains and culture conditions. The bacterial strains used in this study are listed in Table 1. *S. aureus* was cultured in tryptic soy broth (TSB) to stationary phase (18 h) at 37°C with shaking (180 rpm). *S. gordonii* and *E. faecalis* were grown in Todd-Hewitt broth (THB) supplemented with 1% (wt/vol) yeast extract (THB-Y) at 37°C, statically in 5% CO₂. *E. coli* was grown in lysogeny broth (LB) or Terrific broth (TB) (1.2% [wt/vol] tryptone, 2.4% [wt/vol] yeast extract, 0.5% glycerol, 0.17 M KH₂PO₄, 0.72 M K₂HPO₄) for protein expression at 37°C with shaking (180 rpm). Media were supplemented with antibiotics as required. When appropriate, bacteria were grown on Columbia blood agar (CBA) made with 5% defibrinated sheep blood.

PrecA-gfp fluorescent reporter assay. As detailed previously (38), promoter-reporter gene constructs in the JE2 background were used to directly assess the expression of *recA*. Antibiotic 2-fold

TABLE 1 Bacterial strains used in this study

Strain	Description ^a	Reference or source
<i>Staphylococcus aureus</i>		
SH1000	<i>rsbU</i> ⁺ derivative of laboratory strain 8325-4	80
SH1000 <i>rexB</i> ::Tn	SH1000 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> ; Ery ^r	38
SH1000 <i>rexB</i> ::Tn <i>pitet</i> empty	SH1000 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> with the integrated <i>pitet</i> empty plasmid; Ery ^r	38
SH1000 <i>rexB</i> ::Tn <i>pitet rexAB</i>	SH1000 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> with integrated <i>pitet</i> with AHT-inducible <i>rexB</i> ; Ery ^r	38
JE2	Derivative of CA-MRSA USA300 LAC, cured of plasmids	24
JE2 <i>rexB</i> ::Tn	JE2 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> ; Ery ^r	24
JE2 <i>rexB</i> ::Tn <i>pitet</i> empty	JE2 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> with the integrated <i>pitet</i> empty plasmid; Ery ^r	38
JE2 <i>rexB</i> ::Tn <i>pitet rexAB</i>	JE2 with a <i>bursa aurealis</i> transposon insertion in <i>rexB</i> with integrated <i>pitet</i> with AHT-inducible <i>rexB</i> ; Ery ^r	38
JE2 pCN34 <i>PrecA-gfp</i>	JE2 containing pCN34 with <i>gfp</i> under the control of the <i>recA</i> promoter; Kan ^r	38
JE2 <i>rexB</i> ::Tn pCN34 <i>PrecA-gfp</i>	JE2 <i>rexB</i> ::Tn containing pCN34 with <i>gfp</i> under the control of the <i>recA</i> promoter; Kan ^r	38
<i>Streptococcus gordonii</i>		
DL1 (Challis)	Wild type	81
DL1 Δ <i>rexBA</i>	DL1 with the <i>rexA</i> and <i>rexB</i> genes deleted	This study
<i>Enterococcus faecalis</i>		
CK111(pCF10-101)	Conjugative donor strain with pCF10-101	76
OG1X	Gelatinase deficient	82
OG1X Δ <i>rexBA</i>	OG1X with the <i>rexA</i> and <i>rexB</i> genes deleted	This study
<i>Escherichia coli</i>		
EC1000	Cloning host that provides RepA in <i>trans</i>	83
SoluBL21(DE3)	Derivative of BL21(DE3) for the expression of challenging proteins	Genlantis

^aCA-MRSA, community-acquired methicillin-resistant *S. aureus*.

dilutions were made in flat-bottomed black-walled 96-well plates containing TSB and kanamycin (90 μ g ml⁻¹) and inoculated with a 1/10 dilution of a stationary-phase culture of the reporter strains. Plates were placed into an Infinite M200-Pro microplate reader (Tecan), where cultures were grown for 17 h at 37°C (700 rpm), and both the absorbance at 600 nm (optical density at 600 nm [OD₆₀₀]) and GFP relative fluorescence units (RFU) were measured every 30 min.

OD₆₀₀ data and RFU data were normalized to values for the no-antibiotic controls. To account for differences in cell density, RFU values were normalized by the OD₆₀₀ data at each time point.

Neutrophil phagocytosis and measurement of DNA damage. Whole human blood (15 ml) was collected from individual healthy donors in EDTA-treated tubes (BD Biosciences) and layered over 20 ml of room-temperature Polymorph prep (Alere Limited) before centrifugation at 500 $\times$ g for 45 to 60 min (brake off, 30°C) until a clear separation of red blood cells (RBCs), peripheral blood mononuclear cells (PBMCs), and polymorphonuclear leukocytes (PMNs) (or neutrophils) was seen. The PBMCs were discarded, and the PMNs were transferred to a fresh centrifuge tube. Hanks' balanced salt solution (HBSS) was added to the PMNs to a total volume of 50 ml, and cells were pelleted at 500 $\times$ g for 10 min (brake off, 30°C). The cells were resuspended in 3 ml of HBSS, counted using a hemocytometer, and adjusted to 5 $\times$ 10⁶ cells ml⁻¹ in HBSS containing 10% human serum, 0.1 mM calcium, and 0.1 mM magnesium. To the neutrophil suspension, 5 $\times$ 10⁶ cells ml⁻¹ of bacteria (stationary- or exponential-phase TRITC-stained bacteria) were added. The bacterium-neutrophil suspension was then incubated at 37°C with tumbling. At each time point (0.5, 1, 2, and 3 h), 100 to 150 μ l was taken and resuspended in 4% paraformaldehyde (PFA) in PBS for a minimum of 1 h. Before analysis by flow cytometry, samples were washed and resuspended in PBS. Samples were analyzed on a FACSAria or LSRFortessa flow cytometer (BD Biosciences), and at least 100,000 events were captured, except for bacterium-only samples, where at least 50,000 events were captured. Green fluorescence (GFP positive bacteria) was detected at 530 (30) nm, and TRITC labeling was detected at 586 nm (16) nm. Full gating strategies are detailed in Fig. S6 and Fig. S7 in the supplemental material. Measurements of phagocytosis include both ingested bacteria and those bound to the surface of the neutrophil, based on that of Surewaard et al. (39).

Determination of MICs. MICs were determined using a serial broth dilution protocol as described previously (71). Bacteria were diluted to 1 $\times$ 10⁵ CFU ml⁻¹ and incubated in flat-bottomed 96-well plates with a range of antibiotic concentrations for 17 h at 37°C under static conditions (aerobic, anaerobic, or 5% CO₂). The MIC was defined as the lowest concentration at which no growth was observed.

Whole-blood and hydrogen peroxide survival assays. Bacteria were washed twice with PBS and adjusted to 10⁶ CFU ml⁻¹ in HBSS, and 10⁴ CFU (10 μ l) were used to inoculate 90 μ l of freshly donated human blood (collected in EDTA-treated tubes; BD Biosciences) or freshly diluted H₂O₂ (10 mM in PBS) in 96-well plates. Ethical approval for drawing and using human blood was obtained from the Regional Ethics Committee and the Imperial NHS Trust Tissue Bank (REC Wales approval no. 12/WA/0196 and

TABLE 2 Primers used in this study

Oligonucleotide	Sequence (5'–3') ^a
rexB-F BamHI	CCAGGATCCGATGACATTACATGCTTATTTAGG
rexA-R Sall	GCGGTGACCTATAGTTGCAATGTACC
StreptII rexAB SDM-F	ACAATTCCAGAGAAACCACAAGGCGTGATTGGACTGACGCGCAATGGC
StreptII rexAB SDM-R	AGCGCTTTTTTCGAAGTGGGGTGGCTCCACATCTATTGCTACACCCC
Thr rexAB SDM-F	CCCCGAGTTCGAAAAAAGCGCTGGCCTGGTGCCGCGCGGCAGCGGCACAATTCCAGAGAAACCACAAGG
Thr rexAB SDM-R	CCTGTGGTTTCTCTGGAATTGTGCCGCTGCCGCGCGGCACCAGGCCAGCGCTTTTTTCGAAGTGGGG
Sg.rexAB.F1	CAGCAGGACAAGGGAAGTG
Sg.rexAB.R1	TTTTGTTCATAGCCTGCCTTTCTCTAG
Sg.rexAB.F2	AGGTCCCTAGAAGGAAGCATCTGAGTTG
Sg.rexAB.R2	CGTCGAGCACTAGTCTCG
ermAM.SgF	AAGGCAGGCTATGAACAAAAATATAAAATATTCTCA
ermAM.SgR	ATGCTTCCTTCTAGGGACCTCTTTAGCTCC
Ef.rexAB.pheF1	TGACGTCGACGCGTCTGACGAGTACGCTAAAACGCTAGAAGC
Ef.rexAB.pheR1	TAAATCTACTGACACTCATGGTTTCAC
Ef.rexAB.pheF2	GAGTGTGCGAGTAGATTTTATAAAGTAGAAAAAATTAGAAG
Ef.rexAB.pheR2	GCTTAGCATGCCATGGTCTTAATACTTCGGTGATTGG

^aUnderlined regions indicate base pair overlaps for stitch PCR, and italicized regions indicate restriction endonuclease sites for ligation into pCJ47.

ICHTB HTA license no. 12275). In some assays, blood was pretreated for 10 min with diphenyleneiodonium (DPI) (50 μ M) or an equivalent volume of DMSO as a solvent control. After 6 h of incubation, bacterial survival was determined by CFU counts in blood-bacterium mixtures on CBA plates. For H₂O₂ assays, survival was measured after 1 h at 37°C (static) in the dark. Survival for both assays was calculated as a percentage of the number of bacteria in the starting inoculum.

Neutrophil survival assay. Neutrophils were adjusted to 5×10^6 cells ml⁻¹ in HBSS containing 10% human serum, 0.1 mM calcium, and 0.1 mM magnesium. Stationary-phase bacterial cultures were washed in PBS, and 1×10^6 CFU were added to the neutrophil suspension (multiplicity of infection [MOI] of 1:5) to a total volume of 1 ml. Neutrophils were treated for 10 min prior to the addition of bacteria with either DPI (50 μ M) or an equivalent volume of DMSO (solvent control), as needed. The bacterium-neutrophil suspension was subsequently incubated at 37°C with tumbling. At relevant time points (0.5, 1, 2, and 3 h), 50 μ l of the suspension was transferred to a 96-well plate and serially diluted 10-fold in PBS up to a 10^{-3} dilution. All dilutions (including neat) were then plated onto CBA and incubated for 24 h at 37°C before counting. Survival was calculated as a percentage of the number of bacteria in the starting inoculum.

Murine systemic infection model. Animal work was conducted in accordance with the Animals (Scientific Procedures) Act 1986 outlined by United Kingdom Home Office regulations. Work was approved by the United Kingdom Home Office after ethical approval by the Imperial College Animal Welfare and Ethical Review Body (AWERB). Six- to eight-week-old female C57BL/6 mice (Charles River) were infected via the intraperitoneal route with the wild type or the *rexB*:Tn mutant in the JE2 and SH1000 backgrounds. Stationary-phase bacterial cultures were washed twice with PBS and adjusted to 10^7 CFU ml⁻¹. Subsequently, 400 μ l (4×10^6 CFU) of the washed bacterial suspensions was injected into the peritoneal cavity of each mouse (5 mice for each strain; 20 in total). After 6 h, the mice were humanely sacrificed by cervical dislocation, and death was confirmed by severing the femoral artery. The peritoneal cavity was washed with PBS to release the bacteria, and CFU counts were determined by plating onto tryptic soy agar (TSA). Sample size was determined prior to the experiment using power analysis based on *in vitro* data (72). Tubes containing the bacterial suspensions were blinded before the start of the experiment. Mice were randomly allocated to group cages, and each group was randomly allocated to a treatment. According to Home Office regulations, any animals that displayed two or more of the following symptoms were humanely killed using a schedule 1 method and excluded from the study: shivering, hunched posture, reduced movement, cyanosis, circling, or difficulty breathing.

Murine skin infection model. Six- to eight-week-old mice were subcutaneously infected with 2×10^6 CFU of exponential-phase cultures of *S. aureus* as previously described (73). Skin infection animal work was performed according to the *Guide for the Care and Use of Laboratory Animals* of the National Institutes of Health (74), the Animal Welfare Act, and U.S. Federal law. The protocol was approved by the Animal Care and Use Committee of Rutgers New Jersey Medical School.

Construction of Δ rexBA mutants of *E. faecalis* and *S. gordonii*. A Δ rexBA mutant was generated in *S. gordonii* by in-frame allelic replacement with the erythromycin resistance determinant *ermAM*. Flanking regions directly upstream and downstream of the *rexBA* operon were amplified by PCR from *S. gordonii* DL1 genomic DNA with primer pairs Sg.rexAB.F1/Sg.rexAB.R1 and Sg.rexAB.F2/Sg.rexAB.R2 (Table 2), respectively, while the *ermAM* cassette was amplified from plasmid pVA838 (75) using primer pair *ermAM*.SgF/*ermAM*.SgR (Table 2). The resulting amplicons were then joined via 20-bp overlapping regions by stitch PCR using primers Sg.rexAB.F1/Sg.rexAB.R2 and transformed into *S. gordonii*. Erythromycin-resistant transformants were confirmed by sequencing, and the strain was designated UB3018.

A *rexBA* mutant was generated in *E. faecalis* by markerless exchange using a 2-step homologous recombination approach, as previously described (76). In brief, flanking regions directly upstream and downstream of the *rexBA* operon were amplified by PCR from *E. faecalis* OG1X genomic DNA with primer pairs Ef.rexAB.pheF1/Ef.rexAB.pheR1 and Ef.rexAB.pheF2/Ef.rexAB.pheR2 (Table 2), respectively, and then

joined via 20-bp overlapping regions by stitch PCR using primer pair Ef.rxAAB.pheF1/Ef.rxAAB.pheR2. The resultant amplicon was cloned into donor plasmid pCJ47 (76) via the unique restriction sites PstI and NcoI to generate pCJ47-rxAAB, propagated in *E. coli* EC1000, and then introduced into conjugative donor strain *E. faecalis* CK111(pCF10-101) by electroporation. Plasmid pCJ47-rxAAB was transferred to *E. faecalis* OG1X by conjugation. Transconjugants carrying the integrated plasmid were confirmed by colony PCR, before counterselection based on the *P-pheS** marker was used to identify secondary recombinants in which the integrated plasmid had been excised and lost, leaving the desired Δ rxAAB allele. This was confirmed by sequencing, and the strain was designated UB2948.

Construction of an *S. aureus* RexAB expression vector. Cloning of the *rexA* and *rexB* genes from *S. aureus* was achieved by PCR from wild-type genomic JE2 DNA using the *rexB*-F BamHI and *rexA*-R Sall primers listed in Table 2, which allowed the amplification of the *rexBA* operon immediately flanked by suitable restriction endonuclease recognition sequences (BamHI and Sall). The *rexA* and *rexB* genes were inserted into the pET28b⁺ vector (Novagen) using standard cloning techniques, and site-directed mutagenesis (SDM) was performed to insert Strep-tag II and a thrombin site in front of the *rexA* gene. This enabled RexA and RexB proteins to be detected individually via an N-terminal His₆ tag for RexB (His₆ from the pET28b⁺ vector) and N-terminal Strep-tag II for RexA. PCR primers for SDM are listed in Table 2. DNA sequencing was used on the pET28b⁺ *rexBA* expression plasmid to confirm that the sequences of the entire *rexA* and *rexB* genes, tags, and promoter regions were as expected.

Expression and purification of recombinant *S. aureus* RexAB. Cells from single colonies of *E. coli* SoluBL21(DE3), freshly transformed with the pET28b⁺ *rexAB* expression plasmid coding for N-terminally Strep-tag II-tagged RexA and N-terminally His₆-tagged RexB, were used to inoculate a starter culture grown overnight in LB supplemented with 50 μ g ml⁻¹ of kanamycin. The starter culture was diluted to an OD₆₀₀ of 0.05 into 4 liters of TB containing 50 μ g ml⁻¹ of kanamycin. Cells were grown at 37°C with shaking at 180 rpm until an OD₆₀₀ of 0.5 was reached, prior to induction with 1 mM isopropyl- β -D-thiogalactoside (IPTG). Following induction, the temperature was reduced to 20°C, and cultures were further incubated for 20 h. Cells were harvested by centrifugation at >10,000 $\times$ g at 4°C for 30 min, and pellets were resuspended in 100 ml of a solution containing 50 mM Tris and 150 mM NaCl (pH 7.5).

For protein purification, cells were disrupted by sonication, and cell debris was cleared by centrifugation at 32,000 $\times$ g at 4°C for 30 min. The resulting supernatant was added to 5 ml of Chelating Sepharose Fast Flow resin (GE Healthcare), which had been loaded with 0.1 M NiCl₂ and equilibrated with 100 ml of a solution containing 50 mM Tris, 150 mM NaCl, and 20 mM imidazole (pH 7.5). The supernatant-resin mixture was left at 4°C overnight with gentle stirring to optimize the binding of the His-tagged protein to the nickel-charged resin. Next, the mixture was washed nine times with 40 ml of a solution containing 50 mM Tris, 150 mM NaCl, and 20 mM imidazole (pH 7.5) and once with 20 ml of a solution containing 50 mM Tris, 150 mM NaCl, and 70 mM imidazole (pH 7.5). His-tagged protein was eluted with a solution containing 50 mM Tris, 150 mM NaCl, and 150 mM imidazole (pH 7.5); each 10-ml fraction was tested with Bradford reagent (Bio-Rad) for protein content until no more protein could be detected. Fractions containing protein were pooled, buffer exchanged, and concentrated using an Amicon 100-kDa-cutoff concentrator (Merck Millipore); RexAB is >250 kDa. The concentrating device was centrifuged at 2,000 $\times$ g (4°C), and the protein solution was exchanged into a solution containing 50 mM Tris and 150 mM NaCl (pH 7.5) by four serial concentration and redilution steps. The total protein concentration was quantified using the Pierce bicinchoninic acid (BCA) protein assay kit (Thermo Fisher Scientific) according to the manufacturer's instructions, and the presence of intact recombinant RexAB was confirmed via SDS-PAGE and Western blot analysis.

Nuclease and helicase activity assays. The nuclease and helicase activities of RexAB were measured to confirm the AddAB-like activity in our recombinant RexAB protein. Staphylococcal DNA was amplified from the JE2 whole genome by colony PCR using the primer pair Chi control F (5'-TCAGTGAATTAGATGATTCGC-3') and Chi control R (5'-TTCATACGTATGAATGTTATTTGC-3'), where the amplicon lacked a Chi site region to be used as the DNA substrate in these assays.

Reactions were set up with either nuclease assay buffer (25 mM Tris-acetate [pH 7.5], 2 mM Mg acetate, 1 mM dithiothreitol [DTT]) or helicase assay buffer (25 mM Tris-acetate [pH 7.5], 0.25 mM Mg acetate, 1 mM DTT), along with 5 ng μ l⁻¹ of DNA, 1 mM ATP or an equivalent volume of nuclease-free water, and either 20 nM or 50 nM recombinant RexAB for the nuclease and helicase assays, respectively. Additionally, for the helicase assay, 2 μ M SSB protein was added to each sample.

Samples were incubated statically at 37°C, and at 0, 5, 15, 30, 60, and 120 min, 5 μ l was removed and pipetted into 20 μ l of STEB buffer (40% [wt/vol] sucrose, 100 mM Tris-HCl, 10 mM EDTA, 0.5 mg/ml bromophenol blue [pH 8]) to stop the reaction. Twenty microliters of chloroform-isoamyl alcohol (24:1) was added to each tube, vortexed for 10 s, and centrifuged for 2 min at 17,000 $\times$ g to remove the protein and any compounds used for inhibition. The aqueous (upper blue) phase was loaded onto a 1% (wt/vol) agarose gel prepared in Tris-borate-EDTA (TBE) buffer, and electrophoresis was carried out at 85 V for 1 h. The gels were subsequently stained with SYBR Safe DNA gel stain (Invitrogen) at a 1/10,000 dilution in TBE buffer for 2 h with rocking and visualized using a Gel Doc EZ imager (Bio-Rad). The band intensity was quantified using ImageJ software. For nuclease activity, values were normalized to those of the no-ATP controls at 0 h. For helicase activity, values were normalized to those of an ssDNA control that lacked the RexAB protein, which had been heated at 95°C for 2 min to denature the dsDNA and allow the SSB protein to bind and stabilize the two ssDNA strands.

Measurement of bacterial growth. To measure the growth of *S. aureus*, bacterial cultures were first grown to stationary phase in TSB at 37°C (180 rpm) and then inoculated at a 1/50 dilution for growth curves or a 1/10 dilution for growth inhibition assays (supplementary) into a flat-bottomed 96-well plate

(200- μ l total volume) and placed into a POLARstar Omega plate reader (BMG Labtech). Bacteria were grown for 17 h at 37°C (700 rpm), and the absorbance at 600 nm was measured every 30 min.

Hemolytic activity. The hemolytic activity of culture supernatants was determined as described previously (77). Briefly, stationary-phase *S. aureus* cultures were pelleted for 5 min at $17,000 \times g$, and 400 μ l of the supernatant was pipetted into microcentrifuge tubes. An equal volume of 2% defibrinated sheep blood in PBS was added, and the mixture was incubated statically for 1 h at 37°C. Fresh TSB containing 2% defibrinated sheep blood was used as a negative control. After incubation, the samples were centrifuged for 5 min at $500 \times g$, and 200 μ l of the supernatant was transferred into a flat-bottomed 96-well plate. The absorbance of released hemoglobin was measured at 540 nm using a POLARstar Omega plate reader (BMG Labtech). Percent hemolysis was calculated relative to the wild-type control.

Staphyloxanthin extraction and quantification. To extract and quantify staphyloxanthin, *S. aureus* stationary-phase cultures grown in TSB were harvested by centrifugation at $17,000 \times g$ for 2 min. Staphyloxanthin was extracted by incubating the culture in methanol at 42°C for 30 min. Cells were pelleted by centrifugation at $17,000 \times g$ for 2 min, and 100 μ l of the supernatant was transferred into a flat-bottomed 96-well plate. The released staphyloxanthin was quantified by measuring the absorbance at 462 nm using a POLARstar Omega plate reader (BMG Labtech).

Catalase activity. To determine the level of catalase activity, cultures grown overnight were washed three times in PBS, and 10^7 CFU were inoculated into 1 ml of 100 μ M H_2O_2 (diluted in PBS). Samples were incubated at 37°C, protected from light, for 15 min. Two hundred microliters of the sample was centrifuged at $17,000 \times g$ for 3 min, and 20 μ l of the supernatant was added to a 96-well microtiter plate. PBS containing no bacteria was used as a negative control.

Catalase activity was determined indirectly by measuring the concentration of H_2O_2 over time using the Pierce quantitative peroxide assay kit (aqueous-compatible formulation; Thermo Fisher Scientific). According to the manufacturer's instructions, reagents A and B were mixed at a ratio of 1:100, and 200 μ l was added to each sample in the 96-well plate. The plate was then incubated for 30 min at room temperature, and the absorbance was measured at 595 nm using an iMark microplate reader (Bio-Rad). The H_2O_2 concentration was determined using a standard curve of known concentrations (up to 1 mM).

Computational analyses. Multiple-sequence alignments were generated using Clustal Omega via the EMBL-EBI Web server (78). Parameters were left in their default settings with the exception of alignment "order," which was set to consider the input order. Protein structures were predicted using the Phyre2 protein fold recognition server (84). Phyre2 structural models were viewed and manipulated using PyMOL molecular graphics system version 2.3 (Schrödinger).

Statistical analyses. Data are presented as the means or medians from three or more independent experiments and were analyzed by Student's *t* test (two tailed, unpaired, and assuming equal variances), one-way analysis of variance (ANOVA), or two-way ANOVA corrected for multiple comparisons, as described in the figure legends. For each experiment, "n" refers to the number of independent biological replicates. CFU counts from murine experiments are presented as the values obtained from each animal, and significance was assessed using the Mann-Whitney test. A *P* value of <0.05 was considered significant between data points (GraphPad Prism 7 for Windows).

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

FIG S1, TIF file, 0.3 MB.

FIG S2, TIF file, 0.4 MB.

FIG S3, TIF file, 0.6 MB.

FIG S4, TIF file, 0.6 MB.

FIG S5, TIF file, 0.3 MB.

FIG S6, TIF file, 0.5 MB.

FIG S7, TIF file, 0.4 MB.

TABLE S1, DOCX file, 0.01 MB.

ACKNOWLEDGMENTS

A.M.E. and R.S.C. acknowledge funding from Shionogi & Co., Ltd. A.M.E. also acknowledges support from the National Institute for Health Research (NIHR) Imperial Biomedical Research Centre (BRC). K.P.H. is supported by a Ph.D. scholarship funded by a Medical Research Council award to the Centre for Molecular Bacteriology and Infection (MR/J006874/1). D.P. is supported by National Institutes of Health grants R01HL134870 and R21AI153646. D.A.I.M. acknowledges funding from the Medical Research Council (career development award MR/M009505/1). All authors acknowledge the provision of strains by the Network on Antimicrobial Resistance in *Staphylococcus aureus* (NARSA) Program under NIAID/NIH contract no. HHSN272200700055C. The funders had no role in the study design, interpretation of the findings, or the writing of the manuscript.

REFERENCES

- Lenhart JS, Schroeder JW, Walsh BW, Simmons LA. 2012. DNA repair and genome maintenance in *Bacillus subtilis*. *Microbiol Mol Biol Rev* 76: 530–564. <https://doi.org/10.1128/MMBR.05020-11>.
- Baharoglu Z, Mazel D. 2014. SOS, the formidable strategy of bacteria against aggressions. *FEMS Microbiol Rev* 38:1126–1145. <https://doi.org/10.1111/1574-6976.12077>.
- Tong SYC, Davis JS, Eichenberger E, Holland TL, Fowler VG. 2015. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 28: 603–661. <https://doi.org/10.1128/CMR.00134-14>.
- Kreisel K, Boyd K, Langenberg P, Roghmann MC. 2006. Risk factors for recurrence in patients with *Staphylococcus aureus* infections complicated by bacteremia. *Diagn Microbiol Infect Dis* 55:179–184. <https://doi.org/10.1016/j.diagmicrobio.2006.01.021>.
- Rigby KM, DeLeo FR. 2012. Neutrophils in innate host defense against *Staphylococcus aureus* infections. *Semin Immunopathol* 34:237–259. <https://doi.org/10.1007/s00281-011-0295-3>.
- Buvelot H, Posfay-Barbe KM, Linder P, Schrenzel J, Krause K-H. 2017. *Staphylococcus aureus*, phagocyte NADPH oxidase and chronic granulomatous disease. *FEMS Microbiol Rev* 41:139–157. <https://doi.org/10.1093/femsre/fuw042>.
- Ellson CD, Davidson K, Ferguson GJ, O'Connor R, Stephens LR, Hawkins PT. 2006. Neutrophils from p40^{phox}−/− mice exhibit severe defects in NADPH oxidase regulation and oxidant-dependent bacterial killing. *J Exp Med* 203:1927–1937. <https://doi.org/10.1084/jem.20052069>.
- Rogers DE, Tompsett R. 1952. The survival of staphylococci within human leukocytes. *J Exp Med* 95:209–230. <https://doi.org/10.1084/jem.95.2.209>.
- Melly MA, Thomison JB, Rogers DE. 1960. Fate of staphylococci within human leukocytes. *J Exp Med* 112:1121–1130. <https://doi.org/10.1084/jem.112.6.1121>.
- Hirsch J, Cohn Z. 1960. Degranulation of polymorphonuclear leucocytes following phagocytosis of microorganisms. *J Exp Med* 112:1005–1014. <https://doi.org/10.1084/jem.112.6.1005>.
- Babior BM. 1987. The respiratory burst oxidase. *Trends Biochem Sci* 12:241–243. [https://doi.org/10.1016/0968-0004\(87\)90118-6](https://doi.org/10.1016/0968-0004(87)90118-6).
- Fridovich I. 1995. Superoxide radical and superoxide dismutases. *Annu Rev Biochem* 64:97–112. <https://doi.org/10.1146/annurev.bi.64.070195.000525>.
- Hampton MB, Kettle AJ, Winterbourn CC. 1996. Involvement of superoxide and myeloperoxidase in oxygen-dependent killing of *Staphylococcus aureus* by neutrophils. *Infect Immun* 64:3512–3517. <https://doi.org/10.1128/IAI.64.9.3512-3517.1996>.
- Repine JE, Fox RB, Berger EM. 1981. Hydrogen peroxide kills *Staphylococcus aureus* by reacting with staphylococcal iron to form hydroxyl radical. *J Biol Chem* 256:7094–7096.
- Imlay JA, Chin SM, Linn S. 1988. Toxic DNA damage by hydrogen peroxide through the Fenton reaction *in vivo* and *in vitro*. *Science* 240:640–642. <https://doi.org/10.1126/science.2834821>.
- Lanza F. 1998. Clinical manifestation of myeloperoxidase deficiency. *J Mol Med (Berl)* 76:676–681. <https://doi.org/10.1007/s001090050267>.
- Parry MF, Root RK, Metcalf JA, Delaney KK, Kaplow LS, Richar WJ. 1981. Myeloperoxidase deficiency. Prevalence and clinical significance. *Ann Intern Med* 95:293–301. <https://doi.org/10.7326/0003-4819-95-3-293>.
- Ben-Ari J, Wolach O, Gavrieli R, Wolach B. 2012. Infections associated with chronic granulomatous disease: linking genetics to phenotypic expression. *Expert Rev Anti Infect Ther* 10:881–894. <https://doi.org/10.1586/eri.12.77>.
- Pollock JD, Williams DA, Gifford MA, Li LL, Du X, Fisherman J, Orkin SH, Doerschuk CM, Dinuer MC. 1995. Mouse model of X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide production. *Nat Genet* 9:202–209. <https://doi.org/10.1038/ng0295-202>.
- Imlay JA. 2003. Pathways of oxidative damage. *Annu Rev Microbiol* 57:395–418. <https://doi.org/10.1146/annurev.micro.57.030502.090938>.
- Chang W, Small DA, Toghrol F, Bentley WE. 2006. Global transcriptome analysis of *Staphylococcus aureus* response to hydrogen peroxide. *J Bacteriol* 188:1648–1659. <https://doi.org/10.1128/JB.188.4.1648-1659.2006>.
- Friedberg EC, Wood RD, Walker GC, Schultz RA, Siede W, Ellenberger T. 2006. DNA repair and mutagenesis, 2nd ed. ASM Press, Washington, DC.
- Fang FC. 2011. Antimicrobial actions of reactive oxygen species. *mBio* 2:e00141-11. <https://doi.org/10.1128/mBio.00141-11>.
- Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A genetic resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. *mBio* 4:e00537-12. <https://doi.org/10.1128/mBio.00537-12>.
- Liu GY, Essex A, Buchanan JT, Datta V, Hoffman HM, Bastian JF, Fierer J, Nizet V. 2005. *Staphylococcus aureus* golden pigment impairs neutrophil killing and promotes virulence through its antioxidant activity. *J Exp Med* 202:209–215. <https://doi.org/10.1084/jem.20050846>.
- Painter KL, Hall A, Ha KP, Edwards AM. 2017. The electron transport chain sensitizes *Staphylococcus aureus* and *Enterococcus faecalis* to the oxidative burst. *Infect Immun* 85:e00659-17. <https://doi.org/10.1128/IAI.00659-17>.
- Malachowa N, Whitney AR, Kobayashi SD, Sturdevant DE, Kennedy AD, Braughton KR, Shabb DW, Diep BA, Chambers HF, Otto M, DeLeo FR. 2011. Global changes in *Staphylococcus aureus* gene expression in human blood. *PLoS One* 6:e18617. <https://doi.org/10.1371/journal.pone.0018617>.
- Buscher K, Wang H, Zhang X, Striowski P, Wirth B, Saggu G, Lütke-Enking S, Mayadas TN, Ley K, Sorokin L, Song J. 2016. Protection from septic peritonitis by rapid neutrophil recruitment through omental high endothelial venules. *Nat Commun* 7:10828. <https://doi.org/10.1038/ncomms10828>.
- Zuñiga-Castillo J, Romero D, Martínez-Salazar JM. 2004. The recombination genes *addAB* are not restricted to Gram-positive bacteria: genetic analysis of the recombination initiation enzymes RecF and AddAB in *Rhizobium etli*. *J Bacteriol* 186:7905–7913. <https://doi.org/10.1128/JB.186.23.7905-7913.2004>.
- Rocha EPC, Cornet E, Michel B. 2005. Comparative and evolutionary analysis of the bacterial homologous recombination systems. *PLoS Genet* 1:e15. <https://doi.org/10.1371/journal.pgen.0010015>.
- Chedin F, Kowalczykowski SC. 2002. A novel family of regulated helicases/nucleases from Gram-positive bacteria: insights into the initiation of DNA recombination. *Mol Microbiol* 43:823–834. <https://doi.org/10.1046/j.1365-2958.2002.02785.x>.
- Cromie GA. 2009. Phylogenetic ubiquity and shuffling of the bacterial RecBCD and AddAB recombination complexes. *J Bacteriol* 191: 5076–5084. <https://doi.org/10.1128/JB.00254-09>.
- Wigley DB. 2013. Bacterial DNA repair: recent insights into the mechanism of RecBCD, AddAB and AdnAB. *Nat Rev Microbiol* 11:9–13. <https://doi.org/10.1038/nrmicro2917>.
- Drlica K. 1999. Mechanism of fluoroquinolone action. *Curr Opin Microbiol* 2:504–508. [https://doi.org/10.1016/S1369-5274\(99\)00008-9](https://doi.org/10.1016/S1369-5274(99)00008-9).
- Lee Y-J, Park S-J, Ciccone SLM, Kim C-R, Lee S-H. 2006. An *in vivo* analysis of MMC-induced DNA damage and its repair. *Carcinogenesis* 27: 446–453. <https://doi.org/10.1093/carcin/bgi254>.
- Yeeles JTP, Dillingham MS. 2007. A dual-nuclease mechanism for DNA break processing by AddAB-type helicase-nucleases. *J Mol Biol* 371: 66–78. <https://doi.org/10.1016/j.jmb.2007.05.053>.
- Painter KL, Strange E, Parkhill J, Bamford KB, Armstrong-James D, Edwards AM. 2015. *Staphylococcus aureus* adapts to oxidative stress by producing H₂O₂-resistant small-colony variants via the SOS response. *Infect Immun* 83:1830–1844. <https://doi.org/10.1128/IAI.03016-14>.
- Clarke RS, Bruderer MS, Ha KP, Edwards AM. 2019. RexAB is essential for the mutagenic repair of DNA damage caused by co-trimoxazole. *Antimicrob Agents Chemother* 63:e00944-19. <https://doi.org/10.1128/AAC.00944-19>.
- Seward B, de Haas CJC, Vervoort F, Rigby KM, DeLeo FR, Otto M, van Strijp JAG, Nijland R. 2013. Staphylococcal alpha-phenol soluble modulins contribute to neutrophil lysis after phagocytosis. *Cell Microbiol* 15:1427–1437. <https://doi.org/10.1111/cmi.12130>.
- Selton-Suty C, Célaré M, Le Moing V, Doco-Lecompte T, Chirouze C, lung B, Strady C, Revest M, Vandenesch F, Bouvet A, Delahaye F, Alla F, Duval X, Hoen B, AEPEI Study Group. 2012. Prevalence of *Staphylococcus aureus* in infective endocarditis: a 1-year population-based survey. *Clin Infect Dis* 54:1230–1239. <https://doi.org/10.1093/cid/cis199>.
- Holland TL, Baddour LM, Bayer AS, Hoen B, Miro JM, Fowler VG. 2016. Infective endocarditis. *Nat Rev Dis Primers* 2:16059. <https://doi.org/10.1038/nrdp.2016.59>.
- Hooper DC. 2001. Mechanisms of action of antimicrobials: focus on

- fluoroquinolones. Clin Infect Dis 32:59–S15. <https://doi.org/10.1086/319370>.
43. Tai PC, Davis BD. 1979. Triphasic concentration effects of gentamicin on activity and misreading in protein synthesis. Biochemistry 18:193–198. <https://doi.org/10.1021/bi00568a029>.
 44. Gourley CR, Negretti NM, Konkel ME. 2017. The food-borne pathogen *Campylobacter jejuni* depends on the AddAB DNA repair system to defend against bile in the intestinal environment. Sci Rep 7:14777. <https://doi.org/10.1038/s41598-017-14646-9>.
 45. Amundsen SK, Fero J, Hansen LM, Cromie GA, Solnick JV, Smith GR, Salama NR. 2008. *Helicobacter pylori* AddAB helicase-nuclease and RecA promote recombination-related DNA repair and survival during stomach colonization. Mol Microbiol 69:994–1007. <https://doi.org/10.1111/j.1365-2958.2008.06336.x>.
 46. Cano DA, Pucciarelli MG, García-Del Portillo F, Casadesús J. 2002. Role of the RecBCD recombination pathway in *Salmonella* virulence. J Bacteriol 184:592–595. <https://doi.org/10.1128/jb.184.2.592-595.2002>.
 47. Voyich JM, Braughton KR, Sturdevant DE, Whitney AR, Saïd-Salim B, Porcella SF, Long RD, Dorward DW, Gardner DJ, Kreiswirth BN, Musser JM, DeLeo FR. 2005. Insights into mechanisms used by *Staphylococcus aureus* to avoid destruction by human neutrophils. J Immunol 175:3907–3919. <https://doi.org/10.4049/jimmunol.175.6.3907>.
 48. Kobayashi SD, Braughton KR, Palazzolo-Ballance AM, Kennedy AD, Sampaoio E, Kristosturyan E, Whitney AR, Sturdevant DE, Dorward DW, Holland SM, Kreiswirth BN, Musser JM, DeLeo FR. 2010. Rapid neutrophil destruction following phagocytosis of *Staphylococcus aureus*. J Innate Immun 2:560–575. <https://doi.org/10.1159/000317134>.
 49. Schlosser-Silverman E, Elgrably-Weiss M, Rosenshine I, Kohen R, Altuvia S. 2000. Characterization of *Escherichia coli* DNA lesions generated within J774 macrophages. J Bacteriol 182:5225–5230. <https://doi.org/10.1128/jb.182.18.5225-5230.2000>.
 50. Mertens K, Lantsheer L, Ennis DG, Samuel JE. 2008. Constitutive SOS expression and damage-inducible AddAB-mediated recombinational repair systems for *Coxiella burnetii* as potential adaptations for survival within macrophages. Mol Microbiol 69:1411–1426. <https://doi.org/10.1111/j.1365-2958.2008.06373.x>.
 51. Cirz RT, Jones MB, Gingles NA, Minogue TD, Jarrahi B, Peterson SN, Romesberg FE. 2007. Complete and SOS-mediated response of *Staphylococcus aureus* to the antibiotic ciprofloxacin. J Bacteriol 189:531–539. <https://doi.org/10.1128/JB.01464-06>.
 52. Standish AJ, Weiser JN. 2009. Human neutrophils kill *Streptococcus pneumoniae* via serine proteases. J Immunol 183:2602–2609. <https://doi.org/10.4049/jimmunol.0900688>.
 53. Richardson AR, Libby SJ, Fang FC. 2008. A nitric oxide-inducible lactate dehydrogenase enables *Staphylococcus aureus* to resist innate immunity. Science 319:1672–1676. <https://doi.org/10.1126/science.1155207>.
 54. Kelley WL. 2006. Lex marks the spot: the virulent side of SOS and a closer look at the LexA regulon. Mol Microbiol 62:1228–1238. <https://doi.org/10.1111/j.1365-2958.2006.05444.x>.
 55. Erill I, Campoy S, Barbé J. 2007. Aeons of distress: an evolutionary perspective on the bacterial SOS response. FEMS Microbiol Rev 31:637–656. <https://doi.org/10.1111/j.1574-6976.2007.00082.x>.
 56. Varhimo E, Savijoki K, Jalava J, Kuipers OP, Varmanen P. 2007. Identification of a novel streptococcal gene cassette mediating SOS mutagenesis in *Streptococcus uteris*. J Bacteriol 189:5210–5222. <https://doi.org/10.1128/JB.00473-07>.
 57. Vickerman MM, Heath DG, Clewell DB. 1993. Construction of recombination-deficient strains of *Streptococcus gordonii* by disruption of the *recA* gene. J Bacteriol 175:6354–6357. <https://doi.org/10.1128/jb.175.19.6354-6357.1993>.
 58. Savijoki K, Ingmer H, Frees D, Vogensen FK, Palva A, Varmanen P. 2003. Heat and DNA damage induction of the LexA-like regulator HdiR from *Lactococcus lactis* is mediated by RecA and ClpP. Mol Microbiol 50:609–621. <https://doi.org/10.1046/j.1365-2958.2003.03713.x>.
 59. Boutry C, Delplace B, Clippe A, Fontaine L, Hols P. 2013. SOS response activation and competence development are antagonistic mechanisms in *Streptococcus thermophilus*. J Bacteriol 195:696–707. <https://doi.org/10.1128/JB.01605-12>.
 60. Dybvig K, Hollingshead SK, Heath DG, Clewell DB, Sun F, Woodard A. 1992. Degenerate oligonucleotide primers for enzymatic amplification of *recA* sequences from Gram-positive bacteria and mycoplasmas. J Bacteriol 174:2729–2732. <https://doi.org/10.1128/jb.174.8.2729-2732.1992>.
 61. Latorre M, Galloway-Peña J, Roh JH, Budinich M, Reyes-Jara A, Murray BE, Maass A, González M. 2014. *Enterococcus faecalis* reconfigures its transcriptional regulatory network activation at different copper levels. Metallomics 6:572–581. <https://doi.org/10.1039/c3mt00288h>.
 62. Latorre M, Quenti D, Travisany D, Singh KV, Murray BE, Maass A, Cambiazo V. 2018. The role of Fur in the transcriptional and iron homeostatic response of *Enterococcus faecalis*. Front Microbiol 9:1580. <https://doi.org/10.3389/fmicb.2018.01580>.
 63. Permina EA, Mironov AA, Gelfand MS. 2002. Damage-repair error-prone polymerases of eubacteria: association with mobile genome elements. Gene 293:133–140. [https://doi.org/10.1016/S0378-1119\(02\)00701-1](https://doi.org/10.1016/S0378-1119(02)00701-1).
 64. Jackson SP. 2002. Sensing and repairing DNA double-strand breaks. Carcinogenesis 23:687–696. <https://doi.org/10.1093/carcin/23.5.687>.
 65. Dziegielewska B, Beerman TA, Bianco PR. 2006. Inhibition of RecBCD enzyme by antineoplastic DNA alkylating agents. J Mol Biol 361:898–919. <https://doi.org/10.1016/j.jmb.2006.06.068>.
 66. Karu AE, Linn S. 1972. Uncoupling of the *recBC* ATPase from DNase by DNA crosslinked with psoralen. Proc Natl Acad Sci U S A 69:2855–2859. <https://doi.org/10.1073/pnas.69.10.2855>.
 67. Wilkinson M, Troman L, Wan Nur Ismah WA, Chaban Y, Avison MB, Dillingham MS, Wigley DB. 2016. Structural basis for the inhibition of RecBCD by Gam and its synergistic antibacterial effect with quinolones. Elife 5:e22963. <https://doi.org/10.7554/elife.22963>.
 68. Sun D, Hurlley LH. 1992. Structure-activity relationships of (+)-CC-1065 analogs in the inhibition of helicase-catalyzed unwinding of duplex DNA. J Med Chem 35:1773–1782. <https://doi.org/10.1021/jm00088a012>.
 69. Amundsen SK, Spicer T, Karabulut AC, Londoño LM, Eberhart C, Fernandez Vega V, Bannister TD, Hodder P, Smith GR. 2012. Small-molecule inhibitors of bacterial AddAB and RecBCD helicase-nuclease DNA repair enzymes. ACS Chem Biol 7:879–891. <https://doi.org/10.1021/cb3000018x>.
 70. Lim CSQ, Ha KP, Clarke RS, Gavin LA, Cook DT, Hutton JA, Sutherland CL, Edwards AM, Evans LE, Tate EW, Lanyon-Hogg T. 2019. Identification of a potent small-molecule inhibitor of bacterial DNA repair that potentiates quinolone antibiotic activity in methicillin-resistant *Staphylococcus aureus*. Bioorg Med Chem 27:114962. <https://doi.org/10.1016/j.bmc.2019.06.025>.
 71. Wiegand I, Hilpert K, Hancock REW. 2008. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc 3:163–175. <https://doi.org/10.1038/nprot.2007.521>.
 72. Faul F, Erdfelder E, Lang A-G, Buchner A. 2007. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods 39:175–191. <https://doi.org/10.3758/bf03193146>.
 73. Jacquet R, LaBauve AE, Akoolo L, Patel S, Alqarzaee AA, Wong Fok Lung T, Poorey K, Stinear TP, Thomas VC, Meagher RJ, Parker D. 2019. Dual gene expression analysis identifies factors associated with *Staphylococcus aureus* virulence in diabetic mice. Infect Immun 87:e00163-19. <https://doi.org/10.1128/IAI.00163-19>.
 74. National Research Council. 2011. Guide for the care and use of laboratory animals, 8th ed. National Academies Press, Washington, DC.
 75. Macrina FL, Evans RP, Tobian JA, Hartley DL, Clewell DB, Jones KR. 1983. Novel shuttle plasmid vehicles for *Escherichia-Streptococcus* transgeneric cloning. Gene 25:145–150. [https://doi.org/10.1016/0378-1119\(83\)90176-2](https://doi.org/10.1016/0378-1119(83)90176-2).
 76. Kristich CJ, Chandler JR, Dunny GM. 2007. Development of a host-genotype-independent counterselectable marker and a high-frequency conjugative delivery system and their use in genetic analysis of *Enterococcus faecalis*. Plasmid 57:131–144. <https://doi.org/10.1016/j.plasmid.2006.08.003>.
 77. Pader V, James EH, Painter KL, Wigneshweraraj S, Edwards AM. 2014. The Agr quorum-sensing system regulates fibronectin binding but not hemolysis in the absence of a functional electron transport chain. Infect Immun 82:4337–4347. <https://doi.org/10.1128/IAI.02254-14>.
 78. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol 7:539. <https://doi.org/10.1038/msb.2011.75>.
 79. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robertse B, Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova O, Brover V, Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM, Galdorf T, Gupta T, Haft D, Hatcher E, Hlavina W, Joardar VS, Kodali VK, Li W, Maglott D, Masterson P, McGarvey KM, Murphy MR, O'Neill K, Pujar S, Rangwala SH, Rausch D, Riddick LD, Schoch C, Shkeda A, Storz SS, Sun H, Thibaud-Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C, Webb D, Wu W, Landrum MJ, Kimchi A, et al. 2016. Reference sequence

- (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res* 44:D733–D745. <https://doi.org/10.1093/nar/gkv1189>.
80. Horsburgh MJ, Aish JL, White IJ, Shaw L, Lithgow JK, Foster SJ. 2002. SigmaB modulates virulence determinant expression and stress resistance: characterization of a functional *rsbU* strain derived from *Staphylococcus aureus* 8325-4. *J Bacteriol* 184:5457–5467. <https://doi.org/10.1128/jb.184.19.5457-5467.2002>.
 81. Pakula R, Walczak W. 1963. On the nature of competence of transformable streptococci. *J Gen Microbiol* 31:125–133. <https://doi.org/10.1099/00221287-31-1-125>.
 82. Ike Y, Craig RA, White BA, Yagi Y, Clewell DB. 1983. Modification of *Streptococcus faecalis* sex pheromones after acquisition of plasmid DNA. *Proc Natl Acad Sci U S A* 80:5369–5373. <https://doi.org/10.1073/pnas.80.17.5369>.
 83. Leenhouts K, Buist G, Bolhuis A, ten Berge A, Kiel J, Mierau I, Dabrowska M, Venema G, Kok J. 1996. A general system for generating unlabelled gene replacements in bacterial chromosomes. *Mol Gen Genet* 253: 217–224. <https://doi.org/10.1007/s004380050315>.
 84. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 Web portal for protein modeling, prediction and analysis. *Nat Protoc* 10:845–858. <https://doi.org/10.1038/nprot.2015.053>.
 85. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. 2000. The Protein Data Bank. *Nucleic Acids Res* 28:235–242. <https://doi.org/10.1093/nar/28.1.235>.
 86. Saikrishnan K, Yeeles JT, Gilhooly NS, Krajewski WW, Dillingham MS, Wigley DB. 2012. Insights into Chi recognition from the structure of an AddAB-type helicase-nuclease complex. *EMBO J* 31:1568–1578. <https://doi.org/10.1038/emboj.2012.9>.