See, E. J., Caskey, F., Johnson, D. W., & al., E. (2020). Availability, coverage, and scope of health information systems for kidney care across world countries and regions. *Nephrology Dialysis Transplantation*, 2020, [gfaa343]. <https://doi.org/10.1093/ndt/gfaa343>

Peer reviewed version

Link to published version (if available):
[10.1093/ndt/gfaa343](https://doi.org/10.1093/ndt/gfaa343)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/ndt/advance-article/doi/10.1093/ndt/gfaa343/6044457> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Availability, coverage, and scope of health information systems for kidney care across world countries and regions

Emily J See, MBBS^{1,2}; Aminu K Bello, PhD³; Adeera Levin, FRCPC⁴; Meaghan Lunney, MSc⁵; Mohamed A Osman, MD³; Feng Ye, MSc³; Gloria E Ashuntantang, MD⁶; Ezequiel Bellorin-Font, MD⁷; Mohammed Benghanem Gharbi, MD⁸; Sara Davison, MD³; Mohammad Ghnaimat, FACP⁹; Paul Harden, MD¹⁰; Htay Htay, MBBS¹¹; Vivekanand Jha, PhD^{12,13}; Kamyar Kalantar-Zadeh, MD¹⁴; Peter G Kerr PhD^{15,16}; Scott Klarenbach, MSc³; Csaba P Kovesdy, MD¹⁷; Valerie Luyckx, PhD^{18,19}; Brendon Neuen, MD²⁰; Donal O'Donoghue, MD^{21,22}; Shahrzad Ossareh, MD²³; Jeffrey Perl, MD^{24,25}; Harun Ur Rashid, PhD²⁶; Eric Rondeau, PhD^{27,28}; Saad Syed, BSc³; Laura Sola, MD²⁹; Irma Tchokhonelidze, MD³⁰; Vladimir Tesar, PhD³¹; Kriang Tungsanga, MD^{32,33}; Rumeysa Turan Kazancioglu, MD³⁴; Angela Yee-Moon Wang, PhD³⁵; Chih-Wei Yang, MD³⁶; Alexander Zemchenkov, PhD^{37,38}; Ming-hui Zhao, PhD^{39,40,41,42}; Kitty J Jager, PhD⁴³; Fergus Caskey, MD^{44,45,46}; Vlado Perkovic, PhD⁴⁷; Kailash K Jindal, FRCPC²; Ikechi G Okpechi, PhD^{48,49}; Marcello Tonelli, FRCPC^{50,51}; John Feehally, FRCP⁵²; David C Harris, FRACP⁵³; David W Johnson, PhD^{54,55,56}

¹Department of Intensive Care, Austin Health, Melbourne, Australia; ²School of Medicine, University of Melbourne, Melbourne, Australia; ³Division of Nephrology and Immunology, Department of Medicine, University of Alberta, Edmonton, Alberta, Canada; ⁴Division of Nephrology, Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada; ⁵Department of Community Health Sciences, University of Calgary, Calgary, Alberta, Canada; ⁶Faculty of Medicine and Biomedical Sciences, Yaounde General Hospital, University of Yaounde I, Yaounde, Cameroon; ⁷Division of Nephrology and Hypertension, Department of Medicine, Saint Louis University, Saint Louis, Missouri,

USA; ⁸Urinary Tract Diseases Department, Faculty of Medicine and Pharmacy of Casablanca, University Hassan II of Casablanca, Casablanca, Morocco; ⁹Nephrology Division, Department of Internal Medicine, The Specialty Hospital, Amman, Jordan; ¹⁰Oxford Kidney Unit, Oxford University Hospitals NHS Foundation Trust, Oxford, UK; ¹¹Department of Renal Medicine, Singapore General Hospital, Singapore; ¹²George Institute for Global Health India, New Delhi, India; ¹³Nuffield Department of Clinical Medicine, University of Oxford, Oxford, UK; ¹⁴Division of Nephrology and Hypertension, University of California Irvine Medical Center, Orange, California, USA; ¹⁵Department of Nephrology, Monash Medical Centre, Monash Health, Clayton, Victoria, Australia; ¹⁶Department of Medicine, Monash University, Clayton, Victoria, Australia; ¹⁷University of Tennessee Health Science Center, Memphis, Tennessee, USA; ¹⁸Institute of Biomedical Ethics and the History of Medicine, University of Zurich, Zurich, Switzerland; ¹⁹Renal Division, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; ²⁰The George Institute for Global Health, Newtown, New South Wales, Australia; ²¹Salford Royal NHS Foundation Trust, Salford, UK; ²²University of Manchester, Manchester, UK; ²³Division of Nephrology, Department of Medicine, Hasheminejad Kidney Center, Iran University of Medical Sciences, Tehran, Iran; ²⁴Division of Nephrology, St. Michael's Hospital and the Keenan Research Centre in the Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Ontario, Canada; ²⁵Division of Nephrology, Department of Medicine, University of Toronto, Toronto, Ontario, Canada; ²⁶Department of Nephrology, Kidney Foundation Hospital and Research Institute, Dhaka, Bangladesh; ²⁷Intensive Care Nephrology and Transplantation Department, Hopital Tenon, Assistance Publique-Hopitaux de Paris, Paris, France; ²⁸Sorbonne Université, Paris, France; ²⁹Dialysis Unit, CASMU-IAMPP, Montevideo, Uruguay; ³⁰Nephrology Development Clinical Center, Tbilisi State Medical University, Tbilisi, Georgia; ³¹Department of Nephrology, General University Hospital, Charles University, Prague, Czech

Republic; ³² Department of Medicine, Faculty of Medicine, King Chulalong Memorial Hospital, Chulalongkorn University, Bangkok, Thailand; ³³ Bhumirajanagarindra Kidney Institute, Bangkok, Thailand; ³⁴ Division of Nephrology, Bezmialem Vakif University, Istanbul, Turkey; ³⁵ Department of Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong; ³⁶ Kidney Research Center, Department of Nephrology, Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Taoyuan, Taiwan; ³⁷ Department of Internal Disease and Nephrology, North-Western State Medical University named after I.I. Mechnikov, Saint Petersburg, Russia; ³⁸ Department of Nephrology and Dialysis, Pavlov First Saint Petersburg State Medical University, Saint Petersburg, Russia; ³⁹ Renal Division, Department of Medicine, Peking University First Hospital, Beijing, China; ⁴⁰ Key Lab of Renal Disease, Ministry of Health of China, Beijing, China; ⁴¹ Key Lab of Chronic Kidney Disease Prevention and Treatment, Ministry of Education of China, Beijing, China; ⁴² Peking-Tsinghua Center for Life Sciences (CLS), Beijing, China; ⁴³ ERA-EDTA Registry, Department of Medical Informatics, Academic Medical Center, Amsterdam Public Health Research Institute, University of Amsterdam, Amsterdam, the Netherlands; ⁴⁴ UK Renal Registry, Learning and Research, Southmead Hospital, Bristol, UK; ⁴⁵ Population Health Sciences, University of Bristol, Bristol, UK; ⁴⁶ The Richard Bright Renal Unit, Southmead Hospital, North Bristol NHS Trust, Bristol, UK; ⁴⁷ The George Institute for Global Health, University of New South Wales Sydney, Sydney, New South Wales, Australia; ⁴⁸ Division of Nephrology and Hypertension, University of Cape Town, Cape Town, South Africa; ⁴⁹ Kidney and Hypertension Research Unit, University of Cape Town, Cape Town, South Africa; ⁵⁰ Department of Medicine, University of Calgary, Calgary, Alberta, Canada; ⁵¹ Pan-American Health Organization/World Health Organization's Collaborating Centre in Prevention and Control of Chronic Kidney Disease, University of Calgary, Calgary, Alberta, Canada; ⁵² University of Leicester, Leicester, UK; ⁵³ Centre for Transplantation and Renal

Research, Westmead Institute for Medical Research, University of Sydney, Sydney, New South Wales, Australia; ⁵⁴Centre for Kidney Disease Research, University of Queensland, Brisbane, Australia; ⁵⁵Translational Research Institute, Brisbane, Australia; ⁵⁶Metro South and Ipswich Nephrology and Transplant Services (MINTS), Princess Alexandra Hospital, Brisbane, Australia

Corresponding author

Professor David Johnson

Department of Nephrology

Level 2, ARTS Building

Princess Alexandra Hospital

199 Ipswich Road, Woolloongabba, QLD 4102, Australia

Tel: +61 7, Fax: +61 7

Email: david.johnson2@health.qld.gov.au

Word count:

Abstract: 268

Body: 3252

Figures: 4

Tables: 3

ABSTRACT

Background: Health information systems (HIS) are fundamental tools for the surveillance of health services, estimation of disease burden, and prioritisation of health resources. Several gaps in the availability of HIS for kidney disease were highlighted by the first iteration of the Global Kidney Health Atlas.

Methods: As part of its second iteration, the International Society of Nephrology conducted a cross-sectional global survey between July and October 2018 to explore the coverage and scope of HIS for kidney disease, with a focus on kidney replacement therapy (KRT).

Results: Of 182 invited countries, 154 countries responded to questions on HIS (85% response rate). KRT registries were available in almost all high income countries, but few low income countries, while registries for non-dialysis chronic kidney disease (CKD) or acute kidney injury (AKI) were rare. Registries in high income countries tended to be national, in contrast to registries in low income countries, which often operated at local or regional levels. Although cause of end-stage kidney disease, modality of KRT, and source of kidney transplant donors were frequently reported, few countries collected data on patient-reported outcome measures and only half of low income countries recorded process-based measures. Almost no countries had programs to detect AKI and practices to identify CKD targeted individuals with diabetes, hypertension, and cardiovascular disease, rather than members of high-risk ethnic groups.

Conclusions: These findings confirm significant heterogeneity in the global availability of HIS for kidney disease and highlight important gaps in their coverage and scope, especially in low income countries and across the domains of AKI, non-dialysis CKD, patient-reported outcomes, process-based measures and quality indicators for KRT service delivery.

What is already known about this subject:

- Health information systems are fundamental tools for the surveillance of health services, estimation of disease burden, and prioritisation of healthcare resources
- The first iteration of the Global Kidney Health Atlas highlighted significant variability in the availability of health information systems for kidney disease within and between countries
- Coverage of health information systems for kidney disease was particularly poor in low resource settings and across the domains of acute kidney injury and non-dialysis chronic kidney disease

What this study adds:

- This study goes beyond an examination of the availability of health information systems to explore in detail the collection and reporting of clinical outcomes, process-based measures, and quality indicators for service delivery, especially as they relate to kidney replacement therapy.
- Registries for kidney replacement therapy remain poorly available in low income countries, while registries for acute kidney injury and non-dialysis chronic kidney disease are scarce across all income groups. National registries are largely limited to high income countries, while registries in low income countries tend to operate at local or regional levels.
-
- Major gaps in the scope of registries include infrequent measurement and reporting of patient-reported outcome measures, process-based measures, and quality indicators for delivery of kidney replacement therapy. Fewer low income countries collected these metrics compared to high income countries.

What impact this may have on practice or policy:

- Establishing and strengthening health information systems for kidney disease should be a national and international priority
- Incorporating patient-reported outcomes, process-based measures and quality metrics into existing health information system infrastructure may improve the quality of care for patients with kidney disease
- Low and lower middle income countries may require provision of financial support, technical expertise, and mentorship by established systems to ensure success

Keywords: chronic kidney disease; end-stage kidney disease; health information systems; kidney replacement therapy; registries

INTRODUCTION

Health information systems (HIS) play an essential role in health service monitoring and benchmarking by ensuring the timely availability of robust and reliable health data.

Comprising registries, disease surveillance systems, facility databases and electronic health records, these systems can be used to quantify disease burden and outcomes,^{1,2} evaluate the quality and safety of care,^{3,4} develop and implement health-related policies,⁵⁻⁸ identify differences between and within health services, and recognize areas that incur unnecessary health care costs.^{2,9-13} Increasingly, HIS are also being utilized in the health research setting, where they have been successfully integrated into cohort studies and clinical trials to increase efficiency and reduce expenditures.¹⁴⁻¹⁶

Variability in the availability of HIS for kidney disease within and between countries was reported by the first iteration of the Global Kidney Health Atlas (GKHA), especially in low resource settings.¹⁷ This finding was not unexpected given the costs and resources required to establish and maintain such systems and is supported by earlier studies, which found that almost half of HIS for kidney disease were based in Europe.¹⁸⁻²⁰ The scope and content coverage of HIS for kidney disease has not been previously studied, but may be another important source of heterogeneity. Therefore, the objectives of this study were two-fold: first, to perform a temporal comparison of the availability, nature of provider participation and geographic coverage of kidney disease HIS with the 2017 GKHA; second, to comprehensively examine their scope, with respect to measurement and reporting of clinical outcomes, process-based measures, and quality indicators.

METHODS

The second iteration of the GKHA was a multinational cross-sectional survey conducted by the International Society of Nephrology (ISN), which was administered electronically to representatives of all 182 countries with ISN affiliate societies between July and October 2018. Countries were grouped by World Bank income group classification. A comprehensive description of the sampling approach, survey development, data handling and statistical analysis is available in the GKHA methodology paper.^{21,22} In short, a systematic search of the published literature and other data sources (including annual reports of kidney registries and databases) was undertaken to estimate the burden of KRT. Then, a survey questionnaire of three key opinion leaders from each country was carried out, to collect details on national kidney care practices. Further assistance from international contacts, collaborators, ISN leaders, and regional board members was also sought to review the consistency of regional data. The project was approved by the University of Alberta research ethics committee (protocol number: PR000063121).

The questionnaire was designed in five modules that corresponded to each of the six building blocks of health systems towards universal health coverage. One module related specifically to HIS, which included registries, quality indicators, and disease detection programs. Compared to the first iteration of the GKHA, the current survey focused in more detail on the global capacity and readiness to deliver kidney replacement therapy (KRT). Countries were asked to provide data on the availability and geographic coverage (national, regional, or local) of their kidney disease registries, as well as the nature of provider participation (mandatory or voluntary). Information pertaining to the scope and content of registries was also requested, with specific questions tailored to each registry type: dialysis (etiology of end-stage kidney disease [ESKD], KRT modality, process-based measures, hospitalizations, patient-reported

outcome measures, mortality); transplantation (etiology of ESKD, transplant source, process based measures, hospitalizations, patient-reported outcome measures, mortality); non-dialysis chronic kidney disease ([CKD] coverage of all CKD stages or restriction to stages 4 to 5), and acute kidney injury ([AKI] risk factors, etiology, incidence, hospitalizations, requirement for KRT, mortality). Patient-reported outcome measures were defined as health outcomes reported directly by the patient without interpretation of their response by others. They included satisfaction, fatigue, pain and quality of life. Process-based measures referred to markers of anemia, bone disease, and blood pressure control.

The proportion of centres in each country which used a HIS to measure and report quality indicators for KRT was also explored. For hemodialysis and peritoneal dialysis, these indicators included patient-reported outcome measures (e.g. fatigue, quality of life, satisfaction, pain), blood pressure, small solute clearance (e.g. Kt/V, creatinine clearance), hemoglobin/hematocrit, bone mineral markers (e.g. calcium, phosphate, parathyroid hormone), technique survival, and patient survival. For kidney transplantation, these indicators included patient reported outcome measures, delayed graft function, rejection rates, renal allograft function, graft survival and patient survival.

To examine local practices for the screening and detection of kidney disease, the survey also explored whether specific ethnic groups were considered to be at increased risk for AKI and CKD, and whether detection programs based on national policy and/or guidelines existed. Countries were asked about whether cases of AKI or CKD were identified through a “reactive” approach (identification through routine practice), an “active, routine” approach (active screening of high-risk populations during routine health encounters), or an “active, specific” approach (active screening of high-risk populations through specific screening

processes). Specific screening practices for CKD in the following high-risk groups were also covered: hypertension, diabetes, cardiovascular disease, autoimmune disease, elderly, urological disorders, chronic users of nephrotoxic agents, high-risk ethnic groups, and family history of CKD.

RESULTS

Among 182 invited countries, 160 countries (88%) participated in the GKHA survey, comprising 97.8% of the world's population. Of these, 154 countries responded to the questions about HIS (85%). Non-respondent countries were evenly distributed across regions and income groups and mostly represented smaller countries (Supplementary Table S1). Between the first and second iterations of the GKHA, the number of respondent countries providing information on HIS increased by 37. Complete details on the response rate and population coverage of the survey have been published elsewhere.²²

Availability of renal registries

The survey demonstrated wide variation in the availability of registries within and between ISN regions and World Bank income groups (Table 1, Figure 1). Individual results according to country are presented in Supplementary Table S2. Despite finding that the majority of countries had dialysis registries (n = 101, 66%) and more than half had registries for transplantation (n = 88, 57%), few low income countries had dialysis registries (n = 4, 18%) and none had registries for transplantation. Just 13 countries (8%) had registries for AKI, while non-dialysis CKD registries were only available in 19 countries (12%). There were no obvious differences in the availability of AKI or CKD registries between high and low income countries. Notably, few countries in South Asia had registries of any type. These figures are comparable to those reported by the first iteration of the GKHA, suggesting minimal improvement in registry availability in the two years between surveys.

Provider participation in registries

The nature of provider participation varied by registry type and by World Bank income group (Table 2). Compared to the first iteration of the GKHA, more countries in this survey

mandated provider participation in dialysis registries, including all low income countries (n = 4, 100%) and two-thirds of high income countries (n = 30, 63%). Similarly, the percentage of countries with mandatory provider participation in transplantation registries also increased between surveys, both in lower-middle (n = 5, 50%) and high income (n = 37, 73%) countries. For AKI registries, provider participation was mandatory in 8 of 13 countries (62%); among the 19 countries with non-dialysis CKD registries, participation was mandatory in 10 countries (53%). Participation was voluntary for all available registries in South Asian countries.

Geographic coverage of registries

Nearly all dialysis registries were national, especially in high (n = 47, 98%) and upper-middle (n = 28, 90%) income countries (Table 3). Among low income countries, two (50%) had national dialysis registries and one (25%) had a local dialysis registry. For transplantation, national registries were more commonly found in high (n = 43, 84%) or upper-middle (n = 26, 96%) income countries, rather than lower-middle income countries (n = 6, 60%). Less than half of all AKI registries were national (n = 6, 46%), compared to more than two-thirds of non-dialysis CKD registries (n=13, 68%) . The percentage of countries with national non-dialysis CKD registries in this survey (n=13, XX%) was much lower than the 89% of countries reported in the previous GKHA, suggesting that new non-dialysis CKD registries have tended to be regional or local.

Content coverage of registries

Questions pertaining to the content of registries were included for the first time in this iteration of the GKHA. Almost all dialysis registries collected data on the modality of KRT (n = 96, 95%), as well as the etiology of ESKD (n = 91, 90%), and patient mortality (n = 83,

82%; Figure 2). When analyzed by income group, fewer low income countries collected these metrics compared to high income countries. Process-based measures for dialysis were recorded by registries in most high income countries ($n = 40$, 83%), but by only half of registries in low income countries ($n = 2$). Patient-reported outcome measures for dialysis were recorded by one quarter of countries overall ($n = 24$, 24%).

With respect to transplantation registries, the majority of countries captured the donor source ($n = 85$, 97%), and most collected data on patient mortality ($n = 76$, 86%) and ESKD etiology ($n = 75$, 85%; Figure 3). Less than half of countries reported process-based measures for transplantation ($n = 37$, 42%) or hospitalizations ($n = 36$, 41%), and less than one-quarter of countries recorded patient-reported outcome measures for transplantation ($n = 18$, 20%), with no difference according to World Bank income group.

Most countries with AKI registries collected data on its etiology ($n = 10$, 77%), incidence ($n = 9$, 69%), requirement for KRT ($n = 9$, 69%), and patient mortality ($n = 9$, 69%). Fewer countries with AKI registries reported AKI-related hospitalizations ($n = 7$, 54%) or risk factors ($n = 6$, 46%). Where available, non-dialysis CKD registries generally covered the whole spectrum of CKD ($n = 12$, 63%), with registries in 5 countries (26%) covering only advanced CKD (i.e., stages 4 to 5).

Measurement and reporting of quality indicators for kidney replacement therapy

Data pertaining to measurement and reporting of key quality indicators for KRT were also collected for the first time in this survey. Commonly measured and reported indicators for hemodialysis service delivery included blood pressure and hemoglobin/hematocrit, followed by small solute clearance, technique survival and patient survival (Supplementary figure S1).

The proportion of centres increased from low to high income countries across all parameters. With regard to peritoneal dialysis service delivery, blood pressure and hemoglobin/hematocrit were also frequently measured and reported, as was patient survival (Supplementary figure S2). Technique survival and small solute clearance were not reported by many countries, especially those in low income settings. For countries in which kidney transplantation was available, patient survival, graft survival, and kidney allograft function were measured and reported more often than rejection and delayed graft function (Supplementary figure S3). Patient-reported outcome measures for hemodialysis, peritoneal dialysis and transplantation were infrequently measured and reported by all countries, but especially in low and lower-middle income settings.

Identification of high-risk groups

Information on how countries identified and/or screened patients for AKI and CKD was also collected. Just 6 countries (4%) had AKI detection programs in place and 35 countries (23%) had programs to detect CKD. Where available, the most common method of AKI detection was active screening, either through routine health encounters ($n = 4$, 67%), specific screening processes ($n = 3$, 50%) and/or a reactive approach ($n = 2$, 33%). High income countries tended to implement CKD detection programs through a reactive approach ($n = 8$, 47%) or by active screening at routine health encounters ($n = 7$, 41%). The only low income country with a CKD detection program also performed active screening during routine health encounters.

Very few low income countries identified ethnic groups at higher risk for AKI ($n = 3$, 14%) or CKD ($n = 0$, 0%), compared to over one-third of high income countries ($n = 21$ for AKI, 38%; $n = 20$ for CKD, 36%). Individuals with diabetes mellitus, hypertension, cardiovascular

disease, and urological disorders were more frequently screened for CKD than those of older age or high-risk ethnic group, those with a family history of CKD, and chronic users of nephrotoxic medications.

DISCUSSION

This study goes beyond an examination of the availability of HIS for kidney disease to explore the measurement and reporting of clinical outcomes, process-based measures, and quality indicators for service delivery, especially as they relate to KRT. Compared to the first iteration of the GKHA, it covered more countries and captured trends over time, including the presence of new registries. The findings of this study highlight key deficiencies in the availability, coverage and scope of HIS for kidney disease. Dialysis and transplantation registries remained poorly available in low income countries, while AKI and non-dialysis CKD registries continued to be scarce across all income groups. Although an increasing number of countries mandated provider participation, national registries were still largely limited to the high income setting. Major gaps in the scope of registries included the measurement and reporting of patient-reported outcomes, process-based measures, and quality indicators for KRT service delivery. Almost no countries had AKI detection programs in place, while CKD detection programs were generally implemented through routine health encounters and targeted individuals with diabetes mellitus, hypertension and cardiovascular disease, rather than those from high risk ethnic groups.

These findings have important implications for both low and high income countries.^{18,19} First, the persistently low prevalence of KRT registries in low income countries limits their access to local data on the epidemiology of ESKD and the outcomes of KRT, which impedes the delivery of accessible and cost-effective kidney care.^{8,23,24} Second, significant gaps in the availability of AKI and non-dialysis CKD registries in both low and high income countries are concerning because both conditions occur commonly and are associated with substantial morbidity, mortality, resource consumption, and health care costs.^{25,26} AKI and non-dialysis CKD registries could provide useful information to monitor important trends in disease

incidence, identify novel risk factors, and document their natural history. They could also be used to anticipate the future demands on health resources relating to the development of ESKD, and to audit compliance with recommended management. Third, the paucity of countries incorporating patient-reported outcomes, process-based measures, and quality indicators into their HIS leads to a lost opportunity to better understand the patient experience and perception of the quality of care. Patient-reported outcomes are increasingly recognized as important targets for therapeutic interventions,²⁷ while process-based measures and quality indicators are useful tools for auditing the standard of care, benchmarking health services, and identifying opportunities for improvement.

Addressing these gaps requires identification and prioritization of the areas of greatest need.²⁸ Because KRT is associated with substantial health care costs and significant variability in outcomes, establishing dialysis and transplantation registries is of high importance, since they could allow more efficient use of resources. For low income countries, the cost of developing such registries could be minimized by either incorporating them into existing registries, expanding the coverage of local registries to a national scale, or adapting established models for disease surveillance to local settings.^{29,30} Increasing the availability of AKI and non-dialysis CKD registries could be achieved inexpensively through the use of electronic health records and administrative databases since both have a computable phenotype; linkage to hospitalization and mortality data could also be achieved using existing infrastructure.^{31,32} In light of the potential cost associated with measuring and collecting patient-reported outcomes and process based measures, identification of the outcomes of greatest importance to patients and clinicians is first necessary, which has been one of the key objectives of the Standardized Outcomes in Nephrology (SONG) initiative.³³ Where possible, validated tools should be used

to ensure that the collected data are meaningful and comparable. The specific quality metrics selected for inclusion in the HIS should be individualized based on local relevance.

Prior to implementing changes to HIS infrastructure, consideration must be given to ensuring their quality, accuracy and security. This requires the availability of an adequate regulatory framework, trained and skilled personnel, financial and logistic support, and a functioning information technology system. Data collection methods should be continuously monitored to assess quality, and instruments for secure data storage, processing and analysis must be in place.¹⁸ HIS oversight should be transparent, and any output should be accessible to individuals, clinicians, governing bodies and policy makers. Simple systems that do not overburden staff, but that allow comparisons between and within health services would be of particular benefit.^{34–36}

Establishing such systems in low income countries presents a unique challenge.^{13,37,38} Infrastructure for the diagnosis and screening of kidney disease may be unavailable or centralized in major towns, limiting rural residents' access to testing. Collection of longitudinal data concerning disease prevalence and outcomes could be impeded by the migratory nature of some populations or by a lack of national census data for use as a reference population. A lack of trained personnel (i.e., data collectors, coders, analysts, epidemiologists) could also pose a barrier, and ongoing infrastructure and resource limitations could compromise HIS sustainability.^{29,39} Strategies to address these challenges must be considered prior to the development or expansion of HIS in low resource settings.

This study has several key strengths. The GKHA is the most comprehensive report of global readiness and capacity for kidney care, and provides important and unique insights into the

availability, coverage and scope of HIS for kidney disease. The survey was based on a well-validated framework for assessing care capacity that leverages the widely applied World Health Organization (WHO) health system building blocks. Regional and national stakeholders with knowledge of local contexts carefully reviewed responses to ensure data accuracy. Participating countries represented nearly the entire global population, with excellent coverage across all ISN regions and World Bank income levels. Despite these strengths, this study also has certain limitations. A total of 15% of countries were missing, which may have impacted on the representativeness of the data or introduced bias. Population surveys are prone to subjective responses (i.e., social desirability bias) and depend on respondents' knowledge and experiences. Additional aspects of HIS, such as quality, validity and regulation were beyond the scope of this study, but are important to consider when assessing a HIS's value.

In summary, significant gaps remain in the global coverage of HIS for kidney disease, particularly in low income countries, across the domains of AKI, non-dialysis CKD, and in the measurement and reporting of patient-reported outcomes, process-based measures, and quality indicators for KRT service delivery. Establishing and strengthening HIS should be a priority at both national and global levels to enable accurate estimation of the burden of kidney disease and to ensure that available resources are appropriately channelled to areas of greatest need. Financial and technical support for HIS in low and lower-middle income countries should be accelerated, and a collaborative approach with oversight and mentoring by established systems should be implemented to increase their likelihood of success. Integration of patient-reported outcomes, process-based measures, and quality metrics into new and existing HIS could drive important improvements in the quality of kidney care.

CONFLICT OF INTEREST

David Johnson has received consultancy fees, research grants, speaker's honoraria and travel sponsorships for Baxter Healthcare and Fresenius Medical Care. He is supported by a National Health and Medical Research Council Practitioner Fellowship. Htay Htay has received research grants from Johnson & Johnson Company and Singhealth new investigator grant (NIG). Angela Yee-Moon Wang has received speaker honoraria and Travel grants from Sanofi Renal and Fresenius Kabi and research grant from Sanofi Renal and Baxter Healthcare. All other authors have no conflict of interest to declare.

ACKNOWLEDGEMENTS

We thank Sandrine Damster (Research Project Manager at the ISN) and Alberta Kidney Disease Network staff (G. Houston, S. Szigety, S. Tiv) for helping to organize and conduct the survey and providing project management support; Jo-Ann Donner (Awards, Endorsements, ISN-ANIO Programs Coordinator) for helping with the manuscript management and submission process; ISN staff (Charu Malik, Claire Van Der Linden, Luisa Straniand, Luca Segantini) for their support; the GKHA steering committee, executive committee of the ISN, ISN regional leadership and the leaders of the ISN Affiliate Societies at the regional and country levels for their help, particularly with identification of survey respondents and data acquisition, which ensured the success of this initiative. None of the persons acknowledged received compensation for their role in the study.

FUNDING/SUPPORT

This work was supported by the International Society of Nephrology (grant RES0033080 to the University of Alberta).

Role of the funder/sponsor: The ISN provided administrative support for the design and implementation of the study and data collection activities. The authors were responsible for data management, analysis and interpretation, as well as manuscript preparation, review and approval, and the decision to submit the manuscript for publication.

REFERENCES

1. Larsson S, Lawyer P, Garellick G, Lindahl B, Lundström M. Use of 13 disease registries in 5 countries demonstrates the potential to use outcome data to improve health care's value. *Health Aff.* 2012;31:220-227.
2. Parkin D, Bray F, Ferlay J, Pisani P. Estimating the world cancer burden: Globocan 2000. *Int J Cancer.* 2001;94:153-156.
3. Han W, Sharman R, Heider A, Maloney N, Yang M, Singh R. Impact of electronic diabetes registry "Meaningful Use" on quality of care and hospital utilization. *J Am Med Informatics Assoc.* 2015;23:1-7.
4. Hoque DME, Kumari V, Ruseckaite R, Romero L, Evans SM. Impact of clinical registries on quality of patient care and health outcomes: protocol for a systematic review. *BMJ Open.* 2016;6:e010654.
5. McDonald SP. Australia and New Zealand Dialysis and Transplant Registry. *Kidney Int Suppl.* 2015;5:39-44.
6. Wilcox N, McNeil JJ. Clinical quality registries have the potential to drive improvements in the appropriateness of care. *Med J Aust.* 2016;205:21-26.
7. Nwomeh BC, Lowell W, Kable R, Haley K, Ameh EA. History and development of trauma registry: lessons from developed to developing countries. *World J Emerg Surg.* 2006;1:32.
8. Counil É, Cherni N, Kharrat M, Achour A, Trimech H. Trends of incident dialysis patients in Tunisia between 1992 and 2001. *Am J Kidney Dis.* 2008;51:463-470.
9. Collins AJ, Foley RN, Gilbertson DT, Chen S-C. United States Renal Data System public health surveillance of chronic kidney disease and end-stage renal disease. *Kidney Int Suppl.* 2015;5:2-7.
10. Zoccali C, Kramer A, Jager K. The databases: renal replacement therapy since 1989 -

- the European Renal Association and European Dialysis and Transplant Association (ERA-EDTA). *Clin J Am Soc Nephrol*. 2009;4:S18-S22.
11. Jin D-C. Dialysis registries in the world: Korean Dialysis Registry. *Kidney Int Suppl*. 2015;5:8-11.
 12. Leung CB, Cheung WL, Li PKT. Renal registry in Hong Kong-the first 20 years. *Kidney Int Suppl*. 2015;5:33-38.
 13. Jedy-Agba EE, Oga EA, Odutola M, et al. Developing national cancer registration in developing countries – case study of the Nigerian national system of cancer registries. *Front Public Heal*. 2015;3:1-10.
 14. Lim TO, Goh A, Lim YN, Morad Z. Use of renal registry data for research, health-care planning and quality improvement: what can we learn from registry data in the Asia-Pacific region? *Nephrology*. 2008;13:745-752.
 15. Hanafusa N, Nakai S, Iseki K, Tsubakihara Y. Japanese society for dialysis therapy renal data registry - a window through which we can view the details of Japanese dialysis population. *Kidney Int Suppl*. 2015;5:15-22.
 16. Jager KJ, Wanner C. Fifty years of ERA-EDTA Registry—a registry in transition. *Kidney Int Suppl*. 2015;5:12-14.
 17. Bello AK, Levin A, Tonelli M, et al. Assessment of Global Kidney Health Care Status. *JAMA*. 2017;317:1864.
 18. Black N, Barker M, Payne M. Cross sectional survey of multicentre clinical databases in the United Kingdom. *BMJ*. 2004;328:1478.
 19. Hanna TP, Kangolle ACT. Cancer control in developing countries: using health data and health services research to measure and improve access, quality and efficiency. *BMC Int Heal Hum Rights*. 2010;10:24.
 20. Liu FX, Rutherford P, Smoyer-Tomic K, Prichard S, Laplante S. A global overview of

- renal registries: a systematic review. *BMC Nephrol.* 2015;16:31.
21. Bello AK, Johnson DW, Feehally J, et al. Global Kidney Health Atlas (GKHA): design and methods. *Kidney Int Suppl.* 2017;7:145-153.
 22. Bello AK, Levin A, Lunney M, et al. Status of care for end stage kidney disease in countries and regions worldwide: international cross sectional survey. *BMJ.* 2019;367:15873.
 23. Barsoum RS. Overview: end-stage renal disease in the developing world. *Artif Organs.* 2002;26:737-746.
 24. Boutayeb A, Boutayeb S. The burden of non communicable diseases in developing countries. *Int J Equity Health.* 2005;4:2.
 25. Hoste EAJ, Kellum JA, Selby NM, et al. Global epidemiology and outcomes of acute kidney injury. *Nat Rev Nephrol.* 2018;14:607-625.
 26. Honeycutt AA, Segel JE, Zhuo X, Hoerger TJ, Imai K, Williams D. Medical costs of CKD in the Medicare population. *J Am Soc Nephrol.* 2013;24:1478-1483.
 27. Breckenridge K, Bekker HL, Gibbons E, et al. How to routinely collect data on patient-reported outcome and experience measures in renal registries in Europe: an expert consensus meeting. *Nephrol Dial Transplant.* 2015;30:1605-1614.
 28. Evans SM, Scott IA, Johnson NP, Cameron PA, McNeil JJ. Development of clinical-quality registries in Australia: the way forward. *Med J Aust.* 2011;194:360-363.
 29. Razeen Davids M, Eastwood JB, Selwood NH, et al. A renal registry for Africa: first steps. *Clin Kidney J.* 2016;9:162-167.
 30. United States Renal Data System. 2014 USRDS annual data report: epidemiology of kidney disease in the United States. National Institutes of Health. In: National Institute of Diabetes and Digestive and Kidney Diseases; 2014.
 31. Lucas H. Information and communications technology for future health systems in

- developing countries. *Soc Sci Med*. 2008;66:2122-2132.
32. Drawz PE, Archdeacon P, McDonald CJ, et al. CKD as a Model for Improving Chronic Disease Care through Electronic Health Records. *Clin J Am Soc Nephrol*. 2015;10:1488-1499.
 33. Tong A, Craig JC, Nagler E V., et al. Composing a new song for trials: the Standardized Outcomes in Nephrology (SONG) initiative. *Nephrol Dial Transplant*. 2017;32:1963-1966.
 34. Schena FP. Epidemiology of end-stage renal disease: international comparisons of renal replacement therapy. *Kidney Int*. 2000;57:S39-S45.
 35. Jager KJ, Zoccali C. Quality of care in end-stage renal disease: the importance of comparing “apples with apples.” *Nephrol Dial Transplant*. 2008;23:1116.
 36. Kjellstrand CM. International comparisons of dialysis survival are meaningless to evaluate differences in dialysis procedures. *Death Hemodial Prev or Inevitable?*. Published online 1994:55-68.
 37. Valsecchi MG, Steliarova-Foucher E. Cancer registration in developing countries: luxury or necessity? *Lancet Oncol*. 2008;9:159-167.
 38. Rastogi T, Hildesheim A, Sinha R. Opportunities for cancer epidemiology in developing countries. *Nat Rev Cancer*. 2004;4:909-917.
 39. Morsy A, Lim TO, Varatharajan S, Lim JY. National registries in developing countries: understanding construction challenges and implementation steps. *5th Cairo Int Biomed Eng Conf CIBEC*. Published online 2010:75-78.

FIGURE LEGENDS

Figure 1. Percentage of countries with registries for acute kidney injury, non-dialysis chronic kidney disease, dialysis and transplantation, by World Bank income group.

Figure 2. Percentage of countries with dialysis registries covering specific content, by World Bank income group.

Figure 3. Percentage of countries with transplantation registries covering specific content, by World Bank income group.

Figure 4. Percentage of countries adopting specific practices to identify chronic kidney disease among individuals in high-risk groups, by International Society of Nephrology region.

Table 1. Availability of registries for acute kidney injury, chronic kidney disease, dialysis and transplantation, by International Society of Nephrology region (n = 154).

	AKI	CKD	Dialysis	Transplantation
Region	N (%)	N (%)	N (%)	N (%)
Overall	13 (8)	19 (12)	101 (66)	88 (57)
ISN region				
Africa	4 (10)	3 (7)	18 (44)	6 (15)
Eastern & Central Europe	2 (11)	1 (5)	17 (89)	17 (89)
Latin America	1 (6)	5 (28)	14 (78)	13 (72)
Middle East	1 (9)	1 (9)	6 (55)	8 (73)
NIS & Russia	1 (14)	2 (29)	5 (71)	6 (86)
North America	0 (0)	0 (0)	4 (44)	3 (33)
North & East Asia	1 (14)	2 (29)	7 (100)	7 (100)
OSEA	1 (7)	2 (13)	10 (67)	8 (53)
South Asia	0 (0)	0 (0)	2 (29)	1 (14)
Western Europe	2 (10)	3 (15)	18 (90)	19 (95)

AKI: acute kidney injury; CKD: chronic kidney disease; ISN: International Society of Nephrology; NIS: Newly Independent States; OSEA.

Table 2. Provider participation in registries for acute kidney injury, chronic kidney disease, dialysis and transplantation, by World Bank income group (n = 154 countries).

	AKI	CKD	Dialysis	Transplantation
Participation	N (%)	N (%)	N (%)	N (%)
Mandatory				
Overall	8 (62)	10 (53)	60 (59)	57 (65)
Low income	2 (100)	2 (100)	4 (100)	0 (0)
Lower-middle income	2 (50)	3 (75)	10 (56)	5 (50)
Upper-middle income	0 (0)	1 (50)	16 (52)	15 (56)
High income	4 (80)	4 (36)	30 (63)	37 (73)
Voluntary				
Overall	5 (38)	8 (42)	36 (37)	26 (30)
Low income	0 (0)	0 (0)	0 (0)	0 (0)
Lower-middle income	2 (50)	0 (0)	7 (39)	3 (30)
Upper-middle income	2 (100)	1 (50)	11 (35)	9 (33)
High income	1 (20)	7 (64)	18 (38)	14 (27)
Missing or unknown				
Overall	0 (0)	1 (5)	5 (5)	5 (6)
Low income	0 (0)	0 (0)	0 (0)	0 (0)
Lower-middle income	0 (0)	1 (25)	1 (6)	2 (20)
Upper-middle income	0 (0)	0 (0)	4 (13)	3 (11)
High income	0 (0)	0 (0)	0 (0)	0 (0)

AKI: acute kidney injury; CKD: chronic kidney disease.

Table 3. Geographic coverage of registries for acute kidney injury, chronic kidney disease, dialysis and transplantation, by World Bank income group (n = 154 countries)

	AKI	CKD	Dialysis	Transplantation
Geographic Coverage	N (%)	N (%)	N (%)	N (%)
National				
Overall	6 (46)	13 (68)	90 (89)	75 (85)
Low income	2 (100)	1 (50)	2 (50)	0 (0)
Lower-middle income	0 (0)	2 (50)	13 (72)	6 (60)
Upper-middle income	0 (0)	1 (50)	28 (90)	26 (96)
High income	4 (80)	9 (82)	47 (98)	43 (84)
Regional				
Overall	2 (15)	4 (21)	15 (15)	9 (10)
Low income	0 (0)	0 (0)	0 (0)	0 (0)
Lower-middle income	2 (50)	2 (50)	4 (22)	2 (20)
Upper-middle income	0 (0)	0 (0)	2 (6)	0 (0)
High income	0 (0)	2 (18)	9 (19)	7 (14)
Local				
Overall	4 (31)	7 (37)	13 (13)	11 (13)
Low income	0 (0)	1 (50)	1 (25)	0 (0)
Lower-middle income	2 (50)	2 (50)	4 (22)	3 (30)
Upper-middle income	1 (50)	1 (50)	2 (6)	1 (4)
High income	1 (20)	3 (27)	6 (13)	7 (14)
Missing or unknown				
Overall	1 (8)	0 (0)	13 (13)	11 (13)
Low income	0 (0)	0 (0)	1 (25)	0 (0)
Lower-middle income	0 (0)	0 (0)	0 (0)	0 (0)
Upper-middle income	1 (50)	0 (0)	0 (0)	0 (0)
High income	0 (0)	0 (0)	0 (0)	0 (0)

AKI: acute kidney injury; CKD: chronic kidney disease. N.B. Some countries have registries with more than one type of geographic coverage.

Figure 1. Percentage of countries with registries for acute kidney injury, non-dialysis chronic kidney disease, dialysis and transplantation, by World Bank income group (n = 154 countries).

Figure 2. Percentage of countries with dialysis registries covering specific content, by World Bank income group (n = 101 countries).

Figure 3. Percentage of countries with transplantation registries covering specific content, by World Bank income group (n = 88 countries).