et al. (2021). The association between personality disorder traits and suicidality following sudden bereavement: a national cross-sectional survey. *Journal of Personality Disorders*, 35(6), 917-935.
https://doi.org/10.1521/pedi_2021_35_520

Publisher's PDF, also known as Version of record

Link to published version (if available):
[10.1521/pedi_2021_35_520](https://doi.org/10.1521/pedi_2021_35_520)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Guilford Press at [10.1521/pedi_2021_35_520](https://doi.org/10.1521/pedi_2021_35_520). Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The association between personality disorder traits and suicidality following sudden bereavement: a national cross-sectional survey

Sedigheh Zabihi¹

Rebecca Jones¹

Paul Moran²

Michael King¹

David Osborn^{1,3}

Alexandra Pitman^{1,3}

¹ UCL Division of Psychiatry, Maple House, 149 Tottenham Court Rd, London W1T 7NF

² Centre for Academic Mental Health, Department of Population Health Sciences, Bristol Medical School, University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK

³ Camden and Islington NHS Foundation Trust, St Pancras Hospital, London NW1 0PE, UK

Corresponding author: a.pitman@ucl.ac.uk

Abstract

Personality disorder is associated with increased risk of suicidal behaviour. We aimed to investigate the association between number of personality disorder traits and suicidality risk following sudden bereavement. Our secondary analysis of cross-sectional data on 3,167 young UK-based adults who had experienced sudden bereavement investigated the association between number of traits (measured using a standardised screening instrument) and post-bereavement suicide attempt and suicidal ideation. Using multivariable logistic regression we found a linear relationship between number of traits and suicide attempt (adjusted odds ratio [AOR]=1.36, 95% CI: 1.23 to 1.49) and suicidal ideation (AOR=1.31, 95% CI: 1.25 to 1.38) following bereavement. This represented an increase in odds by 36% and 31% respectively for each additional personality trait. Our study suggests that individuals with a greater number of traits suggestive of a personality disorder diagnosis are at increased risk of suicidality after a negative life event.

Keywords:

suicide attempt; suicidal ideation; bereavement; personality traits; personality disorder

Introduction

Suicide is a complex public health problem, with a range of risk factors, and great etiological heterogeneity (Turecki & Brent, 2016). Globally, the suicide rate is 10.5 deaths per 100,000 population, and it is estimated that a person dies by suicide every 40 seconds (WHO, 2016). Suicide is the second leading cause of death among young adults in the UK (ONS, 2016) and the third leading cause of death in 15 to 19 year-olds globally (WHO, 2019). Suicide attempts are 20 to 30 times more prevalent than completed suicides (Turecki & Brent, 2016), and also represent a major public health problem, being a risk factor for repetition (Ribeiro et al., 2016). Although it is very difficult in clinical practice to predict suicide and suicide attempt, improved recognition and understanding of clinical, psychological, sociological, and biological factors holds the potential to help detect individuals at higher risk and tailor support to their needs (Turecki & Brent, 2016).

Risk factors for suicide and suicidal behaviour include psychiatric disorders (Cavanagh, Carson, Sharpe, & Lawrie, 2003; Nock et al., 2008; Cash & Bridge, 2009), sociodemographic factors such as low socioeconomic status, white ethnicity and male gender (Beautrais, 2000; Bilsen, 2018; Goldman-Mellor et al., 2014), and negative life events, such as loss of a relationship, a job, or housing (Beck, Brown, & Steer, 1989; Turecki & Brent, 2016). These risk factors are likely to mediate their effect through personality factors, particularly in people diagnosed with a personality disorder. For example, diagnoses of borderline personality disorder and antisocial personality disorder are characterized by aggressive and impulsive traits (Turecki & Brent, 2016), and people diagnosed with borderline personality disorder have an increased risk of suicide (Black, Blum, Pfohl, & Hale, 2004) and suicide attempt (Ansell et al, 2015).

Personality disorder is described broadly as an enduring and pervasive pattern of emotional and cognitive difficulties that affect the way in which a person relates to others or

understand themselves (Winsper et al., 2020). The pooled prevalence of personality disorders in the general population has been estimated at 8%, and up to 10% in high-income countries (Winsper et al., 2020), but there are few robust estimates of the prevalence of individual traits characteristic of a diagnosis of personality disorder. A systematic review of studies conducted to date investigating the association between personality traits and suicidality identified that neuroticism and extroversion are the strongest predictors of suicidal ideation, suicide attempts, and suicide (Brezo, Paris, and Turecki (2006a). Higher levels of impulsive-aggressive traits are associated with younger age of death by suicide (McGirr et al., 2008). Other personality traits specifically associated with suicidality in young people include neuroticism (Beautrais, Joyce, & Mulder, 1999) and impulsivity (Brezo et al., 2006b). However, most of this work has tended to quantify the contribution of individual personality traits to suicidality rather than exploring the cumulative contribution of traits (Brezo, Paris, and Turecki (2006a).

We investigated whether there was an independent association between the number of traits suggestive of personality disorder and risk of suicidality in young adults; the age group representing the peak incidence of suicidal ideation and behaviour (Turecki & Brent, 2016). We chose to focus on young people who have experienced the loss of a close contact, because this represents a significant loss of social support during a vulnerable period, setting young people apart from their peers. We chose to analyze data on a population sample of young adults, rather than restricting our focus to those with a diagnosis of a personality disorder. We hypothesized that there would be a linear association between number of personality disorder traits and the probability of suicidal behaviour following traumatic bereavement. We also hypothesized that this association would be stronger for people who were bereaved by suicide rather than other sudden deaths, and for people who described themselves as very close to the deceased.

Methods

Sample

We conducted a secondary analysis of data from the UCL Bereavement Study; a cross-sectional study investigating risk of suicide following sudden bereavement (Pitman, Osborn, Rantell, & King, 2016). Sampling methods have been described previously (Pitman, Osborn, Rantell, & King, 2016). In brief, in 2010, all 164 higher education institutions (HEI) in the UK were invited to participate in an online survey to investigate the impact of sudden bereavement on mental health. This was judged to be the best means of accessing hard-to-reach groups, whilst avoiding the biases associated with recruiting a help-seeking sample (Pitman, Osborn, & King, 2015). A total of 37/164 (23%) of HEIs agreed to participate by sending all their students and staff individual emails, with an embedded survey link. Ten of the 37 chose to modify sampling, due to the sensitivities of the topic, by advertising the study on their weekly email digests, advertising the survey on both the student and staff intranet, or sending the survey solely to students. Participating HEIs represented a range of agricultural, performing arts, and academic institutions, providing broad geographic and socioeconomic representation, and accessed a sampling frame of 659,572 bereaved and non-bereaved staff and students. There was no accurate way of measuring response rate as the denominator of bereaved people in this sample was not ascertainable using routine data or survey methods.

The survey's inclusion criteria were respondents aged between 18 and 40 years who had experienced the sudden bereavement of a close friend or relative, with the loss occurring after they were aged ten years. Participants self-reported their exposure to bereavement by suicide, sudden natural causes (e.g. coronary death), or sudden un-natural causes (e.g., accidental death). The exclusion criteria were exposure to bereavement before the age of ten, in order to reduce recall bias and capture an adult level of cognitive processing of life events.

Of the total of 3,686 respondents to the UCL Bereavement Survey who consented to participate and specified mode of traumatic bereavement, we included in our secondary analysis the 3,167 (86%) individuals who had provided data on personality characteristics, suicidal ideation and suicide attempt (Figure 1).

Ethical approval

All participants provided online informed consent. The study protocol was approved by the UCL Research Ethics Committee in 2010 (reference: 1975/002).

Measures

The online survey questionnaire collected quantitative data on sociodemographic and clinical characteristics.

Exposure: personality disorder traits

Personality disorder traits were assessed using the self-report Standardized Assessment of Personality - Abbreviated Scale (SAPAS); an eight-item screen for identifying a probable diagnosis of personality disorder, based on DSM-IV-TR criteria (Germans, Van Heck, Moran, & Hodiament, 2008). These items cover: difficulty making and keeping friends; identifying as a loner; difficulty trusting other people; tendency to lose one's temper easily; tendency to impulsivity; tendency to be a worrier; tendency to depend on others; and tendency to be a perfectionist (see Box 1). The SAPAS was originally validated in a psychiatric population (Moran et al., 2003) and has also been validated for use in general population samples (Fok et al., 2015). Furthermore, other work has confirmed the clinical utility of the SAPAS, with SAPAS scores being prospectively linked with future functioning and clinical impairment, including general health status (Fok et al., 2014), response to psychological treatment (Crawford et al., 2009; Mars et al., 2020), and treatment response in depression (Gorwood et

al., 2010; Bukh et al., 2010). We used the population-derived cut point of four to indicate high probability of a diagnosis of personality disorder (Fok et al., 2015).

Outcomes: Suicidal behaviour

We measured self-reported suicide attempt and self-reported suicidal ideation since exposure to bereavement, captured using the survey questions “*Have you ever made an attempt to take your life, by taking an overdose of tablets or in some other way?*” and “*Have you ever thought of taking your life, even though you would not actually do it?*”. The wording of these questions was taken from the Adult Psychiatric Morbidity Survey (APMS), a regular national survey of the mental health of the general population in England, providing population norms (McManus, Meltzer, Brugha, Bebbington, & Jenkins, 2009). In the case of affirmative responses for either question, a qualifier determined whether the attempt/ideation had occurred before, after, or both before and after the bereavement.

Covariates

On *a priori* grounds, we defined a set of six covariates as potential confounders: gender (Hawton, 2000; Lippa, 2010; Trull, Jahng, Tomko, Wood, & Sher, 2010), age (Moran et al., 2012; Shah, 2007), socioeconomic status (Rehkopf & Buka, 2006; Jonassaint, Siegler, Barefoot, Edwards, & Williams, 2011), pre-bereavement suicidal and non-suicidal self-harm (Ribeiro et al., 2016; Whitlock et al., 2013; Brezo et al., 2006a; Ansell et al., 2015), pre-bereavement depression (Fergusson, Beautrais, & Horwood, 2003), and perceived social support (Tuisku et al., 2014). We measured these as follows:

- gender: using a self-report binary variable
- age: continuous measure
- socio-economic status: five category variable based on assigning own occupation (for staff) or the occupation of a parent or other source of financial support (for students)

using the National Statistics Socio-economic Classification (NS-SEC) of skill level and skill content, based on the Office for National Statistics Standard Occupational Classification 2010 (ONS, 2010), collapsing social classes 1.1 and 1.2 into category 1, and social classes 5 - 9 into category 5.

- pre-bereavement suicidal and non-suicidal self-harm: based on responses to the question on suicide attempt described above, and to a question on non-suicidal self-harm (“*Have you ever deliberately harmed yourself in any way but not with the intention of killing yourself?*”) (McManus, Meltzer, Brugha, Bebbington, & Jenkins, 2009), qualified to include only episodes occurring before the bereavement
- pre-bereavement depression, using the two item Composite International Diagnostic Interview (CIDI) screen for lifetime depression (Robins et al., 1988), qualified by whether the onset of these core symptoms of low mood and anhedonia occurred before or after the sudden bereavement, to derive a binary measure
- perceived social support, using a seven item scale taken from the 1987 Health and Lifestyle survey in England (Cox et al., 1987) and also used in the APMS, in which scores are summed to create a three category variable: no lack of perceived social support, moderate lack of perceived social support, and severe lack of perceived social support

We measured three putative modifiers of the association between the exposure and outcomes: gender, mode of bereavement (suicide *versus* non-suicide loss), and closeness to the deceased. We predicted that the magnitude of the association would be greater in females, the suicide-bereaved, and those most closely attached to the deceased. The latter two variables were measured as follows:

- mode of bereavement, ascertained based on responses to the question “*Since you were aged 10 have you experienced a sudden bereavement of someone close to you due to*

any of the following: (1) sudden natural death (eg. cardiac arrest, epileptic seizure, stroke); (2) sudden unnatural death (eg. road crash, murder or manslaughter, work accident); (3) suicide?". Those who had experienced bereavement by suicide as well as another bereavement by a different cause were classified as suicide-bereaved.

- closeness to the deceased, using an unvalidated Likert-type scale derived from previous bereavement research (Bailley, Kral, & Dunham, 1999), rated from 1 ('Not close at all') to 5 ('As close as any relationship I've had before or since') and dichotomized into 'quite close' (1-3) and 'very close' (4-5).

We also measured the following descriptive characteristics:

- post-bereavement suicidal thoughts: using a question from the standardized APMS survey based on the question "*Have you ever thought of taking your life, even if you would not really do it?*" (McManus, Meltzer, Brugha, Bebbington, & Jenkins, 2009), qualified by whether this was after the bereavement
- post-bereavement non-suicidal self-harm: using a question from the standardized APMS survey based on the question "*Have you ever deliberately harmed yourself in any way but not with the intention of killing yourself?*" (McManus, Meltzer, Brugha, Bebbington, & Jenkins, 2009), qualified by whether this was after the bereavement
- incident depression post-bereavement: using the two item Composite International Diagnostic Interview (CIDI) screen for lifetime depression (Robins et al., 1988) as above, qualified by whether the onset was after the sudden bereavement

Missing data for model covariates varied from <0.5% (for gender, pre-bereavement depression, pre-bereavement self-harm, pre-bereavement suicide attempt) to 3% (for socio-economic status).

Statistical Analysis

We described the sample's demographic and clinical characteristics by exposure group, using Chi-squared tests (categorical variables) and one-way analysis of variance (continuous variables). For descriptive purposes, we classified exposure in our sample using three categories: individuals scoring 0 traits on the SAPAS, those scoring 1 to 3 traits, and those scoring 4 or more traits (denoting the threshold for a possible diagnosis of personality disorder). However, for our main regressions, we used the SAPAS as a continuous measure.

We investigated the relationship between number of personality disorder traits (continuous measure) and our two binary outcomes using multivariable logistic regression models with HEI as random effect, to take into account any clustering effect at the HEI level. We adjusted models for the six confounders described above, adding three blocks sequentially for socio-demographic variables, clinical variables, and social support.

To assess whether gender, mode of bereavement (suicide *versus* non-suicide loss), or closeness to the deceased (very close *versus* quite close) modified the association between personality disorder traits and suicidality we specified separate statistical models with interactions between number of personality disorder traits and each of the putative effect modifiers.

To check for deviations from a linear association between number of personality traits and post-bereavement suicide attempt and ideation (May & Bigelow, 2006) we further included a quadratic term for exposure in relation to both outcomes.

We tested for a threshold effect of high probability of a diagnosis of personality disorder by adding an indicator variable to statistical models for each outcome using the standard population-derived cut point of four SAPAS personality traits (Fok et al, 2015) and used likelihood ratio tests to compare these models to nested models without this indicator variable.

We also tested for a threshold effect at adjacent cut points by repeating these analyses with the cut point as three and five SAPAS personality traits respectively.

As a final exploratory analysis, we ran logistic regressions to test for an association of each of the eight individual SAPAS personality disorder traits with outcomes (see Supplementary online table).

We used complete case analysis, such that participants with missing data on any of the covariates were excluded from final models. All analyses were conducted using STATA version 15 (StataCorp, 2017).

Results

Sample Characteristics

The majority of the sample endorsed one or more traits on the SAPAS (3,038/3,167; 96%), and 11% (347/3,167) endorsed one trait only. A third endorsed four or more traits, reaching the threshold for possible diagnosis of personality disorder (37%), whilst 59% (1,863/3,167) endorsed 1 to 3 traits (Table 1). The majority of the sample were female (81%), white (90%), and aged 23 or more. The highest educational attainment of most participants was A Level or equivalent or degree level qualifications. The majority had been bereaved due to sudden natural death (61%) and by the death of a blood relative (71%). A total of 208 participants (7%) had attempted suicide since the bereavement and 1,512 (48%) reported post-bereavement suicidal ideation. When comparing groups defined by number of SAPAS personality traits, there were no group differences in age, self-defined ethnicity, kinship, closeness to the deceased, or mode of bereavement. Participants endorsing four or more SAPAS personality traits had a higher prevalence of depression, of pre-bereavement non-suicidal self-harm, and of post-bereavement incident depression.

In descending order, the frequency of each SAPAS personality trait endorsed in this sample was: tendency to be a worrier (72%); tendency to being a perfectionist (63%); tendency to impulsivity (43%); difficulty trusting other people (33%); tendency to lose one's temper easily (28%); tendency to depend on others (28%); difficulty making and keeping friends (22%); and identifying as a loner (22%). The total number of personality traits endorsed on the SAPAS was normally distributed (Figure 2).

The association between personality traits endorsed on SAPAS and probability of suicidality after traumatic bereavement

We found evidence to support an association between number of traits endorsed on a screen for possible personality disorder and post-bereavement suicide attempt, both in unadjusted (odds ratio [OR]=1.48, 95% CI: 1.35 to 1.61; $p<0.0001$) and adjusted (adjusted odds ratio [AOR]=1.36; 95% CI: 1.23 to 1.49; $p<0.0001$) models (see Table 2). There was a linear relationship between number of personality disorder traits and the log odds of suicide attempt ($p=0.8$ for non-linearity), with each additional trait increasing the odds of post-bereavement suicide attempt by 48% (unadjusted) or 36% (adjusted model). When adding a quadratic term to our model for suicide attempt there was no evidence for deviation from linearity, whether in an unadjusted ($p=0.795$) or adjusted ($p=0.971$) model.

We also found evidence to support an association between number of personality disorder traits and post-bereavement suicidal ideation, both in unadjusted (OR=1.39, 95% CI: 1.32 to 1.46; $p<0.0001$) and adjusted (OR= 1.31, 95% CI: 1.25 to 1.38; $p<0.0001$) models (see Table 2). There was also a linear relationship between number of personality disorder traits and probability of suicidal thoughts ($p=0.2$ for non-linearity), with each additional trait increasing the odds of post-bereavement suicide ideation by 39% (unadjusted) or 31% (adjusted model). When adding a quadratic term to the model for suicidal ideation there was no evidence for deviation from linearity, whether in an unadjusted ($p=0.158$) or adjusted ($p=0.142$) model.

Effect modification

We did not find evidence that gender modified these adjusted associations, whether in relation to suicide attempt ($p=0.1$) or suicidal ideation ($p=0.5$). We also found no evidence that exposure to suicide bereavement (versus other losses) modified adjusted associations, whether in relation to suicide attempt ($p=0.3$) or suicidal ideation ($p=0.6$). Finally, we found no evidence that reporting emotional closeness to the deceased modified adjusted associations, either in relation to suicide attempt ($p=0.4107$) or suicidal ideation ($p=0.5221$).

Test for model fit

There was no evidence that the addition to final models of binary indicator variables denoting high probability of having a diagnosis of personality disorder (with number of SAPAS personality disorder traits as the main exposure) improved the fit of these models. This was the case when using binary variables based on the conventional cut point of four traits or more, and when using cut points of three and as five traits. These findings provided no evidence to support the existence of threshold effects.

Associations of individual personality disorder traits with outcomes

Our exploratory analysis of individual personality disorder traits with suicide attempt after sudden bereavement (see Supplementary Table) found that, when adjusted for the six covariates as above, only two of these personality traits (being a perfectionist and depending on others) were not associated with post-bereavement suicide attempt (see Supplementary Table). When this exploratory analysis was conducted with post-bereavement suicidal thoughts as the outcome, only one trait (being a perfectionist) was not associated with suicidal thoughts. The traits most strongly associated with risk of suicide attempt were: describing oneself as a loner (AOR=2.11, 95% CI: 1.54 to 2.90; $p<0.0001$), and difficulty in making or keeping friends (AOR=1.88, 95% CI: 1.36 to 2.59; $p<0.0001$). The traits most strongly associated with risk of

suicidal ideation were: describing oneself as a loner (AOR=1.97, 95% CI: 1.64 to 2.37, $p<0.0001$), and a tendency to lose one's temper (AOR=1.74, 95% CI: 1.12 to 1.51, $p<0.0001$).

Discussion

Main Findings

We found evidence to support a linear relationship between the number of personality disorder traits and probability of suicidal behaviour after an adverse life event. For each additional trait endorsed, odds of attempting suicide and of suicidal thoughts after sudden bereavement increased by 36% and 31% respectively. These associations held even after taking into account group differences in socio-demographic and clinical characteristics, and perceived social support after bereavement, and were not modified by gender, mode of bereavement, or closeness to the deceased. This last finding is surprising, given that sensitivities to abandonment are features of borderline personality disorder (WHO, 1993), and in view of its association with suicidality (Black, Blum, Pfohl, & Hale, 2004). Further exploratory work to describe the additive effect of specific combinations of traits suggestive of personality disorder would be clinically useful, but would require larger, longitudinal, representative samples.

Results in the context of other studies

To our knowledge only one previous study has investigated the association of severity of personality disorder traits with suicidality. This too was cross-sectional, and relied on a clinical sample of emergency department attenders, using a categorical exposure (no personality disorder, 'simple' personality disorder, 'diffuse' personality disorder), and found an association of severity of personality disorder with number of past suicide attempts but not severity or lethality of attempts (Blasco-Fontecill et al., 2009). However, we believe the current study to be the first to investigate the association between the number of traits suggestive of a diagnosis of personality disorder and suicidal behaviour, as well as to explore the modifying

effect of gender, closeness, and mode of bereavement. Other studies have tended to capture personality using a range of dimensions or sub-types in assessing associations with suicidality (Brezo, Paris, and Turecki, 2006a).

In relation to the differential associations of each personality disorder trait with suicidality, our exploratory findings were in keeping with work showing that the traits consistently associated with suicidal behaviours are neuroticism (Ortigo et al., 2009; Yen et al., 2009; Beautrais, Joyce, & Mulder, 1999) and impulsivity (Brezo et al., 2006b). Our finding that impulsivity was less strongly associated with suicidality than factors such as being a worrier is in keeping with longitudinal research finding that neuroticism is a stronger predictor of suicide attempt than disinhibition and impulsivity (Yen et al., 2009). Our finding of no association of perfectionism with suicidality conflicts with other work (Hewitt, Flett, & Turnball-Donovan, 1992; Smith et al., 2017) but as an exploratory analysis this may be a chance finding.

Strengths and limitations

We analyzed data from a large, UK-wide sample of 3,167 bereaved adults using a brief, validated personality measure, which reduced question burden on respondents compared to lengthy diagnostic assessments. We tested specific hypotheses reflecting gaps in the personality disorder literature and a focus on young adults, and our statistical models were adjusted for pre-selected potential confounders. In view of the survey's sampling methods, we acknowledge the potential for male non-response bias, and selection bias of white, highly educated adults from higher socio-economic groups. Our use of a validated brief screen for personality disorder traits allowed us to gather data on a large population-based sample. However, by virtue of its brevity, use of a personality disorder screen may have introduced some measurement error. We also acknowledge that perfectionism is a common trait among students (Halgin & Leahy, 1989), and the prevalence of perfectionism is likely to be higher

than in non-HEI samples. Our inclusion of survey participants who had complete data on bereavement type, personality traits, suicidality, and all covariates meant that our sample may not be representative of all those exposed to sudden bereavement. These factors, and the survey's age restrictions, suggest that the study's findings may be generalizable only to highly educated young white women in the UK. This is underlined by the rate of suicide in the UK student population, 4.7 deaths per 100,000 HEI students, (ONS, 2018) being significantly lower than that for non-student peers of the same age (Gunnell, Caul, Appleby, John, & Hawton, 2020). Our focus on exposure to sudden bereavement is also not representative of all adverse life events in this age group.

Our measures of suicidality and depression were potentially subject to recall bias, and that used to measure pre-bereavement depression was a brief screening tool, which may have resulted in over- or under-adjustment for past depression in multivariable models. We used a standardised measure of suicidal ideation, with population norms, but this did not specify the degree of planning or intent inherent to the suicidal thoughts. As this was a cross-sectional study, it was not possible to definitively ascertain the temporal sequence of outcomes, including whether suicidal behaviour following bereavement had preceded the emergence of personality disorder traits. The dataset also lacked measures of the number of attempts or episodes of suicidal intent, or the intentions associated with them. Finally, as this was a secondary analysis of existing data, we were unable to control for a number of important unmeasured potential confounders. These include coping styles and problem-solving skills (Connor-Smith & Flachsbart, 2007), which may be associated with negative outcomes after bereavement (Stroebe et al, 2007), and resilience (Nrugham, Holen, & Sund, 2010).

Clinical and research implications

Our findings from a sample of young adults suggest that young bereaved adults with personality difficulties are a vulnerable population. We know that young people have a

tendency not to seek help when in distress (Biddle, Donovan, Sharp, & Gunnell, 2007) and that those who attempt suicide are more likely to have persistent mental and physical health and social problems (Goldman-Mellor et al., 2014). Young adults are a priority group within suicide prevention strategies due to concerns about their risk of suicide (Department of Health, 2012). Our findings suggest that, following bereavement, young people endorsing a greater number of personality disorder traits fare worse in terms of their future mental health, with higher rates of incident depression and incident suicidality. This finding is in keeping with other longitudinal studies (Moran et al., 2016), and highlights a need for practitioners to tailor bereavement or crisis support packages to those with a greater degree of underlying personality difficulties, as they are a high-risk population. It may be important to profile the personality characteristics of people who experience sudden bereavement, to identify those with personality difficulties and the associated risk of suicidality. Improving our clinical response to such markers of risk in the bereaved population is important because of the established association of sudden bereavement with risk of suicidality and psychiatric morbidity (Ajdacic-Gross et al., 2008; Daniel, Goldston, Erkanli, Heilbron, & Franklin, 2017; Guldin et al., 2017; Qin & Mortensen, 2003), particularly in young people, and after suicide loss (Erlangsen et al., 2017; Guldin et al., 2017; Hawton & Rodham, 2006; Pitman, Osborn, King, & Erlangsen, 2014).

Very little research has explored how personality factors influence adaptation to a life event such as bereavement, despite this being likely to influence coping style, depending on the nature of attachment to the deceased (Stroebe, Schut, & Stroebe, 2007). It is possible that a greater range of personality difficulties hampers ability to cope with grief, and that this discourages others from offering support. It is also possible that loss of a friend or relative (regardless of how close they were) induces emotional dysregulation through abandonment. All of these factors might increase the risk of suicidality, particularly in the context of feeling

lonely or set apart from one's peers. However, to develop these hypotheses there is a need for qualitative work to enrich our understanding of how people with traits suggestive of personality disorder experience sudden loss. Furthermore, there is a need for longitudinal studies to investigate the course of suicidality following bereavement and how this may vary in relation to interpersonal difficulties.

Conclusion

This analysis of British survey data on people who have experienced sudden bereavement found a linear association between number of personality disorder traits and probability of suicidal ideation and suicide attempt after a negative life event. This association was not modified by gender, cause of death, nor degree of closeness to the deceased. Our findings suggest that personality assessment may be clinically useful for bereavement counsellors, crisis teams, and general practitioners after a traumatic life event to gain a sense of a bereaved person's risk of suicidality and the way they respond to others, in order to help tailor appropriate support.

Table 1: Sample characteristics in relation to number of personality traits measured on SAPAS

Characteristics	0 personality traits on SAPAS (n= 129, 4%)	1-3 personality traits on SAPAS (n= 1863, 59%)	4 or more personality traits on SAPAS (n= 1175, 37%)	Total participants (n=3167)	P-value
Demographic and social variables					
Gender					
Female	91 (71)	1511 (81)	977 (83)	2579 (81)	0.01
Male	38 (29)	351 (19)	198 (17)	587 (18)	
Missing	0	1 (<1)	0	1 (<1)	
Age (years); mean (SD)	26.50 (7)	25.14 (6)	24.81 (6)	25.07 (6)	0.3
Ethnicity					
White	118 (91)	1671 (90)	1064 (91)	2853 (90)	0.65
Non-White	11 (9)	189 (10)	111 (9)	311 (10)	
Missing	0	3 (<1)	0	3 (<1)	
Highest educational attainment					
No academic qualification	0	1 (<1)	2 (<1)	3 (<1)	<0.0001
Maximum GCSE level	4 (3)	31 (2)	15 (1)	50 (2)	
Maximum A level	43 (33)	728 (39)	549 (47)	1320 (42)	
Maximum university degree	50 (38)	695 (37)	409 (35)	1153 (36)	
Maximum post-graduate	32 (25)	402 (22)	200 (17)	634 (20)	
Socioeconomic status					
Social class 1 (least deprived)	43 (33)	578 (31)	303 (26)	924 (29)	0.004
Social class 2	41 (32)	607 (33)	385 (33)	1033 (33)	
Social class 3	14 (11)	222 (12)	139 (12)	375 (12)	
Social class 4	9 (7)	93 (5)	47 (4)	149 (5)	
Social class 5 (most deprived)	18 (14)	311 (17)	265 (23)	594 (19)	
Missing	4 (3)	52 (3)	36 (3)	92 (3)	

Work status	Working	37 (29)	410 (22)	197 (17)	644 (20)	0.001
	Studying	51 (40)	931 (50)	604 (51)	1586 (50)	
	Both working & studying	40 (31)	515 (28)	365 (31)	920 (29)	
	Neither	1 (1)	7 (<1)	9 (1)	17 (1)	
Social support	No lack of perceived support	98 (76)	1197 (64)	542 (46)	1837 (58)	<0.0001
	Moderate lack of perceived support	25 (19)	452 (24)	369 (31)	846 (27)	
	Severe lack of perceived support	6 (5)	214 (11)	264 (22)	484 (15)	
Clinical variables						
Depression prior to loss	Yes	6 (5)	294 (16)	323 (28)	626 (20)	<0.0001
	Missing	0	2 (<1)	2 (<1)	4 (<1)	
Non-suicidal self-harm prior to loss	Yes	16 (12)	338 (18)	358 (30)	712 (22)	<0.0001
	Missing	0	6 (<1)	1 (<1)	7 (<1)	
Suicide attempts after bereavement	Yes	0	80 (4)	128 (11)	208 (7)	<0.0001
	Missing	0	0	0	0	
Suicidal thoughts after bereavement	Yes	39 (23)	748 (40)	725 (62)	1512 (48)	<0.0001
	Missing	0	0	0	0	
Non-suicidal self-harm after bereavement	Yes	17 (13)	344 (18)	368 (31)	729 (23)	<0.0001
	Missing	0	6 (<1)	2 (<1)	8 (<1)	
Incident depression after bereavement	Yes	30 (23)	545 (29)	460 (39)	1035 (33)	<0.001
	Missing	0	2 (<1)	2 (<1)	4 (<1)	
Bereavement variables						
Closeness to the deceased	Very close	98 (76)	1379 (74)	865 (74)	2342 (74)	0.07
	Quite close	28 (22)	476 (25)	303 (26)	807 (25)	
	Missing	3 (2)	8 (0.5)	7 (0.5)	18 (0.5)	

Kinship to the deceased					
Blood relative	91 (71)	1308 (70)	854 (73)	2253 (71)	0.54
Non-blood relative	38 (29)	547 (29)	318 (28)	903 (28)	
Missing	0	8 (0.5)	3 (0.2)	11 (0.5)	
Type of bereavement					
Natural sudden death	76 (59)	1129 (61)	733 (62)	1938 (61)	0.28
Unnatural sudden death	30 (23)	411 (22)	222 (19)	663 (21)	
Suicide	23 (18)	323 (17)	220 (19)	566 (18)	

Key:

All statistics are presented as n (%) unless otherwise stated. P values are for group comparisons excluding missing values, using a two-sided significance threshold of $p=0.05$.

SAPAS = Standardised Assessment of Personality – Abbreviated Scale

Table 2. Estimates of the association between number of traits measured using SAPAS and post-bereavement suicidality

Outcome	Odds ratio (95% CI)	P-value
Post-bereavement suicide attempt		
Unadjusted model	1.48 (1.35 - 1.60)	<0.0001
Adjusted for age, gender, socio-economic status	1.47 (1.34 - 1.60)	<0.0001
Adjusted for age, gender, socio-economic status; pre-bereavement depression, pre-bereavement (suicidal and non-suicidal) self-harm	1.42 (1.29 - 1.55)	<0.0001
Final model: Adjusted for above variables and perceived social support	1.36 (1.23 - 1.49)	<0.0001
Post-bereavement suicidal ideation		
Unadjusted model	1.39 (1.32 - 1.46)	<0.0001
Adjusted for age, gender, socio-economic status	1.38 (1.32 - 1.45)	<0.0001
Adjusted for age, gender, socio-economic status; pre-bereavement depression, pre-bereavement (suicidal and non-suicidal) self-harm	1.36 (1.30 - 1.43)	<0.0001
Final model: Adjusted for above variables and perceived social support	1.31 (1.25 - 1.38)	<0.0001

Key:

All models used complete case analysis of n=3,065 individuals

Declaration of interest: All authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank all the HEIs from England, Wales, Northern Ireland, and Scotland that consented to participate, listed below, and all the bereaved individuals who took time to respond to the online survey.

Participating HEIs: Bishop Grosseteste University; Bournemouth University; Royal Central School of Speech and Drama; City University; Cranfield University; Courtauld Institute; De Montfort University; University of Greenwich; King's College London; Liverpool Institute for Performing Arts; Liverpool John Moores University; London Metropolitan University; Norwich University of the Arts; Royal Veterinary College; School of Oriental and African Studies; St George's, University of London; Staffordshire University; Trinity Laban Conservatoire of Music and Dance; UCL; University of Suffolk; University of Bedfordshire; University of Chester; University of Cumbria; University of Leeds; University of Liverpool; University of Oxford; University of Southampton; University of Worcester; University of Westminster; Queen Margaret University; Heriot-Watt University; Scottish Agricultural College (now part of Scotland's Rural University College); University of Dundee; Cardiff University; Cardiff Metropolitan University (formerly University of Wales Institute Cardiff); Queen's University Belfast; University of Ulster.

Financial Support

The UCL Bereavement Study survey was funded by a Medical Research Council Population Health Scientist Fellowship Award to AP (G0802441). AP, MK and DO are supported by the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) at University College London Hospitals (UCLH). DO is also part supported by the NIHR

Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Bart's Health NHS Trust. PM is part-funded by the NIHR Biomedical Research Centre at University Hospitals Bristol NHS Foundation Trust and the University of Bristol. The funders had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, and approval of the manuscript. The views expressed are those of the authors and not necessarily those of the funders.

Ethical standards

The author asserts that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Ajdacic-Gross, V., Ring, M., Gadola, E., Lauber, C., Bopp, M., Gutzwiller, F., & Rössler, W. (2008). Suicide after bereavement: An overlooked problem. *Psychological Medicine*, 38(5), 673-676. doi:10.1017/S0033291708002754
- Ansell, E. B., Wright, A. G. C., Markowitz, J. C., Sanislow, C. A., Hopwood, C. J., Zannarini, M. C., Yen, S., Pinto, A., McGlashan, T.H., Grilo, C. M. (2015). Personality disorder risk factors for suicide attempts over 10 years of follow-up. *Personality Disorders: Theory, Research, and Treatment*, 6(2), 161-167. doi:10.1037/per0000089
- Bailey, S. E., Kral, M. J. and Dunham, K. (1999). Survivors of Suicide Do Grieve Differently: Empirical Support for a Common Sense Proposition. *Suicide and Life-Threatening Behavior*, 29: 256-271. doi:10.1111/j.1943-278X.1999.tb00301.x
- Beautrais, A.L., Joyce, P.R., Mulder, R.T. (1999). Personality Traits and Cognitive Styles as Risk Factors for Serious Suicide Attempts among Young People. *Suicide and Life-Threatening Behaviour*, 29(1), 37-47. doi:10.1111/j.1943-278X.1999.tb00761.x
- Beautrais, A. L. (2000). Risk Factors for Suicide and Attempted Suicide among Young People. *Australian & New Zealand Journal of Psychiatry*, 34(3), 420–436. doi:10.1080/j.1440-1614.2000.00691.x
- Beck, A. T., Brown, G., & Steer, R. A. (1989). Prediction of eventual suicide in psychiatric inpatients by clinical ratings of hopelessness. *Journal of Consulting and Clinical Psychology*, 57(2), 309–310. Doi: 10.1037/0022-006X.57.2.309
- Biddle, L., Donovan, J., Sharp, D. and Gunnell, D. (2007). Explaining non-help-seeking amongst young adults with mental distress: a dynamic interpretive model of illness behaviour. *Sociology of Health & Illness*, 29: 983-1002. doi:10.1111/j.1467-9566.2007.01030.x
- Bilsen J. (2018). Suicide and Youth: Risk Factors. *Frontiers in psychiatry*, 9, 540. doi:10.3389/fpsyt.2018.00540
- Black, D.W., Blum, N., Pfohl, B., Hale, N. (2004). Suicidal Behaviour in Borderline Personality Disorder: Prevalence, Risk Factors, Prediction, and Prevention. *Journal of Personality Disorders*, 18(Suicide and Borderline Personality Disorder), 226-239. doi: 10.1521/pedi.18.3.226.35445
- Blasco-Fontecilla, H., Baca-Garcia, E., Devic, K., Perez-Rodriguez, M.M., Saiz-Gonzalez, M.D., Saiz-Ruiz, J., Oquendo, M.A., de Leon, J. (2009). Severity of personality disorders and suicide attempt. *Acta Psychiatrica Scandinavica*, 119(2), 149-155. doi:10.1111/j.1600-0447.2008.01284.x
- Brezo, J., Paris, J., Turecki, G. (2006a). Personality traits as correlates of suicidal ideation, suicide attempts, and suicide completions: a systematic review. *Acta Psychiatrica Scndinavica*, 113(3), 180-206. doi: 10.1111/j.1600-0447.2005.00702.x
- Brezo, J., Paris, J., Tremblay, R., Vitaro, F., Zoccolillo, M., Hébert, M., & Turecki, G. (2006b). Personality traits as correlates of suicide attempts and suicidal ideation in young adults. *Psychological Medicine*, 36(2), 191-202. doi:10.1017/S0033291705006719

- Bukh, J. D., Bock, C., Vinberg, M., Gether, U., & Kessing, L. V. (2010). Clinical utility of Standardised Assessment of Personality - Abbreviated Scale (SAPAS) among patients with first episode depression. *Journal of affective disorders*, 127(1-3), 199–202. doi:10.1016/j.jad.2010.05.023
- Cash, S. J., & Bridge, J. A. (2009). Epidemiology of youth suicide and suicidal behaviour. *Current opinion in Pediatrics*, 21(5), 613–619. doi:10.1097/MOP.0b013e32833063e1
- Cavanagh, J.T., Carson, A.J., Sharpe, M., Lawrie, S.M. (2003). Psychological autopsy studies of suicide: a systematic review. *Psychological Medicine*, 33(3), 395-405. doi:10.1017/S0033291702006943
- Connor-Smith, J.K., Flachsbart, C. (2007). Relations between personality and coping: A meta-analysis. *Journal of Personality and Social Psychology*, 93(6), 1080-1107. doi: 10.1037/0022-3514.93.6.1080
- Cox, B. D., Blaxter, M., Buckle, A. L. J., Fenner, N. P., Golding, J. F., Gore, M., ... & Wadsworth, M. E. J. (1987). The health and lifestyle survey. Preliminary report of a nationwide survey of the physical and mental health, attitudes and lifestyle of a random sample of 9,003 British adults. *Health Promotion Research Trust*.
- Crawford, M. J., Price, K., Gordon, F., Jossion, M., Taylor, B., Bateman, A., Fonagy, P., Tyrer, P., & Moran, P. (2009). Engagement and retention in specialist services for people with personality disorder. *Acta psychiatrica Scandinavica*, 119(4), 304–311. doi:10.1111/j.1600-0447.2008.01306.x
- Daniel, S. S., Goldston, D. B., Erkanli, A., Heilbron, N., & Franklin, J. C. (2017). Prospective Study of Major Loss Life Events and Risk for Suicidal Thoughts and Behaviours Among Adolescents and Young Adults. *Suicide & life-threatening behavior*, 47(4), 436–449. doi:10.1111/sltb.12305
- Department of Health. (2012) Preventing Suicide in England: A cross-government outcomes strategy to save lives. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/430720/Preventing-Suicide-.pdf
- Erlangsen, A., Runeson, B., Bolton, J. M., Wilcox, H. C., Forman, J. L., Krogh, J., Shear, M.K., Nordentoft, M., Conwell, Y. (2017). Association Between Spousal Suicide and Mental, Physical, and Social Health Outcomes: A Longitudinal and Nationwide Register-Based Study. *JAMA psychiatry*, 74(5), 456–464. doi:10.1001/jamapsychiatry.2017.0226
- Fergusson, D., Beautrais, A., & Horwood, L. (2003). Vulnerability and resiliency to suicidal behaviours in young people. *Psychological Medicine*, 33(1), 61-73. doi:10.1017/S0033291702006748
- Fok, M., Hotopf, M., Stewart, R., Hatch, S., Hayes, R., & Moran, P. (2014). Personality disorder and self-rated health: a population-based cross-sectional survey. *Journal of personality disorders*, 28(3), 319–333. doi:10.1521/pedi_2013_27_119

- Fok, M. L., Seegobin, S., Frissa, S., Hatch, S. L., Hotopf, M., Hayes, R. D., & Moran, P. (2015). Validation of the standardised assessment of personality--abbreviated scale in a general population sample. *Personality and mental health*, 9(4), 250–257. doi:10.1002/pmh.1307
- Germans, S., Van Heck, G.L., Moran, P., Hodiament, P.G. (2008). The Self-report standardized Assessment of Personality-abbreviated Scale: Preliminary results of a brief screening test for personality disorders. *Personality and Mental Health*, 2(2), 70-76. doi:10.1002/pmh.34
- Goldman-Mellor, S. J., Caspi, A., Harrington, H., Hogan, S., Nada-Raja, S., Poulton, R., & Moffitt, T. E. (2014). Suicide attempt in young people: a signal for long-term health care and social needs. *JAMA psychiatry*, 71(2), 119–127. doi:10.1001/jamapsychiatry.2013.2803
- Gorwood, P., Rouillon, F., Even, C., Falissard, B., Corruble, E., & Moran, P. (2010). Treatment response in major depression: effects of personality dysfunction and prior depression. *The British journal of psychiatry: the journal of mental science*, 196(2), 139–142. doi:10.1192/bjp.bp.109.067058
- Guldin, M. B., Ina Siegismund Kjaersgaard, M., Fenger-Grøn, M., Thorlund Parner, E., Li, J., Prior, A., & Vestergaard, M. (2017). Risk of suicide, deliberate self-harm and psychiatric illness after the loss of a close relative: A nationwide cohort study. *World psychiatry: official journal of the World Psychiatric Association (WPA)*, 16(2), 193–199. doi:10.1002/wps.20422
- Gunnell, D., Caul, S., Appleby, L., John, A., & Hawton, K. (2020). The incidence of suicide in University students in England and Wales 2000/2001-2016/2017: Record linkage study. *Journal of affective disorders*, 261, 113–120. doi:10.1016/j.jad.2019.09.079
- Halgin, R.P., Leahy, P.M. (1989). Understanding and Treating Perfectionistic College Students. *Journal of Counselling and Development*, 68(2), 222-225. doi:10.1002/j.1556-6676.1989.tb01362.x
- Hawton, K. (2000). Sex and suicide: Gender differences in suicidal behaviour. *British Journal of Psychiatry*, 177(6), 484-485. doi:10.1192/bjp.177.6.484
- Hawton, K., Rodham, K., & Evans, E. (2006). *By Their Own Young Hand: Deliberate Self-Harm and Suicidal Ideas in Adolescents*. London: Jessica Kingsley Publishers.
- Hewitt, P.L., Flett, G.L., Turnbull-Donovan, W. (1992). Perfectionism and suicide potential. *British Journal of Clinical Psychology*, 31(2), 181-190. doi: 10.1111/j.2044-8260.1992.tb00982.x
- Jonassaint, C. R., Siegler, I. C., Barefoot, J. C., Edwards, C. L., & Williams, R. B. (2011). Low life course socioeconomic status (SES) is associated with negative NEO PI-R personality patterns. *International Journal of Behavioral Medicine*, 18(1), 13–21. doi:10.1007/s12529-009-9069-x.
- Lippa, R.A. (2010). Sex Differences in Personality Traits and Gender-Related Occupational Preferences across 53 Nations: Testing Evolutionary and Social-Environmental Theories. *Archives of Sexual Behavior*, 39(3), 619-636. doi:10.1007/s10508-008-9380-7.

Mars, B., Gibson, J., Dunn, B. D., Gordon, C., Heron, J., Kessler, D., Wiles, N., & Moran, P. (2021). Personality difficulties and response to community-based psychological treatment for anxiety and depression. *Journal of affective disorders*, 279, 266–273. doi:10.1016/j.jad.2020.09.115

May, S., & Bigelow, C. (2006). Modeling nonlinear dose-response relationships in epidemiologic studies: statistical approaches and practical challenges. *Dose-response : a publication of International Hormesis Society*, 3(4), 474–490. doi:10.2203/dose-response.003.04.004

McGirr, A., Renaud, J., Bureau, A., Seguin, M., Lesage, A., & Turecki, G. (2008). Impulsive-aggressive behaviours and completed suicide across the life cycle: a predisposition for younger age of suicide. *Psychological medicine*, 38(3), 407–417. doi:10.1017/S0033291707001419

McManus, S., Meltzer, H., Brugha, T., Bebbington, P., & Jenkins, R. (2009) Adult psychiatric morbidity in England - 2007. Results of a household survey. *National Centre for Social Research*. <https://digital.nhs.uk/data-and-information/publications/statistical/adult-psychiatric-morbidity-survey/adult-psychiatric-morbidity-in-england-2007-results-of-a-household-survey>

Moran, P., Leese, M., Lee, T., Walters, P., Thornicroft, G., & Mann, A. (2003). Standardised Assessment of Personality – Abbreviated Scale (SAPAS): Preliminary validation of a brief screen for personality disorder. *British Journal of Psychiatry*, 183(3), 228–232. doi:10.1192/bjp.183.3.228.

Moran, P., Coffey, C., Romaniuk, H., Olsson, C., Borschmann, R., Carlin, J. B., & Patton, G. C. (2012). The natural history of self-harm from adolescence to young adulthood: a population-based cohort study. *Lancet (London, England)*, 379(9812), 236–243. doi:10.1016/S0140-6736(11)61141-0

Moran, P., Romaniuk, H., Coffey, C., Chanen, A., Degenhardt, L., Borschmann, R., & Patton, G. C. (2016). The influence of personality disorder on the future mental health and social adjustment of young adults: a population-based, longitudinal cohort study. *The lancet. Psychiatry*, 3(7), 636–645. doi:10.1016/S2215-0366(16)30029-3

Nock, M., Borges, G., Bromet, E., Alonso, J., Angermeyer, M., Beautrais, A., Bruffaerts, R., Chiu, W.T., de Girolamo, G., Gluzman, S., de Graaf, R., Gureje, O., Haro, J.M., Huang, Y., Karam, E., Kessler, R.C., Lepine, J.P., Levinson, D., Medina-Mora, M.E., Ono, Y., Posada-Villa, J., Williams, D. (2008). Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *British Journal of Psychiatry*, 192(2), 98–105. doi:10.1192/bjp.bp.107.040113

Nrugham, L., Holen, A., Sund, A.M. (2010). Associations between attempted suicide, violent life events, depressive symptoms, and resilience in adolescents and young adults. *The Journal of Nervous and Mental Disease*, 198(2), 131–136. doi: 10.1097/NMD.0b013e3181cc43a2

Office for National Statistics. (2010). Standard Occupational Classification 2010 Volume 3 The National Statistics Socio-economic classification: (Rebased on the SOC2010) User Manual. HMSO. <https://www.ons.gov.uk/file?uri=/methodology/classificationsandstandards/standardoccupatio>

nalclassificationsoc/soc2010/soc2010volume3thenationalstatistics socioeconomicclassificationnssecrebasedonsoc2010/soc2010vol31amendedjanuary2013tcm77179133.pdf

Office for National Statistics. (2016). Suicides in the UK. Retrieved July 20, 2019, from <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/suicidesintheunitedkingdom/2016registrations#suicides-in-the-uk-by-age>

Office for National Statistics. (2018). Estimating suicide among higher education students, England and Wales: Experimental Statistics. Retrieved August 7, 2019, from <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/estimating-suicide-among-higher-education-students-england-and-wales-experimental-statistics/2018-06-25>

Ortigo, K. M., Westen, D., & Bradley, B. (2009). Personality subtypes of suicidal adults. *The Journal of Nervous and Mental Disease*, 197(9), 687–694. doi:10.1097/NMD.0b013e3181b3b13f

Pitman, A., Osborn, D., King, M., & Erlangsen, A. (2014). Effects of suicide bereavement on mental health and suicide risk. *The Lancet Psychiatry*, 1(1), 86-94. doi: 10.1016/S2215-0366(14)70224-X

Pitman, A., Osborn, D., & King, M. (2015). The use of internet-mediated cross-sectional studies in mental health research. *BJPsych Advances*, 21(3), 175-184. doi:10.1192/apt.bp.114.012948

Pitman, A.L., Osborn, D.P.J., Rantell, K., King, M.B. (2016). Bereavement by suicide as a risk factor for suicide attempt: a cross-sectional national UK-wide study of 3432 young bereaved adults. *BMJ Open*, 6(1), e009948. doi:10.1136/bmjopen-2015-009948

Qin, P., Mortensen, P.B. (2003). The impact of parental status on the risk of completed suicide. *Archives of general psychiatry*, 60(8), 797-802. doi:10.1001/archpsyc.60.8.797

Rehkopf, D.H., Buka, S.L. (2006). The association between suicide and the socio-economic characteristics of geographical areas: a systematic review. *Psychological Medicine*, 36(2), 145-157. doi:10.1017/S003329170500588X.

Ribeiro, J. D., Franklin, J. C., Fox, K. R., Bentley, K. H., Kleiman, E. M., Chang, B. P., & Nock, M. K. (2016). Self-injurious thoughts and behaviours as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychological medicine*, 46(2), 225–236. doi:10.1017/S0033291715001804

Robins, L. N., Wing, J., Wittchen, H. U., Helzer, J. E., Babor, T. F., Burke, J., Farmer, A., Jablenski, A., Pickens, R., Regier, D. A. (1988). The Composite International Diagnostic Interview. An epidemiologic Instrument suitable for use in conjunction with different diagnostic systems and in different cultures. *Archives of general psychiatry*, 45(12), 1069–1077. doi:10.1001/archpsyc.1988.01800360017003

Shah, A. (2007). The relationship between suicide rates and age: an analysis of multinational data from the World Health Organization. *International Psychogeriatrics*, 19(6), 1141-1152.

Smith, M.M., Sherry, S.B., Chen, S., Saklofske, D.H., Mushquash, C., Flett, G.L., Hewitt, P.L. (2017). The perniciousness of perfectionism: A meta-analytic review of the

perfectionism-suicide relationship. *Journal of Personality*, 86(3), 522-542. doi: 10.1111/jopy.12333

StataCorp. (2017). *Stata Statistical Software: Release 15*. College Station, TX: StataCorp LLC.

Stroebe, M., Schut, H., Stroebe, W. (2007). Health outcomes of bereavement. *The Lancet*, 370(9603), 1960-1973. doi: 10.1016/S0140-6736(07)61816-9

Trull, T.J., Jahng, S., Tomko, R.L., Wood, P.K., Sher, K.J. (2010). Revised NESARC Personality Disorder Diagnoses: Gender, Prevalence, and Comorbidity with Substance Dependence Disorders. *Journal of Personality Disorders*, 24(4), 412-426. doi:10.1521/pedi.2010.24.4.412.

Tuisku, V., Kiviruusu, O., Pelkonen M., Karlsson, L., Strandholm, T., Marttunen, M. (2014). Depressed adolescents as young adults- Predictors of suicide attempt and non-suicidal self-injury during an 8-year follow-up. *Journal of Affective Disorders*, 152-154, 313-319. doi: 10.1016/j.jad.2013.09.031

Turecki, G., & Brent, D. A. (2016). Suicide and suicidal behaviour. *Lancet*, 387(10024), 1227–1239. doi:10.1016/S0140-6736(15)00234-2

Whitlock, J., Muehlenkamp, J., Eckenrode, J., Purington, A., Baral Abrams, G., Barreira, P., Kress, V. (2013). Non-suicidal self-injury as a gateway to suicide in young adults. *Journal of Adolescent Health*, 52(4), 486-92. doi: 10.1016/j.jadohealth.2012.09.010

Winsper, C., Bilgin, A., Thompson, A., Marwaha, S., Chanen, A., Singh, S., Wang, A., Furtado, V. (2020). The prevalence of personality disorders in the community: A global systematic review and meta-analysis. *The British Journal of Psychiatry*, 216(2), 69-78. doi:10.1192/bjp.2019.166

World Health Organization. (1993) The ICD-10 Classification of Mental and Behavioural Disorders. Diagnostic criteria for research. <https://www.who.int/classifications/icd/en/GRNBOOK.pdf>

World Health Organization. (2016). Suicide rates. Retrieved July 20, 2019, from https://www.who.int/gho/mental_health/suicide_rates/en/

World Health Organization. (2019). Suicide. Retrieved December 12, 2019, from <https://www.who.int/news-room/fact-sheets/detail/suicide>

Yen, S., Shea, M. T., Sanislow, C. A., Skodol, A. E., Grilo, C. M., Edelen, M. O., Stout, R.L., Morey, L.C., Zanarini, M.C., Markowitz, J.C., McGlashan, T.H., Daversa, M.T., Gunderson, J. G. (2009). Personality traits as prospective predictors of suicide attempts. *Acta psychiatrica Scandinavica*, 120(3), 222–229. doi:10.1111/j.1600-0447.2009.013