Harroud, A., Manousaki, D., Butler-Laporte, G., Mitchell, R. E., Davey Smith, G., Richards, J. B., & Baranzini, S. E. (2021). The relative contributions of obesity, vitamin D, leptin and adiponectin to multiple sclerosis risk: a Mendelian randomization mediation analysis. *Multiple Sclerosis Journal*, 27(13), 1994-2000.
<https://doi.org/10.1177/1352458521995484>

Peer reviewed version

Link to published version (if available):
[10.1177/1352458521995484](https://doi.org/10.1177/1352458521995484)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via SAGE Publications at https://journals.sagepub.com/doi/10.1177/1352458521995484?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%20%20pubmed . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The relative contributions of obesity, vitamin D, leptin and adiponectin to multiple sclerosis risk: a Mendelian randomization mediation analysis

Adil Harroud, MD^{1,2}; Despoina Manousaki, MD, PhD^{3,4}; Guillaume Butler-Laporte, MD^{3,5}; Ruth E. Mitchell, PhD^{6,7}; George Davey Smith, DSc^{6,7}; J. Brent Richards, MD, MSc^{3,4,5,8,9,10}; Sergio E. Baranzini, PhD^{1,2,11,12}

- ¹ Department of Neurology, University of California San Francisco, San Francisco, CA, USA.
- ² Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA, USA.
- ³ Centre for Clinical Epidemiology, Department of Epidemiology, Lady Davis Institute for Medical Research, Jewish General Hospital, McGill University, Montreal, Quebec, Canada.
- ⁴ Department of Human Genetics, McGill University, Montreal, Quebec, Canada.
- ⁵ Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Quebec, Canada.
- ⁶ MRC Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Bristol, United Kingdom.
- ⁷ Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom.
- ⁸ Department of Medicine, McGill University Montreal, Quebec, Canada.
- ⁹ Department of Epidemiology, Biostatistics, Occupational Health, McGill University, Montreal, Quebec, Canada.
- ¹⁰ Department of Twin Research and Genetic Epidemiology, King's College London, London, United Kingdom.
- ¹¹ Institute for Human Genetics, University of California San Francisco, San Francisco, CA, USA.
- ¹² Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, CA, USA.

Corresponding author:

Sergio E. Baranzini, Department of Neurology, Weill Institute of Neuroscience, University of California San Francisco, 675 Nelson Rising Lane, San Francisco, CA 94158, USA

Email: sergio.baranzini@ucsf.edu

Tel: 415-502-6865; Fax: 415-476-5229

Keywords: Multiple sclerosis, Mendelian randomization, obesity, vitamin D, genetic epidemiology

Abstract word count: ~~204~~199

Manuscript word count: ~~1,718~~2,285

Number of figures: 2

Number of tables: 1

Number of references: ~~27~~36

Supplementary File: 1 (Supplementary Tables S1-~~3~~8)

Abstract

Background: Obesity is associated with increased risk of multiple sclerosis (MS); however, the underlying mechanisms remain unclear.

Objective: To determine the extent to which decreased vitamin D bioavailability and altered levels of adiponectin and leptin mediate the association between obesity and MS.

Methods: We performed Mendelian randomization (MR) analyses to estimate the effects on MS of body mass index (BMI), 25-hydroxyvitamin D (25OHD), adiponectin and leptin levels in a cohort of 14,802 MS cases and 26,703 controls. We then estimated the proportion of the effect of obesity on MS explained by these potential mediators.

Results: Genetic predisposition to higher BMI was associated with increased MS risk (OR=1.~~3340~~ per SD, 95%-CI 1.~~0916~~-1.6~~37~~), while higher 25OHD levels reduced odds of MS (OR=0.72 per SD, 95%-CI 0.60-0.87). In contrast, we observed no effect of adiponectin or leptin. In MR mediation analysis, 5.~~24~~% of the association between BMI and MS was attributed to obesity lowering 25OHD levels (95%-CI 0.~~34~~%-~~to 30.531.0~~%).

Conclusions: This study found that a minority of the increased risk of MS conferred by obesity is mediated by lowered vitamin D levels, while leptin and adiponectin had no effect. Consequently, vitamin D supplementation would only modestly reverse the effect of obesity on MS.

Introduction

An increasing body of evidence supports a role for obesity in the development of multiple sclerosis (MS).^{1, 2} However, the mechanisms underlying this association remain unclear. A commonly hypothesized pathway is through vitamin D deficiency, another established MS risk factor,² as obesity decreases the bioavailability of 25-hydroxyvitamin D (25OHD).³ While appealing, these proposed mediating effects of 25OHD have yet to be confirmed or quantified. Obesity is also characterized by a chronic low-grade inflammatory state, driven in part by hormones and cytokines secreted by adipose tissue, such as leptin and adiponectin. These have been implicated in MS pathogenesis and shown to modulate experimental autoimmune encephalomyelitis disease course (reviewed in ⁴). However, it remains unclear whether they have a causal role in the disease process or contribute to mediating the association with obesity.

Establishing the relative contributions of obesity and vitamin D deficiency in particular to MS risk has major public health implications, as their prevalence in the US is respectively 41.6% and 39.6%.^{5, 6} Furthermore, if a large proportion of the association between obesity and MS is explained by lowered 25OHD levels, then vitamin D supplementation at a population level or in high-risk individuals could become a viable strategy to mitigate the effects of obesity on MS.

To address these questions, we undertook a Mendelian randomization (MR) approach, which uses natural genetic variation as a proxy for an exposure to estimate its effect on an outcome. MR greatly reduces confounding since allelic variants influencing different exposures are randomly allocated at conception.⁷ The fact that genotypes are not modifiable by disease onset also limits reverse causation.⁸ This makes MR well suited for mediation analysis.⁹ In this study, we first

estimated ~~the effect of each of~~ whether genetic predisposition towards higher 25OHD, leptin and adiponectin levels ~~on-influence the risk of~~ MS, then took forward significantly associated traits into a two-step MR mediation analysis to determine their contribution to the association between obesity and MS.

Methods

Data sources

We identified single-nucleotide polymorphisms (SNPs) for BMI and each potential mediator using large-scale genome-wide association studies (GWAS) as shown in the **Table**.¹⁰⁻¹³ As genetic estimates for 25OHD were derived from the UK Biobank cohort,¹¹ we used a BMI GWAS that did not include this cohort¹⁰ ~~to avoid participant overlap in the mediation analysis, which can lead to inflated type 1 error rates to avoid bias from sample overlap~~.^{8, 14} For MS susceptibility, summary genetic estimates were obtained from the discovery cohort of the latest International MS Genetics Consortium meta-analysis, which included 14,802 MS cases and 26,703 controls as previously described.¹⁵ To prevent confounding through population stratification, all genetic estimates were from individuals of European ancestry (white British group for the UK Biobank¹¹) and subsequently adjusted for genetic principal components.

We ensured that genetic variants were independent ($r^2 < 0.01$) by using the 1000 Genomes European reference panel¹⁶ and PLINK v1.9¹⁷ ~~(clump command within 10,000 kb distance)~~,¹⁶ except for the 25OHD levels variants for which we used conditionally independent estimates from a COJO analysis.¹¹ When genetic ~~instruments-variants~~ were missing from one of the datasets, we identified proxy SNPs

in linkage disequilibrium (LD; $r^2 > 0.6$) using the same reference panel.¹⁶ We excluded variants within the major histocompatibility complex (MHC) region, as it is strongly associated with MS risk and exhibits complex ~~linkage disequilibrium~~LD which renders it susceptible to bias from pleiotropy. Genetic variants were aligned to the forward strand. For GWAS not originally reported on the Genome Reference Consortium Human Build 37 (BMI, leptin and adiponectin), forward strand alleles for palindromic SNPs were inferred using minor allele frequencies up to 0.42.

The inclusion of genetic variants with smaller effects on the exposure can lead to weak instrument bias, which attenuates MR estimates towards the null. For each phenotype, we evaluated instrument strength using the F-statistic (two-sample conditional F-statistic F_{TS} for multivariable MR), with values greater than 10 indicating adequately strong instruments. As a sensitivity analysis, we also measured the effect of BMI on the risk of MS using the latest GWAS meta-analysis by Yengo and colleagues¹⁸ (n=681,275) which included UK Biobank participants and thus was not used for the mediation analysis, as discussed above. This study identified 941 near-independent SNPs, of which 548 were included after filtering out MHC variants, marginal effects below genome-wide significance and correlated variants.

Statistical analysis

We first carried out univariable inverse-variance weighted (IVW) MR to examine the effect on MS of genetically determined BMI, 25OHD, leptin and adiponectin levels individually. For each genetic variant, the effect on MS was estimated using the SNP effect coefficients via the ratio method, with standard errors derived using the delta method.¹⁹ These individual MR estimates were combined into a summary measure using random-effect inverse-variance weighted meta-analysis. In addition, we

applied the MR-Egger and weighted median MR methods to assess for potential bias from pleiotropic effects, whereby genetic instruments-variants affect the outcome independent of the risk factor.⁸ We also performed the MR-Pleiotropy RESidual Sum and Outlier (MR-PRESSO) outlier test to identify and remove potentially pleiotropic variants.²⁰

As leptin and adiponectin displayed no measurable effect on MS risk, only 25OHD levels were-was taken forward into the mediation analysis. We measured the indirect effect of BMI on MS mediated by 25OHD levels using the product of coefficients method via two-step MR (**Figure 1**).⁹ This involved first estimating the effect of BMI on 25OHD levels, then multiplying this by the effect of 25OHD levels on MS risk adjusting for BMI using regression-based multivariable MR.²¹ The multivariable MR included genetic effects of both 25OHD levels and BMI for variants associated with either phenotype. The proportion mediated was estimated by dividing this-the indirect effect by the total effect of BMI on MS. Outlier-corrected estimates were used for univariable analyses.

To further ensure that our estimates were not biased by pleiotropy, we repeated the mediation analysis using only-6 genetic variants for 25OHD levels identified in a previous GWAS,²² most of which have a well-defined role in vitamin D synthesis (*DHCR7/NADSYN1* [rs12785878]; *CYP2R1* [rs10741657]), transportation (*GC* [rs3755967]), or degradation (*CYP24A1* [rs17216707]). Lastly, we calculated the effect of BMI on MS risk adjusting for 25OHD levels using multivariable MR.

All statistical analyses were performed in R (version 3.6.0). We used the MendelianRandomization²³ (version 0.4.2), TwoSampleMR²⁴ (version 0.5.4) and MVMR (version 0.2)²⁵ R packages. The alpha-level for statistical significance was set to 0.05. The data sources used in this study obtained

informed consent from all participants. Separate institutional review board approval was not required for this study.

Results

Effects of vitamin D, leptin and adiponectin levels on MS

Figure 2 shows the total effect on odds of MS for BMI, 25OHD levels, leptin and adiponectin levels as estimated by univariable MR. A significant effect was ~~only~~ observed for BMI (odds ratio [OR]=1.~~3340~~ per standard deviation [SD] increase in BMI, 95% CI ~~1.09 to 1.63~~, $P=4.8 \times 10^{-3}$ ~~1.16 to 1.67~~, $P=3.12 \times 10^{-4}$) and 25OHD levels (OR=0.72 per SD increase in log-transformed 25OHD, 95% CI 0.60 to 0.87, $P=6.21 \times 10^{-4}$). To guide clinical interpretation, the SD for BMI was 4.7 kg/m²,¹⁰ while for 25OHD it corresponded to a 29.2 nmol/L increase for individuals with a baseline level of 50 nmol/L. Additional equivalences for clinically relevant vitamin D thresholds are presented in the **Supplementary Table S1**. In sensitivity analyses, the MR-Egger intercept was centered around zero for both phenotypes (BMI: intercept=-0.0064, 95% CI ~~-0.020 to 0.0080~~ ~~-0.007 to -0.17~~, $P=0.4350$; 25OHD levels: intercept=0.006, 95% CI ~~-0.003 to 0.005~~ ~~to -0.003~~, $P=0.22$). **Figure 2** shows that the MR-Egger regression coefficient, ~~and~~ weighted median ~~and MR-PRESSO outlier-corrected~~ estimates were consistent with the main IVW analysis. ~~F-statistics were greater than 10 for all phenotypes (range 22.2-167.9) and are reported in the F-statistics for each phenotype are reported in the~~ **Supplementary Table S2**. ~~The effect of BMI on MS risk was replicated using the larger set of 548 variants from the latest BMI GWAS meta-analysis (OR=1.29, 95% CI 1.15 to 1.46, $P=2.6 \times 10^{-5}$).~~

Mediation of the effect of BMI on MS by vitamin D

Higher BMI was associated with lower 25OHD levels (IVW: $\beta=-0.082$ per SD unit increase in BMI, 95% CI -0.118 to -0.046, $P=7.6871 \times 10^{-6}$; outlier-corrected: $\beta=-0.081$, 95% CI -0.111 to -0.051,

$P=2.6 \times 10^{-7}$). In contrast, genetically increased 25OHD levels did not influence BMI ($\beta=0.027$, 95% CI -0.018 to 0.073, $P=0.24$). In line with this, the effect of 25OHD levels on MS remained robust after adjusting for BMI in multivariable MR (OR=0.81, 95% CI 0.68 to 0.97, $P=0.02$)~~OR=0.80, 95% CI 0.67-0.96, $P=0.01$~~ . The slightly weaker adjusted compared to unadjusted estimate is due to the reduced number of variants ($N_{\text{SNPs}}=78$) also present in the BMI dataset (**Supplementary Table S3**). The proportion of the effect of BMI on MS mediated by 25OHD levels was 5.2% (95% CI 0.3% to 31.0%)~~5.4% (95% CI 0.4% to 30.5%)~~. Using only the 6 previously identified SNPs for 25OHD,²² we obtained similar estimates of the mediated proportion (5.9%, 95% CI 0.3% to 36.3%)~~(6.4%, 95% CI 0.7% to 33.7%)~~.

In multivariable MR, an effect of higher BMI on MS persisted after adjusting for 25OHD levels (OR=1.28 per SD unit increase in BMI, 95% CI 1.05-1.55, $P=0.01$), consistent with incomplete mediation (Supplementary Table S3). The two-sample conditional F-statistic F_{TS} was 77.5 and 33.0 for 25OHD levels and BMI respectively.

None of the individual variants associated with BMI, 25OHD, leptin or adiponectin levels were genome-wide significant in the discovery cohort of the MS susceptibility GWAS (Supplementary Tables S4-7). Of the genes reported to be associated with MS (based on distance from their corresponding variants), 2 overlap with 25OHD levels (*CYP24A1* and *TNFAIP8*) and 1 with BMI (*ADCY3*). While the lead variants for MS and those phenotypes were not in LD (Supplementary Table S8), a previous study reported a coding variant in *CYP24A1* which strongly reduces 25OHD levels and also increases risk of MS.²⁶ *CYP24A1* encodes an enzyme that catalyzes the inactivation of calcidiol and calcitriol.

Discussion

This study provides genetic support for a ~~causal~~ role of increased BMI and lowered 25OHD levels in the development of MS, using updated genetic estimates from large-scale cohorts. When considering both phenotypes in a mediation framework, we estimated that ~~only~~ 5.42% (and up to a third) of the association between obesity and MS susceptibility can be explained by lowered vitamin D levels. In contrast, the results show no effect on the risk of the disease for lifelong genetically related increases in leptin or adiponectin levels. Therefore, the majority of the effect of obesity remains unexplained.

Previous MR studies on the effects of 25OHD levels and BMI on MS risk have found directionally consistent results with overlapping estimate confidence intervals.²⁷⁻³¹ Similarly, our previous MR study of adiponectin levels using an independent MS genetic cohort found no effect on MS risk.³² While some observational studies have described increased leptin and lowered adiponectin levels in MS compared to controls, these have generally been small and included prevalent MS cases, making them susceptible to reverse causality.⁴ That said, small effects by those cytokines on MS risk may still exist given their wider estimate confidence intervals.

~~Despite the large body of observational studies on vitamin D deficiency and obesity in MS, to our knowledge none has addressed the question of mediation between these risk factors.~~ A previous study in pediatric-onset MS and another in adult-onset MS reported independent effects of genetically determined BMI and 25OHD levels using multivariable regression.^{28, 31} However, they did

not quantify the mediated effects and employed an approach which may be more susceptible to measurement error in the intermediate phenotype. Moreover, both studies used genetic estimates for 25OHD previously adjusted for BMI, making them ~~##~~less suited to assess mediation of obesity.

A major strength of this work is the MR approach which helps overcome many of the challenges facing traditional mediation analysis by reducing bias from unmeasured confounding, reverse causality and measurement error. It also enabled us to estimate the effects of each phenotype using large-scale genetic studies totaling more than 800,000 participants, while alleviating the need for ~~all~~ direct measurements of those phenotypes to be performed in a single cohort. We also acknowledge a number of limitations. First, potential bias from pleiotropy cannot be entirely excluded. While the genetic variants included are reliably associated and predictive of their respective phenotypes, their functional consequences (or those of strongly correlated variants in the same region) are not known in most cases and may include pleiotropic effects. These cannot be directly tested; ~~h~~However, our sensitivity analyses and the consistent estimates obtained using the reduced set of SNPs mapped to genes with well-characterized roles in vitamin D metabolism decrease the likelihood of pleiotropic bias. Second, a binary outcome can lead to biased mediation estimates due to the non-collapsibility of odds ratio, although this is lessened by the use of product of the coefficients method and the rare prevalence of MS.⁹ Third, MR estimates the risk associated with lifelong differences in the exposure, and as such may not fully capture effects that are time-specific. Fourth, although the risk of weak instrument bias is low given the F-statistics, the lower variance explained for leptin, as well as the smaller GWAS sample size for leptin and adiponectin, have contributed to reduced statistical power compared to other phenotypes, as indicated by the wider confidence intervals. As such, small effects by those cytokines in MS risk may still exist. That said, the genetic variants used in this study

adequately capture differences in BMI in early adulthood,³³ the time period where obesity is most strongly associated with MS risk, while for vitamin D deficiency associations have been reported from in utero through early adulthood.² Lastly, we assume that the associations between MS, obesity and 25OHD levels are linear (assumptions supported by previous studies³⁴⁻³⁶) and without interaction. Confirmation of these findings in observational studies with direct phenotype measurement, studies investigating potential interactions between BMI and 25OHD levels, and exploration of the mechanistic pathways underlying the association between obesity and MS are needed.

Conclusion

This MR study found that a small proportion of the effect of obesity on the risk of MS is mediated by decreasing levels of vitamin D, while leptin and adiponectin had no measurable effect on MS susceptibility. This suggests that widespread vitamin D supplementation would only lead to modest reduction in the association between obesity and MS, most of which remains unexplained.

Acknowledgments: We wish to kindly thank the IMSGC, GIANT and ADIPOGen consortia for access to their summary statistics data. Part of this work was conducted using the UK Biobank resource.

Declaration of Conflicting Interests: B.R. reported receiving compensation for consulting fees from GlaxoSmithKline and Deerfield Capital outside the submitted work. S.E.B. reported receiving compensation for consulting fees from EMD Serono, Novartis, Merck & Co and Sanofi-Aventis outside of the submitted work. No other disclosures were reported.

Funding: This study was supported by the NMSS-ABF Clinician Scientist Development Award from the National Multiple Sclerosis Society and the Multiple Sclerosis Society of Canada (FAN-1808-32256 to A.H.), by the Medical Research Council Integrative Epidemiology Unit (MC_UU_00011/1 to R.E.M.), and the National Institutes of Health (NIH) (R01NS099240 to S.E.B). Dr Baranzini holds the Distinguished Professorship in Neurology I and is the Heidrich Family and Friends Endowed Chair in Neurology at UCSF. The Richards research group is supported by the Canadian Institutes of Health Research (grants 365825; 409511), the Lady Davis Institute of the Jewish General Hospital, the Canadian Foundation for Innovation, the NIH Foundation, Cancer Research UK, Genome Québec, the Public Health Agency of Canada and the Fonds de Recherche Québec Santé (FRQS). Dr Richard is supported by a FRQS Clinical Research Scholarship. Support from Calcul Québec and Compute Canada is acknowledged. TwinsUK is funded by the Wellcome Trust, Medical Research Council, European Union, the National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London.

References

1. Harroud A and Richards JB. Mendelian randomization in multiple sclerosis: A causal role for vitamin D and obesity? *Mult Scler*. 2018; 24: 80-5.
2. Olsson T, Barcellos LF and Alfredsson L. Interactions between genetic, lifestyle and environmental risk factors for multiple sclerosis. *Nat Rev Neurol*. 2017; 13: 25-36.
3. Vimalaswaran KS, Berry DJ, Lu C, et al. Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Med*. 2013; 10: e1001383.
4. Keyhanian K, Saxena S, Gombolay G, Healy BC, Misra M and Chitnis T. Adipokines are associated with pediatric multiple sclerosis risk and course. *Mult Scler Relat Disord*. 2019; 36: 101384.
5. Forrest KY and Stuhldreher WL. Prevalence and correlates of vitamin D deficiency in US adults. *Nutr Res*. 2011; 31: 48-54.
6. Hales CM, Fryar CD, Carroll MD, Freedman DS and Ogden CL. Trends in Obesity and Severe Obesity Prevalence in US Youth and Adults by Sex and Age, 2007-2008 to 2015-2016. *JAMA*. 2018; 319: 1723-5.
7. Smith GD, Lawlor DA, Harbord R, Timpson N, Day I and Ebrahim S. Clustered environments and randomized genes: a fundamental distinction between conventional and genetic epidemiology. *PLoS Med*. 2007; 4: e352.
8. Hemani G, Bowden J and Davey Smith G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum Mol Genet*. 2018; 27: R195-R208.
9. Carter AR, Gill D, Davies NM, et al. Understanding the consequences of education inequality on cardiovascular disease: mendelian randomisation study. *BMJ*. 2019; 365: l1855.

10. Locke AE, Kahali B, Berndt SI, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature*. 2015; 518: 197-206.
11. Manousaki D, Mitchell R, Dudding T, et al. Genome-wide Association Study for Vitamin D Levels Reveals 69 Independent Loci. *Am J Hum Genet*. 2020; 106: 327-37.
12. Dastani Z, Hivert MF, Timpson N, et al. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet*. 2012; 8: e1002607.
13. Kilpelainen TO, Carli JF, Skowronski AA, et al. Genome-wide meta-analysis uncovers novel loci influencing circulating leptin levels. *Nat Commun*. 2016; 7: 10494.
14. Burgess S, Davies NM and Thompson SG. Bias due to participant overlap in two-sample Mendelian randomization. *Genet Epidemiol*. 2016; 40: 597-608.
15. International Multiple Sclerosis Genetics C. Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science*. 2019; 365.
16. The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*. 2015; 526: 68-74.
17. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007; 81: 559-75.
18. Yengo L, Sidorenko J, Kemper KE, et al. Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet*. 2018; 27: 3641-9.
19. Thompson JR, Minelli C and Del Greco MF. Mendelian Randomization using Public Data from Genetic Consortia. *Int J Biostat*. 2016; 12.

20. Verbanck M, Chen CY, Neale B and Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet.* 2018; 50: 693-8.
21. Burgess S, Dudbridge F and Thompson SG. Re: "Multivariable Mendelian randomization: the use of pleiotropic genetic variants to estimate causal effects". *Am J Epidemiol.* 2015; 181: 290-1.
22. Jiang X, O'Reilly PF, Aschard H, et al. Genome-wide association study in 79,366 European-ancestry individuals informs the genetic architecture of 25-hydroxyvitamin D levels. *Nat Commun.* 2018; 9: 260.
23. Yavorska OO and Burgess S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. *Int J Epidemiol.* 2017; 46: 1734-9.
24. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife.* 2018; 7.
25. Sanderson E, Davey Smith G, Windmeijer F and Bowden J. An examination of multivariable Mendelian randomization in the single-sample and two-sample summary data settings. *Int J Epidemiol.* 2019; 48: 713-27.
26. Manousaki D, Dudding T, Haworth S, et al. Low-Frequency Synonymous Coding Variation in CYP2R1 Has Large Effects on Vitamin D Levels and Risk of Multiple Sclerosis. *Am J Hum Genet.* 2017; 101: 227-38.
27. Mokry LE, Ross S, Ahmad OS, et al. Vitamin D and Risk of Multiple Sclerosis: A Mendelian Randomization Study. *PLoS Med.* 2015; 12: e1001866.

28. Jacobs BM, Noyce AJ, Giovannoni G and Dobson R. BMI and low vitamin D are causal factors for multiple sclerosis: A Mendelian Randomization study. *Neurol Neuroimmunol Neuroinflamm.* 2020; 7.
29. Mokry LE, Ross S, Timpson NJ, Sawcer S, Davey Smith G and Richards JB. Obesity and Multiple Sclerosis: A Mendelian Randomization Study. *PLoS Med.* 2016; 13: e1002053.
30. Gianfrancesco MA, Glymour MM, Walter S, et al. Causal Effect of Genetic Variants Associated With Body Mass Index on Multiple Sclerosis Susceptibility. *Am J Epidemiol.* 2017; 185: 162-71.
31. Gianfrancesco MA, Stridh P, Rhead B, et al. Evidence for a causal relationship between low vitamin D, high BMI, and pediatric-onset MS. *Neurology.* 2017; 88: 1623-9.
32. Devorak J, Mokry LE, Morris JA, et al. Large differences in adiponectin levels have no clear effect on multiple sclerosis risk: A Mendelian randomization study. *Mult Scler.* 2017; 23: 1461-8.
33. Brandkvist M, Bjorngaard JH, Odegard RA, Asvold BO, Sund ER and Vie GA. Quantifying the impact of genes on body mass index during the obesity epidemic: longitudinal findings from the HUNT Study. *BMJ.* 2019; 366: l4067.
34. Munger KL, Chitnis T and Ascherio A. Body size and risk of MS in two cohorts of US women. *Neurology.* 2009; 73: 1543-50.
35. Munger KL, Levin LI, Hollis BW, Howard NS and Ascherio A. Serum 25-hydroxyvitamin D levels and risk of multiple sclerosis. *JAMA.* 2006; 296: 2832-8.
36. Wortsman J, Matsuoka LY, Chen TC, Lu Z and Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr.* 2000; 72: 690-3.

Figure Legends

Figure 1. Directed acyclic graph of the MR mediation analysis.

Increased BMI may affect the risk of MS through lowering 25OHD levels (indirect pathways in red), or independently from 25OHD (pathway in blue). The indirect effect can be calculated by multiplying α times β , where α is the effect of BMI on 25OHD levels, and β the effect of 25OHD levels on MS adjusted for BMI using multivariable MR. The proportion mediated can be estimated by dividing the indirect effect by the total effect of BMI on MS. 25OHD=25-hydroxyvitamin D; MS = multiple sclerosis; SNP = single nucleotide polymorphism

Figure 2. Forest plot showing the MR estimates investigating the effect of BMI and its potential mediators on the risk of MS.

There were too few variants to apply the MR-Egger method for leptin levels, and no outliers were identified by MR-PRESSO except for BMI. The unit change for each phenotype associated with reported odds ratio can be found in **Table 1**. 25OHD=25-hydroxyvitamin D; CI=confidence intervals; IVW=inverse-variance weighted; MR=Mendelian randomization; MS=multiple sclerosis; N_{SNPs}=number of single nucleotide polymorphisms included in each analysis; OR=Odds ratio.

Tables

Table 1. Summary of the phenotypes and summary genetic data used as exposures in the MR analyses

Phenotype	Sample size	N _{SNPs} ^a	Units	Data source
BMI	322,154	74	SD increase in BMI, adjusted for age, age ² and sex	GIANT consortium ¹⁰
25OHD	401,460	138	SD increase in standardized log-transformed levels, adjusted for vitamin D supplementation, age, sex, season of measurement and assessment center (as proxy for latitude)	UK Biobank ¹¹
Adiponectin	38,276	11	Unit increase in log-transformed levels, adjusted for age, sex and BMI	ADIPOGen consortium ¹²
Leptin	52,126	4	Unit increase in log-transformed levels, adjusted for age, age ² and sex	Kilpeläinen et al., 2016 ¹³

^a Genome-wide significant SNPs, after exclusions due to linkage disequilibrium, MHC region or palindromic SNPs. For 25OHD levels, this represents the number of conditionally independent variants from COJO analysis. BMI=body mass index; MHC=major histocompatibility complex; MR=Mendelian randomization; SD=standard deviation; SNPs=single nucleotide polymorphisms