Winters, M., Lyng, K. D., Holden, S., Lura, C. B., Welton, N. J., Caldwell, D. M., Vicenzino, B., Weir, A., & Rathleff, M. S. (2021). Infographic. Comparative effectiveness of treatments for patellofemoral pain: a living systematic review with network meta-analysis. *British Journal of Sports Medicine*, 55(22), 1311-1312. <https://doi.org/10.1136/bjsports-2021-104360>

Publisher's PDF, also known as Version of record

License (if available):
CC BY-NC

Link to published version (if available):
[10.1136/bjsports-2021-104360](https://doi.org/10.1136/bjsports-2021-104360)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Infographic. Comparative effectiveness of treatments for patellofemoral pain: a living systematic review with network meta-analysis

Marinus Winters ¹, Kristian Damgaard Lyng,¹ Sinead Holden ^{1,2}, Carolina Bryne Lura,¹ Nicky J Welton,³ Deborah M Caldwell,³ Bill Vicenzino ⁴, Adam Weir ^{5,6,7}, Michael Skovdal Rathleff^{1,8}

INTRODUCTION

Patellofemoral pain (PFP) is one of the most common knee complaints and negatively impacts daily living and quality of life.¹ Despite its high prevalence, managing PFP is often challenging, with poor prognosis and a large variety of treatment strategies available.^{2,3}

THE NEED FOR A LIVING SYSTEMATIC REVIEW AND NETWORK META-ANALYSIS (NMA)

There are many systematic reviews that have investigated different head-to-head comparisons for treatment of PFP. However, the comparative effectiveness of *all* available treatments for PFP has never been investigated. This makes it challenging to decide on the most appropriate treatment. Additionally, systematic reviews need to be updated regularly to reflect the most recent and up-to-date research. A *living* systematic review with a network NMA allows for the simultaneous comparison of all available treatment strategies for PFP.⁴ It supports clinicians with an up-to-date and comprehensive overview of the comparative effectiveness of treatments.⁴

In this paper,⁵ published in the *British Journal of Sports Medicine*, we performed a living systematic review with an NMA to evaluate the comparative effectiveness of all available treatments for patients with a clinical diagnosis of PFP. We included randomised control trials investigating the effectiveness of treatments in patients with PFP. Our main outcome was 'any improvement' on the Global Rating of Change Scale.

We showed that compared with all other treatments, 'wait and see' was the least effective approach at 3 months. Education in combination with a physical treatment (exercise, orthoses or patellar taping/mobilisation or a combination of these treatments) was more likely to be effective at 3 months compared with all other treatments alone. Education alone is comparable to education in combination with a physical treatment at 12-month follow-up.

Infographic

FUTURE TREATMENT OF PFP

We highlight the comparative effectiveness of different treatments for PFP in our living systematic review with NMA. When new evidence emerges, the NMA will be updated to provide clinicians with a contemporary overview of the best available evidence for the treatment of PFP. This paper can be freely accessed on <https://bjsm.bmj.com/content/55/7/369>, and future updates will be available at <https://www.generalpractice.aau.dk/research/Center-for-Musculoskeletal-Health/mangement-patellofemoral-pain>, which also contains ideas on how to deliver treatments presented in the NMA to ensure our NMA supports clinical practice.

RATIONALE FOR INFOGRAPHIC

The rationale for this infographic is to further increase the attention of the study's findings. By creating an infographic, we hope to aid clinical practice and research translation in general. Our infographic includes a link to our living synthesis. We hope our infographic will support clinicians managing PFP now and in the future.

¹Center for General Practice, Aalborg University, Aalborg, Denmark

²SMI, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

³Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

⁴School of Health and Rehabilitation Sciences: Physiotherapy, University of Queensland, Brisbane, Queensland, Australia

⁵Sports Groin Pain Centre, Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar

⁶Department of Orthopaedics, Erasmus MC University Medical Center for Groin Injuries, Rotterdam, The Netherlands

⁷Sports Medicine and Exercise Clinic Haarlem (SBK), Haarlem, The Netherlands

⁸Department of Clinical Medicine, Aalborg University, Aalborg University Hospital, Aalborg, Denmark

Correspondence to Dr Marinus Winters, Research Unit for General Practice in Aalborg, Department of Clinical Medicine, Aalborg Universitet Det Sundhedsvidenskabelige Fakultet, Aalborg, Denmark; marinuswinters@hotmail.com

Twitter Marinus Winters @marinuswinters, Kristian Damgaard Lyng @_Kristianlyng, Sinead Holden @Sinead_Holden and Bill Vicenzino @Bill_Vicenzino

Contributors KDL produced the infographic with revisions provided by MW, SH, CBL, NJW, DMC, BTW, AW and MSR. All authors read and consented to the content of the infographic.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

MW and KDL contributed equally.

MW and KDL are joint first authors.

To cite Winters M, Lyng KD, Holden S, *et al.* *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2021-104360

Accepted 30 June 2021

Br J Sports Med 2021;0:1–2.
doi:10.1136/bjsports-2021-104360

ORCID iDs

Marinus Winters <http://orcid.org/0000-0001-5742-7441>

Sinead Holden <http://orcid.org/0000-0002-7314-2152>

Bill Vicenzino <http://orcid.org/0000-0003-0253-5933>

Adam Weir <http://orcid.org/0000-0003-0861-662X>

REFERENCES

- 1 Coburn SL, Barton CJ, Filbay SR, *et al.* Quality of life in individuals with patellofemoral pain: a systematic review including meta-analysis. *Phys Ther Sport* 2018;33:96–108.
- 2 Rathleff MS, Rathleff CR, Olesen JL, *et al.* Is knee pain during adolescence a self-limiting condition? prognosis of Patellofemoral pain and other types of knee pain. *Am J Sports Med* 2016;44:1165–71.
- 3 Lankhorst NE, van Middelkoop M, Crossley KM, *et al.* Factors that predict a poor outcome 5–8 years after the diagnosis of patellofemoral pain: a multicentre observational analysis. *Br J Sports Med* 2016;50:881–6.
- 4 Winters M, de Vos R-J, van Middelkoop M, *et al.* Stay alive! What are living systematic reviews and what are their advantages and challenges? *Br J Sports Med* 2021;55:519–20.
- 5 Winters M, Holden S, Lura CB, *et al.* Comparative effectiveness of treatments for patellofemoral pain: a living systematic review with network meta-analysis. *Br J Sports Med* 2021;55:369–77.