Feng, Q., Wong, S. H., Zheng, J., Yang, Q., Sung, J. J., & Tsoi, K. K. (2021). Intake of processed meat, but not sodium, is associated with risk of colorectal cancer: Evidence from a large prospective cohort and two-sample Mendelian randomization. *Clinical Nutrition*, 40(7), 4551-4559. <https://doi.org/10.1016/j.clnu.2021.05.036>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.clnu.2021.05.036](https://doi.org/10.1016/j.clnu.2021.05.036)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Elsevier at [10.1016/j.clnu.2021.05.036](https://doi.org/10.1016/j.clnu.2021.05.036). Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Intake of processed meat, but not sodium, is associated with risk of colorectal cancer:
evidence from a large prospective cohort and two-sample Mendelian randomization

Qi Feng¹, Sunny H Wong², Jie Zheng³, Qian Yang³, Joseph JY Sung², Kelvin KF Tsoi^{1, 4*}

1. Division of Epidemiology, School of Public Health and Primary Care, The Chinese
University of Hong Kong

2. Department of medicine & Therapeutics, Faculty of Medicine, The Chinese University of
Hong Kong

3. MRC Integrative Epidemiology Unit, Bristol Population Health Science Institute, The
University of Bristol

4. SH Big Data Decision Analytics Research Centre, The Chinese University of Hong Kong

*: corresponding author (Kelvin KF Tsoi, Division of Epidemiology, School of Public Health
and Primary Care, The Chinese University of Hong Kong; Email: kelvintsoi@cuhk.edu.hk)

Abstract

Background & Aims: Processed meat and high sodium intake are common in Western diet.

The objective was to examine their independent effects on the risk of colorectal cancer (CRC).

Methods: We performed both observational analysis with UK Biobank and genetic analysis with Mendelian randomization (MR). The 24-hour urinary sodium (UNa) and reported intake of processed meat were fitted on incident CRC by multivariable Cox proportional hazard model, adjusted for covariates, such as age, gender, family history, etc. Different sodium measures were used for sensitivity analyses. Two-sample MR analyses were performed using summary data from genome-wide association studies of UNa and CRC. Multivariable MR was adjusted for body mass index.

Results: We included 415 524 eligible participants from UK Biobank. During a median follow-up of 11.1 years, 2 663 participants were diagnosed with CRC. High intake of processed meat independently increased risk of CRC by 23% (HR 1.23; 95% CI: 1.03 to 1.46), but 24-hour UNa was not significantly associated with CRC (HR 0.96; 95% CI: 0.87 to 1.06). Furthermore, MR also showed little evidence for the effect of UNa on CRC (OR 1.02; 95% CI: 0.11 to 9.42). Sensitivity analyses showed consistent results across different measurements of sodium intake.

Conclusions: Intake of processed meat had an independent effect on the risk of CRC, but the risk was not associated with sodium level. Reduction of processed meat intake may be an effective strategy for CRC prevention, while sodium reduction should still be recommended to achieve other health benefits.

Keywords: Sodium intake; urinary sodium; processed meat; colorectal cancer; UK Biobank; Mendelian randomization

Introduction

Colorectal cancer (CRC) is one of the most prevalent cancers worldwide. GLOBOCAN 2018 estimated CRC contributed 10.2% of cancer incidence and 9.2% of cancer mortality among all cancer types [1]. Numerous lifestyle factors have been noted to increase CRC risk, including obesity, smoking, physical inactivity, insufficient intake of fresh vegetables and fruits, etc. [2–4]. Excessive consumption of processed meat has been regarded as a convincing cause of CRC [5]. Increased CRC risk has been observed in people with high processed meat intake consistently in large prospective cohorts [6,7] and meta-analyses [8,9].

Since high sodium intake has shown both direct and indirect carcinogenic effects on gastric cancer [10–13], studies have hypothesized similar effects on other gastrointestinal cancers, such as CRC, but current evidence remains unclear. Two case-control studies found increased CRC risk in people with high dietary sodium intake [14] and in people who usually add extra salt to food [15], while other studies failed to observe similar significant associations [16,17]. Although a recent qualitative systematic review found seven studies with inconsistent results, the majority favored a positive association [18]. However, the methodological quality of these studies was limited by small sample size, unclear temporality and residual confounding.

Processed meat generally has high sodium concentration [19]. Intake of processed meat is closely correlated with sodium intake and is a big contributor for total sodium intake in typical Western pattern diet [20,21]. This complicated the relationship between processed meat intake, sodium intake and CRC risk. It remains unclear whether high sodium intake contributes to the effect of processed meat intake on CRC. The objective of this study was to examine the independent effects of processed meat intake and sodium intake on CRC; the potential results could be used to inform primary prevention of CRC. We aimed to provide both observational

and genetic evidence (Figure 1) [22]. The observational evidence was generated from UK Biobank cohort, and the genetic evidence from Mendelian randomization (MR) analysis.

Method

This study included two independent parts of data analysis, that is an observational analysis using UK biobank and a genetic analysis using MR. The overall methodology flowchart was shown in Figure 1.

a) Observational analysis

Data source and participants

UK Biobank is a large prospective population-based cohort. Half million participants aged 40 to 69 years old were recruited during year 2006 to 2010. Baseline data were collected from each participant, using questionnaire, physical examination and biochemical analysis of blood, urine and saliva samples. The baseline questionnaire collected information on sociodemographic characteristics, family history, health status, comorbidities, medication use, reproductive and environmental exposures and lifestyle (smoking, drinking, physical activity, dietary factors, use of nutrition supplements, etc.). Physical examination measured anthropometric traits (e.g., body weight, body height, waist circumference, etc.) and blood pressure, all with standardized procedures. Clinical outcomes during follow-up were extracted from cancer and death registries. UK Biobank was approved by the North West Multicenter Research Ethics Committee, the National Information Governance Board for Health and Social Care in England and Wales, and Community Health Index Advisory Group in Scotland. All participants have provided informed consents. The UK Biobank protocol is available online (<https://www.ukbiobank.ac.uk/wp-content/uploads/2011/11/UK-Biobank-Protocol.pdf>). We excluded those who were lost to follow-up (n = 1 298), pregnant at baseline (n = 370), reported

clinically-diagnosed cancer (other than non-melanoma skin cancer) at baseline (n = 41 248) and had missing data on exposure and other covariates (n = 44 054). Therefore, among all 502 494 participants, 415 524 were eligible and included (Figure 1).

Exposure

The primary exposure variables were 24-hour urinary sodium (UNa) and processed meat intake. Because dietary sodium intake was not directly measured in UK Biobank, we used 24-h UNa as its substitute, as used in previous studies [23,24]. 24-h UNa was estimated from spot UNa, urinary creatinine (UCr), body mass index (BMI) and age using the gender-stratified INTERSALT equation, which was developed and validated in Western population [25]. It has been shown to be less biased compared to other equations [26,27], and has been used in UK Biobank data previously [28]. Intake of processed meat (such as bacon, ham, sausages, meat pies, kebabs, burgers, chicken nuggets) was measured from baseline questionnaire and coded as a categorical variable (never, $\leq$ once a week and $\geq$ twice a week).

For sensitivity analyses, we used other substitute measures for dietary sodium intake, including spot UNa, urinary sodium-to-potassium (UNa/UK) ratio, urinary sodium-to-creatinine (UNa/UCr) ratio and adding salt to food. We dichotomized continuous 24-h UNa, spot UNa, UNa/UK ratio and UNa/UCr ratio using their median values as cutoff. Urinary sodium and potassium were measured with ion selective electrode analysis from participants' urine sample using Beckman Coulter AU5400, while UCr was measured with enzymatic analysis. Adding salt to food was measured with a question ("Do you add salt to your food?") in baseline questionnaire and categorized as never/rarely vs. usually/always.

Outcomes

The primary outcome was incident CRC (ICD-10 code: C18 - C20) during follow-up, while the secondary outcomes included incident colon cancer (ICD-10 code: C18) and incident rectal cancer (ICD-10 code: C19 - C20). Outcome data were extracted via linkage to cancer registries: the Medical Research Information Services for participants in England and Wales, and the Information Services Division for participants in Scotland. The time to outcome was calculated from the attendance date at the baseline assessment center to the date of outcome diagnosis, the date of death or the last date of follow-up (31 March 2016 for England and Wales, 31 October 2015 for Scotland), whichever was the earliest.

Covariate

We included sociodemographic, lifestyle, health status, medication use and dietary factors as covariates. Sociodemographic variables included age, gender, ethnicity (White, Black, Asian, Mixed and others), Townsend deprivation index (in quintiles), educational attainment (below college, college and above). Townsend deprivation index is a measure of material deprivation, calculated for each area based on the national census data, and each participant was assigned a Townsend deprivation index score corresponding to the area in which their postcode was located at baseline. CRC family history referred to the presence of diagnosed CRC on the participant's mother, father or siblings. Smoking and drinking status were coded as never, previous or current regular users. Physical activity was categorized to low, moderate and high levels based on their weekly metabolic equivalent values measured with the International Physical Activity Questionnaire. BMI was assessed in physical examination and calculated as body weight in kilogram divided by height square in meter. Estimated glomerular filtration rate (eGFR) was calculated based on serum cystatin C concentration with the gender-specific CKD-EPI cystatin C equations [29], as used in previous research [30]. Regular use of aspirin and/or ibuprofen was self-reported. The weekly intake frequencies of processed meat, fresh fruits,

fresh vegetables, cooked vegetables, oily fish, red meat and cereal (dietary fiber) were measured with baseline questionnaire. Intakes of unprocessed beef, lamb/mutton and pork were combined as red meat intake [31]. All answers of “prefer not to say”, “do not know”, “unknown” were combined as a category of “unknown”.

Statistical analysis

For descriptive statistics, mean with standard deviation was used for continuous variables and frequency with proportion was for categorical variables. We compared the 24-h UNa across different levels of processed meat intake, and evaluated their association in a multivariable logistic regression with processed meat intake as independent variable and 24-h UNa as dependent variable, adjusted for age, gender, ethnicity, smoking, drinking, physical activity, BMI, eGFR, intake of fresh fruits, fresh vegetables, cooked vegetables, oily fish, red meat and cereal.

We quantified the association between the exposure and CRC risk using multivariable Cox proportional hazard model, stratified by gender, ethnicity and age (< 50, 50 - 60, $\geq$ 60 years old). Assumption of hazard proportionality was examined with scaled Schoenfeld residual plot, and was satisfied in this study. We fitted a first model with only processed meat intake, a second model with only 24-h UNa, and a third model with both processed meat intake and 24-h UNa. The above three models were adjusted for multiple covariates (educational attainment, Townsend deprivation index, CRC family history, physical activity, smoking, drinking, regular use of aspirin/ibuprofen, BMI, eGFR, intake of fresh fruits, fresh vegetables, cooked vegetables, oily fish, red meat and cereal), to reduce potential confounding biases. In sensitivity analysis, we replaced 24-h UNa with other sodium intake measures (spot UNa, UNa/UK ratio, UNa/UCr ratio and adding salt to food).

173

174 *b) Two-sample MR*

175 MR is a study design to make causal inference between an exposure and an outcome using
176 genetic variants as instrumental variables [32]. We adopted two-sample MR approach [33] to
177 examine the association between UNa and risk of CRC, colon cancer and rectal cancer.
178 Genome wide association (GWA) studies of urine sodium, CRC, colon cancer and rectal cancer
179 were identified to perform the MR analysis. GWA analysis is a statistical technique to
180 investigate the association between a phenotype (e.g., a disease) and multiple genetic variants
181 (usually millions of single nucleotide polymorphisms (SNPs)). Data on the effects of SNPs on
182 phenotype were used for data analysis.

183

184 *Data sources*

185 We identified 50 SNPs associated with log-transformed spot UNa at a significance level of 5
186 $\times 10^{-8}$ from a GWA study of 446 237 unrelated European-ancestry individuals [34]. The 50
187 SNPs are located in different genes, distributed independently, and explain totally 6.4% of the
188 trait variance (Supplementary table 1). GWA analyses of CRC, colon cancer and rectal cancer
189 were conducted in 135 538 individuals from FinnGen cohort (1 573 CRC cases, 968 colon
190 cancer cases and 521 rectal cancer cases) [35].

191

192 Since obesity was associated with both urinary sodium and CRC [2,36], multivariable MR
193 adjusted for BMI was performed. A GWA study of 322 154 European-ancestry individuals
194 from the Genetic Investigation of Anthropometric Traits (GIANT) consortium identified 98
195 SNPs significantly associated with BMI, which explained totally 2.5% of the trait variance [37].
196 For sensitivity analysis, similar to the observational analysis, we used UNa/UK ratio and
197 UNa/UCr ratio as substitute measures for UNa. The GWA study of UNa/UK ratio and

UNa/UCr ratio was conducted in 327 616 unrelated European-ancestry individuals, which identified 8 and 12 SNPs for UNa/UK ratio and UNa/UCr ratio, respectively [38]. All the above GWA studies were adjusted for age, sex, ancestry principle components and array information. Summary-level statistics from the GWA studies were extracted for analysis (Figure 1). Processed meat intake was not included in multivariable MR because there is no relevant GWA study on it so far.

Statistical analysis

We firstly matched the 50 UNa-associated SNPs with the summary-level statistics of the outcome GWA studies in univariable MR. When a SNP could not be matched, we searched for its potential proxies (with $r^2 \geq 0.80$, distance $\leq 500\ 000$ bytes; <https://ldlink.nci.nih.gov/?tab=ldproxy>); in case of no eligible proxy, we removed the SNP from analysis. Summary-level statistics across different GWA studies were harmonized so that their effect estimates were aligned on the same allele [39]. The strength of the SNPs as instrumental variables was measured with F statistics, with a value > 10 suggesting good instrument strength [40]. For multivariable MR, we matched the UNa-associated SNPs and the BMI-associated SNPs with the outcome GWA studies and applied the same proxy-searching strategy.

Inverse-variance weighted method with random-effects model was used as primary analysis in univariable and multivariable MR. For sensitivity analysis, weighted median and MR-Egger methods were used in univariable MR, while only MR-Egger method was used in multivariable MR (Figure 1) [41]. Presence of pleiotropy was examined with MR-Egger intercept test, with a p value < 0.05 suggesting pleiotropy. MR-PRESSO was planned to examine the effect of horizontal pleiotropy, in case of any [42]. Additional sensitivity analyses was performed by (1)

using UNa/UK ratio and UNa/UCr ratio as sodium measure and (2) removing the SNPs that were associated with risk factors for the outcomes (e.g., obesity) at GWA significance level (5×10^{-8}) in manually searching PhenoScanner v2 database [43] (<http://www.phenoscanter.medschl.cam.ac.uk/>).

The effect measure in observational analysis was hazard ratio (HR) and its 95% confidence interval (CI), while the effect measure in MR was odds ratio (OR) and its 95% CI. A statistically significant effect was suggested by a CI not including null effect value of 1. The interpretation of HR was the excessive outcome risk for high 24-h UNa compared to that for low 24-h UNa. The interpretation of OR should be the excessive outcome risk for each 1 standard deviation increment in log-transformed spot UNa. Data analysis was performed in R environment, with “rms” (version 5.1-4), “MendelianRandomization” (version 0.4.2) and “MVMR” (version 0.2) [44] packages.

Results

The baseline characteristics of the 415 524 eligible participants were shown in Table 1. The mean age was 56.3 years old and 53.2% were female. Almost 95% of them were White. The mean BMI was 27.4 kg/m². The mean values for 24-h UNa, log transformed spot UNa, UNa/UK ratio and UNa/UCr ratio were 2.8, 4.2, 1.4 and 10.7, respectively. About 44.5% of participants usually/always added salt to their food. Over 30% had processed meat more than twice a week.

During a median follow-up of 11.1 (interquartile range 10.4 to 11.8) years, 2 663 participants was diagnosed with CRC (1756 colon cancer cases, 907 rectal cancer cases). Compared with the overall cohort, CRC cases were more like to be male, older, White, affluent, previous

smokers, and had lower eGFR, more frequent red meat and processed meat intake and adding salt to food, but less frequent consumption of fresh vegetables and cereal.

Processed meat intake was highly associated with 24-h UNa, spot UNa, UNa/UK ratio, UNa/UCr ratio and adding salt to food. Frequent processed meat consumer ($\geq$ twice a week) had increased risk of high 24-h UNa (OR 1.21 (95%CI 1.17 to 1.25)), high spot UNa (1.45 (1.41 to 1.49)), high UNa/UK ratio (1.38 (1.34 to 1.42)), high UNa/UCr ratio (1.11 (1.08 to 1.14)) and were more likely to add salt to food (1.64 (1.59 to 1.68)) (Table 2).

When fitting processed meat intake and 24-h UNa in separate models, increased CRC risk was observed in high processed meat intake (HR 1.14 (0.97 to 1.34) and 1.23 (1.03 to 1.46) for $\leq$ once/week and $\geq$ twice/week, respectively), but not in high 24-h UNa (0.96 (0.87 to 1.06)).

When fitting processed meat intake and the 24-h UNa in the same model, we observed an independent effect of processed meat intake (1.14 (0.97 to 1.34) and 1.23 (1.03 to 1.46) for $\leq$ once/week and $\geq$ twice/week, respectively), but still not of 24-h UNa (0.96 (0.87 to 1.05)). The effect size of processed meat intake remained similar between models (Table 3). Regarding the subtypes of CRC, neither processed meat intake nor 24-h UNa was associated with colon cancer, but processed meat intake $\geq$ twice/week was associated with rectal cancer (1.45 (1.06 to 1.98)) (Supplementary table 2). When using other sodium intake measures (spot UNa, UNa/UK ratio, UNa/UCr ratio and adding salt to food) in sensitivity analyses, the results remained similar, that is, no evidence of association between the sodium intake measures with CRC or its subtypes (Supplementary table 2).

In Mendelian randomization, we used 48 SNPs (F statistic = 611.25) and 125 SNPs (28 UNa-associated SNPs and 97 BMI-associated SNPs, F statistic = 14.75 and 34.70 for UNa and BMI,

respectively) in univariable and multivariable MR, respectively. We found little evidence for an association between UNa and CRC in both univariable MR (0.89 (0.21 to 3.74)) and multivariable MR (1.43 (0.30 to 6.81)) (Table 3, Figure 2). MR-Egger intercept tests showed no evidence of pleiotropy. UNa showed similarly insignificant effects on colon cancer and rectal cancer (Supplementary table 3; Supplementary figure 1). Using UNa/UK ratio and UNa/UCr ratio also demonstrated similarly null evidence for associations (Supplementary table 3). Removing the 22 SNPs associated with obesity-related traits also yielded similar results (Supplementary table 4).

Table 1: Baseline characteristics of the UK Biobank cohort

	Overall cohort (n=415 524)	Participants with colorectal cancer (n=2 663)
Female gender (n, %)	221 102 (53.2)	1 108 (41.6)
Age (mean, SD)	56.3 (8.1)	60.6 (6.6)
Ethnicity (n, %)		
White	392 008 (94.7)	2 563 (96.8)
Black	6 519 (1.6)	21 (0.8)
Asian	9 410 (2.3)	32 (1.2)
Mixed and others	6 175 (1.5)	33 (1.2)
Townsend deprivation (n, %)		
Q1 (most affluent) (mean = -4.52)	103 837 (25.0)	695 (26.1)
Q2 (mean = -2.90)	103 923 (25.0)	669 (25.1)
Q3 (mean = -0.97)	103 874 (25.0)	648 (24.3)
Q4 (most deprived) (mean = 3.19)	103 890 (25.0)	651 (24.4)
Qualification (n, %)		
College or above	135 554 (39.7)	778 (37.7)
Below college	206 228 (60.3)	1 286 (62.3)
Body mass index (mean, SD)	27.4 (4.8)	27.9 (4.6)
Family history of colorectal cancer	49 295 (11.9)	433 (16.3)

Regular user of aspirin/ibuprofen	123 406 (29.7)	831 (31.2)
eGFR (mean mL/min/1.73m ² , SD)	94.2 (9.0)	90.8 (8.4)
Smoking (n, %)		
Never smoker	228 260 (55.1)	1 236 (46.6)
Previous smoker	142 272 (34.4)	1 157 (43.6)
Current smoker	43 496 (10.5)	260 (9.8)
Drinking (n, %)		
Never drinker	17 885 (4.3)	88 (3.3)
Previous drinker	14 365 (3.5)	92 (3.5)
Current drinker	382 811 (92.2)	2 478 (93.2)
Physical activity level (n, %)		
Low	62719 (18.6)	419 (19.5)
Moderate	137295 (40.7)	886 (41.2)
High	137191 (40.7)	848 (39.4)
Sodium level		
Spot urinary sodium (mean µmol/L, SD)	77.87 (44.6)	78.73 (43.0)
Log-transformed spot urinary sodium (mean, SD)	4.18 (0.6)	4.2 (0.6)
24-h urinary sodium (mean g/d, SD)	2.76 (0.5)	2.67 (0.5)
Urinary sodium/potassium ratio (mean, SD)	1.44 (0.9)	1.41 (0.9)
Urinary sodium/creatinine ratio (mean, SD)	10.7 (7.1)	10.5 (6.2)
Adding salt to food (n, %)		
Never/rarely	230 638 (55.5)	1 424 (53.5)
Usually/always	184 826 (44.5)	1 239 (46.5)
Red meat intake frequency (mean, SD)	2.12 (1.45)	2.29 (1.53)
Processed meat (n, %)		
Never	38 194 (9.2)	174 (6.5)
≤ 1 time/week	246 588 (59.5)	1 531 (57.6)
≥ 2 times/week	129 810 (31.3)	954 (35.9)
Fresh fruit intake (n, %)		
< 1 time/week	38 692 (9.4)	270 (10.2)
1-2 times/week	225 826 (54.6)	1 417 (53.5)
3-4 times/week	117 663 (28.4)	760 (28.7)
≥ 5 times/week	31 626 (7.6)	203 (7.7)

Fresh vegetable intake (n, %)		
< 1 time/week	59 360 (14.5)	413 (15.8)
1-2 times/week	222 627 (54.3)	1 419 (54.2)
3-4 times/week	87 767 (21.4)	543 (20.7)
≥ 5 times/week	40 302 (9.8)	243 (9.3)
Cooked vegetable intake (n, %)		
< 1 time/week	20 650 (5.0)	132 (5.0)
1-2 times/week	191 569 (46.7)	1 187 (45.3)
3-4 times/week	156 241 (38.1)	1 022 (39.0)
≥5 times/week	41 792 (10.2)	277 (10.6)
Oily fish (n, %)		
< 1 time/week	183 609 (44.4)	1 103 (41.7)
1 time/week	155 718 (37.7)	1050 (39.7)
>1 times/week	73 784 (17.9)	495 (18.7)
Cereal (n, %)		
< 1 time/week	71 497 (17.3)	501 (18.8)
1-2 times/week	46 373 (11.2)	296 (11.1)
3-4 times/week	53 853 (13.0)	344 (12.9)
≥5 times/week	242 342 (58.5)	1 517 (57.1)

283

284

285

286

287

288

289

290

291

292

293

294 Table 2: Association between processed meat intake with sodium intake in UK Biobank

	Processed meat intake: Never	Processed meat intake: ≤ 1/week	Processed meat intake: ≥ 2/week
N	38 194	246 588	129 810
24-hour urinary sodium			
Low (n, %)	21 728 (56.9)	128 167 (52.0)	57 474 (44.3)
High (n, %)	16 466 (43.1)	118 421 (48.0)	16 466 (12.7)
Mean (SD)	2.7 (0.5)	2.7 (0.5)	2.8 (0.5)
OR (95%CI)**	1	1.01 (0.98, 1.04)	1.21 (1.17, 1.25)
Spot Urinary sodium			
Low (n, %)	23 354 (61.2)	130 657 (53.0)	53 173 (41.0)
High (n, %)	14 840 (38.8)	115 931 (47.0)	76 637 (59.0)
Mean (SD)*	4.0 (0.7)	4.1 (0.6)	4.3 (0.6)
OR (95%CI)**	1	1.18 (1.15, 1.22)	1.45 (1.41, 1.49)
Urinary sodium/potassium ratio			
Low (n, %)	21 366 (55.9)	129 233 (52.5)	56 631 (43.7)
High (n, %)	1 673 (43.8)	116 825 (47.5)	72 909 (56.3)
Mean (SD)	1.3 (0.9)	1.4 (0.9)	1.6 (0.9)
OR (95%CI)**	1	1.12 (1.09, 1.14)	1.38 (1.34, 1.42)
Urinary sodium/creatinine ratio			
Low (n, %)	18 043 (47.2)	124 822 (50.6)	64 633 (49.8)
High (n, %)	20 151 (52.8)	121 766 (49.4)	65 177 (50.2)
Mean (SD)	11.3 (6.8)	10.6 (7.4)	10.7 (6.4)
OR (95%CI)**	1	0.94 (0.92, 0.97)	1.11 (1.08, 1.14)
Adding salt to food			
Never/rarely (n, %)	24 390 (63.9)	141 426 (57.3)	64 417 (49.6)
Usually/always (n, %)	13 794 (36.1)	105 418 (42.7)	65 381 (50.4)
OR (95%CI)**	1	1.29 (1.26, 1.33)	1.64 (1.59, 1.68)

295 *: showing log-transformed spot urinary sodium.

296 **: odds ratio (95% confidence interval), estimated with processed meat intake as independent

297 variable while sodium intake measures and salt intake behaviour as dependent variable, adjusted for

298 age, sex, ethnicity, smoking, drinking, physical activity, body mass index, regular use of

aspirin/ibuprofen, estimated glomerular filtration rate, intake of fresh fruit, fresh vegetable, cooked vegetables, red meat, oily fish and cereal.

Table 3: Observational analysis and genetic analysis with Mendelian randomization (MR) for processed meat intake and urinary sodium on colorectal cancer

	Processed-meat Intake ($\geq 2/\text{week}$ vs. never)	Urinary Sodium
Observational Analysis*:		(24-hour urinary sodium)
Model 1	1.23 (1.03, 1.46)	Not Included
Model 2	Not Included	0.96 (0.87, 1.06)
Model 3	1.23 (1.03, 1.46)	0.96 (0.87, 1.05)
Genetic Analysis ⁺ :		(Log-transformed spot urinary sodium)
Univariable MR [¶]	NA	0.89 (0.21, 3.74)
Multivariable MR [§]	NA	1.43 (0.30, 6.81)

*Models were stratified by age, sex, ethnicity, and adjusted for education attainment, Townsend deprivation index, colorectal cancer family history, regular use of aspirin/ibuprofen, IPAQ physical activity status, smoking, drinking, body mass index, eGFR, fresh fruit intake, fresh vegetable intake, cooked vegetable intake, cereal intake, red meat and oily fish;

⁺: Inverse-variance weighted method;

[¶]: including 48 SNPs, F statistics = 611.25, p for MR-Egger intercept test = 0.32;

[§]: including 125 SNPs (28 UNa SNPs + 97 BMI SNPs), F statistics = 14.75 and 34.70 for UNa and BMI), p for MR-Egger intercept test = 0.10;

NA: not applicable;

SNP: single nucleotide polymorphism.

Discussion

This study shows evidence that processed-meat intake, but not sodium intake, is associated with an increased risk of CRC. It provided both epidemiological and genetic evidence that the effect of sodium intake on CRC was minimal; therefore, reducing intake of processed meat is a more important strategy for CRC prevention.

There has been plentiful evidence for the association between processed meat intake and CRC risk. Vieira *et al.* found that 100 g/d increment of processed meat intake increased CRC risk by 18% in a meta-analysis of 10 studies with 10 356 participants [9]. Handel *et al.* observed high intake of processed meat increased CRC risk by 13% in a more recent meta-analysis of 15 cohorts [8]. Previous analysis with UK biobank also demonstrated similarly 20-25 g/d higher intake of processed meat increased CRC risk by 18-19% [6,7]. An early pooled analysis of UK dietary cohort consortium failed to observe a significant association, but that was probably due to small number of CRC cases during follow-up [45]. As for the CRC subtypes, inconsistency remains across studies. Händel *et al.* found significant associations with both colon cancer and rectal cancer, and the effect was stronger on colon cancer than the other (excessive risk 25% vs. 18%) [8]. Vieira *et al.* found significantly positive association with colon cancer, but not rectal cancer [9], opposite to the findings in this study. Therefore, future research is merited to delineate its effect on CRC subtypes.

Although the intakes of dietary sodium and processed meat are highly correlated as shown in this study and previous evidence [21], sodium intake did not demonstrate significant effects on CRC or its subtypes as processed meat intake did. Even if the insignificance here was caused by the limited number of events and a true association existed, the effect size should be actually minor, as the HR point estimates are very close to one. The validity of the results is enhanced

by the use of the largest sample size so far and the careful controlling for a range of confounders. The null effect was similarly observed in sensitivity analyses using other substitutes for sodium intake and MR analysis, as well as previous studies, such as a case-control study of 2 226 Western people [16] and a prospective cohort of 77 500 Japanese people [17]. Although other studies found a significant association, such as Kune *et al* [14], Alegria-Lertxundi *et al* [15] and Yakoob *et al* [18], it was likely due to uncontrolled confounding. Future studies on this topic should be evidence-based, carefully designed with larger sample size, higher statistical power and more effective measures for controlling confounding, to reduce research waste [46].

The findings indicated that sodium intake had null carcinogenic effect on CRC, while processed meat had independent carcinogenic effect, to which its high sodium concentration made little contribution. Carcinogenesis of processed meat is believed to be caused by its mutagenic chemicals, including heterocyclic amines, polycyclic aromatic hydrocarbons, haem iron, N-nitroso compounds, etc. [5,47]. Recent evidence has found that high sodium diet was associated with cancer at gastrointestinal tract, especially stomach [12,13,48]. In the case of stomach cancer, high sodium causes direct gastric epithelial cell damage and weakens mucous barrier, thus indirectly enhancing the carcinogenesis of mutagenic compounds and *Helicobacter pylori* infection [49–52]. However, colorectal epithelia are able to actively maintain intra-cellular and inter-cellular sodium homeostasis, via amiloride-sensitive epithelial sodium channel (ENaC), mineralocorticoid receptor, etc. [53,54]. *In vivo* and *in vitro* studies showed that high sodium diet down-regulates expression and activation of mineralocorticoids receptor and ENaC subunits, thus decreasing sodium absorption and consequently maintaining sodium balance [55]. These mechanisms may help prevent or mitigate the potential harmful effects of high sodium in colon and rectum as it does to gastric epithelia. Nevertheless, high

sodium may still cause colon inflammation and exacerbate colitis as shown in animal research [56].

The findings provided further evidence that reduction of processed meat intake may be a potentially effective strategy for primary prevention of CRC, but sodium reduction may not. Nevertheless, we are not encouraging high sodium intake, because no evidence shows it can decrease CRC risk. More importantly, high sodium intake is a well-known risk factor for multiple health outcomes, such as hypertension [57], metabolic syndrome [58], cardiovascular diseases [24] and mortality [59]. Randomized controlled trials have demonstrated the health benefits of dietary sodium reduction [60,61]. Future studies are warranted to investigate how to improve general population's compliance and adherence to intake reduction of sodium and processed meat.

One of the strengths of our study is that we used the large UK Biobank cohort in observational analysis and large GWA studies in genetic analysis, which generated consistent results of null effect of sodium intake on CRC. We examined the independent effect of processed meat intake and urinary sodium, after adjusting for multiple covariates, including family history, use of aspirin, intake of red meat, fruits and vegetables, etc. In MR analysis, using SNPs as instrumental variables mainly relies on three assumptions: (1) the SNPs are associated with the exposure, (2) the SNPs should be independent of unobserved confounders after conditioning on observed confounders, and (3) associated with the outcome only through the exposure. For assumption (1) and (2), our SNPs are associated with UNa, forming strong instrumental variables, and not associated with the outcomes. For assumption (3), we performed multivariable MR adjusted for BMI, and sensitivity analysis by removing SNPs that are

potentially associated with confounders, such as various obesity measures, which yielded consistent findings.

This study has some limitations. First, we used 24-h UNa as a substitute for dietary sodium intake. Direct measurement of sodium intake is difficult and unavailable in UK Biobank, while 24-h UNa is a commonly acceptable substitute [23,24]. Using sodium intake behavior and other urinary sodium measures in sensitivity analyses showed similar results, which further enhanced the reliability of our results. Second, UK biobank only measured the frequency of food intake at baseline assessment, which may not completely represent the quantity consumed and introduce information bias. Third, we observed wide CIs under MR-Egger method in MR analyses, which may suggest underpower. However, MR-Egger method is often underpowered in studies and thus used as secondary sensitivity analysis [62], but the results were qualitatively consistent with the primary analysis of inverse-variance weighted method. Fourth, we did not specifically adjust for macronutrients (e.g., fat, protein, carbohydrates); instead, we adjusted for intakes of food (such as fruit, vegetables, red meat, fish, cereal), as they are the major sources of macronutrients as well as micronutrients. Fifth, although we have adjusted for multiple covariates in analyzing UK Biobank data, residual confounding is still possible, as in other observational studies, but its consistency with MR results increased its validity. Sixth, in multivariable MR analysis, we did not adjust for processed meat intake due to lack of relevant GWA study.

Conclusion

Although sodium intake and processed meat intake are highly correlated, processed meat intake is an independent risk factor for CRC risk, while sodium intake is not. Reducing intake of processed meat may be an effective measure for CRC prevention. Dietary sodium reduction

419 should still be recommended to achieve health benefits in blood pressure, metabolic syndrome,
420 cardiovascular diseases, etc. Future research is merited to study how to improve compliance
421 and adherence to intake reduction of processed meat and sodium.

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444 **Acknowledgement**

445 We thank the participants and researchers in UK Biobank study and FinnGen cohort. This study
446 was conducted under the UK Biobank approved application (ID 65563).

447

448 **Declaration**

449 **Conflict of interests:** None to declare.

450 **Funding statement:** This research did not receive any specific grant from funding agencies in
451 the public, commercial, or not-for-profit sectors.

452 **Data availability:** The UK Biobank individual data are accessible under application at
453 <https://www.ukbiobank.ac.uk/>. The FinnGen GWA summary data of colorectal cancer, colon
454 cancer and rectal cancer are accessible under application at <https://www.finnngen.fi/en>. The
455 BMI GWA summary data is accessible at GIANT consortium
456 [https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_file](https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files)
457 [s](https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files). The GWA summary data of urinary sodium, urinary sodium-to-potassium ratio and urinary
458 sodium-to-creatinine ratio can be accessed via the methods described in the original
459 publications.

460 **Analysis code availability:** Available under reasonable request.

461 **Author contribution:** QF conceived the research idea. QF and KKFT designed the study. JZ
462 and QY provided methodological guidance on Mendelian randomization. QF analyzed the data
463 and draft the manuscript. QF and KKTF interpreted the results. All authors critically reviewed
464 the manuscript.

References

- [1] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424. <https://doi.org/10.3322/caac.21492>.
- [2] Brenner H, Kloor M, Pox CP. Colorectal cancer. *Lancet* 2014;383:1490–502. [https://doi.org/10.1016/S0140-6736\(13\)61649-9](https://doi.org/10.1016/S0140-6736(13)61649-9).
- [3] Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut* 2017;66:683–91. <https://doi.org/10.1136/gutjnl-2015-310912>.
- [4] Weitz J, Koch M, Debus J, Höhler T, Galle PR, Büchler MW. Colorectal cancer. *Lancet* 2005;365:153–65. [https://doi.org/10.1016/S0140-6736\(05\)17706-X](https://doi.org/10.1016/S0140-6736(05)17706-X).
- [5] World Cancer Research Fund, American Institute for Cancer Research. Diet, nutrition, physical activity and colorectal cancer (revised 2018). n.d.
- [6] Bradbury KE, Murphy N, Key TJ. Diet and colorectal cancer in UK Biobank: a prospective study. *Int J Epidemiol* 2020;49:246–58. <https://doi.org/10.1093/ije/dyz064>.
- [7] Knuppel A, Papier K, Fensom GK, Appleby PN, Schmidt JA, Tong TYN, et al. Meat intake and cancer risk: prospective analyses in UK Biobank. *Int J Epidemiol* 2020:dyaa142. <https://doi.org/10.1093/ije/dyaa142>.
- [8] Händel MN, Rohde JF, Jacobsen R, Nielsen SM, Christensen R, Alexander DD, et al. Processed meat intake and incidence of colorectal cancer: a systematic review and meta-analysis of prospective observational studies. *Eur J Clin Nutr* 2020;74:1132–48. <https://doi.org/10.1038/s41430-020-0576-9>.
- [9] Vieira AR, Abar L, Chan DSM, Vingeliene S, Polemiti E, Stevens C, et al. Foods and beverages and colorectal cancer risk: a systematic review and meta-analysis of cohort

489 studies, an update of the evidence of the WCRF-AICR Continuous Update Project. *Ann*
 490 *Oncol* 2017;28:1788–802. <https://doi.org/10.1093/annonc/mdx171>.
 491 [10] D’Elia L, Galletti F, Strazzullo P. Dietary salt intake and risk of gastric cancer. *Cancer*
 492 *Treat Res* 2014;159:83–95. https://doi.org/10.1007/978-3-642-38007-5_6.
 493 [11] Peleteiro B, Barros S, Castro C, Ferro A, Morais S, Lunet N. Worldwide burden of gastric
 494 cancer in 2010 attributable to high sodium intake in 1990 and predicted attributable
 495 burden for 2030 based on exposures in 2010. *Br J Nutr* 2016;116:728–33.
 496 <https://doi.org/10.1017/S0007114516002518>.
 497 [12] D’Elia L, Rossi G, Ippolito R, Cappuccio FP, Strazzullo P. Habitual salt intake and risk of
 498 gastric cancer: a meta-analysis of prospective studies. *Clin Nutr* 2012;31:489–98.
 499 <https://doi.org/10.1016/j.clnu.2012.01.003>.
 500 [13] Umesawa M, Iso H, Fujino Y, Kikuchi S, Tamakoshi A, JACC Study Group. Salty food
 501 preference and intake and risk of gastric cancer: The JACC study. *J Epidemiol*
 502 2016;26:92–7. <https://doi.org/10.2188/jea.JE20150023>.
 503 [14] Kune GA, Kune S, Watson LF. Dietary sodium and potassium intake and colorectal
 504 cancer risk. *Nutr Cancer* 1989;12:351–9.
 505 <https://doi.org/10.1080/01635588909514036>.
 506 [15] Alegria-Lertxundi I, Aguirre C, Bujanda L, Fernández FJ, Polo F, Ordovás JM, et al. Gene–
 507 diet interactions in colorectal cancer: Survey design, instruments, participants and
 508 descriptive data of a case–control study in the Basque Country. *Nutrients*
 509 2020;12:2362. <https://doi.org/10.3390/nu12082362>.
 510 [16] Negri E, La Vecchia C, D’Avanzo B, Gentile A, Boyle P, Franceschi S. Salt preference and
 511 the risk of gastrointestinal cancers. *Nutr Cancer* 1990;14:227–32.
 512 <https://doi.org/10.1080/01635589009514097>.

513 [17] Takachi R, Inoue M, Shimazu T, Sasazuki S, Ishihara J, Sawada N, et al. Consumption of
514 sodium and salted foods in relation to cancer and cardiovascular disease: the Japan
515 Public Health Center-based Prospective Study. *Am J Clin Nutr* 2010;91:456–64.
516 <https://doi.org/10.3945/ajcn.2009.28587>.

517 [18] Yakoob MY, Baig-ansari N. Dietary sodium (salt) intake and risk of colorectal cancer: A
518 systematic review. *Curr Dev Nutr* 2019;3:nzz030.P05-039-19.
519 <https://doi.org/10.1093/cdn/nzz030.P05-039-19>.

520 [19] Petit G, Jury V, Lamballerie M, Duranton F, Pottier L, Martin J. Salt intake from
521 processed meat products: Benefits, risks and evolving practices. *Compr Rev Food Sci*
522 *Food Safety* 2019;18:1453–73. <https://doi.org/10.1111/1541-4337.12478>.

523 [20] Ni Mhurchu C, Capelin C, Dunford EK, Webster JL, Neal BC, Jebb SA. Sodium content of
524 processed foods in the United Kingdom: analysis of 44,000 foods purchased by 21,000
525 households. *Am J Clin Nutr* 2011;93:594–600.
526 <https://doi.org/10.3945/ajcn.110.004481>.

527 [21] Inguglia ES, Zhang Z, Tiwari BK, Kerry JP, Burgess CM. Salt reduction strategies in
528 processed meat products – A review. *Trends Food Sci Technol* 2017;59:70–8.
529 <https://doi.org/10.1016/j.tifs.2016.10.016>.

530 [22] Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *Int J*
531 *Epidemiol* 2016;45:1866–86. <https://doi.org/10.1093/ije/dyw314>.

532 [23] Huang L, Crino M, Wu JHY, Woodward M, Barzi F, Land M-A, et al. Mean population
533 salt intake estimated from 24-h urine samples and spot urine samples: a systematic
534 review and meta-analysis. *Int J Epidemiol* 2016;45:239–50.
535 <https://doi.org/10.1093/ije/dyv313>.

- 536 [24] Mente A, O'Donnell M, Rangarajan S, Dagenais G, Lear S, McQueen M, et al.
537 Associations of urinary sodium excretion with cardiovascular events in individuals with
538 and without hypertension: a pooled analysis of data from four studies. *Lancet*
539 2016;388:465–75. [https://doi.org/10.1016/S0140-6736\(16\)30467-6](https://doi.org/10.1016/S0140-6736(16)30467-6).
- 540 [25] Brown IJ, Dyer AR, Chan Q, Cogswell ME, Ueshima H, Stamler J, et al. Estimating 24-
541 hour urinary sodium excretion from casual urinary sodium concentrations in Western
542 populations: the INTERSALT study. *Am J Epidemiol* 2013;177:1180–92.
543 <https://doi.org/10.1093/aje/kwt066>.
- 544 [26] Cogswell ME, Wang CY, Chen TC, Pfeiffer CM, Elliott P, Gillespie CD, et al. Validity of
545 predictive equations for 24-h urinary sodium excretion in adults aged 18-39 y. *Am J Clin*
546 *Nutr* 2013;98:1502–13. <https://doi.org/10.3945/ajcn.113.059436>.
- 547 [27] de Boer IH, Kestenbaum B. Invited commentary: Quantifying salt in urine--a complex
548 solution. *Am J Epidemiol* 2013;177:1193–5. <https://doi.org/10.1093/aje/kwt064>.
- 549 [28] Elliott P, Muller DC, Schneider-Luftman D, Pazoki R, Evangelou E, Dehghan A, et al.
550 Estimated 24-hour urinary sodium excretion and incident cardiovascular disease and
551 mortality among 398 628 individuals in UK Biobank. *Hypertension* 2020;76:683–91.
552 <https://doi.org/10.1161/HYPERTENSIONAHA.119.14302>.
- 553 [29] Inker LA, Schmid CH, Tighiouart H, Eckfeldt JH, Feldman HI, Greene T, et al. Estimating
554 glomerular filtration rate from serum creatinine and cystatin C. *N Eng J Med*
555 2012;367:20–9. <https://doi.org/10.1056/NEJMoa1114248>.
- 556 [30] Lees JS, Welsh CE, Celis-Morales CA, Mackay D, Lewsey J, Gray SR, et al. Glomerular
557 filtration rate by differing measures, albuminuria and prediction of cardiovascular
558

disease, mortality and end-stage kidney disease. *Nat Med* 2019;25:1753–60.
<https://doi.org/10.1038/s41591-019-0627-8>.

[31] Bradbury KE, Young HJ, Guo W, Key TJ. Dietary assessment in UK Biobank: an evaluation of the performance of the touchscreen dietary questionnaire. *J Nutr Sci* 2018;7:e6. <https://doi.org/10.1017/jns.2017.66>.

[32] Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet* 2014;23:R89–98.
<https://doi.org/10.1093/hmg/ddu328>.

[33] Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genetic Epidemiology* 2013;37:658–65. <https://doi.org/10.1002/gepi.21758>.

[34] Pazoki R, Evangelou E, Mosen-Ansorena D, Pinto RC, Karaman I, Blakeley P, et al. GWAS for urinary sodium and potassium excretion highlights pathways shared with cardiovascular traits. *Nat Commun* 2019;10:3653. <https://doi.org/10.1038/s41467-019-11451-y>.

[35] FinnGen study. FinnGen Biobank: Access results n.d.
https://www.finngen.fi/en/access_results (accessed October 5, 2020).

[36] Feng Q, Yuan S, Yang Q, Lu Y, Loos R, Li GH, et al. Causal associations between urinary sodium with body mass, shape and composition: a two-sample Mendelian randomization study. *Sci Rep* 2020;10:17475. <https://doi.org/10.1038/s41598-020-74657-x>.

[37] Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015;518:197–206.
<https://doi.org/10.1038/nature14177>.

- [38] Zanetti D, Rao A, Gustafsson S, Assimes TL, Montgomery SB, Ingelsson E. Identification of 22 novel loci associated with urinary biomarkers of albumin, sodium, and potassium excretion. *Kidney Intl* 2019;95:1197–208. <https://doi.org/10.1016/j.kint.2018.12.017>.
- [39] Hartwig FP, Davies NM, Hemani G, Davey Smith G. Two-sample Mendelian randomization: avoiding the downsides of a powerful, widely applicable but potentially fallible technique. *Int J Epidemiol* 2016;45:1717–26. <https://doi.org/10.1093/ije/dyx028>.
- [40] Pierce BL, Ahsan H, Vanderweele TJ. Power and instrument strength requirements for Mendelian randomization studies using multiple genetic variants. *Int J Epidemiol* 2011;40:740–52. <https://doi.org/10.1093/ije/dyq151>.
- [41] Zheng J, Baird D, Borges M-C, Bowden J, Hemani G, Haycock P, et al. Recent developments in Mendelian randomization studies. *Curr Epi Rep* 2017;4:330–45. <https://doi.org/10.1007/s40471-017-0128-6>.
- [42] Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018;50:693–8. <https://doi.org/10.1038/s41588-018-0099-7>.
- [43] Kamat MA, Blackshaw JA, Young R, Surendran P, Burgess S, Danesh J, et al. PhenoScanner V2: an expanded tool for searching human genotype-phenotype associations. *Bioinformatics* 2019;35:4851–3. <https://doi.org/10.1093/bioinformatics/btz469>.
- [44] Sanderson E, Spiller W, Bowden J. Testing and correcting for weak and pleiotropic instruments in two-sample multivariable Mendelian randomisation. *BioRxiv* 2020;2020.04.02.021980. <https://doi.org/10.1101/2020.04.02.021980>.

606 [45] Spencer EA, Key TJ, Appleby PN, Dahm CC, Keogh RH, Fentiman IS, et al. Meat, poultry
607 and fish and risk of colorectal cancer: pooled analysis of data from the UK dietary
608 cohort consortium. *Cancer Causes Control* 2010;21:1417–25.
609 <https://doi.org/10.1007/s10552-010-9569-7>.

610 [46] Lund H, Brunnhuber K, Juhl C, Robinson K, Leenaars M, Dorch BF, et al. Towards
611 evidence based research. *BMJ* 2016;i5440. <https://doi.org/10.1136/bmj.i5440>.

612 [47] Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, et al.
613 Carcinogenicity of consumption of red and processed meat. *Lancet Oncol*
614 2015;16:1599–600. [https://doi.org/10.1016/S1470-2045\(15\)00444-1](https://doi.org/10.1016/S1470-2045(15)00444-1).

615 [48] Li J, Sun F, Guo Y, Fan H. High-salt diet gets involved in gastrointestinal diseases
616 through the reshaping of gastroenterological milieu. *Digestion* 2019;99:267–74.
617 <https://doi.org/10.1159/000493096>.

618 [49] Thapa S, Fischbach LA, Delongchamp R, Faramawi MF, Orloff M. Association between
619 Dietary Salt Intake and Progression in the Gastric Precancerous Process. *Cancers*
620 (Basel) 2019;11. <https://doi.org/10.3390/cancers11040467>.

621 [50] Kato S, Tsukamoto T, Mizoshita T, Tanaka H, Kumagai T, Ota H, et al. High salt diets
622 dose-dependently promote gastric chemical carcinogenesis in *Helicobacter pylori*-
623 infected Mongolian gerbils associated with a shift in mucin production from glandular
624 to surface mucous cells. *Int J Cancer* 2006;119:1558–66.
625 <https://doi.org/10.1002/ijc.21810>.

626 [51] Thapa S, Fischbach LA, Delongchamp R, Faramawi MF, Orloff M. The Association
627 between Salt and Potential Mediators of the Gastric Precancerous Process. *Cancers*
628 (Basel) 2019;11. <https://doi.org/10.3390/cancers11040535>.

629 [52] Tamura M, Matsui H, Nagano YN, Kaneko T, Indo HP, Majima HJ, et al. Salt is an
630 oxidative stressor for gastric epithelial cells. *J Physiol Pharmacol* 2013;64:89–94.

631 [53] Malsure S, Wang Q, Charles R-P, Sergi C, Perrier R, Christensen BM, et al. Colon-specific
632 deletion of epithelial sodium channel causes sodium loss and aldosterone resistance. *J*
633 *Am Soc Nephrol* 2014;25:1453–64. <https://doi.org/10.1681/ASN.2013090936>.

634 [54] Sandle GI. Salt and water absorption in the human colon: a modern appraisal. *Gut*
635 1998;43:294–9. <https://doi.org/10.1136/gut.43.2.294>.

636 [55] Lienhard D, Lauterburg M, Escher G, Frey FJ, Frey BM. High salt intake down-regulates
637 colonic mineralocorticoid receptors, epithelial sodium channels and 11 β -
638 hydroxysteroid dehydrogenase type 2. *PLoS ONE* 2012;7:e37898.
639 <https://doi.org/10.1371/journal.pone.0037898>.

640 [56] Aguiar SLF, Miranda MCG, Guimarães MAF, Santiago HC, Queiroz CP, Cunha P da S, et
641 al. High-salt diet induces IL-17-dependent gut inflammation and exacerbates colitis in
642 mice. *Front Immunol* 2017;8:1969. <https://doi.org/10.3389/fimmu.2017.01969>.

643 [57] Graudal N, Hubeck-Graudal T, Jürgens G, Taylor RS. Dose-response relation between
644 dietary sodium and blood pressure: a meta-regression analysis of 133 randomized
645 controlled trials. *Am J Clin Nutr* 2019;109:1273–8.
646 <https://doi.org/10.1093/ajcn/nqy384>.

647 [58] Jürgens G, Graudal NA. Effects of low sodium diet versus high sodium diet on blood
648 pressure, renin, aldosterone, catecholamines, cholesterols, and triglyceride. *Cochrane*
649 *Database Syst Rev* 2004:CD004022.
650 <https://doi.org/10.1002/14651858.CD004022.pub2>.

- [59] Graudal N, Jürgens G, Baslund B, Alderman MH. Compared with usual sodium intake, low- and excessive-sodium diets are associated with increased mortality: a meta-analysis. *Am J Hypertens* 2014;27:1129–37. <https://doi.org/10.1093/ajh/hpu028>.
- [60] He FJ, Li J, MacGregor GA. Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ* 2013;346:f1325–f1325. <https://doi.org/10.1136/bmj.f1325>.
- [61] Huang L, Trieu K, Yoshimura S, Neal B, Woodward M, Campbell NRC, et al. Effect of dose and duration of reduction in dietary sodium on blood pressure levels: systematic review and meta-analysis of randomised trials. *BMJ* 2020;368:m315. <https://doi.org/10.1136/bmj.m315>.
- [62] Pagoni P, Dimou NL, Murphy N, Stergiakouli E. Using Mendelian randomisation to assess causality in observational studies. *Evid Based Mental Health* 2019;22:67–71. <https://doi.org/10.1136/ebmental-2019-300085>.

675 Figure legend

676 Figure 1: The flowchart of method framework and data sources

677 GWAS: Genome-wide association study. MR: Mendelian Randomization. BMI: Body Mass Index.

678 GIANT consortium: Genetic Investigation of Anthropometric Traits consortium. SNP: Single

679 Nucleotide Polymorphism.

680