Chu, A. L., Hickman, M., Steel, N., Jones, P., Davey Smith, G., & Khandaker, G. (2021). Inflammation and Depression: A Public Health Perspective. *Brain, Behavior, and Immunity*, 95, 1-3.
<https://doi.org/10.1016/j.bbi.2021.04.015>

Peer reviewed version

License (if available):
CC BY

Link to published version (if available):
[10.1016/j.bbi.2021.04.015](https://doi.org/10.1016/j.bbi.2021.04.015)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.bbi.2021.04.015> . Please refer to any applicable terms of use of the publisher.

This research was funded in whole, or in part, by the Wellcome Trust [Grant number 201486/Z/16/Z]. For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. This work is licensed under a Creative Commons Attribution 4.0 International License.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Inflammation and Depression: A Public Health Perspective

Alexander L. Chu^{a,*}, Matthew Hickman^{b,c,d,e}, Nicholas Steel^f, Peter B. Jones^{g,h}, George Davey Smith^{b,e}, Golam M. Khandaker^{e,g,h,i}

Author Institutional Affiliations:

^a Dell Medical School, University of Texas at Austin, Austin, TX, USA

^b Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

^c NIHR Bristol Biomedical Research Centre, University of Bristol, Bristol, UK

^d NIHR Health Protection Research Unit in Behavioural Science and Evaluation, University of Bristol, Bristol, UK

^e MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

^f Norwich Medical School, University of East Anglia, Norwich, UK

^g Department of Psychiatry, University of Cambridge, Herchel Smith Building, Robinson Way, Cambridge, UK

^h Cambridgeshire and Peterborough NHS Foundation Trust, Elizabeth House, Fulbourn Hospital, Cambridge, UK

ⁱ Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

*Corresponding Author:

Alexander L. Chu, MPhil PH

Dell Medical School, University of Texas at Austin

1501 Red River St,

Austin, TX, USA 78712

Email: alexanderchu248@utexas.edu

Keywords: Mental health; Depression; Immunopsychiatry; Inflammation; Public Health; Prevention

Word Count:

- **Abstract:** 130
- **Manuscript:** 999 (excluding section headings, sub-headings, tables and references)

Figure(s): 1

- Figure 1. Target populations for primary and secondary prevention of inflammation-related depression

References: 10

Abstract

The multifaceted role of low-grade systemic inflammation in depression and physical illnesses like cardiovascular disease highlights complex interactions between the body, brain and mind. While current research on inflammation and depression has largely focused on exploring possible disease mechanisms and its therapeutic potential, we seek to broaden the current discussion by introducing a public health perspective. In this Viewpoint, we propose that inflammation and its contributing sources could represent important targets for public health strategies aimed at improving both mental and physical health. We discuss potential universal, selective and indicated primary prevention strategies for inflammation-related depression. We also highlight potential approaches for secondary prevention, including scope for anti-inflammatory treatment and CRP testing for predicting treatment prognosis. Preventive strategies discussed here could also be relevant for other inflammation-mediated mental health conditions.

Inflammation as a public health target for chronic physical and mental illnesses

Low-grade systemic inflammation (hereinafter referred to as inflammation), characterised by persistent sub-acute elevated levels of circulating inflammatory markers like C-reactive protein (CRP) and interleukin-6 (IL-6) (Danesh, 2000), is implicated in the pathogenesis of many chronic physical and mental conditions, including coronary heart disease (CHD), type 2 diabetes mellitus (T2DM), depression and schizophrenia (Berk et al., 2013). Current research has predominantly focused on understanding possible pathophysiological roles of inflammation and its therapeutic potential. In this Viewpoint, we propose that inflammation and its modifiable sources represent important targets for public health interventions aimed at improving physical and mental health. We focus on depression as an exemplar for inflammation-mediated mental disorders given its significant disease burden and strong link with inflammation. However, we acknowledge that our discussion is also relevant for other inflammation-mediated conditions and that other disease mechanisms like hypothalamic-pituitary-adrenal axis dysfunction can work in conjunction with or independently of inflammation.

Evidence triangulation from various approaches support a causal role for inflammation in depression. These include inflammation-induced sickness behaviour resembling somatic/neurovegetative symptoms of depression in rodents and primates (Dantzer et al., 2008), case-control studies reporting elevated concentrations of CRP and other inflammatory mediators in depressed patients (Osimo et al., 2020) and randomised controlled trials (RCTs) reporting antidepressant effects for anti-inflammatory drugs (Kappelmann et al., 2018). Causal inferential techniques like Mendelian randomisation show that genetic variants

regulating levels/activities of CRP and IL-6 are associated with depression (Khandaker et al., 2020)

Recent evidence showing linear trends of increasing depression risk with increasing population-wide CRP levels suggests that inflammation and depression could share a similar relationship like blood pressure and stroke (Ye et al., 2021). This makes inflammation an ideal candidate for public health interventions. Using conservative estimates for prevalence of inflammation (CRP, 3-10mg/L) in depression (16%) and a corresponding risk ratio (1.76), we estimate the population attributable fraction of inflammation for depression to be about 7%, meaning roughly 1 in 15 depression cases may be attributed to inflammation. This potentially equates to millions globally, given an estimated depression lifetime prevalence of 10-20% (Lim et al., 2018).

Primary preventive strategies for inflammation-related depression

Universal, selective and indicated preventive strategies directed towards general, at-risk and high-risk populations, respectively, could be a useful framework for thinking about the primary prevention of inflammation-related depression (Figure 1).

Figure 1. Target populations for primary and secondary prevention of inflammation-related depression

Note: Relative size of each box is not representative of the true size for each respective population group.

The main goal of universal preventive strategies would be to reduce inflammation levels in the general population. Such strategies could involve interventions aimed at achieving regular physical activity, weight reduction and diets with a low dietary inflammatory index. While universal preventive strategies possess the greatest potential in reducing incidence of depression, no RCT, to our knowledge, has investigated the impact of universal preventive interventions targeting inflammation on depression risk. We now require large-scale studies

that examine a) the impact of weight reduction, dietary and other interventions on inflammatory levels in the general population; b) the impact of such interventions on inflammation-related depression incidence.

Selective preventive strategies target at-risk individuals whose risk of depression is higher than the general population and seek to reduce inflammation levels by addressing underlying pro-inflammatory risk factors and promoting relevant protective factors. Typical examples of this approach may include promoting sustained physical activity among patients with chronic diseases and stress management programmes in high-intensity occupational settings. Key issues to consider include determining the appropriate at-risk group and choice of intervention. Because depression risk is higher in at-risk groups compared to those in the general population, assessing the impact of selective preventive interventions on inflammation levels and subsequent depression risk may be a more feasible first step.

Indicated preventive strategies target “high-risk” individuals, i.e., those exhibiting depressive symptoms but are below the threshold for a depression diagnosis. One important goal for indicated preventive strategies is improving detection of high-risk individuals for clinical monitoring, management and/or referral. Whether evidence of inflammation in symptomatic individuals increases risk of progressing to clinical depression requires further evaluation. Adapting existing clinical models like collaborative care could be useful for delivering preventive interventions for symptomatic, help-seeking individuals. Further integration of physical and mental healthcare, including increased mental health input from primary to tertiary care settings, could improve monitoring of depressive symptoms and use of self-help interventions. This approach could be particularly useful for individuals with chronic inflammatory physical illnesses. However, it is worth noting that increasing detection and

monitoring of mildly symptomatic individuals could contribute to the over-medicalisation of depression.

Secondary preventive strategies for inflammation-related depression

Inflammation is associated with poor response to conventional antidepressants and psychotherapies, and it has been proposed that depressed patients with evidence of inflammation may be suitable candidates for anti-inflammatory treatment. Targeting inflammation may also be a valuable target for the secondary prevention of depression, which aims to effectively treat and prevent recurrence (Figure 1). Inflammation could specifically be a useful means for guiding treatment decision and predicting/monitoring response.

Meta-analyses of RCTs have showed that adjunct anti-inflammatory drugs provide greater benefit than antidepressants alone (effect size ~0.35) and that anti-inflammatory drugs improve depressive symptoms in individuals with chronic inflammatory diseases independently of improvements in physical illness (effect size ~0.40). Emerging evidence from RCTs suggest that anti-inflammatory treatment could be useful for depressed individuals with evidence of inflammation (CRP >3 mg/L) (Nettis et al., 2021; Raison et al., 2013). Future large-scale clinical trials are needed to test efficacy of anti-inflammatory drugs as monotherapy or adjunct therapy for depression. Studies are also needed to examine whether assessing inflammatory markers and identifying a subset of newly diagnosed patients with inflammation-related depression could be a cost-effective way of predicting treatment response, finding more suitable, personalised treatment regimens and improving treatment outcomes.

Conclusion

As our understanding of the complex interactions between the body, brain and mind improves, this understanding needs to be matched with a concerted effort to develop not only

new treatments but also public health strategies that recognise the broad-spectrum impact of targeting inflammation and its modifiable sources. By offering an initial primary and secondary prevention framework for inflammation-related depression, we call to attention the need for more interdisciplinary research that will drive integration between physical and mental healthcare and other innovative public health approaches.

Declaration of Interests

All authors do not have any conflicts of interest to report regarding this specific work. MH reports receiving personal fees from MSD Gilead. PBJ is a scientific advisory board chair for Ricordati and a scientific advisory board member for Janssen.

Funding

GMK acknowledges funding support from the Wellcome Trust (grant code: 201486/Z/16/Z), the MQ: Transforming Mental Health (grant code: MQDS17/40), the Medical Research Council UK (grant code: MC_PC_17213 and MR/S037675/1) and the BMA Foundation (J Moulton grant 2019). GDS works in the Medical Research Council Integrative Epidemiology Unit at the University of Bristol (MC_UU_00011/1). PBJ acknowledges funding from the MRC and MQ (grant code: MQDS17/40), programmatic funding from NIHR (RP-PG-0616-20003) and support from the Applied Research Collaboration East of England. MH acknowledges support from the NIHR BRC at Bristol and the NIHR HPRU in Behavioural Science and Evaluation.

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors. The funding sources listed above did not have any role in the writing of this article or in the decision to submit the paper for publication.

Acknowledgements

We are grateful to Mr. Peter Templeton, Executive Director of IfM Education and Consultancy Service, a University of Cambridge knowledge transfer company, and Founder of the mental health charity The Foundation for Young People's Mental Health, for his helpful comments.

References

- Berk, M., Williams, L.J., Jacka, F.N., O’Neil, A., Pasco, J.A., Moylan, S., Allen, N.B., Stuart, A.L., Hayley, A.C., Byrne, M.L., Maes, M., O’Neil, A., Pasco, J.A., Moylan, S., Allen, N.B., Stuart, A.L., Hayley, A.C., Byrne, M.L., Maes, M., 2013. So depression is an inflammatory disease, but where does the inflammation come from? *BMC Med.* 11, 200. doi:10.1186/1741-7015-11-200
- Danesh, J., 2000. Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *BMJ* 321, 199–204. doi:10.1136/bmj.321.7255.199
- Dantzer, R., O’Connor, J.C., Freund, G.G., Johnson, R.W., Kelley, K.W., 2008. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat. Rev. Neurosci.* 9, 46–56. doi:10.1038/nrn2297
- Kappelmann, N., Lewis, G., Dantzer, R., Jones, P.B., Khandaker, G.M., 2018. Antidepressant activity of anti-cytokine treatment: a systematic review and meta-analysis of clinical trials of chronic inflammatory conditions. *Mol. Psychiatry* 23, 335–343. doi:10.1038/mp.2016.167
- Khandaker, G.M., Zuber, V., Rees, J.M.B., Carvalho, L., Mason, A.M., Foley, C.N., Gkatzionis, A., Jones, P.B., Burgess, S., 2020. Shared mechanisms between coronary heart disease and depression: findings from a large UK general population-based cohort. *Mol. Psychiatry* 25, 1477–1486. doi:10.1038/s41380-019-0395-3
- Lim, G.Y., Tam, W.W., Lu, Y., Ho, C.S., Zhang, M.W., Ho, R.C., 2018. Prevalence of Depression in the Community from 30 Countries between 1994 and 2014. *Sci. Rep.* 8, 2861. doi:10.1038/s41598-018-21243-x
- Nettis, M.A., Lombardo, G., Hastings, C., Zajkowska, Z., Mariani, N., Nikkheslat, N., Worrell, C., Enache, D., McLaughlin, A., Kose, M., Sforzini, L., Bogdanova, A., Cleare, A., Young, A.H., Pariante, C.M., Mondelli, V., 2021. Augmentation therapy with

minocycline in treatment-resistant depression patients with low-grade peripheral inflammation: results from a double-blind randomised clinical trial.

Neuropsychopharmacology 46, 939–948. doi:10.1038/s41386-020-00948-6

Osimo, E.F., Pillinger, T., Rodriguez, I.M., Khandaker, G.M., Pariante, C.M., Howes, O.D.,
2020. Inflammatory markers in depression: A meta-analysis of mean differences and
variability in 5,166 patients and 5,083 controls. Brain. Behav. Immun.
doi:10.1016/j.bbi.2020.02.010

Raison, C.L., Rutherford, R.E., Woolwine, B.J., Shuo, C., Schettler, P., Drake, D.F., Haroon,
E., Miller, A.H., 2013. A Randomized Controlled Trial of the Tumor Necrosis Factor
Antagonist Infliximab for Treatment-Resistant Depression. JAMA Psychiatry 70, 31.
doi:10.1001/2013.jamapsychiatry.4

Ye, Z., Kappelmann, N., Moser, S., Smith, G.D., Burgess, S., Jones, P.B., Khandaker, G.M.,
2021. Role of Inflammation in Depression and Anxiety: Tests for Disorder Specificity,
Linearity and Potential Causality of Association in the UK Biobank. medRxiv
2021.02.02.21250987. doi:10.1101/2021.02.02.21250987