Buss, H. L., Mathur, R., White, A. F., & Brantley, S. L. (2010). Iron and phosphorus cycling in deep saprolite, Luquillo Mountains, Puerto Rico. *Chemical Geology*, 269, 52-61.
<https://doi.org/10.1016/j.chemgeo.2009.08.001>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.chemgeo.2009.08.001](https://doi.org/10.1016/j.chemgeo.2009.08.001)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Chemical Geology Manuscript #4604 Revision, Special Volume Honoring Tim Drever

Phosphorus and iron cycling in deep saprolite,

Luquillo Mountains, Puerto Rico

Heather L. Buss^{*1}, Ryan Mathur², Arthur F. White¹, and Susan L. Brantley³

July 14, 2009

* Corresponding Author. Tel: +1-650-329-4420, Fax: +1-650-329-4538, hlbuss@usgs.gov.

1. U.S. Geological Survey, 345 Middlefield Rd., MS 420, Menlo Park, CA 94040

2. Department of Geology, Juniata College, Huntingdon, PA 16652

3. Earth and Environmental Systems Institute, The Pennsylvania State University, University
Park, PA 16802

Abstract

Rapid weathering and erosion rates in mountainous tropical watersheds lead to highly variable soil and saprolite thicknesses which in turn impact nutrient fluxes and biological populations. In the Luquillo Mountains of Puerto Rico, a 5 meter-thick saprolite contains high microorganism densities at the surface and at depth overlying bedrock. We test the hypotheses that the organisms at depth are limited by the availability of two nutrients, P and Fe. Many tropical soils are P limited, rather than N limited, and dissolution of apatite is the dominant source of P. We document patterns of apatite weathering and of bioavailable Fe derived from the weathering of primary minerals hornblende and biotite in cores augered to 7.5 m on a ridgetop as compared to spheroidally weathering bedrock sampled in a nearby roadcut.

Iron isotopic compositions of 0.5 N HCl extracts of soil and saprolite range from about $\delta^{56}\text{Fe} = 0$ to -0.1 ‰ throughout the saprolite except at the surface and at 5 m depth where $\delta^{56}\text{Fe} = -0.26$ to -0.64 ‰. The enrichment of light isotopes in HCl-extractable Fe in the soil and at the saprolite-bedrock interface is consistent with active Fe cycling and consistent with the locations of high cell densities and Fe(II)-oxidizing bacteria, identified previously. To evaluate the potential P-limitation of Fe-cycling bacteria in the profile, solid-state concentrations of P were measured as a function of depth in the soil, saprolite, and weathering bedrock. Weathering apatite crystals were examined in thin sections and an apatite dissolution rate of $6.8 \times 10^{-14} \text{ mol m}^{-2} \text{ s}^{-1}$ was calculated. While surface communities depend on recycled nutrients and atmospheric inputs, deep communities survive primarily on nutrients released by the weathering bedrock and thus are tightly coupled to processes related to saprolite formation including mineral weathering.

While low available P may limit microbial activity within the middle saprolite, fluxes of P from apatite weathering should be sufficient to support robust growth of microorganisms in the deep saprolite.

Keywords: phosphorus; iron isotopes; saprolite; apatite weathering rate; Fe(II)-oxidizing bacteria

1. Introduction

Except where recent landslides have occurred, thick saprolites in the Luquillo Mountains isolate the weathering bedrock from the soil ecosystem. Despite high precipitation ($>4000 \text{ mm y}^{-1}$) in the Rio Icacos watershed of the Luquillo Mountains, most water that reaches the soil moves through the shallow soil layers and the water that does infiltrate to the deep saprolite on the ridges does so slowly: $1.8 - 2.6 \text{ m y}^{-1}$ (Turner, 2001). Established tropical forests are typically characterized by a closed nutrient cycle in which nutrients are tightly cycled within the surface soil and biomass because the underlying substrates are highly weathered and thus nutrient poor (e.g., Bruijnzeel, 1991). The majority of nutrients in precipitation and dry deposition in the Rio Icacos watershed are thus likely rapidly scavenged by the abundant tropical vegetation and soil organisms, leaving little to infiltrate to subsurface communities, as evidenced by very low DOC concentrations in the deep saprolite porewaters (Murphy, 1995). Indeed, tracers such as Sr and Si isotope ratios and Si/Ge ratios indicate that while vegetation and porewaters in the soil and shallow saprolite of the Rio Icacos watershed contain atmospherically derived elements, middle and deep saprolite porewaters (collected from nested suction water samplers) are dominated by elements derived from bedrock weathering (Kurtz and Derry, 2004; Ziegler et al., 2005; Pett-

Ridge et al., 2009; Lugolobi et al., Subm.). Furthermore, the majority of the annual precipitation falls as short, intense storm events during which runoff to the streams occurs through the shallow saprolite only (Lugolobi et al., 2006). Deeper saprolite microbial communities are therefore highly dependent upon weathering bedrock for nutrients (Buss et al., 2005). In the deep saprolite of the Rio Icacos watershed, cell densities increase near the bedrock-saprolite interface where Fe(II)-oxidizing bacteria were detected. Maximum growth rates were estimated for these chemolithoautotrophic microorganisms and for their dependent heterotrophic communities based on the rate of release of Fe(II) from silicate minerals during weathering at the bedrock-saprolite interface (Buss et al., 2005). In Buss et al. (2005), cell densities (live + dead) in the saprolite were estimated by direct counting and from DNA yields. The activity of the microorganisms was not examined and it is possible that the organisms present at depth were merely dormant due to P-limitation. In this paper, we compare P fluxes and P requirements to growth rates based on Fe(II) fluxes to test whether the community at depth in these samples was nutrient (P)-limited or energy (Fe(II))-limited. We also use Fe isotopes to examine the activity of Fe-cycling microorganisms in the saprolite profile.

Phosphorus is an essential macronutrient, vital to all biological systems. Adenosine triphosphate (ATP), one of the most important and abundant biomolecules in nature, is formed during photosynthesis and cellular respiration and functions as an energy shuttle, providing chemical energy for biosynthesis, cell motility, and osmotic transport of molecules through cell membranes (Garrett and Grisham, 1999). Phosphorus is found in DNA and RNA as phosphate ester bridges that link individual nucleotides, in cell membranes as phospholipids, and as a component of many signaling molecules, enzymes, and coenzymes (Frausto da Silva and Williams, 191; Garrett and Grisham, 1999). However, P availability is commonly very low in

tropical soils, a fact that is often attributed to the strong sorption of P to Fe and Al oxides (e.g., Sanchez, 1976; Vitousek, 1984; Olander and Vitousek, 2005). In contrast, tropical soils are typically not N limited (e.g., Vitousek and Sanford, 1986). In the tropical Luquillo Mountains of northeastern Puerto Rico, low rates of NO_3^- leaching and export (McDowell and Asbury, 1994) result primarily from effective N retention via dissimilatory nitrate reduction to ammonium (Templer et al., 2008). High rates of NH_4^+ production via rapid decomposition of soil organic matter also contribute to N availability and retention in tropical soils (Vitousek and Sanford, 1986). In saprolite porewaters in the Rio Icacos watershed, N concentrations increase with depth on both ridgetops and slopes (White et al., 1998) suggesting that, like soils, deep saprolite is also not N limited. It is interesting to note that of the two elements that are most often limiting in terrestrial systems, N and P, only P comes directly from rocks (Wardle et al., 2004).

Pools of P in a forest soil may include primary minerals (predominantly apatite), secondary minerals (associated with Fe or Al oxides or clays), aqueous species in pore waters (inorganic or organic complexes), and solid organic P (contained in biomass and other organic matter). External sources of P to a soil system include bedrock weathering and atmospheric deposition (precipitation and dust). Most of the P associated with secondary minerals, biomass, and organic matter represent internal sources of P that are recycled and transferred between pools on various timescales during the lifetime of an ecosystem. In most soil profiles, P is supplied into the system almost entirely by weathering due to low atmospheric input of P (Walker and Syers, 1976). However, Caribbean island soils receive significant nutrient inputs from African dust and volcanic ash (Graham and Duce, 1982; Heartsill-Scalley et al., 2007; Muhs et al., 2007). Thick, highly weathered saprolites on the ridges of the Luquillo Mountains of Puerto Rico effectively isolate surface soils from weathering bedrock (e.g., Buss et al., 2005), thus atmospheric

deposition may dominate exogenous input of P to the ridgetop soils (Pett-Ridge, 2009). In contrast, mineral weathering is expected to dominate P input to deep saprolite or where saprolite is thin due to topography or erosion. Landslides, which are the dominant mechanism of erosion in the Luquillo Mountains (Simon et al., 1990), increase access to mineral-derived nutrients, notably apatite, which affects the rate of vegetation re-growth on landslide scars (e.g., Zarin and Johnson, 1995; Frizano et al., 2002).

Stable Fe isotopes provide a promising new tool for tracing biogeochemical Fe cycles in soils and weathering systems. Iron isotope systematics have been studied for multiple processes that are relevant to soil systems including i) inorganic dissolution (mobilization of Fe from minerals by H^+), ii) organic dissolution (mobilization of Fe by organic ligands or microorganisms), iii) reductive dissolution (reduction of mineral-bound Fe(III) followed by mobilization of Fe(II)), iv) precipitation of Fe(III)-(hydr)oxides from solution, v) adsorption of Fe onto oxide or clay minerals, and vi) uptake of Fe by plants or microorganisms. Fractionations favoring the release of the light isotope (^{54}Fe) to solution leaving an isotopically heavy solid (enriched in ^{56}Fe) have been documented for organic dissolution of hornblende by bacteria, organic acids, and siderophores (Brantley et al., 2001b; 2004) and for ligand-controlled Fe relocation in soils (Wiederhold et al., 2007b). Organic dissolution of goethite did not produce an isotopic fractionation (Brantley et al., 2004), nor did inorganic dissolution of hornblende or goethite (Brantley et al., 2001b; 2004). Reductive dissolution of goethite, hematite, and ferrihydrite in the presence of bacteria generates a solution that is isotopically light with respect to the starting mineral (e.g., Beard et al., 1999; 2003; Icopini et al., 2004; Crosby et al., 2005; 2007), which leads to a loss of isotopically light Fe from soils (Thompson et al., 2007; Wiederhold et al., 2007b). However, fractionation may occur via isotopic exchange between

aqueous Fe(II) and sorbed Fe(II) , rather than during reductive dissolution, because adsorption of Fe(II) onto goethite preferentially removes ^{56}Fe from solution, with or without bacteria (Brantley et al., 2004; Icopini et al., 2004). Similarly, precipitation of ferrihydrite also concentrates the heavier isotope in a solid product while the solution becomes isotopically lighter (Bullen et al., 2001; Teutsch et al., 2005). Plants that obtain Fe by acidifying the rhizosphere to promote reductive dissolution of Fe(III)-oxides preferentially take up Fe that is isotopically light relative to the surrounding soil (Guelke and Von Blanckenburg, 2007). In contrast, uptake of Fe(III)-siderophore complexes does not result in a fractionation of Fe in grasses that produce phytosiderophores (Guelke and Von Blanckenburg, 2007).

Here we use Fe isotope ratios to investigate controls on the bioavailable fraction of Fe. Iron isotopes are measured in the fraction of Fe liberated from soil and saprolite samples by a dilute HCl leach, which is often associated with the most reactive, or bioavailable, fraction of Fe in soils and sediments (e.g., Fantle and DePaolo, 2004; Buss et al., 2005; Severmann et al., 2006; Wiederhold et al., 2007b). In addition to colloid-bound Fe, organic-bound Fe, and the (typically minute) aqueous Fe(II), it is this “reactive Fe” that is transported from weathering profiles to rivers and oceans and dominates the Fe isotope signatures of these bodies of water (Fantle and DePaolo, 2004). In high silicate soils, Fe within this reactive pool is expected to be isotopically light relative to bedrock and residual material (Fantle and DePaolo, 2004; Wiederhold et al., 2007b), which may indicate active biogeochemical cycling of Fe. For example, complexation of Fe by organic ligands or reductive dissolution of Fe(III)-(hydr)oxides would cause an excursion towards a lighter Fe isotope signature within the reactive Fe pool, while Fe uptake by vegetation would likely yield a heavier Fe isotope signature (Fantle and DePaolo, 2004; Wiederhold et al., 2006; Wiederhold et al., 2007a, b). In soils or regolith where the residual (non-HCl extractable)

Fe is dominated by Fe(III)-hydroxides, isotopic distinction between the two pools may be less than in silicate soils due to rapid isotopic exchange between sorbed Fe(II) and Fe(III)-hydroxides (Pedersen et al., 2005). Here we test the hypothesis that HCl-extractable Fe will be isotopically light in the biologically active zones of the Rio Icacos weathering profile, relative to zones in which biological activity is suppressed due to limited nutrient and energy fluxes.

2. Methods

2.1. Field Site and Sample Collection

The Rio Icacos watershed in the Luquillo Experimental Forest of northeastern Puerto Rico is predominantly located in a lower montane wet colorado forest that receives over 4000 mm of rain annually. The watershed is underlain by the Rio Blanco quartz diorite bedrock, which weathers spheroidally resulting in rounded corestones of about 2 m in diameter, surrounded by 0.2 – 2 m thick zones of concentric, partially weathered rock layers called rindlets (Turner et al., 2003; Buss et al., 2004; Buss et al., 2005; Fletcher et al., 2006; Buss et al., 2008). The rindlet zones are mantled by 200 ka regolith, dated using ^{10}Be (Brown et al., 1995), and is comprised of 2-8 m of saprolite and 0.5-1 m of highly weathered Picacho-Ciales complex soils, previously classified as Ultisols (Boccheciamper et al., 1977), but now considered Inceptisols due to weak B horizon development (USDA NCRS, 2002).

Two 5.7 cm diameter cores of soil and saprolite were collected 1 m apart by hand augering to 7.5 m and 5.0 m depth at the top of the Guaba Ridge in the Rio Icacos watershed (latitude 18°16.903'N, longitude 65°47.418'W), which has been mapped by others (Murphy et al., 1998; White et al., 1998; Turner et al., 2003; Buss et al., 2008). Samples were taken from the auger at 0.15 – 0.3 m depth intervals. The deeper core penetrated the rindlet zone below 5 m

depth. The sampling procedure was described by Buss et al. (2005). Redoximorphic features such as gleying or mottling were not observed in the saprolite samples, but several zones of pink and black Mn-oxide concentrations were identified (Buss et al., 2005). The saprolite is formed on a cuchillo, or knife-edge ridge, and is not significantly influenced by lateral flow of groundwater at depth (White et al., 1998; Lugolobi et al., 2006). The pH of the soil and saprolite porewater is 4.0 – 5.4 (White et al., 1998). The edge of a corestone and the complete 49 cm section of rindlets above the corestone were sampled at a nearby roadcut exposed on Route 191 (latitude 18°16.216'N, longitude 65°47.055'W), preserving the position and orientation of the individual rindlets as described by Buss et al. (2008). The corestone was located beneath about 2 m of saprolite forming a small local topographic high.

2.2. Phosphorus and Apatite Analyses

Phosphorus was measured by inductively coupled plasma atomic emission spectrometry (ICP-AES) after lithium metaborate fusion digestion (SGS Mineral Laboratories, Ontario, Canada) of pulverized and sieved (150 µm) corestone, rindlet, and saprolite samples. Apatite crystals were identified in thin sections of the corestone and rindlets using backscattered electron (BSE) imaging and energy dispersive X-ray spectrometry (EDS) on an FEI Quanta 400 SEM. Point counting of apatite crystals was performed on backscattered images using XT-Docu software (v. 3.2, Soft-Imaging System GmbH, Münster, Germany), which calculates the volume of each user-defined phase based on the backscatter grayscale value. This technique has been used to measure porosity and to point count mineral phases in weathering rocks (Dorn, 1995; Dixon et al., 2006; Buss et al., 2008).

2.3. Fe Isotope Analysis

Iron in soils and sediments exist in many different forms (e.g., as readily exchangeable ions; precipitated as, or bound to, oxide or hydroxide phases; bound to organic matter; in carbonates; or in silicates). These fractions are often extracted sequentially by stronger and stronger leaching procedures. Extractions are favored over total digests because only average isotope ratios of all different Fe phases are obtained from total digests. Extraction of soils and sediments with dilute HCl is expected to liberate readily exchangeable ions, adsorbed Fe, some organic-bound Fe, and poorly crystalline Fe oxides and hydroxides (Wiederhold et al., 2007b) and represent a more reactive, or bioavailable, Fe pool than the bulk solid (Emerson, 2002). Iron isotopes were measured in 0.5 N HCl extracts performed on soil and saprolite samples augered from the Guaba Ridge. Iron isotopes are not fractionated by this extraction procedure: fractionation did not occur during dissolution of goethite or hematite in 0.5 N HCl, even when dissolution was incomplete (Skulan et al., 2002; Wiederhold et al., 2006). Soil and saprolite samples were air dried and gently ground only to break up aggregates before sieving to 2 mm. Samples (0.5 g) were then combined with 10 ml 0.5 N HCl in 15 mL centrifuge tubes and agitated on an end-over-end shaker at room temperature for 17 hours, followed by centrifugation at 4000 x g for 10 minutes (Buss et al., 2005). Supernatants were decanted, centrifuged again, and filtered to 0.4 μ m (SFCA-membrane syringe filter, Cole-Parmer). Concentrations of ferrous iron in the extracts were measured by ultraviolet-visible (UV/Vis) spectrophotometry as described and reported by Buss et al. (2005). Total Fe was measured by inductively coupled

plasma mass spectrometry (ICP-MS) and ferric iron was calculated as the difference between total and ferrous iron concentrations, as detailed in Buss et al. (2005).

Iron was separated from other cations in the HCl extracts following a standard ion exchange column method using AG MP-1 (100-200 mesh) resin (BIORAD) as described by Brantley et al. (2004). Post-purification iron yield was measured volumetrically and found to be better than 95%. Fe isotopes were measured at Washington State University using a multi-collector inductively coupled plasma mass spectrometer (MC-ICP-MS, Neptune®, ThermoFinnigan). The Fe concentration in each sample was adjusted to 3 ppm and a 3 ppm Cu standard was added to each. Sample measurements were bracketed with the standard-sample-standard technique. The mass bias in each sample was corrected using the exponential law (Arnold et al., 2004) and by using the graphical techniques developed by Maréchal et al. (1999). Both methods of data reduction resulted in values that were within the reported 2σ error (± 0.1 ‰). This error was calculated by using multiple measurements of the Fe isotope ratio of goethite throughout four weeks of analysis (sample measured 25 times). The $\delta^{56}\text{Fe}$ (‰) values were calculated relative to the international Fe standard IRMM-014, using the following equation:

$$\delta^{56/54}\text{Fe} = \left(\frac{(^{56}\text{Fe}/^{54}\text{Fe})_{\text{sample}}}{(^{56}\text{Fe}/^{54}\text{Fe})_{\text{IRMM}}} - 1 \right) \times 1000 \quad (1).$$

3. Results

To evaluate mass transfer (loss or gain of an element relative to the parent rock composition) of P in the Rio Icacos weathering profile, we calculate Ti-normalized solid-state

concentrations of P, C_w , which may differ from the measured value, C , due to changes in density and concentrations of other elements during weathering:

$$C_w = C \left(\frac{C_{Ti,p}}{C_{Ti,w}} \right) \quad (2)$$

where $C_{Ti,p}$ (mol kg⁻¹) is the solid-state concentration of Ti in the protolith and $C_{Ti,w}$ is the solid-state concentration of Ti in the weathered material (Ti measured by ICP-AES). Open-system mass transport (e.g., Brimhall and Dietrich, 1987; Anderson et al., 2002) is quantifiable in the Rio Icacos weathering profiles because the protolith is homogenous, of uniform age, and contains a relatively inert component that persists in the weathered material (Chadwick et al., 1990). White et al. (1998) and Buss et al. (2008) identified Ti as relatively immobile in the Rio Icacos saprolite relative to the bedrock based on calculations of very low volumetric strain with respect to Ti. Volumetric strain (here the change in volume due to expansion or collapse), should be near-zero in isovolumetrically weathered material such as saprolite.

The average of five analyses from a single corestone adjacent to the rindlets sampled was used as the protolith composition for C_w calculations of rindlet samples (Buss et al., 2008). The deepest sample (7.24 m) obtained from the augered cores on the Guaba Ridge was used as the protolith for calculations performed on the core samples. Ti concentrations in the bedrock vary slightly throughout the watershed (Turner et al., 2003) and thus it is appropriate to use the least weathered sample obtainable under a given weathering profile as the parent material. Normalized P concentrations (C_w) increase with increasing depth into the regolith, with a dramatic increase at the bottom of the saprolite at 5 m (Figure 1). An increase in P at approximately 2.5 m depth corresponds to a visible, biological, and chemical heterogeneity in the saprolite interpreted to be

a rindlet that has not fully saprolitized: a ‘ghost rindlet’ (Buss et al., 2005). Below 5 m depth (where the auger penetrated the rindlet zone), normalized P concentrations range from 0.01 – 0.02 mol kg⁻¹, which is consistent with P concentrations in the rindlet zone sampled in the roadcut (Figure 2). The gradients in C_w over the augered rindlet zone and the roadcut rindlet zone are equivalent when the rindlet zone thicknesses are scaled so that the two interfaces that delineate the rindlet zone (the saprolite-rindlet and rindlet-bedrock interfaces) coincide for both sample sets (Figure 2).

Dissolution of apatite within the rindlet zone is apparent in thin section as well as in the P elemental concentration profiles of the bulk materials. Loss of apatite occurs with increasing distance from the corestone-rindlet interface (Table 1). Apatite grains in the bedrock exist along grain boundaries of other minerals or as inclusions, mostly within hornblende, but also occasionally within plagioclase or primary Fe oxides. Apatite inclusions are seen to dissolve out of relatively intact crystals such as hornblende or plagioclase (Figure 3). Phosphorus-containing secondary minerals were not detected.

Fe isotopic compositions in the HCl extracts, $\delta^{56}\text{Fe}$, of the soil and saprolite range from about 0 to -0.1 ± 0.1 ‰ throughout most of the profile, consistent with values for igneous rocks (Beard et al., 2003). Only at the surface (above ~ 1.2 m depth) and in the deepest saprolite (at 4.9 m depth) did we observe distinct variation in isotopic compositions: values were observed to be as light as -0.64 and -0.40‰ in those zones respectively (Table 2, Figure 4). These zones of lighter Fe isotope ratios correspond to zones of higher concentrations of extractable Fe, cell numbers, DNA yields, and Fe(II)-oxidizing bacteria (Buss et al, 2005; Figure 4).

4. Discussion

4.1. Phosphorus Flux and Apatite Weathering

Changes in P concentration with depth in regolith generally represent depletion profiles, biogenic profiles, or depletion-addition profiles, (or a combination) (Brantley et al., 2007; Brantley et al., 2008). A depletion profile occurs when P-bearing minerals dissolve and P is leached from the system and a biogenic profile is created when P that is solubilized at depth is transported upward and stored in the upper soil by biological processes. A depletion-addition profile is characterized by mobilization from the upper soil layer and enrichment in lower layers. A fourth type of profile is also possible: an addition profile, if atmospheric deposition or addition of P-rich fertilizers introduces P into the system. The slope of a concentration profile provides information about the rates and mechanisms of the depletion, addition, or biological processes that control the transfer of mass (e.g., White, 2002; Brantley et al., 2008; Buss et al., 2008). In the Rio Icacos watershed, the thicknesses of the weathering profiles are maintained by a positive feedback between weathering and erosion, controlled by the infiltration of aqueous O₂ to the weathering interface (Fletcher et al., 2006). Although the regolith and rindlet zones of the roadcut and augered site differ in thickness, that the concentration profiles of P in these two locations are consistent (that the same amount of P is lost from interface-to-interface, Figure 2) points to the same mechanism of P mass transfer in both profiles.

Phosphorus in soils exists in a variety of forms, many of which are considered to be occluded or only sparingly soluble and thus not readily available to biota. However, these recalcitrant and occluded pools of P (sorbed to Al and Fe oxides, trapped within soil aggregates, or contained within clays or phosphate minerals) are slowly liberated and thus can be considered

bioavailable when integrating over longer (e.g., decadal) timescales (Cumming and Weinstein, 1990; Richter et al., 2006). In addition, P sequestered in Fe(III)-(hydr)oxides can be liberated on much shorter timescales by reductive dissolution in environments subjected to redox fluctuations. In surface soils in the nearby Bisley Research Watersheds, also in the Luquillo Forest, periodic O₂ depletion (in approximately 2 week intervals) related to high biological activity, warm temperatures, and abundant rainfall, causes redox fluctuations that lead to repeated cycles of reductive dissolution and oxidative re-precipitation of Fe(III)-(hydr)oxides (Liptzin and Silver, 2009). The P released during the reductive half-cycle is immediately taken up by the biota, bypassing the “labile” P pool (NaHCO₃-extractable) (Liptzin and Silver, 2009). Although the deep saprolite is not subjected to the environmental fluctuations of the surface soils, microscale redox-cycling at depth remains likely. Indeed, visible Mn-oxide gradients in the saprolite that are coincident with zones of high cell densities indicate local zones of redox-cycling associated with microbial growth (Buss et al., 2005).

In deep saprolite profiles such as on the Guaba Ridge, where atmospheric and recycled (e.g., leaf litter) P are not available, apatite dissolution during bedrock weathering provides a source of P for the deep microbial communities. These communities are directly dependent on the flux of P from the weathering of apatite crystals in the rindlet zone. Laboratory experiments have demonstrated the ability microorganisms to enhance dissolution of apatite to obtain P (e.g., Welch et al., 2002; Schaperdoth et al., 2007) indicating that apatite-bound P can be considered a bioavailable form. In the Rio Icacos watershed, apatite weathers in the rindlet zone, often within rock layers where porosity is insufficient to host microorganisms (Figure 3), thus microbial dissolution of apatite is not considered a dominant process relative to abiotic dissolution. Much of the P that is released by apatite dissolution is lost, as evidenced by the decline in P

concentrations with decreasing depth (Figures 1 and 2). Remaining P is either taken up by microorganisms or mineralized, most likely sorbed to Fe(III)-(hydr)oxides. Phosphorus-containing secondary minerals were not detected in the rindlet zone.

When the concentration of a mobile element (e.g., P) in a soil reflects dissolution without concomitant secondary mineral precipitation, it demonstrates a decrease in the solid-state concentration C_0 at depth z_I to C_w at a shallower depth z_0 (White, 2002) and can be characterized as a depletion profile (Brantley et al., 2008). The slope of such a concentration–depth plot can be related to the weathering velocity, ω , or rate of advance of weathering into unaltered (parent) material, and the weathering rate, R , of the mineral supplying the mobile element. Here C_0 corresponds to the solid-state concentration of the element within the parent rock and C_w is the solid-state concentration of the element in the weathered material at depth z . While such depletion profiles often show curvature (Brantley et al., 2008), they can often be fitted satisfactorily using linear regression to calculate a slope. The reaction rate R ($\text{mol m}^{-2} \text{s}^{-1}$) for a given mineral in a weathering profile is calculated from the elemental distribution in the profile using the following expression from White (2002):

$$R = 10^{-3} \frac{1}{\phi \nu s} \frac{\omega}{b_s} \quad (3)$$

where ϕ is the mass fraction of the mineral in the weathering material (g g^{-1}), ν is the stoichiometric coefficient of the element in the mineral (mol mol^{-1}), s is the specific surface area of the mineral ($\text{m}^2 \text{g}^{-1}$), b_s is the weathering gradient (m kg mol^{-1}), and ω is the weathering velocity (m s^{-1}), which is also known as the weathering advance rate or the regolith propagation rate. The weathering velocity is calculated as the thickness of the profile, Δz , divided by the time required to form the profile, t :

$$\omega = \frac{\Delta z}{t} \quad (4)$$

The methodology of White (2002) permits the calculation of solid-state reaction rates R for minerals in a weathering profile from the elemental distribution in the profile. Here we use Equation 3 (White, 2002) to quantify a weathering rate for apatite across the rindlet zone in the Rio Icacos watershed as a function of the average weathering velocity, $\omega = 43 \text{ m Ma}^{-1}$ (Brown et al., 1995). We consider the gradient in P across the rindlet zone ($b_s = 36.3 \text{ m kg mol}^{-1}$, Figure 2) and mineralogical parameters $\varphi = 0.008 \text{ g g}^{-1}$ for apatite (determined by digital point counting and converted to mass units), and $v = 3 \text{ mol P mol}^{-1}$. We calculate a geometric surface area, $s = 0.023 \text{ m}^2 \text{ g}^{-1}$, by using an average grain size of $80 \text{ }\mu\text{m}$ and a cubic volume (Brantley et al., 1999). The resulting rate of apatite weathering is $6.8 \times 10^{-14} \text{ mol m}^{-2} \text{ s}^{-1}$, or $\log R = -13.2$. This rate is several orders of magnitude slower than apatite dissolution rates measured in the laboratory (Figure 5, Valsami-Jones et al., 1998; Welch et al., 2002; Guidry and Mackenzie, 2003, Schaperdoth et al., Unpub.), which is expected when comparing field rates to laboratory rates (White and Brantley, 2003). The only other field-based rate of apatite weathering that we are aware of is normalized to the volume of soil rather than the specific surface area of the mineral and thus cannot be directly compared (Nezat et al., 2004).

The apatite weathering profile and weathering velocity are directly related to the rate at which P is released from apatite. The flux of an element j , in terms of unit volume: F_j ($\text{mol } j \text{ m}^{-3} \text{ s}^{-1}$), can describe the rate at which apatite-derived P is supplied to the deep saprolite microbial community:

$$F_j = 10^{-3} \frac{\omega \rho}{\varphi b_s} \quad (5)$$

where j is phosphorus and ρ is the density of the weathering material, which ranges from 1.83×10^6 to $2.7 \times 10^6 \text{ g m}^{-3}$ over the rindlet zone (Buss et al., 2008). Using this range of density values yields: $F_P = 8.6 \times 10^{-9}$ to $1.3 \times 10^{-8} \text{ mol P m}^{-3} \text{ s}^{-1}$.

Buss et al. (2005) estimated maximum rates of microbial growth in the deepest saprolite at this site based on the flux of Fe(II) from the weathering bedrock, assuming Fe(II)-oxidizing bacteria as the dominant primary producers. These organisms are chemolithoautotrophs, that is, they use inorganic substrates for energy (e.g., Fe) and fix CO_2 to create organic carbon (autotrophy). The calculated growth rates, β , were $6.9 \times 10^{-8} \text{ mol biomass C m}^{-3} \text{ s}^{-1}$ for Fe(II)-oxidizing bacteria and $3.5 \times 10^{-8} \text{ mol C m}^{-3} \text{ s}^{-1}$ for heterotrophic bacteria. These latter bacteria are dependent on the organic carbon produced by the Fe(II)-oxidizers because the dissolved organic carbon (DOC) in porewaters at these depths is extremely low: $125 \text{ }\mu\text{M}$ (Murphy, 1995) and therefore the flux of DOC from the surface is insignificant relative to the flux of organic carbon produced by autotrophy: $\sim 10^{-7}$ vs $10^{-4} \text{ mol C m}^{-3} \text{ h}^{-1}$, respectively (Buss et al., 2005).

To estimate the flux of P required, F_{req} ($\text{mol P m}^{-3} \text{ s}^{-1}$), to maintain a given growth rate, β ($\text{mol C m}^{-3} \text{ s}^{-1}$), we divide the growth rate by the molar C:P ratio of the cells, X (mol mol^{-1}):

$$F_{req} = \frac{\beta}{X} \quad (6).$$

The molar C:P ratio measured mixed cultures of P-limited lakewater heterotrophic bacteria has been reported to lie in the range 80:1 to 100:1 (Jurgens and Gude, 1990). Other values in the literature range from 57:1 to 100:1 for several individual species of P-limited, non-marine, environmental bacteria (Jurgens and Gude, 1990). Taking a C:P ratio of 100 as a rough estimate, we can use Equation 6 to estimate the flux of P required to support microbial growth at the rates calculated by Buss et al. (2005): $F_{req} = 6.9 \times 10^{-10}$ and $3.5 \times 10^{-10} \text{ mol P m}^{-3} \text{ s}^{-1}$ for the

Fe(II)-oxidizing and heterotrophic communities, respectively. These fluxes are less than the flux of P provided by apatite weathering, F_P (Equation 5). Phosphorus requirements vary among specific organisms and therefore we may be underestimating F_{req} , but because actual growth rates are expected to be lower than the maximum rates estimated by Buss et al. (2005), these results suggest that P is not limiting at the saprolite-bedrock interface.

4.2. Fe Isotopic Composition of the Regolith Profile

The HCl extraction is expected to liberate aqueous Fe; Fe contained within microorganisms and biofilms; and weakly sorbed, amorphous, or poorly crystalline solid Fe (Emerson, 2002; Buss et al., 2005; Wiederhold et al., 2007b). Thus we consider two general reservoirs of Fe in the weathering profile: the extractable reservoir and the non-extractable reservoir, which contains well-crystallized phases including Fe-silicates and goethite. Significant sequestration of Fe in recalcitrant organic material is unlikely because concentrations of organic C are extremely low below 1 m depth (Buss et al., 2005).

An enrichment of light Fe isotopes, such as we measured at the top and bottom of the Rio Icacos weathering profile, is created by preferential input of ^{54}Fe to the extractable reservoir, preferential removal of ^{56}Fe , or both. At the bottom of the saprolite, input to the reservoir of HCl-extractable Fe is dominated by complete and rapid dissolution of hornblende, which occurs only within a 7 cm zone at the rindlet-saprolite interface (Buss et al., 2008). Inorganic dissolution of hornblende does not produce a fractionation in Fe isotopes in laboratory experiments; however, dissolution of hornblende in the presence of organic acids, siderophores, or bacteria preferentially releases ^{54}Fe via a kinetic isotope effect, producing a solution that is as much as

0.8‰ lighter than the mineral (Brantley et al., 2001a; Brantley et al., 2004). Organic dissolution of hornblende is thus consistent with the enrichment in light Fe isotopes and the higher cell densities at the bedrock-saprolite interface. Although the Rio Icacos weathering profile is characterized by oxidation processes (White et al., 1998; Buss et al., 2005; Fletcher et al., 2006; Buss et al., 2008) and neither Fe(III)-reducing bacteria nor redoximorphic features (e.g., mottling) have been detected in the saprolite (Buss et al., 2005), reductive dissolution cannot be entirely ruled out. Microaerophilic, circumneutral Fe(II)-oxidizing bacteria were found in the deep saprolite (Buss et al., 2005), which implies that oxygen may be depleted at depth, possibly within microenvironments. Microscale redox cycling of Fe by bacteria has been demonstrated in circumneutral sediments (Sobolev and Roden, 2002; Roden et al., 2004). Regardless, the rapid dissolution of hornblende clearly dominates Fe input to the extractable reservoir at the bedrock-saprolite interface.

Isotopically light extractable Fe also correlates with a larger reservoir of HCl-extractable Fe and high cell densities in the top meter of the weathering profile, which comprises the soil and rooting zone. Organic dissolution of biotite (and possibly Fe-bearing dust), decomposition of isotopically light vegetation, and microbial or plant-mediated reductive dissolution of Fe(III)-(hydr)oxides may all contribute to inputs of isotopically light Fe to the HCl-extractable reservoir in the surface soil.

The higher concentration of Fe in the HCl-extracts at the top and bottom of the profile relative to the middle saprolite indicates that input to the reservoir of extractable Fe is faster than output from the reservoir in these zones. Organic dissolution of hornblende by organic ligands or bacteria is substantially faster than inorganic dissolution (Kalinowski et al., 2000; Liermann et al., 2000; Brantley et al., 2001b; Brantley et al., 2004; Buss et al., 2007), producing a “pulse” of

isotopically light Fe at the start of the hornblende weathering process. Calculations of C_w for Fe in the saprolite and rindlets (Buss et al., 2005; 2008) are approximately constant with depth, indicating that Fe is conserved within the system. Therefore, the only mechanism for removing Fe from the extractable reservoir is by sequestration within well-crystallized Fe(III)-(hydr)oxides either by re-crystallization of ferrihydrite to form goethite or hematite or possibly diffusion of sorbed Fe into the crystalline phases. The rate of crystallization is a function of the concentration of the ferrihydrite precursor, but can be affected by redox cycling and sorption processes. Repeated redox cycling of Fe has been shown to increase the crystallinity of Fe(III)-(hydr)oxides and decrease the size of the HCl-extractable Fe reservoir (Thompson et al., 2006). However, the formation of goethite can be retarded by the sorption onto ferrihydrite of organic acids including citrate and meso-tartrate, simple sugars including glucose and maltose, and silicate ions (Cornell, 1987). In contrast, adsorption of Fe(II) onto ferrihydrite substantially accelerates the formation of goethite, likely because it can act as a reductant, encouraging dissolution of ferrihydrite (Yee et al., 2006). A sorbed Fe(II) atom oxidizes and precipitates as goethite, while a ferrihydrite Fe(III) atom is reduced and propagates the process. Because the heavier isotope preferentially partitions into the oxidized product during equilibrium isotope exchange between Fe(II) and Fe(III), this process could contribute to the isotopically light signature of the extractable reservoir (Icopini et al., 2004), allowing it to persist for some time after the dissolving hornblende becomes depleted in ^{54}Fe . Although batch experiments demonstrated rapid isotopic equilibration between poorly crystalline and well-crystallized Fe(III)-hydroxides when exposed to Fe(II) (Pedersen et al., 2005), such equilibration is expected to be significantly slower in the hydrologically unsaturated (vadose zone) saprolite. Unlike surface soils, the deep saprolite is not chemically or hydrologically affected by precipitation events or seasonal cycles and thus the chemical

composition, moisture content, O₂ partial pressure, and reaction rates are stable over relatively long timescales in the deep saprolite (White et al., 1998), at least at the bulk sampling scale (i.e., there may be more variability within microenvironments). However, the potential for Fe isotopic transience in the HCl-extractable reservoir due to aqueous Fe(II)-exchange with Fe(III)-hydroxides or repeated redox cycling of Fe (Thompson et al., 2006) should be noted.

Biological activity at the bedrock-saprolite interface and in the surface soil may slow the output of Fe from the extractable reservoir by organic complexation of Fe in solution or sorption of microbial products such as organic acids or sugars onto ferrihydrite. However, once the input of Fe from hornblende dissolution (which becomes isotopically heavier as dissolution progresses) is exhausted, it is only a matter of time before the extractable Fe crystallizes into a non-extractable form, homogenizing the two isotopic pools into a single reservoir. Without the large flux of Fe from the dissolution of hornblende (Buss et al., 2008), active microbial communities can no longer be supported and this is reflected in the low cell densities, extremely low extractable Fe concentration, and near-zero $\delta^{56}\text{Fe}$ within the middle saprolite. In the saprolite, Fe exists within oxide minerals and biotite, which weathers to kaolinite throughout the saprolite (Murphy et al., 1998). The very low concentrations of HCl-extractable Fe in the middle saprolite may also reflect a history of redox cycles at the rindlet-saprolite interface, which can decrease the size of this pool (Thompson et al., 2006) (Figure 1b). It is also relevant to note that redox cycling of Fe increases the bioavailability of P (Liptzin and Silver, 2009), which is often sorbed to Fe(III)-(hydr)oxides in soil and saprolite.

We can estimate the flux of Fe into the extractable reservoir due to hornblende weathering using Equation 5. The gradient, b_s , in total Fe(II) across the zone of hornblende weathering, which occurs over 7 cm at the bedrock-rindlet interface, is $b_s = 0.088 \text{ m kg mol}^{-1}$

(Buss et al., 2008), $\phi = 0.063 \text{ g g}^{-1}$ (White et al., 1998), and $\rho = 1.83 \times 10^6 \text{ g m}^{-3}$ in the deep saprolite (White et al., 1998). The resulting flux, $F_{Fe} = 4.5 \times 10^{-7} \text{ mol Fe m}^{-3} \text{ s}^{-1}$. This rate of input can be compared to the rate of output via goethite crystallization using Equation 5 and the gradient in extractable Fe in the deep saprolite, $b_s = 19.01 \text{ m kg mol}^{-1}$ (Figure 4b), and $\phi = 0.0018 \text{ g g}^{-1}$ (Buss et al., 2005). The resulting Fe flux out, $-F_{Fe} = 7.3 \times 10^{-8} \text{ mol Fe m}^{-3} \text{ s}^{-1}$, is an order of magnitude slower than the Fe flux in due to hornblende dissolution. This result is consistent with the observation that the extractable reservoir reaches the isotopic value of the middle saprolite before (at $\sim 4.6 \text{ m}$ depth) it reaches the concentration value of the middle saprolite (at $\sim 4.0 \text{ m}$ depth).

Enhanced dissolution of hornblende by organic ligands, suggested by the isotopic and concentration gradients in extractable Fe at the bedrock-saprolite interface, indicates an active role for biota in chemical weathering. In addition to increasing the rate of hornblende dissolution, organic ligands produced by microorganisms may increase the rate of other mineral weathering reactions by lowering the pH of the porefluids. Furthermore, microbial consumption of oxygen during respiration may affect the rate of spheroidal fracturing, which is thought to be controlled by the concentration of oxygen in the porewater (Fletcher et al., 2006). These results raise the possibility of a “bio-limited” weathering profile in which a change in the rate of biogeochemical weathering would lead to a change in the total denudation rate. Weathering systems are typically classified as either weathering-limited or transport-limited (Stallard, 1992; Kump et al., 2000; Riebe et al., 2004; West et al., 2005). In a weathering-limited regime, total denudation is controlled by the chemical weathering rate, which is limited by the kinetics of mineral weathering reactions. In contrast, in a transport-limited regime (also called supply-limited), the total denudation rate is controlled by the rate of erosion. In the transport-limited case,

perturbations in the weathering rate do not affect the total denudation rate because the weathering rate is high relative to the erosion rate; development of a thick regolith is a consequence. Because the Rio Icacos weathering profile is consistent with models of transport-limitation (Fletcher et al., 2006; Lebedeva et al., 2007), it is unlikely that microbial influence on the weathering rate would affect the denudation rate or the thickness of the regolith profile studied here. However, it is interesting to note that a weathering-limited profile could also be bio-limited.

5. Conclusions

Iron in HCl extracts of saprolite samples is slightly fractionated relative to igneous rocks through the majority of a saprolite profile overlying quartz diorite in the Luquillo Mountains, Puerto Rico: $\delta^{56}\text{Fe}$ values generally range from approximately 0 to -0.1 ‰. However, HCl-extractable Fe is significantly enriched in the lighter isotope in samples taken from the soil and from the bottom of the saprolite directly overlying the bedrock, where $\delta^{56}\text{Fe} = -0.64$ and -0.40‰, respectively. These zones of lighter Fe isotopic compositions are consistent with the zones of highest biological activity in the saprolite as measured by cell density. Whereas organisms in the soil are generally heterotrophic and can thrive on relatively abundant organic carbon in the soil and porewaters, little to no organic carbon is available at the bedrock-saprolite interface, other than cellular material. Microorganisms at the bottom of the saprolite therefore rely on Fe and P released from the spheroidally weathering bedrock as an energy and nutrient source, respectively, with Fe as the limiting “nutrient”. The bedrock-saprolite interface is characterized by spheroidal fractures that delineate ~2.5 cm-thick rindlets and that comprise ~50

cm thick rindlet sets. Apatite is present in the corestone at parent abundance, but is not measured in the saprolite: we calculate that apatite weathers across the rindlet sets at a rate of 6.8×10^{-14} mol m⁻² s⁻¹. Using growth yields from the literature and weathering release rates of Fe(II), biomass production rates were calculated previously by Buss et al. (2005). The apatite dissolution rate calculated here produces a flux of P sufficient to support these growth rates. Therefore our results suggest that P is not limiting at the saprolite-bedrock interface. These results are consistent with the conclusion, advanced in our earlier paper, that the ecosystem that survives at ~5 m depth in the saprolite directly above the bedrock in the Rio Icacos weathering profile is dependent upon chemolithoautotrophic primary producers. These chemolithoautotrophs are Fe(II) oxidizers and are sustained through Fe released from the weathering bedrock. Active cycling of Fe by microorganisms is reflected in the isotopically light Fe in the HCl extracts from the top and bottom of the weathering profile, although the cycling and fractionation mechanisms may differ between the surface and deep ecosystems. The preferential mobilization of light Fe isotopes at the bedrock-saprolite interface suggests that microorganisms are likely influencing concentrations of Fe and oxygen in the porewater and thus affecting the rates of weathering of Fe-bearing minerals.

Acknowledgements

We thank Lisa Stillings, Aaron Thompson, Tom Bullen, and two anonymous reviewers for thoughtful comments that improved the manuscript, R.C. Fletcher, J. Troester, and M. Rosario-Torres for field assistance and J. Kittleson, G. Icopini, M. Angelone, G. Hart, and J. Vervoort for laboratory and data assistance. Funding provided by DOE grant no. DE-FG02-05ER15675, the Penn State Biogeochemical Research Initiative for Education (BRIE) supported

547 by NSF-IGERT Grant no. DGE-9972759, and the Water Energy and Biogeochemical Budgets
548 Program of the U.S. Geological Survey. H.L. Buss acknowledges fellowship support of the NSF
549 Graduate Research Fellowship Program and the National Academy of Sciences Research
550 Associateship Program.

551 **References**

- 552 Anderson, S.P., Dietrich, W.E. and Brimhall, G.H., 2002. Weathering profiles, mass-balance
553 analysis, and rates of solute loss: Linkages between weathering and erosion in a small,
554 steep catchment. *Geological Society of America Bulletin*, 114: 1143-1158.
- 555 Arnold, G.L., Anbar, A.D., Barling, J. and Lyons, T.W., 2004. Molybdenum isotope evidence for
556 widespread anoxia in mid-Proterozoic oceans. *Science*, 304(5667): 87-90.
- 557 Bandstra, J.Z., Buss, H.L., Campen, R.K., Liermann, L.J., Moore, J., Hausrath, E.M., Navarre-
558 Sitchler, A.K., Jang, J.-H. and Brantley, S.L., 2008. Appendix: Compilation of Mineral
559 Dissolution Rates. In: S.L. Brantley, J.D. Kubicki and A.F. White (Editors), *Kinetics of*
560 *Water-Rock Interaction*. Springer, New York, pp. 737-823.
- 561 Beard, B.L., Johnson, C.M., Cox, L., Sun, H., Nealson, K.H. and Aguilar, C., 1999. Iron isotope
562 biosignatures. *Science*, 285: 1889-1892.
- 563 Beard, B.L., Johnson, C.M., Skulan, J.L., Nealson, K.H., Cox, L. and Sun, H., 2003. Application
564 of Fe isotopes to tracing the geochemical and biological cycling of Fe. *Chem. Geol.*, 195:
565 87-117.

566 Boccheciamp, R.A., Rivera, W.F., Trigo, J.E., Brunet, J.E., Torres, E.O., McKinze, W.E. and
 567 Rivera, L.H., 1977. Soil Survey of the Humacao Area of Eastern Puerto Rico. USDA Soil
 568 Conservation Service., Washington, D.C.

569 Brantley, S.L., Bandstra, J., Moore, J. and White, A.F., 2008. Modelling chemical depeletion
 570 profiles in regolith. *Geoderma*, 145: 494-504.

571 Brantley, S.L., Bullen, T.D. and Liermann, L., 2001a. Fractionation of Fe isotopes by soil
 572 microbes and organic acids. *Geology*, 29: 535-538.

573 Brantley, S.L., Goldhaber, M.B. and Ragnarsdottir, K.V., 2007. Crossing disciplines and scales
 574 to understand the critical zone. *Elements*, 3: 307-314.

575 Brantley, S.L., Liermann, L. and Bullen, T.D., 2001b. Fractionation of Fe isotopes by soil
 576 microbes and organic acids. *Geology*, 29(6): 535 - 538.

577 Brantley, S.L., Liermann, L.J., Guynn, R.L., Anbar, A., Icopini, G.A. and Barling, J., 2004. Iron
 578 isotope fractionation during mineral dissolution with and without bacteria. *Geochim.*
 579 *Cosmochim. Acta*, 68(15): 3189-3204.

580 Brantley, S.L., White, A.F. and Hodson, M., 1999. Specific surface area of primary silicates. In:
 581 B. Jamtveit and P. Meakin (Editors), *Growth and Dissolution in Geosystems*. Kluwer
 582 Academic Publishers, Dordrecht, pp. 291-326.

583 Brimhall, G. and Dietrich, W.E., 1987. Constitutive mass balance relations between chemical
 584 composition, volume, density, porosity, and strain in metasomatic hydrochemical
 585 systems: results on weathering and pedogenesis. *Geochim. Cosmochim. Acta*, 51: 567-
 586 587.

587 Brown, E.T., Stallard, R., Larsen, M.C., Raisbeck, G.M. and Yiou, F., 1995. Denudation rates
588 determined from the accumulation of in situ-produced ^{10}Be in the Luquillo Experimental
589 Forest, Puerto Rico. *Earth and Planetary Science Letters*, 129: 193-202.

590 Bruijnzeel, L.A., 1991. Nutrient input-output budgets of tropical forest ecosystems: A review.
591 *Journal of Tropical Ecology*, 7(1): 1-24.

592 Bullen, T.D., White, A.F., Childs, C.W., Vivit, D.V. and Schulz, M.S., 2001. Demonstration of
593 significant abiotic iron isotope fractionation in nature. *Geology*, 29(8): 699-702.

594 Buss, H.L., Bruns, M.A., Schultz, M.J., Moore, J., Mathur, C.F. and Brantley, S.L., 2005. The
595 coupling of biological iron cycling and mineral weathering during saprolite formation,
596 Luquillo Mountains, Puerto Rico. *Geobiology*, 3(4): 247-260.

597 Buss, H.L., Luttge, A. and Brantley, S.L., 2007. Etch pit formation on iron silicate surfaces
598 during siderophore-promoted dissolution. *Chem. Geol.*, 240: 326-342.

599 Buss, H.L., Sak, P.B., Webb, S.M. and Brantley, S.L., 2008. Weathering of the Rio Blanco
600 quartz diorite, Luquillo Mountains, Puerto Rico: Coupling oxidation, dissolution, and
601 fracturing. *Geochim. Cosmoch. Acta*, 72: 4488-4507.

602 Buss, H.L., Sak, P.B., White, A.F. and Brantley, S.L., 2004. Mineral dissolution at the granite-
603 saprolite interface. In: R.B. Wanty and R.R.I. Seal (Editors), 11th International
604 Symposium on Water-Rock Interaction. Taylor and Francis, Saratoga Springs, NY, pp.
605 819-823.

606 Chadwick, O.A., Brimhall, G.H. and Hendricks, D.M., 1990. From black box to a grey box: a
607 mass balance interpretation of pedogenesis. *Geomorphology*, 3: 369-390.

608 Cornell, R.M., 1987. Comparison and classification of the effects of simple ions and molecules
609 upon the transformation of ferrihydrite into more crystalline products. *Z. Pflanzenernahr.*
610 *Bodenk.*, 150: 304-307.

611 Crosby, H.A., Johnson, C.M., Roden, E.E. and Beard, B.L., 2005. Coupled Fe(II)-Fe(III)
612 electron and atom exchange as a mechanism for Fe isotope fractionation during
613 dissimilatory iron oxide reduction. *Environmental Science & Technology*, 39: 6698-
614 6704.

615 Crosby, H.A., Roden, E.E., Johnson, C.M. and Beard, B.L., 2007. The mechanisms of iron
616 isotope fractionation produced during dissimilatory Fe(III) reduction by *Shewanella*
617 *putrefaciens* and *Geobacter sulfurreducens*. *Geobiology*, 5: 169-189.

618 Cumming, J.R. and Weinstein, L.H., 1990. Utilization of AlPO₄ as a phosphorus source by
619 ectomycorrhizal *Pinus rigida* Mill. seedlings. *New Phytol.*, 116: 99-106.

620 Dixon, J.C., Campbell, S.W., Thorn, C.E. and Darmody, R.G., 2006. Incipient weathering rind
621 development on introduced machine-polished granite discs in an Arctic alpine
622 environment, northern Scandinavia. *Earth Surface Processes and Landforms*, 31: 111-
623 121.

624 Dorn, R., 1995. Digital processing of back-scatter electron imagery: a microscopic approach to
625 quantifying chemical weathering. *Geological Society of America Bulletin*, 107: 725-741.

626 Emerson, D., 2002. Microbial oxidation of Fe(II) and Mn(II) at circumneutral pH. In: D.R.
627 Lovely (Editor), *Environmental Microbe-Metal Interactions*. ASM, Washington D.C.

628 Fantle, M.S. and DePaolo, D.J., 2004. Iron isotopic fractionation during continental weathering.
629 *Earth and Planetary Science Letters*, 228: 547-562.

630 Fletcher, R.C., Buss, H.L. and Brantley, S.L., 2006. A spheroidal weathering model coupling
631 porewater chemistry to soil thicknesses during steady-state denudation. *Earth and*
632 *Planetary Science Letters*, 244: 444-457.

633 Frausto da Silva, J.J.R. and Williams, R.J.P., 191. *The Biological Chemistry of the Elements*.
634 Clarendon Press, Oxford, UK, 561 pp.

635 Frizano, J., Johnson, A.H. and Vann, D.R., 2002. Soil phosphorus fractionation during forest
636 development on landslide scars in the Luquillo Mountains, Puerto Rico. *Biotropica*,
637 34(1): 17-26.

638 Garrett, R.H. and Grisham, C.M., 1999. *Biochemistry*. Harcourt Brace College Publishers,
639 Orlando, 1127 pp.

640 Graham, R.C. and Duce, R.A., 1982. The atmospheric transport of phosphorus to the western
641 North Atlantic. *Atmospheric Environment*, 16(5): 1089-1097.

642 Guelke, M. and Von Blanckenburg, F., 2007. Fractionation of stable iron isotopes in higher
643 plants. *Environmental Science & Technology*, 41: 1896-1901.

644 Guidry, M.W. and Mackenzie, F.T., 2003. Experimental study of igneous and sedimentary
645 apatite dissolution: Control of pH, distance from equilibrium, and temperature on
646 dissolution rates. *Geochim. Cosmochim. Acta*, 67(16): 2949-2963.

647 Heartsill-Scalley, T., Scatena, F.N., Estrada, C., McDowell, W.H. and Lugo, A.E., 2007.
648 Disturbance and long-term patterns of rainfall and throughfall nutrient fluxes in a
649 subtropical wet forest in Puerto Rico. *Journal of Hydrology*, 333: 472-485.

650 Icopini, G.A., Anbar, A.D., Ruebush, S.S., Tien, M. and Brantley, S.L., 2004. Iron isotope
651 fractionation during microbial reduction of iron; the importance of adsorption. *Geology*,
652 32(3): 205-208.

653 Jurgens, K. and Gude, H., 1990. Incorporation and release of phosphorus by planktonic bacteria
654 and phagotrophic flagellates. *Marine Ecology Progress Series*, 59: 271-284.

655 Kalinowski, B.E., Liermann, L.J., Brantley, S.L., Barnes, A. and Pantano, C.G., 2000. X-ray
656 photoelectron evidence for bacteria-enhanced dissolution of hornblende. *Geochim.*
657 *Cosmochim. Acta*, 64(8): 1331-1343.

658 Kump, L.R., Brantley, S.L. and Arthur, M.A., 2000. Chemical weathering, atmospheric CO₂ and
659 climate. *Ann. Rev of Earth and Planetary Sciences*, 28: 611-667.

660 Kurtz, A.C. and Derry, L.A., 2004. Tracing silicate weathering and terrestrial silica cycling with
661 Ge/Si ratios. In: R.B. Wanty and R.R.I. Seal (Editors), *The 11th International Symposium*
662 *on Water-Rock Interaction*. A.A. Balkema, Saratoga Springs, NY, pp. 833-840.

663 Lebedeva, M.I., Fletcher, R.C., Balashov, V.N. and Brantley, S.L., 2007. A reactive diffusion
664 model describing transformation of bedrock to saprolite. *Chem. Geol.*, 244: 624-645.

665 Liermann, L.J., Kalinowski, B.E., Brantley, S.L. and Ferry, J.G., 2000. Role of bacterial
666 siderophores in dissolution of hornblende. *Geochim. Cosmochim. Acta*, 64(4): 587-602.

667 Liptzin, D. and Silver, W.L., 2009. Effects of carbon additions on iron reduction and phosphorus
668 availability in a humid tropical forest soil. *Soil Biol. Biochem.*,
669 doi:10.1016/j.soilbio.2009.05.013.

670 Lugolobi, F., Kurtz, A.C. and Derry, L.A., Subm. Germanium-silicon fractionation in a tropical,
671 granitic weathering environment. *Geochimica Cosmochimica Acta*.

672 Lugolobi, F., Kurtz, A.C. and Salvucci, G.D., 2006. Silica sources and water flowpaths during
673 storm events at Rio Icacos, Puerto Rico, AGU Fall Meeting, San Francisco.

674 Maréchal, C.N., Télouk, P. and Albarède, F., 1999. Precise analysis of copper and zinc isotopic
675 compositions by plasma-source mass spectrometry. *Chem. Geol.*, 156: 251-273.

676 McDowell, W.H. and Asbury, C.E., 1994. Export of carbon, nitrogen, and major ions from three
 677 tropical montane watersheds. *Limnology and Oceanography*, 39(1): 111-125.
 678 Muhs, D.R., Budahn, J.R., Prospero, J.M. and Carey, S.N., 2007. Geochemical evidence for
 679 African dust inputs to soils of western Atlantic islands: Barbados, the Bahamas, and
 680 Florida. *Journal of Geophysical Research*, 112: F02009.
 681 Murphy, S.F., 1995. Weathering of biotite in a tropical forest soil, Luquillo Mountains, Puerto
 682 Rico. M.S. Thesis, The Pennsylvania State University, University Park, PA, 100 pp.
 683 Murphy, S.F., Brantley, S.L., Blum, A.E., White, A.F. and Dong, H., 1998. Chemical weathering
 684 in a tropical watershed, Luquillo Mountains, Puerto Rico; II. Rate and mechanism of
 685 biotite weathering. *Geochim. Cosmochim. Acta*, 62(2): 227-243.
 686 Nezat, C.A., Blum, J.D., Klaue, A., Johnson, C.E. and Siccama, T.G., 2004. Influence of
 687 landscape position and vegetation on long-term weathering rates at the Hubbard Brook
 688 Experimental Forest, New Hampshire, USA. *Geochim. Cosmochim. Acta*, 68(14): 3065-
 689 3078.
 690 Olander, L.P. and Vitousek, P.M., 2005. Short-term controls over inorganic phosphorus during
 691 soil and ecosystem development. *Soil Biol. Biochem.*, 37: 651-659.
 692 Pedersen, H.D., Postma, D., Jakobsen, R. and Larsen, O., 2005. Fast transformation of iron
 693 oxyhydroxides by the catalytic action of aqueous Fe(II). *Geochim. Cosmochim. Acta*, 69:
 694 3967-3977.
 695 Pett-Ridge, J., 2009. Contributions of dust to phosphorus cycling in tropical forests of the
 696 Luquillo Mountains, Puerto Rico. *Biogeochemistry*, 94(1): 63-80.

697 Pett-Ridge, J.C., Derry, L.A. and Kurtz, A.C., 2009. Sr isotopes as a tracer of weathering
 698 processes and dust inputs in a tropical granitoid watershed, Luquillo Mountains, Puerto
 699 Rico. *Geochim. Cosmochim. Acta*, 73: 25-43.

700 Richter, D.D., Allen, H., Li, J., Markewitz, D. and Raikes, J., 2006. Bioavailability of slowly
 701 cycling soil phosphorus: major restructuring of soil P fractions over four decades in an
 702 aggrading forest. *Oecologia*, 150(2): 259-271.

703 Riebe, C.S., Kirchner, J.W. and Finkel, R.C., 2004. Erosional and climatic effects on long-term
 704 chemical weathering rates in granitic landscapes spanning diverse climate regimes. *Earth
 705 and Planetary Science Letters*, 224: 547-562.

706 Roden, E.E., Sobolev, D., Glazer, B. and Luther III, G.W., 2004. Potential for microscale
 707 bacterial Fe redox cycling at the aerobic-anaerobic interface. *Geomicrobiology*, 21(6):
 708 379-391.

709 Sanchez, P.A., 1976. *Properties and Management of Soils in the Tropics*. Wiley, New York, 618
 710 pp.

711 Schaperdoth, I., Liermann, L. and Brantley, S., 2007. The Effect of Polymeric Substances on
 712 Apatite Reactivity in the Presence of a Freshwater Cyanobacterium. *Geomicrobiol. J.*,
 713 24(2): 79-91.

714 Severmann, S., Johnson, C.M., Beard, B.L. and McManus, J., 2006. The effect of early
 715 diagenesis on the Fe isotope compositions of porewaters and authigenic minerals in
 716 continental margin sediments. *Geochim. Cosmochim. Acta*, 70: 2006-2022.

717 Simon, A., Larsen, M.C. and Hupp, C.R., 1990. The role of soil processes in determining
 718 mechanisms of slope failure and hillslope development in a humid-tropical forest, eastern
 719 Puerto Rico. *Geomorphology*, 3: 263-286.

720 Sobolev, D. and Roden, E.E., 2002. Evidence for rapid microscale bacterial redox cycling of iron
 721 in circumneutral environments. *Antonie van Leeuwenhoek*, 81(587-597).

722 Stallard, R.F., 1992. Tectonic processes, continental freeboard, and the rate-controlling step for
 723 continental denudation. In: S.S. Butcher, R.J. Charlson, G.H. Orians and G.V. Wolfe
 724 (Editors), *Global Biogeochemical Cycles*. Academic Press, London, pp. 93-121.

725 Templer, P.H., Silver, W.L., Pett-Ridge, J., DeAngelis, K.M. and Firestone, M.K., 2008. Plant
 726 and microbial controls on nitrogen retention and loss in a humid tropical forest. *Ecology*,
 727 89(11): 3030-3040.

728 Teutsch, N., von Gunten, U., Porcelli, D., Cirpka, O.A. and Halliday, A.N., 2005. Adsorption as
 729 a cause for iron isotope fractionation in reduced groundwater. *Geochimica*
 730 *Cosmochimica Acta*, 17(1): 4175-4185.

731 Thompson, A., Chadwick, O.A., Rancourt, D.G. and Chorover, J., 2006. Iron-oxidized crystallinity
 732 increases during soil redox oscillations. *Geochim. Cosmochim. Acta*, 70: 1710-1727.

733 Thompson, A., Ruiz, J., Chadwick, O.A., Titus, M. and Chorover, J., 2007. Rayleigh
 734 fractionation of iron isotopes during pedogenesis along a climate sequence of Hawaiian
 735 basalt. *Chem. Geol.*, 238: 72-83.

736 Turner, B., 2001. Effects of temperature and climate on chemical weathering in two contrasting
 737 high-rainfall mountainous catchments. Ph.D. Thesis, Pennsylvania State University,
 738 University Park, PA.

739 Turner, B.F., Stallard, R.F. and Brantley, S.L., 2003. Investigation of in situ weathering of quartz
 740 diorite bedrock in the Rio Icacos basin, Luquillo Experimental Forest, Puerto Rico.
 741 *Chem. Geol.*, 202(3-4): 313-341.

742 USDA NCRS, 2002. Soil Survey of Caribbean National Forest and Luquillo Experimental
 743 Forest, Commonwealth of Puerto Rico. USDA, Natural Resources Conservation Service,
 744 Washington D.C., pp. 181.

745 Valsami-Jones, E., Ragnarsdottir, K.V., Putnis, A., Bosbach, D., Kemp, A.J. and Cressey, G.,
 746 1998. The dissolution of apatite in the presence of aqueous metal cations at pH 2-7.
 747 Chem. Geol., 151: 215-233.

748 Vitousek, P.M., 1984. Litterfall, nutrient cycling, and nutrient limitation in tropical forests.
 749 Ecology, 65(1): 285-298.

750 Vitousek, P.M. and Sanford, R.L., Jr., 1986. Nutrient cycling in moist tropical forest. Ann. Rev.
 751 Ecol. Syst., 17: 137-167.

752 Wardle, D.A., Walker, L.R. and Bardgett, R.D., 2004. Ecosystem properties and forest decline in
 753 contrasting long-term chronosequences. Science, 305(5683): 509-513.

754 Welch, S.A., Taunton, A.E. and Banfield, J.F., 2002. Effect of microorganisms and microbial
 755 metabolites on apatite dissolution. Geomicrobiol. J., 19(3): 343-367.

756 West, A.J., Galy, A. and Bickle, M., 2005. Tectonic and climatic controls on silicate weathering.
 757 Earth and Planetary Science Letters, 235: 211-228.

758 White, A.F., 2002. Determining mineral weathering rates based on solid and sout weathering
 759 gradients and velocities: application to biotite weathering in saprolites. Chem. Geol., 190:
 760 69-89.

761 White, A.F., Blum, A.E., Schulz, M.S., Vivit, D.V., Stonestrom, D.A., Larsen, M., Murphy, S.F.
 762 and Eberl, D., 1998. Chemical weathering in a tropical watershed, Luquillo Mountains,
 763 Puerto Rico: I. Long-term versus short-term weathering fluxes. Geochim. Cosmochim.
 764 Acta, 62(2): 209-226.

765 White, A.F. and Brantley, S.L., 2003. The effect of time on the experimental and natural
 766 weathering rates of silicate minerals. *Chem. Geol.*, 202: 479-506.

767 Wiederhold, J.G., Kraemer, S.M., Teutsch, N., Borer, P.M., Halliday, A.N. and Kretzschmar, R.,
 768 2006. Iron isotope fractionation during proton-promoted, ligand-controlled, and reductive
 769 dissolution of goethite. *Environmental Science & Technology*, 40: 3787-3793.

770 Wiederhold, J.G., Teutsch, N., Kraemer, S.M., Halliday, A.N. and Kretzschmar, R., 2007a. Iron
 771 isotope fractionation during pedogenesis in redoximorphic soils. *Soil Sci. Soc. Am. J.*,
 772 71(6): 1840-1850.

773 Wiederhold, J.G., Teutsch, N., Kraemer, S.M., Halliday, A.N. and Kretzschmar, R., 2007b. Iron
 774 isotope fractionation in oxic soils by mineral weathering and podzolization. *Geochimica
 775 et Cosmochimica Acta*, 71: 5821-5833.

776 Yee, N., Shaw, S., Benning, L.G. and Nguyen, T.H., 2006. The rate of ferrihydrite
 777 transformation to goethite via the Fe(II) pathway. *Amer. Mineral.*, 91: 92-96.

778 Zarin, D.J. and Johnson, A.H., 1995. Base saturation, nutrient cation, and organic matter
 779 increases during early pedogenesis on landslide scars in the Luquillo Experimental
 780 Forest, Puerto Rico. *Geoderma*, 65: 317-330.

781 Ziegler, K., Chadwick, O.A., White, A. and Brzezinski, M.A., 2005. $\delta^{30}\text{Si}$ systematics in a
 782 granitic saprolite, Puerto Rico. *Geology*, 33(10): 817-820.

783
 784
 785

786
787

Table 1.
Apatite content of rindlets and corestone

Distance¹ (cm)	Apatite² (vol %)
49.0	nd ³
48.5	nd
48.0	nd
47.5	0.01
33.0	0.14
30.0	nd
28.0	nd
21.0	0.20
16.0	0.35
8.5	0.38
5.0	0.11
4.0	0.36
0	0.62

788
789
790
791
792
793
794
795

¹ Distance from corestone

² Determined by point counting
backscattered electron images of thin
sections using image analysis software.

³ nd = not detected

Table 2.
Fe isotope ratios of HCl extracts

Depth (m)	$\delta^{56}\text{Fe}$ (‰)
0.15	-0.64
0.3	-0.66
0.6	-0.55
0.9	-0.13
1.2	-0.26
2.1	-0.12
2.4	-0.08
2.4	-0.11
3.0	0.00
3.7	-0.11
4.3	-0.10
4.6	0.01
4.9	-0.40

796
797

Figure Captions

Figure 1.

Solid-state, Ti-normalized concentrations of P, C_w , with depth in the weathering profile that was sampled by augering on a ridgetop. The dashed line denotes the bottom of the saprolite. Samples below the saprolite were inferred to be rindlet-zone samples on the basis of texture, color, and chemistry. Error is the standard deviation of the measured concentrations of P for five individual samples of a single corestone. The higher concentrations at ~2.5 m correspond to a visible, chemical, textural, and biological heterogeneity identified by Buss et al. (2005) as a “ghost rindlet”.

Figure 2.

Solid state, Ti-normalized concentrations of P, C_w , in the rindlet and corestone samples collected from a roadcut (open triangles). The left-hand y-axis shows the distance below the saprolite-rindlet interface. A linear fit through the roadcut data is shown, which was used as the gradient in P, b_s , for calculation of the apatite weathering rate and P flux (Equations 3 and 5). The equation for the regression line is $Y = 36.3X - 0.13$, $R^2 = 0.76$. Error bars are the standard deviation of the measured concentrations of P for five individual samples of a single corestone. Solid square symbols are C_w for P in the rindlet zone portion of the augered core (Figure 1) corresponding to the right-hand y-axis, which shows the depth in the augered core.

Figure 3.

Backscattered Electron (BSE) image of a thin section of a rindlet from about 4 cm above the corestone sampled in a road cut. The remnants of a dissolving apatite crystal are visible within an

apatite-shaped hole. While apatite weathers across the entire 49 cm rindlet zone, hornblende only weathers within 7 cm at the saprolite-rindlet interface (Buss et al., 2008). The sample in this image is below the zone of hornblende weathering and thus the hornblende crystal appears pristine.

Figure 4.

a) Isotopic composition of Fe extracted with 0.5 N HCl from the soil and saprolite samples. Error is $2\sigma = 0.1\text{‰}$ calculated for multiple measurements of a goethite standard. **b)** Fe concentrations in 0.5 N HCl extracts (Buss et al., 2005) with a linear regression through concentrations at depths below 4 m representing the gradient in extractable Fe over this zone. Error bars are smaller than most of the symbols. **c)** Cell numbers estimated by direct counting (Buss et al., 2005), and **d)** cell numbers estimated from DNA yields (Buss et al., 2005). Notice the log scale for the cell numbers in c-d. The anomalously high cell density at ~2.2 m in c corresponds to a relatively unweathered zone in the saprolite, thought to be a “ghost rindlet” (Buss et al., 2005).

Figure 5.

Rates of apatite dissolution determined from laboratory experiments (Valsami-Jones et al., 1998; Welch et al., 2002; Guidry and Mackenzie, 2003, Schaperdoth et al. Unpub.) show a linear dependence on pH (Bandstra et al., 2008). The field rate determined in the present study is several orders of magnitude slower than the laboratory rates. Laboratory rates are normalized to BET surface area of the mineral. The field rate is normalized to a geometric mineral surface area. The field pH of 5.4 was measured by White et al. (1998) in porewater on the Guaba Ridge.