Tubiello, F. N., Salvatore, M., F. Ferrara, A., House, J. I., Federici, S., Rossi, S., Biancalani, R., D. Condor Golec, R., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo, P., Schmidhuber, J., Sanz Sanchez, M. J., Srivastava, N., & Smith, P. (2015). The Contribution of Agriculture, Forestry and other Land Use activities to Global Warming, 1990-2012. *Global Change Biology*, 21(7), 2655-2660. <https://doi.org/10.1111/gcb.12865>

Peer reviewed version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1111/gcb.12865](https://doi.org/10.1111/gcb.12865)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Title Page

i. Title:

The Contribution of Agriculture, Forestry and other Land Use activities to Global Warming,
1990-2012: Not as high as in the past

ii. Running Title:

The Contribution of AFOLU to Global Warming

iii. Authors:

Francesco N. Tubiello^{1*}, Mirella Salvatore¹, Alessandro F. Ferrara¹, Simone Rossi^{1,2}, Riccardo Biancalani¹, Rocio D. Condor Golec¹, Sandro Federici¹, Heather Jacobs¹, Alessandro Flammini¹, Paolo Prosperi¹, Paola Cardenas¹, Josef Schmidhuber³, Maria J. Sanz Sanchez⁴, Pete Smith⁵, Jo House⁶, Nalin Srivastava⁷

iv. Institute or Laboratory of Origin:

¹Climate, Energy and Tenure Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome 00153 Italy. Francesco.tubiello@fao.org ph: +39 06 57052169

²European Commission Joint Research Center, Ispra (VA) 28100 Italy. simone.rossi@jrc.europa.eu; ph: +39-0332-78-3080

³Statistics Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome 00153 Italy; josef.schmidhuber@fao.org; ph: +39 06 570 56264

⁴Forest Management Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome 00153 Italy; maria.sanzsanchez@fao.org; ph: +39 06 570 54587

⁵Institute of Biological and Environmental Sciences, School of Biological Sciences, University of Aberdeen, 23 St Machar Drive, Room G45, Aberdeen AB24 3UU, Scotland, UK. pete.smith@abdn.ac.uk, ph: +44 1224 272703

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

⁶School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8 1SS, UK. jo.house@bristol.ac.uk; ph: +44 117 954 5978

⁷IPCC Task Force on National GHG Inventories, IGES, 2108-11 Kamiyamaguchi Hayama, Kanagawa, Japan. shrivastava@iges.or.jp, ph: +81 46 855 3754

v. Corresponding Author:

*Francesco N. Tubiello, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome 00153 Italy.

email: francesco.tubiello@fao.org ph: +39-06-57052169

vi. Keywords: Agriculture, AFOLU, GHG, Emissions, Climate Change, Mitigation

vii. Type of Paper: Original Research

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Abstract

We refine the information available through the IPCC AR5 with regards to recent trends in global GHG emissions from agriculture, forestry and other land uses (AFOLU), including global emissions updates to 2012. By using all three available AFOLU datasets employed for analysis in the IPCC AR5, rather than just one as done in the IPCC AR5 WGIII Summary for Policy Makers, our analyses point to a down-revision of global AFOLU shares of total anthropogenic emissions, while providing important additional information on sub-sectoral trends. Our findings confirm that the share of AFOLU emissions to the anthropogenic total declined over time. They indicate a decadal average of $28.7 \pm 1.5\%$ in the 1990s, $23.6 \pm 2.1\%$ in the 2000s, and an annual value of $21.2 \pm 1.5\%$ in 2010. The IPCC AR5 had indicated a 24% share in 2010. In contrast to previous decades, when emissions from *land use* (land use, land use change and forestry, including deforestation) were significantly larger than those from *agriculture* (crop and livestock production), in 2010 agriculture was the larger component, contributing $11.2 \pm 0.4\%$ of total GHG emissions, compared to $10.0 \pm 1.2\%$ of the land use sector. Deforestation was responsible for only 8% of total anthropogenic emissions in 2010, compared to 17% in the 1990s. Since 2010, the last year assessed by the IPCC AR5, new FAO estimates indicate that land use emissions have remained stable, at about 4.8 in 2012. Emissions minus removals have also remained stable, at $3.2 \text{ Gt CO}_2\text{eq yr}^{-1}$ in 2012. By contrast, agriculture emissions have continued to grow, at roughly 1% annually, and remained larger than the land use sector, reaching $5.4 \text{ Gt CO}_2\text{eq yr}^{-1}$ in 2012. These results are useful to further inform the current climate policy debate on land use, suggesting that as many efforts and resources should be directed to agriculture mitigation in the coming years as they have been devoted to REDD+ in the past decade, considering the increasingly larger emission profile of crop and livestock activities compared to the forestry and land use change sector.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

1. Introduction

Agriculture, Forestry and Other Land Use (AFOLU) activities generate greenhouse gas emissions by sources as well as removals by sinks, through the oxidation and fixation of organic matter *via* photosynthesis, followed by loss from decomposition and fire, including complex microbial processes associated with human management and disturbance of ecosystems. They comprise mainly non-CO₂ emissions by agriculture (CH₄ and N₂O), and CO₂ emissions by sources and removals by sinks by land use and land use change.

Assessing historical and current trends of AFOLU emissions, including *agriculture* (crop and livestock production) and *land use* (land use land use change and forestry) dynamics at global and regional level, is important both for science and climate policy. On the science side, better data help to characterize anthropogenic forcing of the atmosphere, while providing useful constraints to carbon cycle assessments. On the climate policy side, better data provide support for concerted global action, allowing for transparent exchange of information between parties, based on more accurate greenhouse gas inventories for reporting under the United Nations Convention on Climate Change (UNFCCC). More reliable data, for instance, provided the operational basis for implementation of the Kyoto Protocol First Commitment Period, the adoption of the Bali Action Plan, and the Cancun Agreements. Better AFOLU data specifically contributed to raising awareness on the need to reducing emissions from deforestation and forest degradation (REDD+) (FAO, 2011). Indeed, early quantifications of the share of deforestation to total anthropogenic emissions helped to make the case for forests and the land use sector being as an effective, short-term climate change mitigation option, compared to more complex and costly long-term measures in other sectors, including agriculture (Stern, 2007).

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Despite the high relevance of this information, quantification of AFOLU emissions sources and sinks remains uncertain, even at the global level. The recent IPCC WGIII Summary for Policy Makers (SPM) states that AFOLU contributed 24% of total anthropogenic emissions in 2010 (IPCC, 2014). This estimate however was based on only one of three global emission data sources available in the IPCC AR5 WGIII AFOLU chapter. These included the EDGAR database, which was the single data source used in the IPCC AR5 SPM, and additionally the FAOSTAT database and the Houghton dataset (Smith et al., 2014).

We show in this paper that different absolute values and relative shares of emissions from agriculture and land use are obtained by integrating valuable information from the FAO and Houghton's available estimates, in addition to EDGAR.

2. Materials and Methods

Three global AFOLU datasets used in the IPCC WGIII chapter 11 (Smith et al., 2014) are considered herein: FAOSTAT (2014), EDGAR (2013) and the bookkeeping method of Houghton et al. (2012). These data sources robustly represent the range of AFOLU data used in the IPCC AR5 WGIII for global and regional emission analyses. Specifically, EDGAR data were used as a reference for estimating total anthropogenic emissions and perform cross-sectoral synthesis, while Houghton's and FAO data were used for more in-depth analyses of the AFOLU sector. Three data combinations were considered herein as follows: i) EDGAR only (E); ii) FAOSTAT only (F); and iii) FAOSTAT for agriculture and peat degradation and fire, plus Houghton for forest and land use change (H). The three combinations correspond to data analyses reported in the IPCC AR5 WGIII SPM (E) and in the IPCC AR5 WGIII Ch. 11, Section 11.2.2 (H) and 11.2.3 (F).

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

The FAOSTAT Emissions database covers all agriculture, forestry and other land use activities and their associated emissions of CO₂, CH₄ and N₂O, following IPCC 2006 Guidelines at Tier 1 and land approach 1 (Tubiello et al., 2013; FAO, 2014). Emissions are estimated for nearly 200 countries, for the reference period 1961-2012 (agriculture) and 1990-2012 (FOLU), based on activity data collected by FAO. Emissions and removals are calculated both within and across land categories, including effects of crop and livestock management, land use and land conversion. FAOSTAT includes estimates of emissions from biomass fires, peatland drainage and fires, based on geo-spatial information, as well as forest carbon stock changes based on national-level FAO Forest Resources Assessment data (FAO, 2010). FOLU carbon balances in FAOSTAT should only be intended as first-order estimates of underlying complex dynamics arising from afforestation/reforestation, degradation, re-growth and harvest activities. The FAOSTAT emission estimates are at the basis of regular FAO annual emission updates for AFOLU (FAO, 2014).

The EDGAR database covers all IPCC sectors (Energy, Transport, Industry, Buildings, and AFOLU), applying IPCC guidelines for emission estimation (EDGAR, 2013). Several methodological differences with FAOSTAT and Houghton exist, which result in differences in emission estimates. The latter are quantified in the results section below. Methodological differences are as follows. First, for agriculture, EDGAR follows the Revised 1996 IPCC Guidelines (IPCC, 1997) rather than the 2006 IPCC Guidelines used by FAOSTAT, with repercussions that will be discussed in the results section. Second, and more importantly, for FOLU EDGAR does not follow either IPCC Guidelines or standard carbon cycle approaches (Le Quere et al., 2013), in contrast to the FAOSTAT and Houghton data. Specifically, EDGAR emissions estimates for forest and forest conversion are termed collectively *Forest Fires and Decay*. While the associated sub-category *Forest Fires* contains data taken directly from the *Global Fire Emissions Database* (GFED version 2; Van der Werf

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

et al., 2006), lack of transparent information in relevant EDGAR literature and associated metadata makes the interpretation of the *Forest Decay* emission estimates difficult. Apparently the EDGAR *Forest Decay* category covers emissions from the decay of dead biomass left over after fire damage. These are then used as proxy to estimate overall deforestation emissions, by applying unpublished scaling factors directly to the relevant GFED forest fire data. Such an approximation is incorrect however, because not all carbon losses from deforestation are associated to the use of fire.

Another area of inconsistency with both the IPCC guidelines and FAOSTAT is the fact that EDGAR includes in AFOLU the emissions from energy use in agriculture, while they should be reported in the energy sector.

The Houghton dataset is based on a bookkeeping approach for estimating CO₂ emissions due to land use and land use change, such as deforestation and afforestation, forest degradation, wood harvest and shifting cultivation (Houghton et al., 2012). The method uses parameter values for biomass stock, biomass growth and decay rates, taken from published national inventories and the FAO Forest Resource Assessment (FAO, 2010). In the IPCC AR5, the Houghton dataset provides the basis for the central estimates of FOLU emissions for the carbon budget (Cias et al., 2013; Smith et al., 2014). Furthermore, this dataset is at the core of the Global Carbon Project's annual carbon emissions updates for land use change (Le Quere et al., 2013; Friedlingstein et al., 2011).

Units

AFOLU activities emit all three major GHGs, as discussed above. The analyses that follow use CO₂ equivalent (CO₂eq) as a common metric for comparison across emission

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

categories, obtained by using the 100-year global warming potentials specified by the current UNFCCC reporting procedures for national GHG inventories¹ (IPCC, 1997 and 2006).

Uncertainty

While CO₂ emissions related to fossil fuel carry relatively low uncertainties, around $\pm 10\%$, uncertainties in AFOLU are much larger. Uncertainty for global emission estimates from crop and livestock carry $\pm 30\%$ uncertainty ranges (e.g., Tubiello et al., 2013). Uncertainties in FOLU are even larger, with a $\pm 50\%$ uncertainty range (le Qure et al., 2013; Pongratz et al., 2014). All data presented in this paper are characterized by the above uncertainties. The ranges indicated in the results section below, however, refer to the variability of mean estimates across the three datasets used for analysis, rather than to the above uncertainties.

¹ GWP-CO₂ = 1; GWP-CH₄ = 21, GWP-N₂O = 310

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

3. Results and Discussion

EDGAR total AFOLU emission estimates were 12.7 Gt CO₂eq yr⁻¹ on average over the period 2000-2009 (2000s hereafter), and 11.9 Gt CO₂eq yr⁻¹ for the year 2010. By contrast, estimates made with FAOSTAT and Houghton data show the same total AFOLU decadal average emissions of 9.9 Gt CO₂eq yr⁻¹ for 2000s, and a range of 9.5-10.1 Gt CO₂eq yr⁻¹ in 2010. The two datasets thus exhibit substantial differences with EDGAR, producing estimates that are lower than EDGAR's by nearly 3 Gt CO₂eq yr⁻¹ for the 2000s decadal estimates, and about 2 Gt CO₂eq yr⁻¹ for 2010.

The aim of this paper is therefore to investigate the reasons for these differences, in an effort to improve the characterization of AFOLU emissions and their contribution to total anthropogenic forcing. We include analysis of recent emission data, published by FAOSTAT and the Global Carbon Project up to the year 2012, in order to gain further insights into the trends identified by the IPCC AR5 WGIII up to the year 2010.

5.1 Differences between EDGAR and FAOSTAT/Houghton

We identified four major differences between EDGAR and FAOSTAT/Houghton, as follows.

First, as discussed in the methodological section above, EDGAR reports excess agriculture emissions, by counting under AFOLU the emissions arising from energy use in agriculture. These represent an excess of 0.6 Gt CO₂eqyr⁻¹ in 2010, which should be subtracted from the 11.9 Gt CO₂eqyr⁻¹EDGAR estimates published in IPCC AR5 WGIII Section 11.2.2. Reallocation to the energy sector is needed in order to be consistent with existing IPCC guidelines and UNFCCC reporting rules. This results in revised total AFOLU emissions by EDGAR of 11.3 Gt CO₂eqyr⁻¹ in 2010.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Second, additional differences between EDGAR and FAOSTAT agriculture estimates were identified within the IPCC AR5 (Tubiello et al., 2013; Smith et al., 2014). These differences, which are behind the 5.2-5.8 Gt CO₂eq yr⁻¹ emission range identified by the IPCC AR5 WGIII Ch. 11, were further analyzed herein. They arose mainly from rice and savannah burning emissions estimates, explaining 0.3 Gt CO₂eq yr⁻¹ and 0.2 Gt CO₂eq yr⁻¹, respectively, of the 0.6 Gt CO₂eq yr⁻¹ gap between the two databases. Rice differences were explained by the use of different parameters linked to different IPCC Guidelines between EDGAR (IPCC, 1997) and FAOSTAT (IPCC, 2006). Differences in estimates for savannah burning were related instead to different Tier approaches between FAOSTAT and the GFED3 database used by EDGAR (Rossi et al, 2014, Submitted).

The third and more significant difference identified between EDGAR, FAOSTAT and the Houghton datasets is in estimated FOLU emissions. These generate a range of emission estimates of 4.2-5.5 Gt CO₂ eq yr⁻¹ for the year 2010 (Tab. 1), with Houghton and EDGAR providing the lower and higher estimates, respectively, and FAOSTAT estimates in the middle of the range. The reason for this large difference across databases, about 1.3 Gt CO₂eq yr⁻¹, was more difficult to assess, compared to those observed for the agriculture sector discussed above. On the one hand, differences between Houghton and FAOSTAT were easily explained by the different Tier approaches between FAOSTAT and the GFED3 database used by Houghton for forest and peat fires (Rossi et al, 2014, Submitted). On the other, differences between EDGAR and the other two databases could not be resolved. Indeed, IPCC AR5 WGIII Section 11.2.3 had already acknowledged the significant differences between EDGAR and FAOSTAT/Houghton for FOLU, noting that the EDGAR estimates could only be considered approximations of overall land-based CO₂ fluxes. Additional analyses we made over a test period 2001-2010, focusing on comparing the EDGAR sub-category *Forest Decay*, and potentially relevant FAOSTAT and Houghton categories net forest conversion and forest

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

carbon stock change, failed to add insight into the observed differences. As discussed in the methodology section, a lack of needed details from published sources for EDGAR limited our ability to tease out plausible explanations for the substantial differences in estimates. This large unexplained gap added to the serious methodological issues raised above with regards to the EDGAR category *Forest Decay*. These findings thus suggest that, at a minimum, the EDGAR database should not be used as a single source for analyzing FOLU emissions, as was done instead in the IPCC AR5 SPM.

Aggregate AFOLU emission estimates

Based on the above results and conclusions, we re-estimated total AFOLU emissions and their share to total anthropogenic emissions, by jointly analyzing the three datasets used in IPCC WGIII AR5 (Fig. 1). The aggregated results confirmed the historical AFOLU trends reported by IPCC.

Decadal average agriculture emissions grew, from 4.6-5.1 (4.8 ± 0.3) Gt CO₂ eq yr⁻¹ in the 1990s, to 5.0-5.5 (5.1 ± 0.3) Gt CO₂ eq yr⁻¹ in the 2000s, reaching 5.4 ± 0.3 Gt CO₂ eq yr⁻¹ in 2010 (Tab. 1). Overall, the results confirmed a growth trend in agriculture emissions of about 1% over the last two decades (FAO, 2014). Recent updates by FAO, to 5.4 Gt CO₂ eq yr⁻¹ in 2012, show that this trend is continuing past 2010, the last year assessed by IPCC, with emission growth driven primarily by livestock and synthetic fertilizer use (FAOSTAT, 2014).

Decadal average FOLU emissions declined, from a range of 5.7-6.8 (6.4 ± 0.6) Gt CO₂ eq yr⁻¹ in the 1990s, to a range of 4.9-6.5 (5.5 ± 0.9) Gt CO₂ eq yr⁻¹ in the 2000s, reaching 4.9 ± 0.6 Gt CO₂ eq yr⁻¹ in 2010 (Tab. 1). As for agriculture estimates, the three IPCC WGIII sources confirmed a marked decline of FOLU emissions over the last two decades already noted in the IPCC AR5, and due largely to a slow-down in deforestation rates.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Furthermore, recent FAO updates indicated that total FOLU emissions, as well as emissions from net forest conversion, which is a proxy for deforestation, remained stable in both 2011 and 2012 at their 2010 levels of 4.8 and 3.8 Gt CO₂eq yr⁻¹, respectively (FAOSTAT, 2014). The same trend was reflected in recent updates of FOLU emissions minus removals, resulting in a range of 2.9-3.2 Gt CO₂eq yr⁻¹ in 2012 (FAOSTAT, 2014; Global Carbon Project, 2014).

As a result of the above opposite trends between agriculture and land use emissions, total AFOLU emissions declined slightly, from 10.3-11.9 (11.1±0.8) Gt CO₂ eq yr⁻¹ in the 1990s to 9.9-12.0 (10.6±1.2) Gt CO₂ eq yr⁻¹ in the 2000s, reaching 10.3±0.9 Gt CO₂ eq yr⁻¹ in 2010 (Tab. 1).

Aggregate AFOLU share of total anthropogenic emissions

Using the above aggregate AFOLU emission results, together with EDGAR's estimates of non-AFOLU sectors, we produced three different estimates of total anthropogenic emissions, corresponding to a range of 47.7- 49.0 Gt CO₂eq yr⁻¹ in 2010. It was thus possible to assess the shares of AFOLU to total anthropogenic emissions, for each of the three databases considered (Tab. 2). The resulting aggregated analysis confirmed that the share of AFOLU decreased over time, while the use of the three datasets allowed us to further refine the recently published IPCC AR5 findings.

Results (Tab. 3) indicated that the decadal average share of AFOLU decreased from 28.7±1.5% in the 1990s, to 23.6±2.1% in the 2000s, and reached an annual value of 21.2±1.5% in 2010, lower than the 24% reported by the IPCC AR5 WGIII SPM. Similarly, results confirmed that both the share of agriculture and FOLU decreased over time (Tab. 3). The agriculture share decreased less than the AFOLU aggregate, from a decadal average of 12.3±0.6% in 1990s, to 11.5±0.3% in 2000s, reaching an annual value of 11.2±0.4% in 2010.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

The FOLU share decreased substantially more than the AFOLU aggregate, from a decadal average of $16.1 \pm 1.1\%$ in 1990s, to $12.2 \pm 1.7\%$ in 2000s, reaching an annual value of $10.0 \pm 1.2\%$ in 2010. Fig. 2 provides a visual representation of the time dependency of these figures.

Finally, using the explicit information provided by the FAOSTAT database, our results further indicated that the share of deforestation to total anthropogenic emissions was a mere 7.9% in 2010. This share was much smaller than the 20% value cited by Stern (2007) when proposing REDD+ as an effective global mitigation option. Data shown in Tab. 2 for EDGAR 1990s estimates, further suggest that it is likely that the widely cited 20% estimate for deforestation alone, was in fact the total FOLU emission share (17.1%) mentioned in the 2007 IPCC AR4 WGIII SPM Report (IPCC, 2007). Similarly to the 2014 SPM, the 2007 SPM had also relied on EDGAR-only emissions estimates.

7. Conclusions

The findings of this work confirm general known trends in AFOLU emissions over the last two decades, but add additional new estimates that further quantify the continuous increase of agriculture emissions and the substantial decrease of emissions from forest and other land use. These absolute trends, plus the fact that energy emissions continue to grow faster than those from AFOLU, result in decreasing shares of AFOLU compared to total anthropogenic emissions over time.

By aggregating all three AFOLU databases available in IPCC AR5 WGIII, this work was able to further refine critical trends in AFOLU. Results indicated that the share of AFOLU emissions to total anthropogenic emissions in 2010 was 21% rather than 24% and that agriculture (11%) has become as large a contributor to global emissions as forest and land use

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

combined (10%). Furthermore, this analysis added new estimates for AFOLU up to the year 2012, to confirm trends that the IPCC AR5 had analyzed only until 2010.

Despite the large uncertainties that characterize emissions estimates in AFOLU, these findings are important for several reasons.

First, they highlight difficulties in the use of a widely applied database of emissions estimates, EDGAR, indicating specific problems in its FOLU emission estimates. As a result, the findings of this work warn against the use of this database as the sole source of FOLU data.

Second, they show that crop and livestock activities have become the dominant source of AFOLU emissions, indicating the need to increase attention to this sector, in parallel to the ongoing international REDD+ efforts to reduce emissions from deforestation and forest degradation. In fact, a large component of FOLU emissions are in any case driven by agriculture, in particular deforestation and peatland degradation.

In conclusion, the analysis presented herein contributes to the global knowledge base on GHG emissions from AFOLU, and points to the importance of developing better integrated and multi-source data for an improved understanding of the role that agriculture, forestry and other land use activities can play, under current and future international climate agreements, towards limiting dangerous anthropogenic interference with the climate system.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

8. Acknowledgements

This work was carried out with generous funding by the Governments of Germany (GCP/GLO/286/GER) and Norway (GCP/GLO/325/NOR) to the “*Monitoring and Assessment of GHG Emissions and Mitigation Potential from Agriculture*” Project of the FAO Climate, Energy and Tenure Division. P. Smith is a Royal Society Wolfson Merit Award holder and his input contributes to the University of Aberdeen Environment and Food Security Theme and to Scotland’s ClimateXChange. J. House was funded by a Leverhulme Research Fellowship. The FAO Statistics Division maintains the FAOSTAT Emissions database with regular program funds allocated through Strategic Objective 6.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

9. References

- Ciais, P., C. Sabine, G. Bala, et al. (2013) Carbon and Other Biogeochemical Cycles. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker, T.F., D. Qin, G.-K. Plattner, et al. (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- EDGAR (2013) Emissions Database for Global Atmospheric Research (EDGAR), available at link <http://edgar.jrc.ec.europa.eu>, accessed on September 2013.
- FAO (2010) Global Forest Resources Assessment, FAO, Rome.
- FAO (2011) Linking sustainability and climate financing: Implications for agriculture. Rome, Italy. <http://www.fao.org/docrep/015/i2480e/i2480e00.pdf>
- FAO (2014) Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks: 1990-2011 Analysis. FAO Statistics Division Working Paper Series, 14/01. UN FAO, Rome, Italy. <http://www.fao.org/docrep/019/i3671e/i3671e.pdf>
- FAOSTAT (2014) FAOSTAT Emissions database. Agriculture: http://faostat3.fao.org/faostat-gateway/go/to/browse/G1/*E and Land Use: http://faostat3.fao.org/faostat-gateway/go/to/browse/G2/*E
- Friedlingstein P, S. Solomon, G.-K. Plattner, R. Knutti, and M. Raupach (2011) Long-term climate implications of twenty-first century options for carbon dioxide emission mitigation. *Nature Climate Change*, **1**, 4457–4461.
- Houghton, R. A. (2003) Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850–2000. *Tellus B*, **55**, 378–390.
- Houghton, R. A., G. R. van der Werf, R. S. DeFries, M. C. Hansen, J. I. House, C. Le Quéré, J. Pongratz, and N. Ramankutty (2012) Chapter G2 Carbon emissions from land use and land-cover change. *Biogeosciences Discuss*, **9**, 835–878.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (Eds), IGES, Hayama, Japan.

IPCC (2007) Summary for Policymakers. In: *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer (eds), Cambridge University Press, Cambridge, UK and New York, NY, USA.

IPCC (2014) Summary for Policymakers. In: *Climate Change 2014, Mitigation. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and J.C. Minx (eds.). Cambridge University Press, Cambridge, UK and New York, NY, USA.

Le Quéré C., Andres R.J., Boden T., et al. (2013) The global CO₂ budget 1959 – 2011. *Earth System Science Data*, **5**, 165-185, doi:10.5194.

Pongratz J. , Reick C. H., Houghton R. A., House and J. I. (2014) Terminology as a key uncertainty in net land use and land cover change carbon flux estimates., *Earth Systems Dynamics*, **5**, 177–195

Rossi, S., Tubiello, F. N., Salvatore, M., Ferrara. A., Rossi. S. (2014) New estimates of greenhouse gas emissions from biomass fires. *Global Change Biol.*, submitted.

Smith, P., Bustamante, M., Ahammad, et al. (2014) Agriculture, Forestry and Other Land Use (AFOLU). In: *Climate Change 2014, Mitigation. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Zwicker and J.C. Minx (eds.). Cambridge University Press, Cambridge, UK and New York, NY, USA

Stern N. (2007) Economics of climate change: the Stern review. Cambridge University Press, Cambridge, UK, 712 pg.

Tubiello, F. N., Salvatore, M., Ferrara. A., Rossi. S., Fitton. N., and Smith. P. (2013) The FAOSTAT database of GHG Emissions from agriculture. *Environmental Research Letters*, **8** doi:10.1088/1748-9326/8/1/015009.

Van der Werf G.R., J.T. Randerson, L. Giglio, et al. (2006) Interannual variability of global biomass burning emissions from 1997 to 2004. *Atmospheric Chemistry and Physics Discussions*, **6**, 3175–3226.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

10. Supporting Information Legends

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

11. Tables

Table 1. Emissions by recent decade and in 2010 (Gt CO₂ eq).

	1990s				2000s				2010			
Database	E	H	F	Mean±SD	E	H	F	Mean±SD	E	H	F	Mean±SD
Agriculture	5.1	4.6	4.6	4.8±0.3	5.5	5.0	5.0	5.1±0.3	5.8	5.2	5.2	5.4±0.3
FOLU	6.8	6.6	5.7	6.4±0.6	6.5	4.9	4.9	5.5±0.9	5.5	4.2	4.9	4.9±0.6
AFOLU	11.9	11.2	10.3	11.1±0.8	12.0	9.9	9.9	10.6±1.2	11.3	9.5	10.1	10.3±0.9

Data sources: E= EDGAR; H=Houghton; F=FAO

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Table 2. Share of AFOLU and its sub-components to total anthropogenic emissions, by decade and for 2010.

	1990s			2000s			2010		
SECTOR	E	H	F	E	H	F	E	H	F
Agriculture	12.9%	11.8%	12.1%	11.8%	11.3%	11.3%	11.6%	11.0%	10.8%
FOLU	17.1%	17.0%	15.1%	14.2%	11.2%	11.2%	11.1%	8.9%	10.1%
AFOLU	30.1%	28.8%	27.1%	26.0%	22.4%	22.4%	22.8%	19.9%	20.9%

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

Table 3. Aggregated share of AFOLU and its sub-components to total anthropogenic emissions, using the three main datasets available in IPCC AR5 WGIII

SECTOR	1990s	2000s	2010
Agriculture	12.3±0.6%	11.5±0.3%	11.2±0.4%
FOLU	16.4±1.1%	12.2±1.7%	10.0±1.2%
AFOLU	28.7±1.5%	23.6±2.1%	21.2±1.5%

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

12. FIGURES LEGENDS

Figure 1. Total AFOLU emissions, with sub-components, for the year 2010, estimated with the three databases used in IPCC AR5.

Figure 2. Historical changes in the share of AFOLU to total anthropogenic emissions, 1990-2010, based on FAOSTAT data for AFOLU and EDGAR data for non-AFOLU sectors.

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

13. FIGURES

FIGURE 1

The Contribution of Agriculture, Forestry and other Land Use activities to Global

Warming, 1990-2010: Not as high as in the past

FIGURE 2