Sagisaka, M., Iwama, S., Ono, S., Yoshizawa, A., Mohamed, A., Cummings, S., Yan, C., James, C., Rogers, S. E., Heenan, R. K., & Eastoe, J. (2013). Nanostructures in Water-in-CO₂ Microemulsions Stabilized by Double-chain Fluorocarbon Solubilizers. *Langmuir*, 29(25), 7618-7628. <https://doi.org/10.1021/la400376g>

Peer reviewed version

License (if available):
Unspecified

Link to published version (if available):
[10.1021/la400376g](https://doi.org/10.1021/la400376g)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Langmuir*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <http://dx.doi.org/10.1021/la400376g>.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Nanostructures in Water-in-CO₂ Microemulsions

Stabilized by Double-chain Fluorocarbon Solubilizers

Masanobu Sagisaka^{1}, Shuho Iwama¹, Shinji Ono¹, Atsushi Yoshizawa¹, Azmi Mohamed², Stephen Cummings³,*

Ci Yan³, Craig James³, Sarah E. Rogers⁴, Richard K. Heenan⁴, and Julian Eastoe^{3}*

¹ Department of Frontier Materials Chemistry, Graduate School of Science and Technology, Hirosaki University,

3 Bunkyo-cho, Hirosaki, Aomori 036-8561, JAPAN

²Department of Chemistry, Faculty of Science and Mathematics, Universiti Pendidikan Sultan Idris
Tanjong Malim Perak 35900 MALAYSIA

³ School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, U.K.

⁴ ISIS-CCLRC, Rutherford Appleton Laboratory, Chilton, Oxon OX11 0QX, U.K.

*To whom all correspondence should be addressed

Masanobu SAGISAKA E-mail: sagisaka@cc.hirosaki-u.ac.jp Phone and Fax: +81-172-39-3569

Julian EASTOE E-mail: Julian.Eastoe@bristol.ac.uk Phone: +44-117-928-9180 Fax: +44-117-925-1295

Abstract

High-pressure small-angle neutron scattering (HP-SANS) studies were conducted to investigate nanostructures and interfacial properties of water-in-supercritical CO₂ (W/CO₂) microemulsions with double-fluorocarbon-tail anionic surfactants, having different fluorocarbon chain lengths and linking groups (glutarate or succinate). At constant pressure and temperature, the microemulsion aqueous cores were found to swell with an increase in water-to-surfactant ratio, W_0 , until their solubilizing capacities were reached. Surfactants with fluorocarbon chain lengths of $n = 4, 6$, and 8 formed spherical reversed micelles in supercritical CO₂ even at W_0 over the solubilizing powers as determined by phase behavior studies, suggesting formation of Winsor-IV W/CO₂ microemulsions and then Winsor-II W/CO₂ microemulsions. On the other hand, a short C2 chain fluorocarbon surfactant analogue displayed a transition from Winsor-IV microemulsions to lamellar liquid crystals at $W_0 = 25$. Critical packing parameters and aggregation numbers were calculated by using area per head group, shell thickness, the core/shell radii determined from SANS data analysis: these parameters were used to help understand differences in aggregation behavior and solubilizing power in CO₂. Increasing the microemulsion water loading led the critical packing parameter to decrease to ~ 1.3 and the aggregation number to increase to > 90 . Although these parameters were comparable between glutarate and succinate surfactants with the same fluorocarbon chain, decreasing the fluorocarbon chain length n reduced the critical packing parameter. At the same time, reducing chain length to 2 reduced negative interfacial curvature, favoring planar structures, as demonstrated by generation of lamellar liquid crystal phases.

Keywords: Supercritical CO₂, Microemulsion, Fluorinated surfactant, Aqueous Core, Micelle, Small-Angle Neutron Scattering

1. Introduction

Supercritical CO₂ (scCO₂) is seen as a promising green solvent in various fields, including organic synthesis, dry cleaning, polymerization, extraction, nanomaterial processing.¹ The reasons why scCO₂ has attracted much attention for those applications are based on CO₂ and supercritical fluid properties, which are, low cost, non-flammability, environmentally benign, natural abundance, high mass transfer, and pressure/temperature-tunable solvency (or CO₂ density). Unfortunately, supercritical CO₂ can dissolve only nonpolar and small molecular mass materials, and common polar or high molecular mass materials always separate from neat scCO₂.² Improving the poor solubility of polar materials is important for developing applications of scCO₂. One of the most promising approaches for enhancing solubility in scCO is to form reversed micelles with high-polarity aqueous cores in the continuous scCO₂ phase, that is, water-in-scCO₂ microemulsions (W/CO₂ μEs).² Since such organized fluids have the attractive characteristics of scCO₂, as well as the solvation properties of water, they have potential as volatile organic compound (VOC)-free and energy-efficient solvents for nano-material synthesis, enzymatic reactions, dry-cleaning, dyeing, preparation of inorganic/organic hybrid materials, and so on².

To be a viable green technology, the amount of surfactant used should be as small as possible, and this needs to be balanced against the need for large interfacial areas in W/CO₂ μEs and appropriate levels of dispersed water for enhanced process efficiencies. One approach to meet these requirements is to explore or develop a highly efficient solubilizers for W/CO₂ μEs. Since 1990, much effort has been directed towards the development of surfactants for W/CO₂ μEs.³⁻³⁰ Three main kinds of surfactant have been examined so far, being hydrocarbon, silicone, and fluorocarbon (FC) surfactants.

Development of CO₂-philic hydrocarbon surfactants has also been conducted for economic and environmental reasons.³⁻⁹ However, most commercial and known hydrocarbon surfactants are insoluble and inactive in scCO₂ systems³. In this regard, it became apparent that conventional surfactant-design theory cannot be applied to W/CO₂ systems directly, and that CO₂-philicity is not directly comparable to

oleo-philicity. Therefore, advancing molecular-design theory for CO₂-philic surfactants has required new directions and paradigms in the field of surfactants.

Only a few hydrocarbon surfactants have so far been reported to stabilize a W/CO₂ μ Es. One of these successful surfactants is the nonionic TMN-6,⁴⁻⁶ which has highly branched alkyl tails and ~8 oxyethylene units: TMN-6 was reported to solubilize water up to a water-to-surfactant molar ratio, W_0 of 30. Custom-made anionic surfactants with highly branched double- or triple-tails (sodium bis(3,5,5-trimethyl-1-hexyl) sulfosuccinate⁷ or (sodium 1,4-bis(neopentyloxy)-3-(neopentyloxycarbonyl)-1,4-dioxobutane-2-sulfonate⁸) were also found to be soluble in scCO₂ and yield transparent single phase W/CO₂ μ Es (Winsor-IV W/CO₂ μ E or **IV μ E**) with W_0 less than 15. Note that the commercial analogue of these surfactants, Aerosol-OT (sodium bis-(2-ethyl-1-hexyl) sulfosuccinate, AOT) is inactive and insoluble scCO₂, hence is ineffective at stabilizing W/CO₂ μ Es. It has been recognized that hydrocarbon surfactants with highly branched tails, especially methyl-branches, can be considered sufficiently CO₂-philic. On the other hand, ester and ether groups have been reported to increase solubility in scCO₂ as well as methyl branches, and highly CO₂-philic, and W/CO₂-interfacially-active copolymers with these groups have been developed and used for emulsification and polymerization.⁹ Unfortunately, an efficient and cost effective hydrocarbon stabilizer for W/CO₂ μ Es, like the AOT used commonly for W/O μ Es¹⁰, has not yet been found.

In the case of silicone surfactants, polydimethylsiloxanes (PDMS) have attracted attention as potential non-fluorinated CO₂-philic. While they are known to be miscible with CO₂, they still require greater pressures to dissolve than is generally needed for fluorocarbon surfactants.¹¹ The majority of siloxane surfactants tested were only found to be soluble at low concentration (~ 0.1 wt%), and none were observed to form microemulsions or generate micellar structures in scCO₂.¹² A copolymer of PDMS_{24-*b*}-EO₂₂ was shown to lower the water - scCO₂ interfacial tension from 20 to 0.2 mN m⁻¹, a PDMS surfactant did however form W/CO₂ emulsions,¹³ which flocculated and coalesced, limiting the stability of these systems. New trisiloxanes ((CH₃)₃SiO)₂Si(CH₃)₂(CH₂)₃(OCH₂CH₂)_{*n*}R were reported to be effective emulsifiers, forming both W/CO₂ emulsions at $n < 7$ and CO₂/W emulsions at $n \geq 7$.¹⁴ This

change in curvature is attributed to variations in hydrophilic-CO₂-philic balance (HCB)¹⁵, which is accompanied by a minimum in interfacial tension between water and CO₂.

In earlier studies, several fluorinated surfactants were found to dissolve in CO₂ and have a high activity at the W/CO₂ interface, suggesting the feasibility of forming W/CO₂ μ Es.^{16,17} Amongst others, surfactants described below are noteworthy for generating W/CO₂ μ Es. Early on a perfluoropolyether (PFPE) surfactant¹⁶ was found to stabilize **IV μ E**, but with only a low $W_0 = 21$ (also expressed as a corrected water-to-surfactant molar ratio by subtracting the low background water solubility in CO₂, $W_0^c = 14$). After that, numerous reports dealing with W/CO₂ μ Es focused on PFPEs.

Another successful class of CO₂-philic surfactants are the HC-FC hybrids, for example sodium 1-pentadecafluoroheptyl-1-octanesulfate (F7H7, (C₇H₁₅)(C₇F₁₅)CHOSO₃Na): these have both a HC and a FC chain in the same molecule. As such F7H7 is able to solubilize up to $W_0=35$ (or $W_0^c=32$) to form stable Winsor-IV W/CO₂ μ Es.¹⁸ Further studies¹⁹, with hybrid surfactants related to F7H7 but with different FC and HC chain lengths, resulted in the formation of Winsor-IV W/CO₂ μ Es for most of the analogues, but unfortunately smaller attainable W_0 values than for F7H7.

Other investigations²⁰⁻²² studied a class of fluorinated AOT analogues, for example sodium bis(1*H*,1*H*,5*H*-octafluoropentyl)-2-sulfosuccinate (di-HCF4), which yield Winsor IV μ Es with $W_0 = 30$ ($W_0^c \sim 20$). In addition, double-FC-tail phosphate surfactants were also found to be efficient μ E stabilizers²³, the most favorable case stabilizing W_0 up to 45.²⁴

Berkowitz et al. and Cummings et al. studied nanostructures of W/CO₂ μ Es with FC and HC surfactants by molecular dynamics simulations. Berkowitz observed a quick self-assembly of the PFPE surfactant spherical reverse micelles over time periods of 5 ns, irrespective of initial conditions. In most cases, the self-assembled PFPE reverse micelles have a spherical shape and properties consistent with SANS results REF. When the FC surfactant is replaced by a HC analogue, the HC assembly contains a region of direct contact between water and carbon dioxide, indicating that HC surfactants are likely to be inappropriate for formation of W/CO₂ μ Es. On the other hand, Cummings reported that the HC tails in scCO₂ had an average of $74 \pm 4\%$ trans bonds (cf. 89% in vacuum) while the FC tails had $91 \pm 2\%$ (cf.

81% in vacuum). This implies that the HC tails assume more contracted conformations in CO₂, indicating CO₂-phobic interactions while the FC tails assume more extended conformations, consistent with favourable CO₂-philic interactions.

Recent molecular simulation studies^{27,28} have elucidated the reasons why FC surfactants can act as efficient W/CO₂ solubilizers; as compared with HC chains, FC groups have (1) a stronger interactions with CO₂ via quadrupolar and dispersion interactions, and (2) weaker FC-FC chain-chain interactions which are down to a weak repulsion, electrostatic in origin. These properties conspire together to give FC surfactant reversed micelles better solvation by CO₂, and this causes lower surfactant interfacial packing densities, and weaker attractive inter-micellar interactions as compared with hydrocarbon surfactant analogues.

To obtain more economic and environmentally friendly W/CO₂ μ Es, the minimum fluorine content necessary to render a surfactant CO₂-philic has been identified by using double-pentyl-tail surfactants with different fluorination levels. It was found that at least two fluorinated carbons (CF₃CF₂-) are required to stabilize μ Es.²⁹ An increase in fluorine level leads to a lower aqueous surface tension at CMC and a lower stabilization cloud pressure for the W/CO₂ μ Es.

Recently, with the aim of optimizing surfactant structure of fluorinated AOT analogues for W/CO₂ μ E, double-FC-tail anionic surfactants with various FC lengths ($n = 4, 6, 8$) and a sulfo-glutarate (n FG(EO)₂) or sulfo-succinate (n FS(EO)₂) head group were synthesized (see **Table 1** for their chemical structures).^{24,30,31} As the structural difference between n FS(EO)₂ and n FG(EO)₂ is only one methylene unit in the group linking the polar ionic head group to the pendant FC tails, these surfactants examined the effect of not only FC length but also the additional methylene spacer. From surface tension measurements on aqueous surfactant solutions, no significant differences were found in CMC, surface tension at CMC and area per head group between n FG(EO)₂ and n FS(EO)₂ at the same FC length n .³¹ However, visual observation and UV-vis spectral measurements with a hydrophilic dye methyl orange (MO) clarified that the solubilizing powers of the glutarates n FG(EO)₂ were higher than those of the succinate analogues n FS(EO)₂: the most efficient found in that study was 4FG(EO)₂ at 75°C (max.

$W_0=80$ in IV μ E) even though it has the shortest FC tails.³¹ Significantly, this is the highest W_0 reported to date for W/CO₂ μ Es. Many earlier papers mentioned that fluorocarbon is one of a few CO₂-philic groups, with longer FC chains promoting the higher the solubilising powers.³² This is an exciting result leading to new approach to equip a fluorine-light surfactant with high solubilizing power. As well as the high solubilizing power, 4FG(EO)₂ was found to give the fastest dissolution into scCO₂, being capable of solubilizing water in just a few seconds, even at high W_0 and close to its maximum solubilizing capacity.³¹ Such a fast solubilization rate is very rare in this field.

It is very interesting to consider how the beneficial properties of 4FG(EO)₂ are related to the molecular structure, especially for advancing surfactant design theory. These excellent solubilizing properties could be related to a low HCB¹⁵, a high critical packing parameter (CPP)³³, weak interactions between tail-tail and tail-head groups^{34,35}, if the spherical reversed micelles supported the formation of IV μ Es.

To further characterize the surfactant, and explore the origins of the super efficiency, this new study reports aggregation behavior and nanostructures of the custom-made double-FC-tail surfactants, n FG(EO)₂ (FC length $n=2, 4, 8$) and n FS(EO)₂ ($n=4, 8$) in W/CO₂ μ Es, by High-pressure Small-Angle Neutron Scattering (HP-SANS). The significance of this study to the field of surfactant science is that optimized, super-efficient, low fluorine content surfactants are now available for stabilization of W/CO₂ μ Es.

2. Experimental Section

2.1. Materials

The surfactants used in this study were sodium 1,5-bis[(1*H*,1*H*,2*H*,2*H*-perfluorobutyl)oxy]-1,5-dioxopentane-2-sulfonate (2FG(EO)₂), sodium 1,5-bis[(1*H*,1*H*,2*H*,2*H*-perfluorohexyl)oxy]-1,5-dioxopentane-2-sulfonate (4FG(EO)₂), sodium 1,5-bis[(1*H*,1*H*,2*H*,2*H*-perfluorodecyl)oxy]-1,5-dioxopentane-2-sulfonate (8FG(EO)₂), and sodium 1,4-bis[(1*H*,1*H*,2*H*,2*H*-perfluorohexyl)oxy]-1,4-dioxobutane-2-sulfonate (4FS(EO)₂), sodium 1,4-bis[(1*H*,1*H*,2*H*,2*H*-perfluorodecyl)oxy]-1,4-dioxobutane-2-sulfonate (8FS(EO)₂). These surfactants except 2FG(EO)₂ were synthesized and evaluated **in terms of** interfacial properties **as described** previously^{24,30,31}. **Two** surfactants were newly synthesized to examine the FC length effect in this study, as shown in **Sec. 2.2**. 1*H*,1*H*,2*H*,2*H*-pentafluoro-1-butanol (SantaCruz) and dimethyl glutaconate (Aldrich) were used without further purification. Reagent grade acetone, dichloromethane, hexane, 1,4-dioxane, toluene, *p*-toluene sulfonic acid monohydrate, and sodium hydrogensulfite were obtained from Wako Pure Chemical Industries and employed as received. Surfactant structures are shown in **table 1** with interfacial properties of aqueous solutions obtained by standard measurements and solubilizing powers in scCO₂.^{24,30,31} As compared to *n*FS(EO)₂, *n*FG(EO)₂ has an extra –CH₂– spacer between the FC chain and the sulfonate group. Ultrapure water with a resistivity of 18.2 MΩ cm, obtained from a Millipore Milli-Q Plus system, was used in the experiments. CO₂ 99.99% purity (Ekika Carbon Dioxide Co., Ltd.) was used. The structures of these steric models and the length of one surfactant molecule in the absence of other molecules were **obtained** using MM2 (Molecular Mechanics program 2) calculations (Chem 3D; CambridgeSoft Corp., Cambridge, MA).

2.2 Synthesis

2.2.1 Synthesis of bis(1*H*,1*H*,2*H*,2*H*-pentafluorobutyl) glutaconate

A mixture of 9.79 g 1*H*,1*H*,2*H*,2*H*-pentafluoro-1-butanol (61.0 mmol), 4.80 g dimethyl glutaconate (30.4 mmol), 1.45 g *p*-toluene sulfonic acid monohydrate (7.4 mmol) in 200 cm³ toluene was refluxed under stirring at 130 °C for 40 h. During the reaction, the methanol liberated was removed azeotropically from the reaction system to shift the equilibrium of the trans-esterification reaction. After the reaction was complete, the mixture was purified by column chromatography with dichloromethane/*n*-hexane (3:1) as a developing solvent and silica gel. Finally, transparent sticky liquid, bis(1*H*,1*H*,2*H*,2*H*-pentafluorobutyl) glutaconate was obtained (yield 3.69 g, 28.8 %): ¹H-NMR (500MHz, CDCl₃, TMS), (δ_H/ppm): 2.40-2.51 (a, m, 4H), 3.28 (c, dd, 2H, *J* = 1.5, 7.2 Hz), 4.41(b, t, 2H, *J* = 5.6 Hz), 4.44 (b', t, 2H, *J* = 5.6 Hz), 5.96 (e, dt, 1H, *J* = 1.6, 15.7 Hz), 7.02 (d, dt, 1H, *J* = 7.2, 15.7 Hz) for C₂F₅CH₂^aCH₂^bOCOCH₂^cCH^d=CH^eCOOCH₂^{b'}CH₂^aC₂F₅; IR (film) ν_{max}/cm⁻¹: 2979, 1749, 1663, 1464, 1348, 1277, 1195, 1159, 1080, 987, 720, 694.

2.2.2 Synthesis of sodium bis(1*H*,1*H*,2*H*,2*H*-pentafluorobutyl)-2-sulfoglutarate (2FG(EO)₂)

Bis(1*H*,1*H*,2*H*,2*H*-pentafluorobutyl) glutaconate (3.29 g, 7.79 mmol) was dissolved in 1,4-dioxane (140 cm³); then, the mixture was heated to 50 °C. A solution of sodium hydrogensulfite (3.60 g, 34.4 mmol) in water (60 cm³) was added to this mixture. The reaction mixture was stirred under reflux for 24 h. After, solvents were evaporated leaving a white solid residue, which was washed with 1,4-dioxane to remove the unreacted diester. The product was Soxhlet extracted with dry acetone to remove excess NaHSO₃ and recrystallized from acetone. It afforded a white powder, 2FG(EO)₂ (with no regioisomers as C₂F₅CH₂CH₂OCOCH₂CH(-SO₃Na)CH₂COOCH₂CH₂C₂F₅), after vacuum drying (yield 3.11 g, 76.0 %); ¹H-NMR (500MHz, CF₃COOD, TMS), (δ_H/ppm): 2.73-2.59 (a, m, 4H), 3.07 (d, dd, 2H, *J* = 6.0, 17.0 Hz), 3.36 (c, dd, 2H, *J* = 7.5, 17.0 Hz), 4.34-4.44 (e, m, 1H), 4.69 (b, t, 4H, *J* = 6.5 Hz) for C₂F₅CH₂^aCH₂^bOCOCH₂^cCH₂^dCH^e(SO₃Na)COOCH₂^bCH₂^aC₂F₅; IR (KBr) ν_{max}/cm⁻¹: 2977, 1736, 1600,

1408, 1348, 1311, 1197, 1080, 800, 720; Elemental analysis for $C_{13}H_{13}O_7F_{10}Na$: found C, 29.1; H, 2.5; S, 7.0. Calcd C, 29.7; H, 2.5.; S, 6.1.

2.3 High-Pressure Small-Angle Neutron Scattering (HP-SANS) measurements and data analysis

Due to the range of neutron wavelengths available, time-of-flight SANS is suitable for studying the shapes and sizes of colloidal systems. High-pressure SANS (HP-SANS) is a particularly important technique for determining surfactant aggregation structures in supercritical CO_2 . The HP-SANS measurements of the D_2O /surfactant/sc CO_2 systems were performed at 45 °C and 350 bar. The LOQ time-of-flight instrument, at the Rutherford Appleton Laboratory at ISIS UK, was used in conjunction with a stirred high-pressure cell (Thar). The path length was 10 mm, the neutron beam diameter was 10 mm. The measurements gave the absolute scattering cross section $I(Q)$ (cm^{-1}) as a function of momentum transfer Q ($\text{\AA}^{-1}$), which is defined as $Q = (4\pi/\lambda)\sin\theta$. The accessible Q range was 0.007-0.22 $\text{\AA}^{-1}$, arising from an incident neutron wavelength of 2.2-10 $\text{\AA}$. The data were normalized for transmission, empty cell, solvent background, and pressure induced changes in cell volume as before.²⁰

Pre-determined amounts of D_2O and surfactant, where the molar ratio of surfactant to CO_2 was fixed at 8×10^{-4} (= 17 mM at the experimental condition), were loaded into the Thar cell. Then, CO_2 (11.3 g), was introduced into the cell by using a high pressure pump, and the surfactant/ D_2O / CO_2 mixture was pressurized up to 350 bar at 45 °C by decreasing inner volume of the Thar cell. With vigorous stirring, visual observation was carried out to identify the mixture to be a transparent single IV μ E or the other turbid phases. Finally, HP-SANS measurements were conducted for not only IV μ Es but also turbid phases. The densities of CO_2 were calculated using the Span-Wagner equation of state (EOS)³⁶.

Neutrons are scattered by short-range interactions with sample nuclei, with the “scattering power” of different components being defined by a scattering-length density (SLD), ρ (cm^{-2}). For CO_2 , $\rho_{CO_2} \sim 2.50 \times \text{mass density} \times 10^{10} \text{ cm}^{-2}$ ³⁷; at the experimental pressure of 350 bar and temperature of 45

°C, the CO₂ density is 0.917 g cm⁻³ so that $\rho_{\text{CO}_2} \sim 2.29 \times 10^{10} \text{ cm}^{-2}$. The scattering length density of surfactant (ρ_{surf}) and D₂O ($\rho_{\text{D}_2\text{O}}$) were obtained using;

$$\rho = \sum_i b_i / V_m \quad (1)$$

b_i are the nuclear scattering lengths as given in the literature³⁸ and V_m is the molecular volume, which can be obtained from the mass density. Mass densities of surfactants were assumed to be 1.0-1.7 g/cm³ as 1.7 g/cm³ for a typical fluorinated compound¹⁷ and 1.0 g/cm³ for a hydrocarbon surfactant. The calculated scattering length densities for $n\text{FS}(\text{EO})_2$ and $n\text{FG}(\text{EO})_2$ were $3.24 \times 10^{10} \text{ cm}^{-2}$ ($n=8$), $2.80 \times 10^{10} \text{ cm}^{-2}$ ($n=4$), and $2.33 \times 10^{10} \text{ cm}^{-2}$ ($n=2$), respectively. The scattering length density of D₂O at 45 °C was calculated to be $\sim 6.32 \times 10^{10} \text{ cm}^{-2}$. Samples in pure CO₂ (11.3 g) were run at the constant molar ratio of surfactant to CO₂ of 8×10^{-4} .

For model fitting data analysis the μE droplets were treated as spherical core-shell particles with a Schultz distribution in core radius.³⁹ Full accounts of the scattering laws are given elsewhere³⁹⁻⁴³. For polydisperse spherical droplets at volume fraction ϕ , radius R_i , volume V_i , and coherent scattering length density ρ_p dispersed in a medium of ρ_m , the normalized SANS intensity $I(Q)$ (cm⁻¹) may be written as

$$I(Q) = \phi (\rho_p - \rho_m)^2 [\sum_i V_i P(Q, R_i) X(R_i)] S(Q, R_{\text{hs}}, \phi_{\text{hs}}) \quad (2)$$

$P(Q, R_i)$ is the single-particle form factor. The Schultz distribution $X(R_i)$ defines the polydispersity using an average radius, R^{av} , and a root-mean-squared deviation, $\sigma = R^{\text{av}}/(Z + 1)^{0.5}$, Z being a width parameter. $S(Q, R_{\text{hs}}, \phi_{\text{hs}})$ is the structure factor, and a hard-sphere model modified for polydispersity was used:^{39,42} the constraints were $\phi_{\text{hs}} = \phi_d$ and $R_{\text{hs}} = R_d^{\text{av}}$ together with the known ρ values for solvents. (The subscripts “ d ” and “ hs ” denote the droplet and hard-sphere, respectively). Using the approach of Ottewill et al., **eq. 2** can be modified to allow for sharp-step shells built onto a spherical core.⁴³

The least-squares FISH program was used to analyze the SANS data.^{19-21,40,41} The fitted parameters are the volume fraction ϕ , the core radius R_c^{av} , polydispersity index σ/R_c^{av} , and shell thickness t_s ; these were initially set at physically reasonable values, radii obtained by preliminary

Guinier analyses R_{sph} , the typical σ/R_c^{av} value (0.15) reported for many common W/O μE systems⁴⁴ and the estimated length of the chains (12.7 Å for $n = 8$, 7.5 Å for $n = 4$, and 4.9 Å for $n = 2$)³², respectively. The sequence of fitting parameters was as follows; firstly, at constant σ/R_c^{av} and t_s values, the other two parameters were adjusted to the experimental data, and then suitable σ/R_c^{av} and t_s were obtained by floating for the best fit (to minimize sum of weighted squared residuals).

3. Results and Discussion

3.1 Characterization of structure in W/CO₂ microemulsions

Previous studies reported solubilizing powers of each surfactant listed in **Table 1**,^{24,30,31} which were measured by visual observation and spectroscopically with the water-soluble dye methyl orange (MO) as an indicator for the μ E water pool. At W_0 lower than the solubilizing power, the water/surfactant/CO₂ mixtures form transparent single-phases W/CO₂ μ E (IV μ Es), but at higher W_0 values turbid W/CO₂ macroemulsions or a precipitate (liquid crystals) are observed.^{24,30-32}

To examine the shape and size of aggregates in surfactant/D₂O/CO₂ mixtures at different W_0 , SANS $I(Q)$ profiles were measured as a function of W_0 at 45 °C and 350 bar, as shown in **Figure 1**, with the fitted $I(Q)$ functions added. SANS profiles can be useful in determining the shape of colloid particles. In the low Q region (typically in the case of droplet microemulsions $< 0.01 \text{ \AA}^{-1}$), the scattering may scale as $I(Q) \sim Q^{-D}$, where D is a characteristic “fractal dimension” for the colloids; hence, the gradient of a log-log plot will be $-D$. In the case of non-interacting spheres, D should be zero in this low Q region, whereas $D = 1$ for rods and 2 for disks.^{30,40-43} For all SANS, profiles except for 2FG(EO)₂ at $W_0 \leq 20$, the gradients in the low Q region of log-log plots were found to be ~ 0 , suggesting the presence of globular (spherical) nanodomains. On the other hand, 2FG(EO)₂ shows negative slopes at $Q < 0.01$ and $W_0 \geq 25$, and shoulders and peaks at Q of 0.02-0.2. These were ascribed to Bragg peaks, suggesting formation of liquid crystal (LC) phases. Earlier lamellar LCs of n FS(EO)₂ in water and/or scCO₂³² and peaks from Miller indices (001), (002) and (003) observed in SANS profiles were identified as lamellar LC phases. Using $d = 2\pi/Q$ the Q max value (001) allows to estimate a layer spacing (d) comprising one 2FG(EO)₂ bilayer and an aqueous layer. The calculated d was 62.5 Å at $W_0=30$, increasing to 110.2 Å at W_0 60. The molecular length of 2FG(EO)₂ is approximately 10 Å by MM2 simulations, so thickness of the bilayers could be ~ 20 Å if CO₂ was not incorporated into the interior, and then thickness of the aqueous layer would be ~ 43 Å at $W_0=30$ and ~ 90 Å at $W_0=60$. It is noticed that doubling the W_0 ratio yielded almost double the aqueous layer thickness, according to a typical swelling behaviour^{45,46} for

lamellar LCs. For lamellar LCs, the relation between d and surfactant volume fraction ϕ_{surf} ($\phi_{\text{surf}} = 1 - \phi_{\text{solvent}}$, ϕ_{solvent} : solvent volume fraction) could be expressed as follows:⁴⁶

$$d = \frac{\delta}{\phi_{\text{surf}}} \quad (3),$$

where δ is a bilayer thickness. On the assumption of absence of CO₂ layer in the lamellar LC, d values calculated by **eq. 3** were 50 Å at $W_0=30$ ($\phi_{\text{surf}}=0.40$) and 80 Å at $W_0=60$ ($\phi_{\text{surf}}=0.25$). The differences of d values from Bragg peak and **eq. 3** are 10-30 Å, implying the presence of CO₂ incorporated into the lamellar LC.

One method to approximate radii from SANS data for the spherical microemulsions is via Guinier plots⁴⁷ ($\log [I(Q)]$ vs Q^2) as shown in supporting information (**Figures SI1-SI3**). In the all plots, linearity was obtained in the appropriate low Q^2 region, and the gradients allowed estimation of a radius of gyration, R_g (the slope = $-R_g^2/3$). This R_g may also be related to a principal sphere radius R_{sph} as $R_g = (3/5)^{0.5} R_{\text{sph}}$.⁴⁷ The values of R_{sph} were calculated and then employed as the starting points for model fit analyses using the full polydisperse Schultz sphere model. The parameter outputs are the average radii for the D₂O cores (R_c^{av}) and reversed micelles ($R_s^{\text{av}} = t_s + R_c^{\text{av}}$), and the polydispersity width (σ/R_c^{av}). These fitted parameters with R_{sph} are shown in **Table 2**. As W_0 increases, the change in intensity at low Q , and crossover at high Q , are characteristic of an increase in droplet size, which is also clear from the fitted radii. The polydispersities σ/R_c^{av} of **IVμEs** at $W_0 \leq 40$ ranged from 0.24 to 0.39, which was 1.2 – 2.0 times larger than typical values for reversed microemulsions.⁴⁴ Such a high polydispersity was reported for W/CO₂ **IVμEs** with the anionic fluorinated double-tail surfactant di-HCF4 and the cationic perfluoropolyether surfactant PFPE-TMMA; the σ/R_c^{av} values being 0.17–0.40 at $W_0 = 5–30$ for di-HCF4²⁰ and for 0.22–0.49 at $W_0 = 19.4–38.1$ for PFPE-TMMA⁴⁸. As the fluorocarbon-hydrocarbon hybrid surfactant F7H7¹⁹ and AOT analogue surfactants (AOK and AO-Vac)⁴⁰ were found to give typical σ/R_c^{av} values (< 0.2) for W/CO₂ **IVμEs**, the origin of these high polydispersities is unlikely to be due to the use of compressed CO₂ fluid (i.e. high diffusivity⁴⁹ increasing frequency of μE-droplet aggregation/separation)^{20,48}. For the surfactant/W/CO₂ mixtures be thermodynamically stable

microemulsions (energy-minimum equilibrium state), longer equilibration might be necessary (> 1 hour) than employed in this study (~5 min owing to expense of SANS time)²⁴, even though clear single phases have already appeared. The limited equilibration time in this experiment may be one reason for the higher polydispersity.

3.2 Adsorption and aggregation Properties of the Double-chain Fluorocarbon Solubilizers in W/CO₂ systems

Table 2 also shows that R_c^{av} and R_s^{av} increased with W_0 , and R_s^{av} was slightly different from R_{sph} in IV μ E phases, namely at W_0 below the maximum solubilizing powers listed in Table 1. To confirm the reliability of the radii obtained by fitting and to estimate an effective area per head group $A_{h,p}$ for n FG(EO)₂ in W/CO₂ IV μ E, Porod plots were generated. Figure 2 shows Porod plots for 4FGEO₂, and those of the other surfactants are displayed in supporting information (Figures SI4 and SI5). For W/O μ E droplets, $A_{h,p}$ can be estimated from high Q SANS data, by assuming a sharp interface and applying the Porod equation⁵⁰:

$$\{I(Q) \cdot Q^4\}_{Q \rightarrow \infty} = 2\pi (\Delta\rho)^2 \Sigma \quad (4)$$

where Σ is the total area per unit volume. Another requirement here is that the cmc (concentration of non-adsorbed free surfactant) in CO₂ is negligible compared with the experimental surfactant concentration; i.e. all N surfactant molecules are at the CO₂-D₂O interface; given those conditions then the area per head group, $A_{h,p} \approx \Sigma/N$. In the Porod plots of Figure 2 at $Q > 0.16 \text{ \AA}^{-1}$, the data are essentially asymptotic, suggesting these approximations are reasonable. The $\Delta\rho$ in eq. 4 was assumed to be $\{(\rho_{D2O} + \rho_{surf})/2 - \rho_{CO2}\}$ because the scattering will arise from not only the D₂O core but also the surfactant shell (ρ_{surf} were $2.33\text{--}3.24 \times 10^{10} \text{ cm}^{-2}$ against $\rho_{CO2} \sim 2.29 \times 10^{10} \text{ cm}^{-2}$) and volume fractions of total surfactant molecular fragments in the W/CO₂ μ E interfacial region were calculated as ~0.5 (see supporting information Fig SI6 and Tables SI1-SI2). The broken lines at each W_0 indicate asymptotes

$\{I(Q) \cdot Q^4\}_{Q \rightarrow \infty}$ giving A_h values as listed in **Table 3**. Except at $W_0 = 60$ which gives rise to a macroemulsion, $A_{h,p}$ values of same surfactant are similar within an experimental uncertainty $\pm 10 \text{ \AA}^2$.

For Porod plots of spherical particles, with these characteristic polydispersities, the position of the first maximum and first minimum typically occurs at $\sim 2.7/R$ and $\sim 4.5/R$ in Q , respectively.⁵⁰ The Porod plots display only the first maxima, but not the first minima clearly. By using the Porod- Q first maxima the μE droplet radii were estimated (**Table 2**); in the table, $R_{p\text{max}}$ means the radius given from the first Porod maximum. As seen in the table R_s^{av} and $R_{p\text{max}}$ were similar at same W_0 , supporting the validity of these analyses.

Figure 3 shows changes in the radii obtained by the theoretical curve fitting for D_2O cores and reversed micelle shells as function of W_0 . When comparing $n\text{FS}(\text{EO})_2$ and $n\text{FG}(\text{EO})_2$, no significant difference in D_2O core radii could be seen at constant W_0 for the IV μEs . Differences in FC length n did not affect the radius in the IV μE region, but at $W_0 \geq 60$ an increase of $\sim 10 \text{ \AA}$ was seen in the radius between $n=4$ and 8 surfactants. This suggests the following: (1) the turbid two-phase system at high W_0 was identified as a Winsor-II $\text{W}/\text{CO}_2\mu\text{E}$, and (2) excess water molecules ($=W_0 -$ the solubilizing power) formed W/CO_2 “macro”-emulsion droplets, and separated from scCO_2 in the absence of stirring whilst the equilibrium reversed micelles still remain in scCO_2 . This behaviour was suggested by changes in UV-vis light absorption of the surfactant/aqueous methyl orange solution/ CO_2 mixtures as a function of W_0 , as seen previously⁵¹. In addition, the linear dependences for radii vs W_0 suggested D_2O molecules are incorporated into spherical μE droplets. This linear behavior for R_c^{av} as a function of W_0 can be used to provide another estimate for the effective head group area using¹⁰;

$$\alpha(p)R_c^{\text{av}} = \frac{3v_w}{A_h}W_0 + \frac{3v_h}{A_h} \quad (5)$$

where v_w is the volume of a water molecule, p is the polydispersity index (σ/R_c^{av}) and $\alpha(p)$ depends on the Schultz distribution $\alpha(p) = 1 + 2p^2$. Therefore, assuming the polydispersity to be independent of W_0 , the broken line for R_c^{av} in **Fig. 3** is expected to have slope which depends on the interfacial area A_h and an intercept which depends on the average head group volume v_h via the head group radius r_h . The head

group areas calculated in this way span 117-129 Å² with an uncertainty of ± 10 Å² (**Table 3**), which is consistent with the previous results obtained by Porod analyses of high Q SANS data in W/CO₂ μ Es. Areas per head group $A_{h,p}$ of 4FG(EO)₂, obtained by Porod analysis were slightly larger than $A_{h,l}$ obtained from the slope of **Fig.3**. The assumption in these calculations for $A_{h,p}$ is that all surfactant molecules are at the W/CO₂ interface, which may not be the case: the computed area will be too large if the solubilities of surfactant in CO₂ and water are high. In a previous study³¹, the dissolution pressures of 0.08 mol% surfactant in scCO₂ were found to decrease at shorter FC chain lengths for n FG(EO)₂ and n FS(EO)₂, suggesting the solubility of 4FG(EO)₂ in pure CO₂ (i.e. number of free 4FG(EO)₂ molecules not at the W/CO₂ interface) is larger than those of longer FC surfactants. In addition, the Kraft temperature shown in **Table 1** shows that 4FG(EO)₂ seems to be more soluble in water, meaning the actual area per head group of 4FG(EO)₂ should be lower than $A_{h,p}$, and could be similar to the $A_{h,l}$.

In previous work²¹, fluorinated double-tail surfactants (di-HCF₄, di-HCF₆ and di-CF₄) having same head groups as n FG(EO)₂ and n FS(EO)₂ were investigated in a similar fashion to as described above, giving 115 ± 5 Å² for A_h (15 °C and 500 bar). Interestingly, and importantly, that value is almost similar as found here, not only for the sulfosuccinate n FS(EO)₂ but also the analogous glutarate surfactant n FG(EO)₂, on account of difference of 30 °C in experimental temperature. For AOT in a range of n -alkanes from propane to decane, a mean value of 72 Å² for A_h was reported.⁴¹ The difference in A_h between W/CO₂ and W/O μ Es suggests that packing density of AOT type surfactants at water surface increases in CO₂ as compared to typical hydrocarbon oil. Earlier papers reported A_h of AOT to be 155 Å² at a W/scCO₂⁵² but 75 Å² at the water/air⁵³ and 72 Å² at the water/alkane interface.⁴¹

Although the values of radii R_c^{av} in **Fig. 3** are approximately half those seen for AOT W/alkane μ Es,¹⁰ this could be accounted for owing to the difference in interfacial packing and the determined A_h values. In **Fig. 3**, intercepts of linear functions of R_c^{av} and R_s^{av} vs W_0 give mean radii of the head group core and the dry reversed micelle, in other words, dimensions of head group (l_h) and overall surfactant molecule (l_{surf}), respectively. The head groups -SO₃Na of n FG(EO)₂ and n FS(EO)₂ displayed the same

radius in dry micelle cores (~ 5.2 Å). The l_{surf} values obtained from **Fig.3** were almost consistent with those calculated by the MM2 calculation, suggesting R_c^{av} and R_s^{av} values in **Fig. 3** to be reasonable: for example, l_{surf} of 8FG(EO)₂ was 17.6 Å from the intercept (radius of dry reversed micelles) of **Fig.3** and 18 Å obtained by MM2 calculation.

By using the intercept for l_h and eq. 5, the effective head group volume was calculated as $v_h = 199\text{--}231$ Å³, and assuming the head group to be spherical, the radius (r_h) would be 3.6–3.8 Å, as summarized in **Table 3**. AOT analogues in W/O μEs were found to have $v_h = 200\text{--}236$ Å³ and $r_h = 3.8$ Å,¹⁰ and these are consistent with the values calculated here for surfactants with identical polar head groups. This suggests the head group volume is not affected by being in supercritical CO₂, making it unlikely the CO₂ penetrates into the polar region of the micellar structure.

These parameters relating to the size of the chain and head group allow estimates of the aggregation number numbers N_{agg} and critical packing parameters (CPP)³³ by using;

$$N_{\text{agg}} = \frac{4\pi(R_c^{\text{av}})^2}{A_h} \quad (6)$$

$$\text{CPP} = \frac{v_c}{A_h \cdot l_c} \quad (7)$$

where v_c is hydrophobic chain volume. If the hydrophobic part is assumed to be a truncated core, the volume should be;

$$v_c = \frac{l_c(A_h + A_c + \sqrt{A_h \cdot A_c})}{3} \quad (8)$$

where A_c is area per hydrophobic chain terminus, obtained using $A_c = 4\pi(R_s^{\text{av}})^2 / N_{\text{agg}}$. To calculate N_{agg} and CPP of surfactants used in this study, A_h values obtained from **eq. 5** were employed here. The calculated N_{agg} and CPP are shown as a function of W_0 in **Figure 4** with previous data for AOT/W/*n*-heptane μEs ¹⁰. As expected from typical behavior of W/O reversed micelle systems^{54,55}, N_{agg} for these W/CO₂ μEs increased with W_0 . For example, N_{agg} was close to 10 at $W_0=10$, but it reached up to ~ 100 and ~ 200 for 4FG(EO)₂ at $W_0=80$ and 8FG(EO)₂ at $W_0=66$, respectively. When the N_{agg} data were compared at a constant W_0 , significant differences were not observed between these FC surfactants. On

the other hand, N_{agg} values of the FC surfactants in scCO₂ μ Es were found to be quite small, one fifth to one sixth of those of AOT/W/*n*-heptane μ Es. Even if compared with AOT in the other organic solvents (eg, $N_{agg} = \sim 90$ at $W_0 = 10$ and ~ 290 at $W_0 = 20$ in isooctane and cyclohexane^{54,55}), the difference in N_{agg} is quite large, resulting from the small R_s^{av} and large A_h for n FG(EO)₂ and n FS(EO)₂ in **IV** μ Es.

The critical packing parameter (CPP)³³ is an indicator of “bulkiness” of a given hydrophobic group compared to the head group, and dictates interfacial curvature of a surfactant assembly. Negative interfacial curvature for a W/O or W/CO₂ μ Es is only stabilized for surfactants with $CPP > 1$, and if $CPP \sim 1$ the interfacial curvature is planar: in other words, the reversed curvature μ E droplets would tend to more planar macroemulsions. At $W_0=10$ in **Fig. 4**, the CPP values are 1.92, 1.76 and 2.57 for 4FG(EO)₂, 4FS(EO)₂ 8FG(EO)₂, respectively. These values were found to decrease with increasing W_0 , and then finally reached constant values of ~ 1.3 at W_0 .

3.3 Effects of Fluorocarbon and Methylene Spacer Lengths on Solubilization

Earlier SANS data for AOT/W/*n*-heptane μEs ¹⁰ were used to calculate CPPs, represented by closed circles in **Fig. 4**. The AOT-analogue glutarate in W/*n*-heptane μEs also had CPP values similar to the regular AOT (e.g. 1.18 at 25 °C and $W_0 = 40$)⁵⁶. These are always smaller than those for the FC-surfactants in scCO_2 at any W_0 , and demonstrate that even at small CPP of ~ 1.15 can stabilize Winsor-IV W/O μEs , whereas such a small CPP is not possible in the W/ CO_2 μEs . It implies **that** other parameters, **for example**, HCB¹⁵, Winsor R^{34,35} and/or W/ CO_2 interfacial tension^{5,52} mainly affect the solubilizing power of the double-FC-tail surfactants in IV μEs .

On the other hand, CPPs of the sulfosuccinate (8FS(EO)₂ and 4FS(EO)₂) and the glutarate (8FG(EO)₂ and 4FG(EO)₂) were found to be similar at same chain lengths *n* in the W/ CO_2 μEs , suggesting that the extra methylene in *n*FG(EO)₂ did not affect CPP: i.e. CPP is not the main parameter explaining differences in solubilizing power. Then, the question “Why can the glutarate surfactants solubilize a higher W_0 than the succinate surfactants can?” still remains unsolved.

The subtle structural change in the head group region (i.e. succinate $\rightarrow$ glutarate) has also been investigated before for hydrocarbon AOT-like surfactants⁵⁶. Interestingly, the addition of just one extra $-\text{CH}_2-$ in the head group had little noticeable effect on most interfacial properties at both air/water and oil/water interfaces; for example, in aqueous systems, the differences in CMC, surface tension and area per head group at CMC were 0.8 mM, < 0.1 mN/m, and $< 7 \text{ \AA}^2$, respectively.⁵⁶ However, a maximum attainable W_0 ratio with the glutarate analogue in W/*n*-heptane μEs was higher by > 10 as compared with the normal succinate (10–40 °C).⁵⁶ Previously studies have been made of succinate and glutarate fluoro-AOT-type analogues in hydrocarbon W/ CO_2 μEs , suggesting that there is a notable enhancement in stability for the glutarate^{21,22} (**i.e. comparing** bis(1H,1H,5Hoctafluoropentyl)-2-sulfosuccinate or di-HCF₄, versus bis(1H,1H,5Hoctafluoropentyl)-2-glutarate or di-HCF₄GLU). However, that work was only limited, comparing the pressure-temperature (*P-T*) phase stability of the W/ CO_2 μEs at one fixed $W_0 = 10$: at 25 °C the phase transition pressure P_{trans} was ~ 12 bar lower for the glutarate analogue (181 vs 193 bar)^{21,22}. The higher solubilizing powers of glutarates have been often reported not only in scCO_2

but also in typical organic solvents. Without a hypothesis based on CPP for the enhanced solubilizing power, there are another two possibilities: the extra methylene spacer results in (1) an increase in structural disorder by lowering molecular symmetry leading to a weaker molecular packing, and (2) a decrease in HCB by increasing hydrophobicity of linking group promoting (possible) CO₂-ester interactions⁵⁷ instead of hydration. These possibilities could be investigated by measuring FT-IR spectra for the ester groups in the two water/scCO₂ mixtures with *n*FG(EO)₂ and *n*FS(EO)₂.⁵⁸

The effect of FC length *n* of the double-FC-tail surfactants on CPP can be seen in **Fig. 4**: longer chains result in larger CPP values. (longer surfactants have larger shells in spite of the core radius being similar, independent of FC length). As listed in **Table 1**, the solubilizing power of *n*FG(EO)₂ at 45 °C increases with an increase in *n*, and this trend would be accelerated by the higher CPP at larger *n* values. On the other hand, the solubilizing power of 4FG(EO)₂ at 75 °C was the highest. An earlier paper^{30,31} discussed the reason why 4FG(EO)₂ is the most efficient solubilizer at temperature > 65 °C, and one of the proposed reasons was formation of bicontinuous microemulsions at high temperatures. Unfortunately, the HP-SANS cell used is restricted, and cannot be used at high temperature, so that bicontinuous microemulsions could not be directly confirmed. However, lamellar LCs were observed in 2FG(EO)₂/W/CO₂ mixtures at $W_0 \geq 25$, raising the possibility of formation of 4FG(EO)₂ bicontinuous microemulsions. In general, a negative curvature of W/O (or W/scCO₂) μ Es approaches zero with decreasing CPP³³ and/or increasing HLB (or HCB)¹⁵, and the microemulsion often turns into a lamellar LC or a bicontinuous system bearing net zero interfacial curvature. In the case of 4FG(EO)₂, the CPP and HCB values should be close to those of 2FG(EO)₂, which prefers zero interfacial curvature, but the ability to lower W/CO₂ interfacial tension could be higher than that of 2FG(EO)₂ owing to the longer FC chains. In addition, affinities of 4FG(EO)₂ tail-to-CO₂ and head-to-water would be enhanced at higher temperature, which would in turn enhance the ability to lower interfacial tension. The discussion about CPP and HCB, suggests that the 4FG(EO)₂ bicontinuous microemulsions, which should require not only zero interfacial curvature but also an extremely low W/CO₂ interfacial tension < 1 mN/m, are highly probable.

4. Conclusions

A previous study found the glutarate headgroup compound 4FG(EO)₂ to be an effective and efficient double-FC-tail surfactant for water-in-scCO₂ microemulsions: this compound can solubilize water of W_0 up to 80 at 75 °C, in spite of the short C₄ FC tails. At high temperature > 65 °C, the solubilizing power was the highest in for FG(EO)₂ analogues with different FC tail lengths ($n = 2, 4, 6$, and 8). Microemulsions in CO₂ are promising solvents for green chemistry, and therefore should be prepared with low levels of surfactant, being also inexpensive and environmentally-benign. Therefore, finding short FC-tail surfactants, which generate high solubilizing power, is key to designing useful CO₂-philic surfactants. Further surfactant structure-performance studies are needed to develop CO₂ as an “environmentally-benign” and “energy-saving” solvent, for applications such as extraction, dyeing, dry cleaning, metal-plating, and organic or nanomaterial synthesis.

This study characterized microemulsions with the double-FC-tail solubilizers n FG(EO)₂ and n FS(EO)₂, and clarified the effects of FC tail and methylene spacer lengths on the nanostructure (radii of D₂O cores and reversed micelle shells, area per head group, aggregation number, and critical packing parameter CPP) of the microemulsions. Through SANS experiments and data analyses, relationships between nanostructure and the surfactant chain length or composition W_0 were revealed: (1) the core and shell radii and aggregation number increased with W_0 , but these were almost independent of addition of extra methylene spacer at same FC length and W_0 ; (2) values of area per head group for each surfactant were almost the same based on the common –SO₃Na head group employed, and (3) the CPPs of FC surfactants decreased for the shorter FC lengths and larger W_0 values. These changes account for the observed transition of microemulsions into lamellar LCs as a function of surfactant structure and microemulsion composition. It was also interesting to note that the nanostructures (core radius) and surfactant properties ($A_{h,l}$, N_{agg} , and CPP) in the CO₂ IVμEs were significantly different from those in typical AOT/W/O μEs at same W_0 , in spite of similar surfactant structure and concentration. Those differences could be down to significant effects of intermolecular interactions between hydrocarbon chains and hydrocarbon solvents, as compared with fluorocarbon chains and CO₂.

Although the reason why 4FG(EO)₂ displays the highest solubilizing power (maximum $W_0=80$) at the high temperatures is still unclear, one conceivable reason is formation of W/CO₂-type bicontinuous microemulsions, and it seems highly probable based on fact (3) mentioned above. If bicontinuous μ Es form in the 4FG(EO)₂ mixtures, this would be first example of single-phase bicontinuous microemulsions for a ternary single-surfactant/W/CO₂ mixture. However, in earlier papers bicontinuous microemulsions were found but in equilibrium with separated water and CO₂ phases⁵⁹ or by using mixed fluorinated surfactants (Zonyl FSH/Zonyl FSN 100 mixtures⁶⁰). High water content bicontinuous microemulsions could represent new generation solvents with unique properties. Further studies on 4FG(EO)₂ and its analogues should be continued to advance surfactant design theory for W/CO₂ μ E and bicontinuous systems.

Acknowledgments

MS thanks the Japan Society for the Promotion of Science (JSPS) for a 1-year fellowship (Excellent Young Researcher Overseas Visit Program) and Hirosaki University. SC thanks the EPSRC for a PhD scholarship under the Next Generation Facility User scheme (EP/F020686). AM thanks the Ministry of Higher Education of Malaysia and Universiti Pendidikan Sultan Idris for the provision of a PhD studentship. We also acknowledge STFC for the allocation of beam time, travel, and consumables grants at ISIS.

This project was supported by JSPS [KAKENHI, Grant-in-Aid for Young Scientists (A), No. 23685034], JST [A-STEP, No. AS231Z02302D], and Leading Research Organizations, namely NSERC, ANR, DFG, RFBR, RCUK and NSF as Partner Organizations under the G8 Research Councils Initiative for Multilateral Research Funding, the Noguchi Institute Foundation, and Nissan Chemical Industries. The EPSRC is thanked for funding under grants EP/C523105/1, EP/F020686 and EP/K020676/1.

Figure captions

Figure 1. SANS profiles for surfactant/D₂O/CO₂ mixtures with various W_0 at 45 °C and 350 bar (CO₂ density = 0.92 g/cm³). Fitted curves were based on a model incorporating a Schultz distribution of polydisperse spheres with a core/shell structure. These SANS profiles are for (a)-(b) 2FG(EO)₂, (c) 4FG(EO)₂, (d) 4FS(EO)₂, (e) 8FG(EO)₂, and (f) 8FS(EO)₂. The molar ratio of the surfactant to CO₂ was fixed at 8×10^{-4} .

Figure 2. Porod plots to obtain μ E droplet radii R_{pmax} and areas per head group $A_{\text{h,p}}$ in D₂O/CO₂ mixtures with 4FG(EO)₂ for several W_0 values at 45 °C and 350 bar. Each arrow shows the first maximum used for calculating R_{pmax} . Broken lines display $\{I(Q) Q^4\}_{Q \rightarrow \infty}$ for calculation of $A_{\text{h,p}}$.

Figure 3. Dependence of droplet radius for the 8FG(EO)₂ D₂O/CO₂ μ Es as a function of W_0 at 45 °C and 350 bar.

Figure 4. Change in aggregation number (N_{agg}) and critical packing parameter (CPP) of n FG(EO)₂ and n FS(EO)₂ in D₂O/CO₂ μ Es or AOT in D₂O/ n -heptane μ Es¹⁰ as a function of W_0 at 45 °C and 350 bar.

References

- (1) Beckman, E. J. Supercritical and Near-Critical CO₂ in Green Chemical Synthesis and Processing. *J. Supercrit. Fluids.* **2004**, 28, 121-191.
- (2) Goetheer, E. L. V.; Vortaman, M. A. G.; Keurentjes, J. T. F. Opportunities for Process Intensification Using Reverse Micelles in Liquid and Supercritical Carbon Dioxide. *Chem. Eng. Sci.* **1999**, 54, 1589-1596.
- (3) Consani, K. A.; Smith, R. D. Observations on the Solubility of Surfactants and Related Molecules in Carbon Dioxide at 50 °C. *J. Supercrit. Fluids* **1990**, 3, 51-65.
- (4) Ryoo, W.; Webber, S. E.; Johnston, K. P. Water-in-Carbon Dioxide Microemulsions with Methylated Branched Hydrocarbon Surfactants. *Ind. Eng. Chem. Res.*, **2003**, 42, 6348-6358.
- (5) Sagisaka, M.; Koike, D.; Mashimo, Y.; Yoda, S.; Takebayashi, Y.; Furuya, T.; Yoshizawa, A.; Sakai, H.; Abe, M.; Otake, K. Water/Supercritical CO₂ Microemulsions with Mixed Surfactant Systems. *Langmuir* **2008**, 24, 10116-10122.
- (6) Sagisaka, M.; Fujii, T.; Koike, D.; Yoda, S.; Takebayashi, Y.; Furuya, T.; Yoshizawa, A.; Sakai, H.; Abe, M.; Otake, K. Surfactant-Mixing Effects on the Interfacial Tension and the Microemulsion Formation in Water/Supercritical CO₂ System. *Langmuir* **2007**, 23, 2369-2375.
- (7) Johnston, K. P.; Cho, D.; DaRocha, S. R. P.; Psathas, P. A.; Ryoo, W.; Webber, S. E.; Eastoe, J.; Dupont, A.; Steytler, D. C. Water in Carbon Dioxide Macroemulsions with a Hydrocarbon Surfactant. *Langmuir* **2001**, 17, 7191-7193.
- (8) Hollamby, M. J.; Trickett, K.; Mohamed, A.; Cummings, S.; Tabor, R. F.; Myakonkaya, O.; Gold, S.; Rogers, S.; Heenan, R. K.; Eastoe, J. Tri-Chain Hydrocarbon Surfactants as Designed Micellar Modifiers for Supercritical CO₂. *Angew. Chem., Int. Ed.* **2009**, 48, 4993-4995.

- (9) Lee, H.; Pack, J. W.; Wang, W.; Thurecht, K. J.; Howdle, S. M. Synthesis and Phase Behavior of CO₂-Soluble Hydrocarbon Copolymer: Poly(Vinyl Acetate-*alt*-Dibutyl Maleate). *Macromolecules* **2010**, *43*, 2276-2282.
- (10) Nave, S.; Eastoe, J.; Heenan, R. K.; Steytler, D.; Grillo, I. What is So Special About Aerosol-OT? 2. Microemulsion Systems. *Langmuir* **2000**, *16*, 8741-8748.
- (11) Hoefling, T. A.; Newman, D. D.; Enick, E. J.; Beckman, E. J. Effect of Structure on the Cloud-Point Curves of Silicone-Based Amphiphiles in Supercritical Carbon Dioxide. *Supercrit. Fluids*, **1993**, *6*, 165-171.
- (12) Psathas, P. A.; da Rocha, S. R. P.; Lee, C. T.; Johnston, K. P.; Lim, K. T.; Webber, S. Water-in-Carbon Dioxide Emulsions with Poly(dimethylsiloxane)-based Block Copolymer Ionomers. *Ind. Eng. Chem. Res.* **2000**, *39*, 2655-2664.
- (13) Johnston, K. P. Block Copolymers as Stabilizers in Supercritical Fluids. *Curr. Opin. Coll. Int. Sci.* **2000**, *5*, 351-356.
- (14) da Rocha, S. R. P.; Dickson, J.; Cho, D.; Rossky, J. P.; Johnston, K. P. Stubby Surfactants for Stabilization of Water and CO₂ Emulsions: Trisiloxanes. *Langmuir* **2003**, *19*, 3114-3120.
- (15) Lee, C. T., Jr.; Psathas, P. A.; Johnston, K. P.; deGrazia, J.; Randolph, T. W. Water-in-Carbon Dioxide Emulsions: Formation and Stability. *Langmuir* **1999**, *15*, 6781-6791.
- (16) Johnston, K. P.; Harrison, K. L.; Klarke, M. J.; Howdle, S. M.; Heitz, M. P.; Bright, F. V.; Carlier, C.; Randolph, T. W. Water-in-Carbon Dioxide Microemulsions: A New Environment for Hydrophiles Including Proteins. *Science* **1996**, *271*, 624-626.
- (17) Zielinski, R. G.; Kline, S. R.; Kaler, E. W.; Rosov, N. A Small-Angle Neutron Scattering Study of Water in Carbon Dioxide Microemulsions. *Langmuir* **1997**, *13*, 3934-3937.

- (18) Niemeyer, E. D.; Bright, F. V. The pH within PFPE Reverse Micelles Formed in Supercritical CO₂. *J. Phys. Chem. B* **1998**, *102*, 1474-1478.
- (19) Eastoe, J.; Bayazit, Z.; Martel, S.; Steytler, D. C.; Heenan, R. K. Droplet Structure in Water-in-CO₂ Microemulsion. *Langmuir* **1996**, *12*, 1423-1424.
- (20) Eastoe, J.; Cazelles, B. M. H.; Steytler, D. C.; Holmes, J. D.; Pitt, A. R.; Wear, T. J.; Heenan, R. K. Water-in-CO₂ Microemulsions Studied by Small-Angle Neutron Scattering. *Langmuir* **1997**, *13*, 6980-6984.
- (21) Eastoe, J.; Downer, A.; Paul, A.; Steytler, D. C.; Rumsey, E.; Penfold, J.; Heenan, R. K. Fluoro-Surfactants at Air/Water and Water/CO₂ Interfaces. *Phys. Chem. Chem. Phys.* **2000**, *2*, 5235-5242.
- (22) Eastoe, J.; Paul, A.; Downer, A.; Steytler, D. C.; Rumsey, E. Effects of Fluorocarbon Surfactant Chain Structure on Stability of Water-in-Carbon Dioxide Microemulsions. Links Between Aqueous Surface Tension and Microemulsion. *Langmuir* **2002**, *18*, 3014-3017.
- (23) Xu, B.; Lynn, G. W.; Guo, J.; Melnichenko, Y. B.; Wignall, G. D.; McClain, J. B.; DeSimone, J. M.; Johnston, C. S., Jr. NMR and SANS Studies of Aggregation and Microemulsion Formation by Phosphorus Fluorosurfactants in Liquid and Supercritical Carbon Dioxide. *J. Phys. Chem. B* **2005**, *109*, 10261-10269.
- (24) Sagisaka, M.; Yoda, S.; Takebayashi, Y.; Otake, K.; Kitiyanan, B.; Kondo, Y.; Yoshino, N.; Takebayashi, K.; Sakai, H.; Abe, M. Preparation of a W/scCO₂ Microemulsion Using Fluorinated Surfactants. *Langmuir* **2003**, *19*, 220-225.
- (25) Lu, L.; Berkowitz, M. L. Molecular Dynamics Simulation of a Reverse Micelle Self Assembly in Supercritical CO₂. *J. Am. Chem. Soc.* **2004**, *126*, 10254-10255.
- (26) Salaniwal, S.; Cui, S. T.; Cummings, P. T.; Cochran, H. D. Self-Assembly of Reverse Micelles in Water/Surfactant/Carbon Dioxide Systems by Molecular Simulation. *Langmuir* **1999**, *15*, 5188-5192.

- (27) Dalvi, V. H.; Srinivasan, V.; Rossky, P. J. Understanding the Effectiveness of Fluorocarbon Ligands in Dispersing Nanoparticles in Supercritical Carbon Dioxide. *J. Phys. Chem. C* **2010**, *114*, 15553-15561.
- (28) Dalvi, V. H.; Srinivasan, V.; Rossky, P. J. Understanding the Relative Effectiveness of Alkanethiol Ligands in Dispersing Nanoparticles in Supercritical Carbon Dioxide and Ethane. *J. Phys. Chem. C* **2010**, *114*, 15562-15573.
- (29) Mohamed, A.; Sagisaka, M.; Guittard, F.; Cummings, S.; Paul, A.; Rogers, S. E.; Heenan, R. K.; Dyer, R.; Eastoe, J. Low Fluorine Content CO₂-Philic Surfactants. *Langmuir* **2011**, *27*, 10562-10569.
- (30) Sagisaka, M.; Iwama, S.; Hasegawa, S.; Yoshizawa, A.; Mohamed, A.; Cummings, S.; Rogers, S. E.; Heenan, R. K.; Eastoe, J. Super-Efficient Surfactant for Stabilizing Water-in-Carbon Dioxide Microemulsions. *Langmuir* **2011**, *27*, 5772-5780.
- (31) Sagisaka, M.; Iwama, S.; Yoshizawa, A.; Mohamed, A.; Cummings, S.; Eastoe, J. An Effective and Efficient Surfactant for CO₂ Having Only Short Fluorocarbon Chains. *Langmuir* **2012**, *28*, 10988-10996.
- (32) Sagisaka, M.; Koike, D.; Yoda, S.; Takebayashi, Y.; Furuya, T.; Yoshizawa, A.; Sakai, H.; Abe, M.; Otake, K. Optimum Tail Length of Fluorinated Double-Tail Anionic Surfactant for Water/Supercritical CO₂ Microemulsion Formation. *Langmuir* **2007**, *23*, 8784-8788.
- (33) Israelachvili, J. N. Measurements of Hydration Forces Between Macroscopic Surfaces. *Chem. Scr.* **1985**, *25*, 7-14.
- (34) Winsor, P. A. Hydrotropy, Solubilisation and Related Emulsification Processes. *Trans. Faraday Soc.* **1948**, *54*, 376-398.
- (35) Harrison, K. L.; Johnston, K. P.; Sanchez, L. C. Effect of Surfactants on the Interfacial-Tension Between Supercritical Carbon-Dioxide and Polyethylene-Glycol. *Langmuir* **1996**, *12*, 2637-2644.

- (36) Span, R.; Wähler, W. A New Equation of State for Carbon Dioxide Covering the Fluid Region from the Triple-Point Temperature to 1100 K at Pressures up to 800 MPa. *J. Phys. Chem. Ref. Data* **1996**, *25*, 1509-1596.
- (37) McClain, J. B.; Londono, D.; Combes, J. R.; Romack, T. J.; Canelas, D. A.; Betts, D. E.; Wignall, G. D.; Samulski, E. T.; DeSimone, J. M. Solution Properties of a CO₂-Soluble Fluoropolymer via Small Neutron Scattering. *J. Am. Chem. Soc.* **1996**, *118*, 917-918.
- (38) Sears, V. F. Neutron Scattering Lengths and Cross Sections. *Neutron News* **1992**, *3*, 26-37.
- (39) Kotlarchyk, M.; Chen, S.-H.; Huang, J. S.; Kim, M. W. Structure of Three-Component Microemulsions in the Critical Region Determined by Small Angle Neutron Scattering Data. *Phys. Rev. A* **1984**, *29*, 2054-2069.
- (40) Eastoe, J.; Gold, S.; Rogers, S.; Wyatt, P.; Steytler, D. C.; Gurgel, A.; Heenan, R. K.; Fan, X.; Beckman, E. J.; Enick, R. M. Designed CO₂-Philes Stabilize Water-in-Carbon Dioxide Microemulsions. *Angew. Chem.* **2006**, *118*, 3757-3759.
- (41) Eastoe, J.; Robinson, B. H.; Young, W. K.; Steytler, D. C. Scattering Studies of Microemulsions in Low-Density Alkanes. *J. Chem. Soc., Faraday Trans.* **1990**, *86*, 2883-2889.
- (42) Hayter, J. B.; Penfold, J. Determination of Micelle Structure and Charge by Neutron Small Angle Scattering. *Colloid Polym. Sci.* **1983**, *261*, 1022-1030.
- (43) Markovic, I.; Ottewill, R. H.; Cebula, D. J.; Field, I.; Marsh, J. Small Angle Neutron Scattering Studies on Non-Aqueous Dispersions of Calcium Carbonate. Part 1. The Guinier Approach. *Colloid Polym. Sci.* **1984**, *262*, 648-656.
- (44) Almgren, M.; Jóhannsson, R.; Eriksson, J. C. Polydispersity of AOT Droplets Measured by Time-Resolved Fluorescence Quenching. *J. Phys. Chem.* **1993**, *97*, 8590-8594.

- (45) Figueiredo Neto, A. M.; Salinas, S. R. A. *The Physics of lyotropic Liquid Crystals, Phase Transition and Structural Properties*; Oxford University Press; New York, 2005.
- (46) Roux, D.; Safinya, C. R. A Synchrotron X-Ray Study of Competing Undulation and Electrostatic Interlayer Interactions in Fluid Multimembrane Lyotropic Phases *J. Phys.* **1988**, *49*, 307-318.
- (47) Guinier, A.; Fournet, G. *Small-Angle Scattering of X-Rays*, Wiley, New York, 1956.
- (48) Lee, C. T.; Psathas, P. A.; Ziegler, K. J.; Johnston, K. P.; Dai, H. J.; Cochran, H. D.; Melnichenko, Y. B.; Wignall, G. D. Formation of Water-in-Carbon Dioxide Microemulsions with a Cationic Surfactant: a Small-Angle Neutron Scattering Study. *J. Phys. Chem. B* **2000**, *104*, 11094-11102.
- (49) Gopalan, A. S.; Wai, C. M.; Jacobs, H. K.; eds., *Supercritical Carbon Dioxide; Separations and Processes*, American Chemical Society, Washington, DC, 2003.
- (50) Porod, G. Die Röntgenkleinwinkelstreuung von dichtgepackten kolloiden Systemen. *Kolloid-Zeitschrift* **1951**, *124*, 83-114.
- (51) Sagisaka, M.; Hino, M.; Oasa, J.; Yamamoto, M.; Yoda, S.; Takebayashi, Y.; Furuya, T.; Yoshizawa, A.; Ochi, K.; Otake, K. Characterization of Water/Supercritical CO₂ Microemulsion by UV-Visible Spectroscopy and Dynamic Light Scattering. *J. Oleo Sci.* **2009**, *58*, 75-83.
- (52) Sagisaka, M.; Fujii, T.; Ozaki, Y.; Yoda, S.; Takebayashi, Y.; Kondo, Y.; Yoshino, N.; Sakai, H.; Abe, M.; Otake, K. Interfacial Properties of Branch-Tailed Fluorinated Surfactants Yielding a Water/Supercritical CO₂ microemulsion. *Langmuir* **2004**, *20*, 2560-2566.
- (53) Antelmi, D. A.; Kikicheff, P. Measurement of the Compressibility Modulus in a Lyotropic Lamellar Phase Stabilized by Undulation Forces. *J. Phys. Chem. B* **1997**, *101*, 8169-8179.
- (54) Izquierdo, C.; Moyá, M. L.; Usero, J. L.; Casado, J. Estimation of the Aggregation Number and Core Radius of Microemulsions. *Monatsh. Chem.* **1992**, *123*, 383-389.

- (55) Kawai, T.; Hamada, K.; Kon-No, K. Formation of AOT Reversed Micelles and W/O Microemulsions. *Bull. Chem. Soc. Jpn.*, **1992**, *65*, 2715-2719.
- (56) Nave, S.; Eastoe, J.; Heenan, R. K.; Steytler, D.; Grillo, I. What Is So Special About Aerosol-OT? Part III – Glutaconate versus Sulfosuccinate Headgroups and Oil-Water Interfacial Tensions. *Langmuir* **2002**, *18*, 1505-1510.
- (57) Kazarian, S. G.; Vincent, M. F.; Bright, F. V.; Liotta, C. L.; Eckert, C. A. Specific Intermolecular Interaction of Carbon Dioxide with Polymers. *J. Am. Chem. Soc.* **1996**, *118*, 1729-1736.
- (58) Zhou, G.-W.; Li, G.-Z.; Chen, W.-J. Fourier Transform Infrared Investigation on Water States and the Conformations of Aerosol-OT in Reverse Microemulsions. *Langmuir* **2002**, *18*, 4566-4571.
- (59) Lee Jr., C. T.; Bhargava, P.; Johnston, K. P. Percolation in Concentrated Water-in-Carbon Dioxide Microemulsions. *J. Phys. Chem. B* **2000**, *104*, 4448-4456.
- (60) Holderer, O.; Klostermann, M.; Monkenbusch, M.; Schweins, R.; Lindner, P.; Strey, R.; Richter, D.; Sottmann, T. Soft Fluctuating Surfactant Membranes in Supercritical CO₂-Microemulsions. *Phys. Chem. Chem. Phys.* **2011**, *13*, 3022-3025.