Mohamed, M. E., Pahirulzaman, K. A. K., & Lazarus, C. M. (2016). Production of 3-Oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic Acid in the Fungus *Aspergillus oryzae*: A Step Towards Heterologous Production of Pyrethrins in Fungi. *Molecular Biotechnology*, 58(3), 172-178. <https://doi.org/10.1007/s12033-015-9911-0>

Peer reviewed version

Link to published version (if available):
[10.1007/s12033-015-9911-0](https://doi.org/10.1007/s12033-015-9911-0)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Humana Press at <http://dx.doi.org/10.1007/s12033-015-9911-0>.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Production of 3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid in the fungus

***Aspergillus oryzae*: a step towards heterologous production of pyrethrins in fungi**

Abstract

Pyrethrins are natural insecticides, which accumulate to high concentrations in pyrethrum (*Chrysanthemum cinerariaefolium*) flowers. Synthetic pyrethroids are more stable, more efficacious and cheaper, but contemporary requirements for safe and environmentally friendly pesticides encourage a return to the use of natural pyrethrins, and this would be favoured by development of an efficient route to their production by microbial fermentation. The biosynthesis of pyrethrins involves ester linkage between an acid moiety (chrysanthemoyl or pyrethroyl, synthesized via the mevalonic acid pathway from glucose), and an alcohol (pyrethrolone). Pyrethrolone is generated from 3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid, which originates from α -linolenic acid via the jasmonic acid biosynthetic cascade. The first four genes in this cascade, encoding lipoxygenase 2, allene-oxide synthase, allene-oxide cyclase 2 and 12-oxophytodienoic acid reductase 3, were amplified from an *Arabidopsis thaliana* cDNA library, cloned in a purpose-built fungal multigene expression vector and expressed in *Aspergillus oryzae*. HPLC-MS analysis of the transgenic fungus homogenate gave good evidence for the presence of 3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid.

Keywords

3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid; *Aspergillus oryzae*; Biosynthesis; Heterologous gene expression; Pyrethrins

Introduction

Pyrethrum (*Chrysanthemum cinerariaefolium*) flowers, and the pyrethrins isolated from them, have been used successfully as household insecticides for the last 150 years. Natural pyrethrins, however, are expensive as well as photosensitive, and their production affects ecosystems by occupying agricultural landscapes (1). It is not surprising, therefore, that the natural compounds have been substantially replaced by synthetic pyrethroids, which are more stable, have greater insecticidal efficacy and are cheaper to produce (2), but recent requirements for safe and environmentally friendly pesticides encourage the re-adoption of natural pyrethrins (3). This is unlikely to happen to any extent if production costs remain high, so development of a cheaper production platform, such as microbial fermentation, would be highly desirable.

Filamentous fungi are seen as attractive heterologous production hosts due to their ability to synthesise large amounts of protein, and to perform post-translational modifications on recombinant proteins. The most commonly used filamentous fungal platforms are *Aspergillus niger*, *Aspergillus oryzae* and *Trichoderma reesei* (4, 5). *A. oryzae* has been used in the food and fermentation industry for several hundred years, and has GRAS (Generally Regarded As Safe) status (6). Its use as a heterologous expression system for recombinant protein production dates back to 1988, and more recently it has been used for the heterologous production of secondary metabolites (6-8). A useful feature of *A. oryzae* is that its genome has been sequenced (9, 10), and established methods for its genetic manipulation have been highly refined (11, 12). Over a decade of improvements to the *A. oryzae* expression system have made it a versatile and high yielding platform for protein and secondary metabolite production (6, 8, 12, 13). It is currently used in the production of biopharmaceuticals such as antibodies and antimicrobial peptides (14-16), and as a research tool to elucidate biosynthetic

pathways of secondary metabolites such as polyketides (17, 18). It has the potential to be used for cost-effective production of pyrethrins by fermentation.

Natural pyrethrins include six major compounds: pyrethrin I and II, cinerin I and II, and jasmolin I and II (Figure 1A) (2). Their biosynthesis involves ester linkage between an acid moiety (chrysanthemoyl or pyrethroyl) and an alcohol (pyrethrolone) (Figure 1B) (19, 20).

Chrysanthemoyl and pyrethroyl moieties are both synthesized via the mevalonic acid pathway from glucose, while pyrethrolone is generated from 3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid (OPC8:0) originating from α -linolenic (ALA) acid via the jasmonic acid biosynthetic cascade (Figure 2) (21).

[Place Figure 1]

[Place Figure 2]

This work reports on the ability of transformed cultures of *A. oryzae* to produce OPC8:0, which represents an important step in production of pyrethrolone, the alcoholic moiety of pyrethrins. Production of this pyrethrolone precursor was achieved using genes from the jasmonate biosynthesis cascade in *Arabidopsis thaliana*.

Materials and Methods

Gene isolation from *Arabidopsis thaliana*

The coding regions of the first four genes in the jasmonic acid biosynthetic cascade were amplified from an *Arabidopsis thaliana* cDNA library using Phusion High Fidelity DNA Polymerase (Thermo Scientific). The genes encoding Lipoxxygenase 2 (ATLOX2), Allene-oxide synthase (ATAOS), Allene-oxide cyclase 2 (ATAOC2) and 12-oxophytodienoic acid reductase 3 (ATOPR3) (Genbank accession numbers NM_114383, X92510, NM_113476, NM_126619, respectively) were amplified using the primers shown in Table 1 and directionally cloned in pENTR/D-TOPO (Life Technologies). DNA sequencing confirmed that all cloned PCR products were mutation free.

Table 1. Primers used in this work. Those with names commencing with "FUN" have 5' extensions corresponding to the cut ends of the vector (in green) to facilitate cloning of PCR products by heterologous recombination in yeast. Underlining indicates the start or (reverse complement of) stop codons of the coding regions amplified.

Primer Name	Primer Sequence
ATLOX2For	5' CACC <u>ATGT</u> ATTGTAGAGAGTCCTTGTC 3'
ATLOX2Rev	5' AGA <u>TCAA</u> ATAGAAATACTATAAGG 3'
ATAOSFor	5' CACC <u>ATGG</u> CTTCTATTTCAACCCCTTTT 3'
ATAOSRev	5' CTC <u>CTA</u> AAAGCTAGCTTTCCTTAAC 3'
ATAOC2For	5' CACC <u>ATGG</u> CTTCTTCAGCAGTGTC 3'
ATAOC2Rev	5' GAA <u>TTAG</u> TTGGTATAGTTACTTAT 3'
ATOPR3For	CACC <u>ATG</u> ACGGCGGCACAAGGGAAC
ATOPR3Rev	CAA <u>TCAG</u> AGGCGGGAAAAAGGAGCC
FUNATLOX2For	5' TACCCAAGCGCTTCGATTAGGAAGTAACCCACC <u>ATGT</u> ATTGTAGAGAGTCCTTGTC 3'
FUNATLOX2Rev	5' TCAGTGTCGAAAGATCCACTAGAGTAAATCAGAT <u>TCAA</u> ATAGAAATACTATAAGG 3'
FUNATAOSFor	5' TAACAGCTACCCCGCTTGAGCAGACATCACACC <u>ATGG</u> CTTCTATTTCAACCCCTTTT 3'
FUNATAOSRev	5' GGGAATGCGGCTCCACAGCTGCAGGCGTAGCTC <u>CTA</u> AAAGCTAGCTTTCCTTAAC 3'
FUNATAOC2For	5' ATCTCAACTCCATCACATACAATCGATCCCACC <u>ATGG</u> CTTCTTCAGCAGTGTC 3'
FUNATAOC2Rev	5' ATTTCACACCGCATACGTCAAAGCAACCATGA <u>TTAG</u> TTGGTATAGTTACTTAT 3'

Fungal expression vector construction

This followed the strategy described in (12) essentially as follows. Multigene expression vector pTAYAGS3tgo (Figure 3A) was cut downstream of each of the three promoters using *AscI*, and three of the cloned coding regions (ATLOX2, ATAOS and ATAOC2) were amplified using primers with 5' extensions corresponding to the cut ends of the vector (Table 1). The digested vector and three PCR products were assembled into a single recombinant plasmid by homologous recombination in yeast strain YPH500 (Agilent Technologies). The resultant pTAYAGS3tgoLSC expression plasmid (Figure 3B) was shuttled to *E. coli* for confirmation and amplification and the ATOPR3 coding region was transferred into the *amyB* expression cassette by the Gateway LR recombination reaction (Life Technologies) to produce pTAYAtgoLSCO (Figure 3C).

[Place Figure 3]

Fungal transformation and fermentation

Transformation of *Aspergillus oryzae* M-2-3 with pTAYA-GS3tgoLSCO was carried out as previously described (13). Transformants were selected and sub-cultured on Czapek Dox Agar then on solid DPY medium (2% dextrin, 1% peptone, 0.5% yeast extract, 0.5% potassium dihydrogen orthophosphate, 0.05% magnesium sulphate, 1.5% agar) for further production of spores, which were used to inoculate 100 mL liquid cultures (DPY with or without supplementation with 1 mg α -linolenic acid). Cultures were shaken for 7 days at 200 rpm and 28 °C.

Metabolite extraction and analysis by LCMS

Liquid cultures were homogenised, acidified with 38% HCl to pH 3 and filtered. The supernatant was extracted twice with 70 ml ethyl acetate, and the organic phase dried with anhydrous MgSO₄ before evaporation to dryness under reduced pressure. The crude extract was dissolved in 10% aqueous methanol and defatted by extraction with hexane. The

methanolic layer was evaporated to dryness and the extract dissolved to a concentration of 10 mg/mL in HPLC grade methanol. Metabolites were analysed using a Waters Platform Liquid chromatography-mass spectrometry system comprising a Waters 600 pump system, Waters 996 diode array detector (detecting between 210 and 400 nm), Macromass Platform LC mass spectrometer (detecting between 150 and 600 m/z units ESI⁺) and Phenomenex Luna 5 μ C 18 (250 $\times$ 4.6 mm, 5 μ m) reverse phase column with water + 0.05% TFA (solution A) and acetonitrile (solution B) as the mobile phase. The injection volume was 30-50 μ l, and elution was done using a non-linear gradient, water/acetonitrile 95/5 % to 5/95 % (0 min, 5% B; 5 min, 5% B; 45 min, 75% B; 46 min, 95% B; 50 min, 95% B; 55 min, 5% B; 60 min, 5%B). Compounds were identified by reference to the Wiley Registry of Mass Spectral Data (10th edition, April 2013), the NIST 11 Mass Spectral Library (NIST11/2011/EPA/NIH) and literature data (22-24).

Results and Discussion

The coding regions of the first four genes required for the biosynthesis of the pyrethrolone moiety of pyrethrins were amplified from an *A. thaliana* cDNA library and cloned. LOX2, AOS, AOC2 and OPR3 are ubiquitous in all plants as the first four enzymes in jasmonate biosynthesis. The coding sequences were inserted into the four expression cassettes of the fungal multigene expression vector pTAYA-GS3tgoLSCO in a two-step process (Figure 3), and used to transform *A. oryzae*. Seven transformants bearing the 4-gene recombinant expression plasmid were obtained and genetically purified by serial sub-culturing on solid selective medium; two were then put into liquid culture for fermentation and extraction of metabolites. LCMS analysis of transgenic fungal extracts, monitored at ES^+ m/z 316 (M^+ + sodium), indicated the presence of a new peak eluting at 9.4 minutes in extracts of both transformants; the data for one transformant and an untransformed (control) culture extract are shown in Figure 4. Submitting the mass spectrum of the newly identified peak to a library search indicated the production of 3-oxo-2-(2'-pentenyl)-cyclopentane-1-octanoic acid (OPC-8:0), the expected product of the combined activities of all four transgenes. This finding was also confirmed by comparing the mass spectrum of the new peak with published literature (23, 24). Figure 4B shows that the transgenic lines were capable of producing OPC-8:0 in the absence of added ALA precursor, but supplementing the culture medium with ALA greatly enhanced production of OPC-8:0 (Figure 4C). Optimisation of the system could therefore include increasing the endogenous ALA content above the maximum reported level of about 3.7 mol % (25). Since linoleic acid accounts for about 60% of the fatty acid pool this could be achieved by over-expression of the native Δ^{15} desaturase or of a foreign enzyme such as that encoded by the *Arabidopsis FAD3* gene (26). No OPC-8:0 was apparent in the extract of the untransformed fungus culture, even on supplementation with ALA (Figure 4A).

[Place Figure 4]

Figure 2 shows that *cis*-jasmonone may also be made by an alternative route from 12-oxophytodienoic acid (OPDA), the immediate precursor of OPC-8:0. This route requires isomerization of OPDA followed by β -oxidation of the iso-OPDA product. While no plant OPDA isomerase enzyme has been identified, an insect glutathione S transferase has been described with this activity (27), and thus the possibility exists of investigating this alternative route by substituting the oxophytodienoic acid reductase (OPR) gene with one for OPDA isomerase. However the expected product would be much more difficult to detect. In contrast our deployment of OPR has yielded the easily detectable OPC-8:0, whose existence testifies to the prior production of the OPDA substrate, and goes a step further along the route to *cis*-jasmonone via the established route.

Successful expression of the four selected genes represents an important step towards heterologous production of pyrethrolone by microbial fermentation. The next step will involve the conversion of OPC-8:0 to *cis*-jasmonone or 7-hydroxy-jasmonic acid (19) by simple decarboxylation with β -oxidation resulting in chain shortening (28). However, while production of pyrethrolone from *cis*-jasmonone or 7-hydroxy-jasmonic acid should involve simple decarboxylation and terminal unsaturation, the enzyme(s) involved have not been identified. Different natural pyrethrins can be made via esterification between pyrethrolone and chrysanthemyl moieties, and it should be possible to produce a chrysanthemoyl moiety by heterologous expression of the chrysanthemyl diphosphate synthase gene isolated previously from the pyrethrum flower (29).

In summary, current global environmental sentiments favour the re-use of natural pyrethrin insecticides over their synthetic pyrethroid counterparts. Thus the low production level of pyrethrins in the natural host, the pyrethrum flower, encourages the idea of producing them heterologously in a fungal host using DNA recombination and metabolic engineering tools. The first four genes responsible for jasmonic acid biosynthesis in *A. thaliana* were expressed

in *A. oryzae* to OPC-8:0, a precursor of the alcoholic moiety of natural pyrethrins. The production of natural pyrethrins in a fungal platform could provide a commercially viable alternative source of these valuable pesticides and offers an approach to protect the environment and plant ecosystems.

Acknowledgment

Conflict of interests

The author declare no conflict of interests

References

1. Matsuo, N., Mori, T. and Matsuda, K. (2012) Pyrethrin Biosynthesis and Its Regulation in *Chrysanthemum cinerariaefolium*, in *Pyrethroids*, vol. 314: Topics in Current Chemistry, Springer Berlin Heidelberg: pp. 73-81.
2. Matsuda, K., Kikuta, Y., Haba, A., Nakayama, K., Katsuda, Y., Hatanaka, A. and Komai, K. (2005) Biosynthesis of pyrethrin I in seedlings of *Chrysanthemum cinerariaefolium*. *Phytochemistry* 66, 1529-1535.
3. Gorbacheva, N. A. and Orlova, A. M. (1999) [Synthetic pyrethroids: their toxicology and metabolism]. *Sud Med Ekspert* 44, 28-31 contd.
4. Nevalainen, K. M. H., Te'o, V. S. J. and Bergquist, P. L. (2005) Heterologous protein expression in filamentous fungi. *Trends in Biotechnology* 23, 468-474.
5. Iwashita, K. (2002) Recent studies of protein secretion by filamentous fungi. *Journal of Bioscience and Bioengineering* 94, 530-535.
6. Sakai, K., Kinoshita, H. and Nihira, T. (2012) Heterologous expression system in *Aspergillus oryzae* for fungal biosynthetic gene clusters of secondary metabolites. *Appl Microbiol Biotechnol* 93, 2011-2022.
7. Mabashi, Y., Kikuma, T., Maruyama, J., Arioka, M. and Kitamoto, K. (2006) Development of a versatile expression plasmid construction system for *Aspergillus oryzae* and its application to visualization of mitochondria. *Biosci Biotechnol Biochem* 70, 1882-1889.
8. Lubertozzi, D. and Keasling, J. D. (2009) Developing *Aspergillus* as a host for heterologous expression. *Biotechnology Advances* 27, 53-75.
9. Machida, M., Yamada, O. and Gomi, K. (2008) Genomics of *Aspergillus oryzae*: learning from the history of Koji mold and exploration of its future. *DNA Res* 15, 173-183.

10. Kobayashi, T., Abe, K., Asai, K., Gomi, K., Juvvadi, P. R., Kato, M., Kitamoto, K., Takeuchi, M. and Machida, M. (2007) Genomics of *Aspergillus oryzae*. *Biosci Biotechnol Biochem* 71, 646-670.
11. Unkles, S., Campbell, E., de Ruiter-Jacobs, Y. J. T., Broekhuijsen, M., Macro, J., Carrez, D., Contreras, R., van den Hondel, C. M. J. J. and Kinghorn, J. (1989) The development of a homologous transformation system for *Aspergillus oryzae* based on the nitrate assimilation pathway: A convenient and general selection system for filamentous fungal transformation. *Molecular and General Genetics MGG* 218, 99-104.
12. Pahirulzaman, K. A., Williams, K. and Lazarus, C. M. (2012) A toolkit for heterologous expression of metabolic pathways in *Aspergillus oryzae*. *Methods Enzymol* 517, 241-260.
13. Heneghan, M. N., Yakasai, A. A., Halo, L. M., Song, Z., Bailey, A. M., Simpson, T. J., Cox, R. J. and Lazarus, C. M. (2010) First heterologous reconstruction of a complete functional fungal biosynthetic multigene cluster. *Chembiochem* 11, 1508-1512.
14. Frenzel, A., Hust, M. and Schirrmann, T. (2013) Expression of recombinant antibodies. *Front Immunol* 4, 217.
15. Mygind, P. H., Fischer, R. L., Schnorr, K. M., Hansen, M. T., Sonksen, C. P., Ludvigsen, S., Raventos, D., Buskov, S., Christensen, B., De Maria, L., Taboureau, O., Yaver, D., Elvig-Jorgensen, S. G., Sorensen, M. V., Christensen, B. E., Kjaerulff, S., Frimodt-Moller, N., Lehrer, R. I., Zasloff, M. and Kristensen, H.-H. (2005) Plectasin is a peptide antibiotic with therapeutic potential from a saprophytic fungus. *Nature* 437, 975-980.
16. Punt, P. J., van Biezen, N., Conesa, A., Albers, A., Manguis, J. and van den Hondel, C. (2002) Filamentous fungi as cell factories for heterologous protein production. *Trends Biotechnol* 20, 200-206.

17. Chen, X. Z., Shen, W., Fan, Y. and Wang, Z. X. (2011) Genomics and metabolic engineering of filamentous fungi in the post-genomics era. *Yi Chuan* 33, 1067-1078.
18. Fisch, K. M., Bakeer, W., Yakasai, A. A., Song, Z., Pedrick, J., Wasil, Z., Bailey, A. M., Lazarus, C. M., Simpson, T. J. and Cox, R. J. (2011) Rational Domain Swaps Decipher Programming in Fungal Highly Reducing Polyketide Synthases and Resurrect an Extinct Metabolite. *Journal of the American Chemical Society* 133, 16635-16641.
19. Matsuda, K. (2012) Pyrethrin biosynthesis and its regulation in *Chrysanthemum cinerariaefolium*. *Top Curr Chem* 314, 73-81.
20. Ramirez, A. M., Saillard, N., Yang, T., Franssen, M. C., Bouwmeester, H. J. and Jongsma, M. A. (2013) Biosynthesis of sesquiterpene lactones in pyrethrum (*Tanacetum cinerariifolium*). *PLoS One* 8, e65030.
21. Delker, C., Stenzel, I., Hause, B., Miersch, O., Feussner, I. and Wasternack, C. (2006) Jasmonate biosynthesis in *Arabidopsis thaliana*--enzymes, products, regulation. *Plant Biol (Stuttg)* 8, 297-306.
22. Aina, T., Matsuura, M. and Kobayashi, Y. (2003) Efficient Total Synthesis of 12-oxo-PDA and OPC-8:0. *The Journal of Organic Chemistry* 68, 7825-7832.
23. Schaller, F. and Weiler, E. W. (1997) Molecular cloning and characterization of 12-oxophytodienoate reductase, an enzyme of the octadecanoid signaling pathway from *Arabidopsis thaliana*. Structural and functional relationship to yeast old yellow enzyme. *J Biol Chem* 272, 28066-28072.
24. Schaller, F. and Weiler, E. W. (1997) Enzymes of octadecanoid biosynthesis in plants--12-oxo-phytyldienoate 10,11-reductase. *European journal of biochemistry / FEBS* 245, 294-299.

25. Sakuradani, E. and Shimizu, S. (2003) Gene Cloning and Functional Analysis of a Second Δ^6 -Fatty Acid Desaturase from an Arachidonic Acid-producing *Mortierella* Fungus. *Bioscience, Biotechnology, and Biochemistry* 67, 704-711.
26. Shah, S., Xin, Z. and Browse, J. (1997) Overexpression of the FAD3 desaturase gene in a mutant of *Arabidopsis*. *Plant Physiol* 114, 1533-1539.
27. Dabrowska, P., Freitak, D., Vogel, H., Heckel, D. G. and Boland, W. (2009) The phytohormone precursor OPDA is isomerized in the insect gut by a single, specific glutathione transferase. *Proc Natl Acad Sci U S A* 106, 16304-16309.
28. Kienow, L., Schneider, K., Bartsch, M., Stuible, H. P., Weng, H., Miersch, O., Wasternack, C. and Kombrink, E. (2008) Jasmonates meet fatty acids: functional analysis of a new acyl-coenzyme A synthetase family from *Arabidopsis thaliana*. *Journal of experimental botany* 59, 403-419.
29. Rivera, S. B., Swedlund, B. D., King, G. J., Bell, R. N., Hussey, C. E., Jr., Shattuck-Eidens, D. M., Wrobel, W. M., Peiser, G. D. and Poulter, C. D. (2001) Chrysanthemyl diphosphate synthase: isolation of the gene and characterization of the recombinant non-head-to-tail monoterpene synthase from *Chrysanthemum cinerariaefolium*. *Proc Natl Acad Sci U S A* 98, 4373-4378.

List of Figures:

Figure 1: Natural pyrethrins. **A:** Names and chemical structures of various natural pyrethrins.

B: Biosynthesis of natural pyrethrins; the reaction involves esterification of an alcohol (pyretholone) with an acid (chrysanthemyl or pyrethroyl moieties).

Figure 2: *cis*-Jasmone biosynthetic cascade from α -linolenic acid. The pathway utilises the activities of four enzymes followed by a series of β -oxidations and dehydration to produce *cis*-jasmone. The alternative iso-OPDA pathway is also illustrated.

Figure 3: Fungal expression vector construction. pTAYAGS3tgo (**A**) was cut downstream of each of the three promoters using *Asc*I, and three amplified coding regions (ATLOX2, ATAOS and ATAOC2) inserted by homologous recombination in yeast to produce pTAYAGS3tgoLSC (**B**). The cloned ATOPR3 coding region was transferred into the *amyB* expression cassette of pTAYAGS3tgoLSC by Gateway LR recombination to produce pTAYAtgoLSCO (**C**).

Figure 4: LCMS analysis of wild-type and transgenic *A. oryzae* culture extracts. The trace was monitored at ES^+ m/z 316. (**A**) Wild-type (untransformed) *A. oryzae* culture extract with added α -linolenic acid. (**B**) Representative *A. oryzae* transformant cultured in the absence of exogenous α -linolenic acid. (**C**) Representative *A. oryzae* transformant cultured with exogenous α -linolenic acid, added as a precursor. The scales of the LCMS traces (boxed and written at larger scale) are different indicating the enhancing effect of exogenous α -linolenic acid. The new peak produced (at 9.4 minutes) is indicated by the dotted box on the trace.

Figure 1

Figure 2

Figure 3

Figure 4