Carswell, A. M., Hill, P., Jones, D., Blackwell, M., Johnes, P. J., & Chadwick, D. (2016). Short-term biotic removal of dissolved organic nitrogen (DON) compounds from soil solution and subsequent mineralisation in contrasting grassland soils. *Soil Biology and Biochemistry*, 96, 82-85. <https://doi.org/10.1016/j.soilbio.2016.01.017>

Peer reviewed version

Link to published version (if available):
[10.1016/j.soilbio.2016.01.017](https://doi.org/10.1016/j.soilbio.2016.01.017)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <http://www.sciencedirect.com/science/article/pii/S0038071716000328>.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Short-term biotic removal of dissolved organic nitrogen (DON) compounds from soil solution and subsequent mineralisation in contrasting grassland soils

A.M. Carswell^{a*}, P.W. Hill^b, D.L. Jones^b, M.S.A. Blackwell^c, P. Johnes^a, D.R. Chadwick^b

^a *School of Geographical Sciences, University of Bristol, University Road, Bristol BS8 1SS, UK*

^b *School of the Environment, Natural Resources & Geography, Bangor University, Gwynedd, LL57 2UW, UK*

^c *Rothamsted Research, Okehampton EX20 2SB, Devon, UK*

Corresponding author: Tel +44 117 928 8808; fax: +44 117 928 7878

e-mail address: alison.carswell@bristol.ac.uk (A.M. Carswell)

ABSTRACT

Cycling of low molecular weight dissolved organic nitrogen compounds constitutes an important component of soil organic matter turnover in soils. We determined how rapidly grassland soils can cycle urea, compared to the amino acid L-alanine, and the peptide L-trialanine. Using naturally occurring concentrations of ^{14}C -labelled compounds the rates of removal from soil solution and subsequent mineralisation were measured. Biotic removal of all three compounds and subsequent mineralisation to CO_2 occurred within minutes. This research has demonstrated, for the first time, the potential for rapid removal of urea at low concentrations by the soil microbial biomass.

Keywords: Dissolved organic matter, Nutrient cycling, Urea, Urine patch

Adverse ecosystem effects of excess nitrogen (N) have been observed globally (Vitousek et al., 2009). Excess N in grasslands, prone to leaching and N emissions, is typically derived from N amendments including organic manures, urine patches and excessive use of synthetic fertilisers. Of these, urea has frequently been examined due to its importance as a fertiliser (IFA, 2014), and its presence in manures and urine (Ball and Ryden, 1984).

Urea is a low molecular weight dissolved organic N (LMW-DON) compound with a C:N molar ratio of 1:2.33, and similar to nitrate and ammonium, is capable of being taken up directly by both plants and microorganisms (Berman and Bronk, 2003; Wang et al., 2008). The extent to which plants can acquire LMW-DON and the degree to which it leaches down the soil profile, is critically dependent on the activity of the soil microbial biomass (SMB; Jones et al., 2013). Recent studies indicate that uptake of LMW-DON by the SMB is frequently driven by carbon (C) rather than by N demand (Farrell et al., 2014). Therefore, the presence of C within urea may drive its rate of removal from grassland soil solutions. Although urease activity in soil (Nielsen et al., 1998; Bolado-Rodríguez et al., 2005), and to a lesser extent urea assimilation by the SMB (Smith et al., 2007) have been investigated, urea removal from the soil solution by the SMB over short time-scales has not.

Here SMB removal of ^{14}C -urea from the soil solution and its subsequent catabolic and anabolic partitioning was examined in three grassland soils. Microbial cycling of urea was directly compared to that of other typical LMW-DON compounds found in soils: the amino acid ^{14}C -L-alanine and the oligopeptide ^{14}C -L-trialanine, whose turnover have been extensively characterised and have also been implicated in direct plant LMW-DON acquisition (Hill et al., 2011; Wilkinson et al., 2014).

Soil was collected from three separate grazed grassland sites in the UK (Table 1). All soils were collected towards the end of the growing season (October), with three independent

replicates collected for each type. Soil cores (10 × 8.5 cm; h × i.d.) were kept intact, at field-moisture, in gas-permeable bags, in the dark at 4°C prior to use.

To characterise LWM-DON in each soil, porewater was obtained from intact soil cores, with the root mat removed, by centrifugation-drainage (Giesler and Lundström, 1993). Soluble N was determined as described by Farrell et al. (2013) and Sullivan and Havlin (1991). All experimentation with ^{14}C -labelled compounds was performed on < 2 mm sieved soil from separately taken soil samples, which had equilibrated to 20°C overnight. The rate of LMW-DON depletion from soil solution was measured according to Hill et al. (2008). Briefly, 1 g soil (dry weight equivalent; DW) was placed in a microcentrifuge tube with a hole pierced in the bottom. This tube was placed inside another microcentrifuge tube. 300 μl of either ^{14}C -labelled urea, L-alanine or L-trialanine (10 μM , 0.9 kBq mL^{-1}) was then applied to the soil surface and allowed to infiltrate the soil (< 2.5 sec, 20°C; associated soil water content increase to 45-52%). This concentration was chosen to reflect the urea and free amino acid concentrations naturally occurring within soil solution (Table 1). At 1, 5, 10, 30 and 60 min after substrate addition, the soil was centrifuged (4000 g, 1 min, 4°C; data presented for 0 min in Figures 1 and 2 are assumed) allowing the soil solution to pass to the lower microcentrifuge tube. The ^{14}C content of the recovered soil solution was determined after addition of Scintisafe3 scintillation cocktail (Fisher Scientific, Loughborough, UK) using a Wallac 1404 liquid scintillation counter (Perkin Elmer Life Sciences, Boston, MA). To assess the mineralisation rate of LMW-DON compounds, 1 g soil was placed in a glass tube through which air was passed before being transferred through 2 successive 0.1 M NaOH traps to capture evolved $^{14}\text{CO}_2$. At 1, 5, 10, 30 and 60 min after ^{14}C -labelled substrate addition (as above), NaOH was replaced and its ^{14}C content determined as above. To separate biotic (e.g. microbial, enzymatic) and abiotic (e.g. sorption) LMW-DON removal processes, the soil solution recovery experiment was also performed on sterilised soil (autoclaved at 121°C, 20 min). Recovery of

^{14}C -labelled compounds from the sterilised soil solutions was used to calculate the theoretical maximum ^{14}C -activity (Hill et al., 2008) that could be recovered following complete mixing of amended ^{14}C -labelled treatments with native soil solution. A two-way ANOVA was used to test for differences and interactions between soils and LMW-DON treatments.

Complete mixing of the applied treatments with native soil solution was not achieved, and after 60 min deviated between 101-153%. Greater than 100% recovery of ^{14}C -compounds was achieved at all incubation periods, thus demonstrating that no retention of ^{14}C -compounds occurred in the sterile soils (Wilkinson et al., 2014; see supplementary information for equations), consequently no evidence of abiotic loss pathways was observed. However autoclaving soils can increase the solubilisation of soil organic matter (SOM; Powlson and Jenkinson, 1976), which may block adsorption sites that would be available in the living soils. Nonetheless autoclaving has been found to be more effective than CHCl_3 fumigation or gamma irradiation at reducing viable cell numbers (Blankinship et al., 2014), and consequently was chosen as the optimal sterilisation method for this study.

All ^{14}C -labelled LMW-DON compounds were rapidly removed from the soil solution (Fig. 1). After 60 min, removal of ^{14}C -L-alanine and ^{14}C -L-trialanine was almost complete, at 98.7 and 99.5% respectively. However, removal of ^{14}C -urea was consistently lower at all incubation periods, and after 60 min was 88.7%. Removal from soil solution followed the series: alanine > trialanine > urea ($p < 0.001$). Across all soils, the half-life of urea, alanine and trialanine in solution was 4.15 ± 0.69 , 0.30 ± 0.04 , 0.94 ± 0.13 min (mean $\pm$ SEM; based on fitting first order single exponential decay to the data), respectively. In contrast, no effect of soils was observed on substrate depletion. This is perhaps unsurprising as cycling of key LMW-DON compounds can be remarkably similar across diverse soils and systems (Jones et al., 2009). Slower uptake of urea relative to the other compounds could be linked to lower transporter expression and affinity within the SMB. It has also been proposed that soil

microorganisms exist in a C-starved state (Hobbie and Hobbie, 2013). Accordingly, in the soils examined here the SMB may have exhibited a preference for compounds with a greater C content and which are commonly present in soil solution (amino acids and peptides are present in rhizodeposits and via protease action on SOM).

In contrast to removal from soil solution, soil affected ($p < 0.001$) the mineralisation of different compounds at 1, 5, 10 and 30 min, but not after 60 min (Fig. 2). The differences between soils may, in part, be due to differences in initial soil water content, which can impact net mineralisation rates (Paul et al., 2003). However, the experiment was performed on field moist soils to represent the same preceding climatic conditions rather than target a specific soil water content (Table 1). ^{14}C -urea had the highest mineralisation rates of the three LMW-DON compounds. This may be attributed to the alternative mineralisation pathway urea can take via the enzyme urease, which is encountered both intra- and extracellularly. Extracellular urease has been shown to account for an average of 46% of total urease activity in a range of soils (Klose and Tabatabai, 1999). This may account for the more rapid mineralisation of urea relative to alanine and trialanine. Although ^{14}C -urea was most rapidly mineralised of the three compounds, only 40-45% of ^{14}C -urea removed from the soil solution was subsequently respired as $^{14}\text{CO}_2$ over 60 min, suggesting that the remaining ^{14}C -urea was assimilated by the SMB (Nielsen et al., 2008). Another intracellular urea pathway is via ATP:urea amidolyase (Cheng et al., 2005; Strope et al., 2011), which produces NH_3 and HCO_3^- via two enzyme (urea carboxylase and allophanate hydrolase) reactions, making it a likely path for urea assimilation in soils.

Although it is widely acknowledged that urea is rapidly mineralised in soils, this is the first time that such rapid removal of urea from the soil solution by the SMB has been reported. This suggests that there will be strong microbial competition for urea in soil which may limit its capture by plant roots when present in low concentrations. The fate of urea-derived NH_4^+

requires further investigation. Assimilation of urea by the SMB at higher concentrations of urea and following application of solid urea as fertiliser requires further work.

Acknowledgements

This work was funded by the UK Natural Environment Research Council (NE/K01093X/1), and the Lawes Trust/University of Reading Studentship scheme.

References

- Ball, P.R., Ryden, J.C., 1984. Nitrogen relationships in intensively managed temperate grasslands. *Plant and Soil* 76, 23-33.
- Berman, T., Bronk, D.A., 2003. Dissolved organic nitrogen: a dynamic participant in aquatic ecosystems. *Aquatic Microbial Ecology* 31, 279-305.
- Blankinship, J.C., Becerra, C.A., Schaeffer, S.M., Schimel, J.P., 2014. Separating cellular metabolism from exoenzyme activity in soil organic matter decomposition. *Soil Biology and Biochemistry* 71, 68-75.
- Bolado-Rodriguez, S., Alonso-Gaite, A., Alvarez-Benedi, J., 2005. Characterization of nitrogen transformations, sorption and volatilization processes in urea fertilized soils. *Vadose Zone Journal* 4, 329-336.
- Cheng, G., Shapir, N., Sadowsky, M.J., Wackett, L.P., 2005. Allophanate hydrolase, not urease, functions in bacterial cyanuric acid metabolism. *Applied and Environmental Microbiology* 71, 4437-4445.
- Farrell, M., Hill, P.W., Farrar, J., DeLuca, T.H., Roberts, P., Kielland, K., Dahlgren, R., Murphy, D.V., Hobbs, P.J., Bardgett, R.D., Jones, D.L., 2013. Oligopeptides represent a preferred source of organic n uptake: a global phenomenon? *Ecosystems* 16, 133-145.

- Farrell, M., Prendergast-Miller, M., Jones, D.L., Hill, P.W., Condron, L.M., 2014. Soil microbial organic nitrogen uptake is regulated by carbon availability. *Soil Biology & Biochemistry* 77, 261-267.
- Giesler, R., Lundstrom, U., 1993. Soil solution chemistry - Effects of bulking soil samples. *Soil Science Society of America Journal* 57, 1283-1288.
- Harrod, T.R., Hogan, D.V., 2008. The soils of North Wyke and Rowden Revised edition of Harrod, T.R. (1981) Soils in Devon V: Sheet SS61 (Chumleigh) Soil Survey Record no. 70.
- Hill, P.W., Farrar, J.F., Jones, D.L., 2008. Decoupling of microbial glucose uptake and mineralization in soil. *Soil Biology & Biochemistry* 40, 616-624.
- Hill, P.W., Farrell, M., Jones, D.L., 2012. Bigger may be better in soil N cycling: Does rapid acquisition of small L-peptides by soil microbes dominate fluxes of protein-derived N in soil? *Soil Biology & Biochemistry* 48, 106-112.
- Hill, P.W., Quilliam, R.S., DeLuca, T.H., Farrar, J., Farrell, M., Roberts, P., Newsham, K.K., Hopkins, D.W., Bardgett, R.D., Jones, D.L., 2011. Acquisition and assimilation of nitrogen as peptide-bound and D-enantiomers of amino acids by wheat. *PLoS ONE* 6, 1-4.
- Hobbie, J.E., Hobbie, E.A., 2013. Microbes in nature are limited by carbon and energy: the starving-survival lifestyle in soil and consequences for estimating microbial rates. *Frontiers in Microbiology* 4, 324.
- IFA., 2014. Production and international trade, IFA Statistics, 2014, online.
- Jones, D.L., Clode, P.L., Kilburn, M.R., Stockdale, E.A., Murphy, D.V., 2013. Competition between plant and bacterial cells at the microscale regulates the dynamics of nitrogen acquisition in wheat (*Triticum aestivum*). *New Phytologist* 200, 796-807.

- Jones, D.L., Kielland, K., Sinclair, F.L., Dahlgren, R.A., Newsham, K.K., Farrar, J.F.,
Murphy, D.V., 2009. Soil organic nitrogen mineralization across a global latitudinal
gradient. *Global Biogeochemical Cycles* 23, GB1016.
- Nielsen, T.H., Bonde, T.A., Sorensen, J., 1998. Significance of microbial urea turnover in N
cycling of three Danish agricultural soils. *FEMS Microbiology Ecology* 25, 147-157.
- Paul, K.I., Polglase, P.J., O'Connell, A.M., Carlyle, J.C., Smethurst, P.J., Khanna, P.K.,
2003. Defining the relation between soil water content and net nitrogen mineralization.
European Journal of Soil Science 54, 39-47.
- Powlson, D.S., Jenkinson, D.S., 1976. Effects of biocidal treatments on metabolism in soil. 2.
Gamma-irradiation, autoclaving, air-drying and fumigation. *Soil Biology and
Biochemistry* 8, 179-188.
- Smith, J.L., Bell, J.M., Bolton, H., Bailey, V.L., 2007. The initial rate of C substrate
utilization and longer-term soil C storage. *Biology and Fertility of Soils* 44, 315-320.
- Strope, P., Nickerson, K., Harris, S., Moriyama, E., 2011. Molecular evolution of urea
amidolyase and urea carboxylase in fungi. *BMC Evolutionary Biology* 11, 1-15.
- Sullivan, D.M., Havlin, J.L., 1991. Flow injection analysis of urea nitrogen in soil extracts.
Soil Science Society of America Journal 55, 109-113.
- Tabatabai, M.A., 1994. Soil enzymes, In: Weaver, R.W., Angel, G.S., Bottomley, P.S. (eds),
Methods of Soil Analysis, part 2: Microbiological and Biochemical Properties,
Madison, USA, Soil Science Society of America.
- Vitousek, P.M., Naylor, R., Crews, T., David, M.B., Drinkwater, L.E., Holland, E., Johnes,
P.J., Katzenberger, J., Martinelli, L.A., Matson, P.A., Nziguheba, G., Ojima, D., Palm,
C.A., Robertson, G.P., Sanchez, P.A., Townsend, A.R., Zhang, F.S., 2009. Nutrient
imbalances: Pollution remains response. *Science* 326, 665-666.

- Wang, W.H., Kohler, B., Cao, F.Q., Liu, L.H., 2008. Molecular and physiological aspects of urea transport in higher plants. *Plant Science* 175, 467-477.
- Wilkinson, A., Hill, P.W., Farrar, J.F., Jones, D.L., Bardgett, R.D., 2014. Rapid microbial uptake and mineralization of amino acids and peptides along a grassland productivity gradient. *Soil Biology and Biochemistry* 72, 75-83.

Table 1 Characteristics of the three grassland soils (upper 10 cm) used in the study. Values represent mean $\pm$ SEM ($n = 3$).

	Dystric Cambisol*	Stagni-vertic Cambisol*	Eutric Cambisol**
Sample location	50°46'N, 3°55'W	50°47'N, 3°57'W	53°14'N, 4°01'W
Soil texture	Loam*	Heavy clay*	Sandy clay loam**
Soil pH	5.2 $\pm$ 0.07	5.3 $\pm$ 0.07	4.8 $\pm$ 0.03
Total soil C (g kg ⁻¹ DW)	21.1 $\pm$ 0.04	29.1 $\pm$ 0.29	27.1 $\pm$ 0.19
Total soil N (g kg ⁻¹ DW)	3.6 $\pm$ 0.02	4.0 $\pm$ 0.02	4.1 $\pm$ 0.04
Soil water (kg kg ⁻¹ DW)	0.19 $\pm$ 0.01	0.35 $\pm$ 0.02	0.29 $\pm$ 0.00
Soil solution free amino acids (μ M)	11.3 $\pm$ 1.08	6.50 $\pm$ 1.18	8.00 $\pm$ 1.48
Soil solution short peptides (<1 kDa; μ M)	153 $\pm$ 47.3	145 $\pm$ 24.7	164 $\pm$ 52.0
Soil solution NO ₃ -N (mg N l ⁻¹)	10.6 $\pm$ 0.92	0.74 $\pm$ 0.23	2.22 $\pm$ 1.21
Soil solution urea (μ M)	6.22 $\pm$ 0.20	7.24 $\pm$ 0.62	15.33 $\pm$ 8.12
Soil solution NH ₄ -N (mg N l ⁻¹)	0.14 $\pm$ 0.02	0.30 $\pm$ 0.07	0.23 $\pm$ 0.05
Soil solution DOC (mg C l ⁻¹)	29.38 $\pm$ 3.25	31.80 $\pm$ 5.51	19.98 $\pm$ 2.77
Soil solution DON (mg N L ⁻¹)	1.94 $\pm$ 0.38	2.17 $\pm$ 0.38	1.54 $\pm$ 1.26
Soil respiration (mg C kg ⁻¹ dry soil h ⁻¹)	0.37 $\pm$ 0.05	0.86 $\pm$ 0.32	0.60 $\pm$ 0.14

Data gained from the literature are marked with either * (described by Harrod and Hogan, 2008) or ** (described by Hill et al., 2012).

Figure legends

Fig. 1. Microbially-mediated depletion of ^{14}C -labelled alanine, trialanine or urea from soil solution in three grassland soils. Data points represent means $\pm$ SEM ($n = 3$).

Fig. 2. Time-dependent cumulative mineralisation of ^{14}C -labelled alanine, trialanine or urea in three grassland soils. Data points represent means $\pm$ SEM ($n = 3$).