Cornthwaite, K., Jefferys, A., Lenguerrand, E., Haase, A., Lynch, M., Johnson, A., Draycott, T., & Siassakos, D. (2016). Pregnancy after weight loss surgery: A commentary. *BJOG: An International Journal of Obstetrics and Gynaecology*, 123(2), 165-170.
<https://doi.org/10.1111/1471-0528.13791>

Peer reviewed version

Link to published version (if available):
[10.1111/1471-0528.13791](https://doi.org/10.1111/1471-0528.13791)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at <http://dx.doi.org/10.1111/1471-0528.13791>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Pregnancy after weight loss surgery: A commentary

Katie Cornthwaite MBBS^{1,2}, Amanda Jefferys BMBS^{1,2}, Erik Lenguerrand PhD²,
Anne Haase PhD², Mary Lynch BSc¹, Andrew Johnson MD³, Tim Draycott MD^{1,2},
Dimitrios Siassakos MD^{1,2}

¹ Women's Health, North Bristol NHS Trust, Southmead Hospital, Bristol, BS10 5NB,
UK

² The University of Bristol, Senate House, Tyndall Avenue, Bristol, BS8 1TH, UK

³ Department of Medicine, North Bristol NHS Trust, Southmead Hospital, Bristol,
BS10 5NB, UK

Corresponding author:

Dr Katie Cornthwaite
Academic Centre for Women's Health,
Chilterns,
Southmead Hospital,
Westbury on Trym,
Bristol,
BS10 5NB,
UK
katiecornthwaite@gmail.com
0044 (0)117 4146835

28 The obesity epidemic poses increasing challenges to maternity services across the
29 UK and the world; almost one fifth of women of childbearing age in the UK are
30 classified obese (BMI ≥ 30) ¹. Optimisation of pre-pregnancy weight can improve
31 maternal and perinatal outcomes and greater numbers are resorting to bariatric
32 surgery pre-pregnancy to aid weight loss. Whilst the negative impact of obesity on
33 maternal and perinatal health is well-established ², there is a dearth of evidence
34 evaluating the effect of previous bariatric surgery on pregnancy. Given the likelihood
35 of rising rates of pregnancies following bariatric surgery, it is essential that both
36 patients and clinicians have evidence-based information to inform care. However, as
37 yet, there are very few robust clinical trials that have directly assessed pregnancy
38 outcomes following bariatric surgery. Consequently, the majority of our
39 understanding comes from case reports and observational studies (Table 1). This
40 commentary reviews recent and emerging evidence for the management of women
41 who are pregnant after bariatric surgery.

42
43 'Bariatric surgery' is an umbrella term that encompasses several different types of
44 weight loss procedures. In the UK, the most commonly performed procedures are
45 Laparoscopic Adjustable Gastric Banding (LAGB), Roux-en-Y gastric bypass, and
46 sleeve gastrectomy ³ (Table 2). Broadly, bariatric procedures can be categorised as
47 malabsorptive, restrictive or a combination of both. Malabsorptive procedures bypass
48 an absorptive part of the gastrointestinal tract, whereas restrictive procedures reduce
49 food intake by limiting gastric volume. Each type of bariatric surgery has specific
50 procedure related metabolic effects and complications. Nutritional deficiencies are
51 more likely after malabsorptive surgery ⁴. It is axiomatic that each method will have
52 specific effects on pregnancy, maternal outcomes and perinatal health.

53
54 Therefore it is unhelpful that most published studies group all bariatric procedures
55 together as a single intervention. There are other problems with the existing data too:

understandably for a previously uncommon intervention many studies to date have necessarily small sample sizes. To improve statistical power a great deal of the research has taken place over a protracted number of years, which makes firm conclusions difficult as there are likely to have been temporal changes in clinical practice and policy. Moreover, many studies have not accounted for important confounders. Finally, there is often a lack of adjustment for socio-demographics factors such as smoking, parity, and age. Other studies have compared outcomes in the same woman in pregnancies before and after bariatric surgery. This approach does not completely disentangle the impact of weight loss per se, from the complex effects of bariatric surgery and is also prone to confounding.

However, some broad conclusions can be drawn from the available data: after bariatric surgery ⁴⁻⁶ there is reduction in obesity-related maternal complications such as gestational diabetes and hypertension. However, the risk remains higher than for non-obese women⁷.

Improvements in some maternal outcomes notwithstanding, there have been concerns raised regarding the negative impact of bariatric surgery on perinatal outcomes, including small for gestational age infants and pre-term birth ^{4,8}. There has also been a suggestion of an increased risk of perinatal mortality ⁶. Moreover, very few studies have exclusively explored the impact of gastric banding on pregnancy, and there has been no randomized controlled trial comparing the management (deflation versus inflation) of the gastric band in pregnancy ⁹.

In addition to general risks, there are other specific complications including maternal intestinal hernias and nutritional deficiencies, as well as a greater risk of birth defects amongst pregnancies following gastric bypass surgery ⁴.

84 We recently completed a national cohort study of Laparoscopic Adjustable Gastric
85 Banding (LAGB) in pregnancy ^{10, 11} to understand the procedure related impact of
86 LAGB on pregnancy outcomes. This study is currently awaiting publication. Data
87 were collected on all pregnancies following LAGB in the UK over a discrete time
88 period using an established surveillance system (UK Obstetric Surveillance System –
89 UKOSS). Uniquely, the study compared outcomes according to index band
90 management (maintenance of band inflation during pregnancy versus band deflation
91 for some or all of pregnancy) and adjusted for important confounding or modifying
92 factors. Outcomes were compared with controls without a gastric band, and national
93 data, allowing comparison with the background population.

94
95 The study ^{10, 11} identified significant variation in band management across the UK,
96 possibly related to the current paucity of evidence to inform decision-making for
97 these women during pregnancy. In this cohort study, inflation of the gastric band was
98 related to a reduction in gestational weight gain, and an associated improvement of
99 some maternal outcomes.

100
101 However, our findings suggest that these maternal benefits may be at the expense of
102 perinatal health. Moreover, some maternal outcomes such as gestational diabetes
103 and caesarean birth were consistently worse in those with a gastric band compared
104 to normal BMI controls, regardless of band management.

105
106 On the basis of our findings, we would not recommend a one-size-fits-all policy
107 during pregnancy, but band adjustments should be made on a case-by-case basis.
108 Irrespective of the band management, pregnancies after bariatric surgery remain
109 high risk and it is essential that women are closely monitored. A multi-professional
110 approach with close involvement of endocrinologists and nutritionists, monitoring of

gestational complications such as diabetes and hypertension, as well as regular fetal growth scans are recommended.

However, clinical judgment alone should not dictate management. It is important for professionals to involve women in decision-making, enabling them to remain in control of their pregnancy and thereby optimising pregnancy and birth experience ¹².

A woman's preference regarding band management is likely to be multifactorial.

Decision-making may not rest solely on the balance of the maternal and perinatal outcomes, but may have a more complex psychological element. Obesity has been associated with high rates of depression and poor body image, and bariatric surgery can lead to improvements in mental health ¹³. However, it is unclear how pregnancy may influence the psychological wellbeing and mood of women with a gastric band, and what psychosocial support, if any, these women may require during pregnancy.

The ENGAGE (ENquiry into women with GAstric banding in pregnancy to Guide management and improve Experience) study aims to explore women's experience of pregnancy following LAGB and assess which factors drive a woman's decision regarding management of the band during pregnancy ¹⁴. Most women interviewed thus far had their bands inserted in the private sector. Preliminary analysis suggests limited information, guidance and support in pregnancy, and need for clarity regarding band management during pregnancy.

A national prospective cohort study, using the same UKOSS methodology, is currently ongoing to better understand the effects of gastric bypass surgery on pregnancy ¹⁵. All women who are pregnant following gastric bypass surgery are included. Epidemiology, management and outcomes, will be evaluated to assess current practice. The findings will help to inform obstetricians, endocrinologists, nutritionists, and bariatric surgeons for pre-pregnancy counseling, suitable monitoring and appropriate nutritional supplementation.

139

140 Furthermore, there is a paucity of data regarding how long to delay pregnancy
141 following weight loss surgery. American guidelines recommend a surgery to
142 conception interval of at least one year ¹⁶ however, there is no firm evidence to
143 support such advice ¹⁷. Data from the UKOSS gastric banding and bypass studies
144 are currently being analysed to investigate the impact of time to conception on
145 maternal and fetal outcomes; the results will be informative.

146

147 In conclusion, the rates of bariatric surgery in women are likely to increase further,
148 but there are very few data available to guide clinicians. Much of the evidence is from
149 registry-based studies and case reports. Furthermore, there has been a tendency to
150 group all bariatric procedures together in studies evaluating associated pregnancy
151 outcomes, despite clear mechanistic differences as well as different procedure-
152 related effects on pregnancy. However, there is emerging evidence that will inform
153 both the understanding of different types of bariatric surgery, and the effect of
154 management options on patient outcome and experience.

155

156 In the meantime, we would advocate multi-professional care for all women who have
157 undergone bariatric surgery (Table 3). Ideally, these women should receive pre-
158 pregnancy counseling; until further advice is available, we would suggest that women
159 delay pregnancy for one year following surgery. Women should be screened and
160 treated for nutritional deficiencies prior to conception ¹⁸ and receive prompt specialist
161 dietary advice during early pregnancy. They should be advised about the associated
162 maternal and fetal risks antenatally. We would recommend low dose aspirin and
163 vitamin supplementation; close blood pressure and urinalysis surveillance; and
164 universal testing of gestational diabetes. Importantly, fetal wellbeing should be
165 carefully monitored with serial growth scans. Clinicians may consider the role of

induction at term to reduce the risk of placental dysfunction and intrauterine fetal death.

Clearly, the findings from ongoing studies will refine these recommendations and provide the clarity clinicians need, and women want, to make appropriate decisions together.

Insert Table 3

Acknowledgements

None

Disclosure of Interests

None

Contribution to Authorship

Katie Cornthwaite conceived the idea, wrote the manuscript, is principal investigator of UKOSS gastric bypass study, and co-investigator of UKOSS gastric banding and ENGAGE studies.

Amanda Jefferys edited the manuscript and is principal investigator of UKOSS gastric banding and ENGAGE studies.

Erik Lenguerrand edited the manuscript and conducted the UKOSS gastric banding analyses.

Anne Haase edited the manuscript and is co-investigator of ENGAGE study.

Mary Lynch edited the manuscript and is co-investigator of ENGAGE study.

Andrew Johnson provided funding for the two UKOSS studies and approved the manuscript.

Tim Draycott edited and approved the manuscript.

Dimitrios Siassakos conceived the idea, co-wrote and edited the manuscript, and supervises the UKOSS gastric bypass, gastric banding and ENGAGE studies.

Ethics Approval

Data for UKOSS gastric banding (NRES 11/SW/0227), UKOSS gastric bypass (NRES 14/LO/0491), and ENGAGE studies (13/SW/0285) were collected with ethics approval

Funding

Bristol Bariatric Pregnancy Research Hub
North Bristol Trust Springboard Fund

References

1. National Obesity Observatory. Bariatric surgery for obesity: Department of Health, 2010. Available at <http://www.noo.org.uk>
2. Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW, et al. Maternal obesity and pregnancy outcome: a study of 287,213 pregnancies in London. *Int J Obes Relat Metab Disord* 2001;25:1175-82.
3. National Bariatric Surgery Registry Report. BOMSS, 2014. Available at http://www.bomss.org.uk/wp-content/uploads/2014/04/Extract_from_the_NBSR_2014_Report.pdf
4. Guelinckx I, Devlieger R, Vansant G. Reproductive outcome after bariatric surgery: a critical review. *Hum Reprod Update*. Oxford University Press; 2009 Mar 1;15(2):189–201.
5. Kjær MM, Nilas L. Pregnancy after bariatric surgery – a review of benefits and risks. *Acta Obstet Gynecol Scand*. Blackwell Publishing Ltd; 2013 Mar 1;92(3):264–71.
6. Johansson K, Cnattingius S, Näslund I, Roos N, Trolle Lagerros Y, Granath F, et al. Outcomes of Pregnancy after Bariatric Surgery. *N Engl J Med*. 2015 Feb 26;372(9):814–24.

- 220 7. Vrebosch L, Bel S, Vansant G, Guelinckx I, Devlieger R. Maternal and
221 Neonatal Outcome After Laparoscopic Adjustable Gastric Banding: a
222 Systematic Review. *Obes Surg*. Springer-Verlag; 2012;22(10):1568–79.
- 223 8. Roos N, Neovius M, Cnattingius S, Lagerros YT, Sääf M, Granath F, et al.
224 Perinatal outcomes after bariatric surgery: nationwide population based
225 matched cohort study. *BMJ*. BMJ Publishing Group Ltd; 2013 Nov
226 12;347:f6460.
- 227 9. Jefferys AE, Siassakos D, Draycott T, Akande VA, Fox R. Deflation of gastric
228 band balloon in pregnancy for improving outcomes. *Cochrane Database of*
229 *Systematic Reviews* 2013, Issue 4. Art. No.: CD010048. DOI:
230 10.1002/14651858.CD010048.pub2.
- 231 10. Cornthwaite K, Jefferys A, Lenguerrand E, Hyde J, Lynch M, Draycott T, et al.
232 One size does not fit all. Management of the laparoscopic adjustable gastric
233 band in pregnancy: a national prospective cohort study. Poster abstract.
234 *Lancet*, Elsevier; 2015 Feb 26;385:S32.
- 235 11. Cornthwaite K, Jefferys A, Lenguerrand E, Hyde J, Lynch M, Draycott T, et al.
236 Striking the balance between maternal and perinatal outcomes for pregnant
237 women with gastric banding: national cohort study (UKOSS). Oral
238 presentation. Annual academic meeting. RCOG. Jan 2015
- 239 12. Waldenström U, Hildingsson I, Rubertsson C, Rådestad I. A Negative Birth
240 Experience: Prevalence and Risk Factors in a National Sample. *Birth*.
241 Blackwell Science Inc; 2004 Mar 1;31(1):17–27.
- 242 13. Herpertz S, Kielmann R, Wolf AM, Langkafel M, Senf W, Hebebrand J. Does
243 obesity surgery improve psychosocial functioning? A systematic review.
244 *International Journal of Obesity*. Nature Publishing Group; 2003 Nov
245 1;27(11):1300–14.
- 246 14. ENGAGE. *ENquiry into women with GAstric banding in pregnancy to Guide*
247 *management and improve Experience*.

248 <http://www.bristol.ac.uk/sps/research/projects/current/engage/> (accessed 31
 249 April 2015)

250 15. UKOSS. *Pregnancy following gastric bypass surgery*.
 251 <https://www.npeu.ox.ac.uk/ukoss/current-surveillance/gby> (accessed 31 April
 252 2015)

253 16. Armstrong C. "ACOG guidelines on pregnancy after bariatric surgery." (2010):
 254 905-906.

255 17. Kjær MM, Nilas L. Timing of Pregnancy After Gastric Bypass—a National
 256 Register-Based Cohort Study. *Obes Surg*. Springer US; 2013;23(8):1281–5.

257 18. O’Kane M, Aasheim E, Barth J, Batterham R, Welbourn R. GP Guidance:
 258 Management of nutrition following bariatric surgery. British Obesity and
 259 Metabolic Surgery Society; Aug 2014

260

261