Bailey, P., Edwards, A., & Courtney, A. (2016). Living kidney donation. *BMJ*, (354), [i4746]. <https://doi.org/10.1136/bmj.i4746>

Publisher's PDF, also known as Version of record

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1136/bmj.i4746](https://doi.org/10.1136/bmj.i4746)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via BMJ at <http://www.bmj.com/content/354/bmj.i4746>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

PRACTICE

CLINICAL UPDATE

Living kidney donation

Pippa Bailey *clinical lecturer in renal medicine*^{1 2}, Anusha Edwards *consultant transplant and endocrine surgeon*², Aisling E Courtney *consultant nephrologist*³

¹School of Clinical Sciences, University of Bristol, Bristol BS2 8DZ, UK; ²Southmead Hospital, North Bristol NHS Trust, Bristol BS10 5NB, UK;

³Belfast City Hospital, Belfast Health and Social Care Trust, Belfast BT9 7AB, UK

Globally each year more than 30 000 people become living kidney donors.¹ In the UK there have been more than 1000 living kidney donations every year since 2009.² There are two types of kidney donation, living and deceased. Living donor kidney transplantation offers the best treatment in terms of life expectancy and quality of life²⁻⁹ for most people with kidney failure (see box 1), the prevalence of which is steadily rising. Living kidney donation is constantly evolving, with new ways of maximising recipient opportunities and increasing information regarding long term outcomes associated with donation. This review presents an overview of current practice covering who can donate, to whom, and the possible impact of donation on the donor's health.

Why do non-specialists need to know about this?

Non-specialists might be approached for information about living kidney donation and need to know where to access up-to-date relevant information. Refer interested potential donors to a living donor kidney transplant centre for advice. General practitioners or family physicians may be asked by a specialist team for information about a potential donor undergoing assessment, or to assist with organisation of investigations in the pre-donation stage. Given that the prevalence of living donors is increasing (currently about 1000 per annum in the UK) doctors are increasingly likely to encounter people who have donated and should be aware of what impact donation may have on their health. Some patients prefer to see their regular general practitioner for long term follow-up after donation.¹⁸

Who can donate a kidney, and to whom?

Most living kidney donors donate to relatives or friends, which is termed directed (or specified) donation. Alternatively, individuals can donate a kidney to a recipient with whom they do not have a pre-existing relationship—termed non-directed altruistic donation (or “Good Samaritan” or unspecified donation).^{19 20} This was formally legalised in the UK in 2007.

Altruistic donors now account for about 10% of all living kidney donations.²

In the UK in 2012 the British Transplantation Society and Human Tissue Authority provided guidance on “directed altruistic donation.” This term is used to describe organ donation that happens either between individuals who have a genetic relationship but no established emotional relationship, or between a donor-recipient pair who had no pre-existing emotional or genetic relationship. This definition was an attempt to overcome some of the confusion brought about by the use of social media to recruit potential living kidney donors, which caused blurring of the lines between directed and non-directed donations. Terminology varies, however, and a working group from the European Society for Organ Transplantation alternatively describes publically solicited donors as “solicited specified donors.”²¹ The stimulus for a potential donor to come forward is the publication of an individual's need for a transplant on social networks or through the media.^{20 22}

Historically, all transplants had to occur between ABO blood group and HLA antigen compatible donor and recipient pairs, to prevent hyperacute rejection of the transplanted kidney. However, donors no longer need to be have such compatibility. Most transplants are still between compatible pairs, but desensitisation techniques have been developed to reduce anti-donor antibody titres in the recipient to allow immunologically incompatible kidney transplants.²³⁻²⁵ It is not always possible to lower antibody titres sufficiently to proceed, and such transplants do carry increased risks to the recipient, particularly of infection and rejection in the early post-transplant period. Long term graft survival, though improving,²⁵ is compromised compared with compatible living donor graft survival. Immunologically matched transplants remain preferable, and in many countries this is achieved via a regional or national living donor kidney exchange scheme (fig 1).²⁶⁻³⁰

What you need to know

- Living donor kidney transplantation is the best treatment for most people with renal failure
- There is no upper age limit and few absolute contraindications to living kidney donation
- A living kidney donor does not always need to be the same blood group as the intended recipient
- Most donors have no long term ill effects from living with a single kidney
- Ensure living kidney donors have annual follow-up including blood pressure assessment, urine analysis for protein, and estimation of renal function

Box 1: Advantages of living donor over deceased donor kidney transplants

- Living donor kidney transplants reduce the number of individuals on waiting lists for deceased donor transplants and offer the possibility of a transplant to more patients, who would otherwise be dialysis dependent¹⁰
- Living donor kidney transplants typically last longer, and recipient survival is greater²⁻¹¹
- Living donor transplants are associated with shorter hospital stays, minimising disruption to recipients' lives
- Planned desensitisation of recipients can occur more easily to allow immunologically incompatible transplants
- The elective nature of the surgery permits transplantation in patients who would be unsuitable for emergency surgery
- The recipient costs of living donor kidney transplants are less, because of shorter hospital stays and the decreased incidence of delayed transplant function and early transplant failure.¹²⁻¹³
- Living donor kidney transplants are more likely to take place before the recipient has started dialysis (pre-emptive). Shorter periods spent on dialysis are associated with less comorbidity and better post-transplant outcomes.¹⁴⁻¹⁵ Pre-emptive transplants are also associated with cost savings from the avoidance of dialysis.¹⁶⁻¹⁷

What motivates people to become living kidney donors?

A thematic synthesis of qualitative studies has described directed and non-directed living kidney donors as being compelled by altruism, inherent responsibility, family expectation, personal benefit, and spiritual beliefs.³¹ Non-directed altruistic donors, much like directed donors, report being motivated by a desire to benefit other people or society as a whole, a compelling desire to donate and recognise personal psychosocial gains.³²⁻³⁴ Where commercial organ trading exists, or where donation is financially incentivised, most unrelated donors report financial motivation.³⁵⁻³⁸ Payment for donation is illegal in almost all countries.³⁹

What are the risks of living kidney donation?**Short term risks**

Mortality in living kidney donation is estimated to be between 0.01 and 0.03%.⁴⁰⁻⁴² A recent systematic review reported that perioperative complications, such as wound infection and bleeding, occur in about 7.3% of cases.⁴¹ Immediately after nephrectomy, the glomerular filtration rate of the donor roughly halves. However, a year after donation, this is expected to increase to 60-70% of pre-donation levels,⁴³ due to adaptive hyperfiltration in the remaining kidney.

Longer term risks

Our knowledge of the long term risks of living kidney donation is incomplete, especially for specific donor subgroups (such as those defined by age, ethnicity, socioeconomic position, and with comorbidities). Observational data are limited by duration of follow-up and identifying an appropriate comparative group. In addition, when living donation was first introduced only relatively young and healthy individuals were accepted for kidney donation. As the criteria for donation have expanded to include older donors and those with comorbidities,⁴⁴ however, the current donor population is now demographically different from the cohort for whom long term follow-up data are

available. This limits the generalisability of findings from studies of early donor populations to currently accepted donors.

Advising donors about risks

In order for a living donor to provide informed consent, potential donors and recipients require information on the known risks, and the limits to our understanding of these risks. When counselling a potential donor, risk must be considered on an individual basis. Establish whether younger, non-white,⁴⁵ overweight individuals or those with pre-existing hypertension or impaired glucose tolerance are prepared to make lifestyle changes to minimise long term risks. A small case-control study of long term risks in US army personnel 45 years after unilateral nephrectomy due to trauma reported that mortality was not increased in servicemen with one kidney compared with age matched controls with two kidneys.⁴⁶ Other small single centre studies have also reported that the survival of living kidney donors is better than that of the general population.⁴⁷⁻⁴⁸

Two recent large cohort studies have reported a higher risk of kidney failure among donors compared with healthy non-donors, but the absolute 15 year reported incidence of kidney failure in both studies was <1%.⁴⁹⁻⁵⁰ These observational studies had limitations, and criticisms have included concerns regarding the comparability of non-donor controls⁵¹⁻⁵³ and relatively short follow-up.⁵⁴

A recent US analysis attempted to estimate an individual's risk of renal failure if they did not donate a kidney, and compared 15-year projections with the observed risk among a large cohort of living kidney donors.⁵⁵ This concluded that the relative risk for donors was 3.5-5.3 times higher than the predicted risks in the absence of donation.⁵⁵ Overall, the absolute risk for donors was <1% over this period, comparable with other observational studies, but the risk was greater in black donors as well as current or former smokers.

Two meta-analyses have suggested that kidney donors may have a small increase in blood pressure⁵⁶ (<6 mm Hg) and in urinary protein,⁵⁷ although the quality of research included in both analyses was reported as poor.⁵⁸ The data on an increased cardiovascular event rate in donors is equivocal.⁴⁰⁻⁶⁰ Although the risks of gestational hypertension and pre-eclampsia seem to be higher in pregnancies among donors than among healthy

non-donors,⁶¹⁻⁶³ adverse outcomes for mother or offspring have not been documented.

Work exploring the mental health of living kidney donors is limited, but a recent case-control study from the Netherlands suggested that donation is not associated with short term changes in mental health.⁶⁴ Multiple studies suggest that the quality of life of most living kidney donors seems to be at least equal to that of the general population⁶⁵⁻⁷² and usually returns to pre-donation levels after donation. A thematic synthesis of qualitative work from Europe, the United States, Canada and Australia, however, identified some negative donation-specific experiences among the overall positive experiences, including a sense of loss, fear, vulnerability, and neglect.³¹

Several studies have reported that potential donors are more willing to accept greater donor risks than potential recipients and transplant professionals.⁷³ Greater risks seem to be accepted when the intended recipient is closely related and when his or her prognosis is poor.^{74 75} In addition, potential donors have been found to be more likely than potential recipients or clinicians to agree that living donation is acceptable when long term donor risks are uncertain.⁷³

In addition to evaluating the risks, consideration of living kidney donation must also involve an evaluation of the benefits to potential donors, which can be substantial, and the harms of non-donation.⁷⁶

How are living donors assessed and how long does it take?

National guidelines exist for living kidney donor evaluation.¹⁸⁻⁸⁰ Details of the guidance differ between countries,⁸⁰ and the International Kidney Disease: Improving Global Outcomes (KDIGO) committee is developing a global, evidence based guideline which is currently available for public consultation.⁸¹

The primary goals of donor evaluation are to determine the suitability of an individual for donation, to ensure a donor is making an informed choice free from coercion or monetary incentive, and to confirm that the kidney is suitable to be transplanted into the intended recipient.

In the UK the living donor evaluation process typically encompasses the steps outlined in figure 2.

The duration of living donor evaluation varies between and within countries.⁸² Potential donors need to be given a suitable period to consider donation, often described as a “cooling off” period.⁸³ The clinical assessments and investigations can be carried out in one day, and some centres in the UK run one-day assessments.⁸⁴ To be a donor, an individual needs to be suitable to undergo surgery under general anaesthesia and to be able to cope with one kidney for the longer term.

Who cannot donate?

Very few absolute contraindications to living kidney donation exist (box 2).⁴⁴

The *United Kingdom Guidelines for Living Donor Kidney Transplantation*¹⁸ specify the minimum measured glomerular filtration rate (mGFR) required for donation in order to ensure that the mGFR of the remaining kidney is predicted to be more than 37.5 mL/min/1.73 m² at 80 years of age. The data on which this was based were limited, and there is variation between UK centres in the measurement of GFR.⁸⁶

In most transplant programmes living kidney donors are required to be over 18 years of age,¹⁸ but cases where adolescents have donated do exist.⁸⁷ Caution is recommended when accepting

donors under 25 years of age as younger people have more time to develop comorbidities and an increased lifetime risk of renal failure.⁵⁵ Women who wish to have children need to be counselled regarding the small increased incidence of gestational hypertension and pre-eclampsia among kidney donors,⁶¹⁻⁶³ and alternative donors are preferred if available. Other relative contraindications to donation include obesity and diabetes, and factors that might affect a potential donor's decision-making capacity (such as cognitive impairment, undertreated psychiatric conditions, or active substance misuse).

In most countries potential donors are prevented from donating if there is evidence of donor coercion, as identified by any member of the donor evaluation team or by an independent assessor or donor advocate.^{20 80}

What is the surgical procedure?

Laparoscopic kidney removal is preferred¹⁸ as it is associated with less pain, shorter hospital stay, and earlier return to normal activities.⁸⁸ Minimally invasive surgical techniques for donor nephrectomy include robot assisted laparoscopy⁸⁹ and a laparoendoscopic single site approach.^{41 90} After surgery, living donors tend to stay in hospital for two to three days, and full recovery is expected within 6-12 weeks.¹⁸

What is the global picture?

Almost 100 of the World Health Organization member states now report transplantation services, but the rates of living kidney donation and the proportion of kidney transplants from living donors vary widely (fig 3 and 4).⁹¹⁻⁹³

Reasons for variation include differences in the underlying prevalence of renal failure, the acceptability and rates of deceased donor transplantation, the healthcare funding for preoperative and postoperative donor care, and differing donor reimbursement or incentives. Iran is currently the only country in the world to have a compensated and government regulated programme for unrelated living donor renal transplantation.³⁹ Payment for donation in most other countries is illegal. However, there remains an international trade in both organs and living donors.^{94 95} The Declaration of Istanbul on organ trafficking and transplant tourism outlines the circumstances in which removing organs from a living or deceased donor are unacceptable and aims to halt unethical transplant activities, including organ trafficking, transplant tourism, and transplant commercialism.⁹⁶

Contributors: PB proposed the topic to the editorial board, and wrote the first draft of the paper. AE and AEC reviewed and revised the drafted paper. All authors approved the final version.

Competing interests: We have read and understood the BMJ policy on declaration of interests, and declare the following interests: AEC is the unpaid chair of the Living Donor Kidney Transplantation 2020 Strategy Implementation Group, a sub-group of the UK's NHS Blood and Transplant Kidney Advisory Group.

Provenance and peer review: Not commissioned; externally peer reviewed.

Patient consent obtained.

- 1 European Directorate for the Quality of Medicines. Newsletter transplant: international figures on donation and transplantation 2014. EDQM vol 20, 2015. https://www.edqm.eu/sites/default/files/newsletter_transplant_2015.pdf.
- 2 NHS Blood and Transplant. Statistics and clinical audit: NHS blood and transplant organ donation and transplantation activity report 2014/15. 2015. <http://www.odt.nhs.uk/uk-transplant-registry/annual-activity-report/>.
- 3 Cecka JM. Living donor transplants. *Clin Transpl* 1995;363-77. PMID:8794280.

Box 2: Absolute contraindications to living kidney donation¹⁸⁻⁸⁵

- Active malignancy* or chronic infection
- Nephrolithiasis secondary to a metabolic abnormality
- Uncontrolled hypertension
- Overt proteinuria, glomerular pathology, or an inadequate glomerular filtration rate (GFR)
- Bilateral renal artery atherosclerosis or fibromuscular dysplasia involving the orifices of both renal arteries
- Sickle cell disease

*In some countries donors with certain types of cancer or successfully treated low grade tumour may be considered for kidney donation (such as small (<4 cm) subcapsular renal cell carcinoma with excision at time of donation and no distant spread, or low grade non-melanoma skin cancer)

Methods box

A PubMed Medline search was undertaken in May 2016 based on the following key words and variants: living, donor(s), donation, transplant(s), kidney, renal. The Cochrane Collaboration database was also screened for relevant articles. The Kidney Disease Improving Global Outcomes (KDIGO) reports *Summary of Published Living Kidney Donor Guidelines*, *Guideline Evidence Report*, and *Outcomes of Living Kidney Donation*, and the draft of *Clinical Practice Guideline on the Evaluation and Follow-up Care of Living Kidney Donors* were reviewed, along with the *United Kingdom Guidelines for Living Donor Kidney Transplantation* and references.

The authors' own knowledge, clinical experience, research, and personal archives of references were used. Research published since article submission was highlighted by reviewers. AE Courtney is chair of the UK NHS Blood and Transplant's Living Donor Kidney Transplantation 2020 Strategy Implementation Group.

Living kidney donation—a donor and family's experience

Our daughter thrived without illness for the first 18 months of her life, but then we noticed a swelling on one side of her stomach and took her to the doctor. A number of scans and x rays later, we were finally given the news that she had completely no function in one kidney and her other kidney was under severe pressure due to hydronephrosis. Our whole family was shocked and traumatised.

With appropriate drains and stents, our daughter had reasonable kidney function for a number of years, but at the age of 11 years her creatinine levels were creeping up and her consultant discussed dialysis. We raised the option of live donation. We had no idea really at this point what it entailed so had a list of questions to ask: how long the process will take, the likelihood of a match, whether it will take straight away, and numerous others. We wanted to do it to get our daughter healthy again without the dialysis if at all possible. Decision made, we wanted to proceed.

My husband was a slightly better match than me, and was put in contact with the living kidney donor coordinator. At the first appointment, we were told that the process takes approximately nine or 10 months, and all I could think of was whether we had this amount of time, as our daughter's kidney was failing and she was determined not to have dialysis if she could avoid it.

A few weeks later my husband had the first of the tests. A number of weeks later the next and so on. We asked whether a lot of these tests could be run on the same day to avoid disruption and minimise impact on his employment. We were informed that unfortunately the process was slow. Our daughter at this stage was losing a lot of weight and her condition was deteriorating quickly.

After the first date was cancelled four days before the surgery, which was devastating, on 8 August 2007 all went ahead. I was surprised at how well my husband looked that afternoon, and at around 4.30 pm our daughter was back onto the intensive care ward and sitting up looking amazing. The kidney worked straight away and all was good. My husband was discharged the following afternoon and returned to work six weeks later. Our daughter experienced instant improvement in her health, and joked about how tanned she looked, to which I replied "That's normal skin tone."

All went well for four years, until a virus known as BK attacked the kidney, and then rejection developed. With appropriate treatment, our daughter's kidney function stabilised but deteriorated again in 2014. She was in need of another kidney. Timing was not good. She was about to enter the final year of her master's degree and was determined to finish it alongside her peers. However, around Christmas we recognised the familiar signs that she had shown many years before—weight loss, vomiting, and extreme tiredness.

In January I rang the living donor coordinator to ask if I could be tested—and was informed that all the tests were now carried out in one day! A few weeks later, I arrived on the ward at 8.30 am for my tests to commence. The coordinator had a list of appointments and went to each department alongside me. By 4.30 pm that afternoon I was on my way home, exhausted but delighted to know that in a week or two we would know for sure if my kidneys were healthy enough for donating. They were, and surgery was booked for 23 March 2015 (when our daughter had no lectures due to the Easter break).

All went well, I returned to work after seven weeks, and our daughter graduated from university three months later. She started a PhD in physics that same year.

As a family having been through the living donation process twice, we would most certainly recommend the one day testing. The process for my husband was long and drawn out and impacted greatly on family and work life. During the time waiting, we were further traumatised watching our daughter become increasingly unwell, and there was little we could do. On the second occasion, it was a full day in hospital, but by the time I was leaving most of the tests had been conducted, we knew things were progressing quickly, and I only had to take one day off work.

Nine years on, and my husband remains well. One year on, I had my review at the hospital and all is good. Most importantly our daughter's results are excellent and she is living life to the full.

We are truly grateful for the staff that cared for us during both these transplants. As a family we thank them all.

Jacqueline Johnston, donated at Belfast City Hospital, Belfast Health and Social Care Trust
Previous accounts of living kidney donation have been published in the BMJ:

Ferriman A. A patient's journey: Becoming a live kidney donor. *BMJ* 2008;336:1374 (<http://www.bmj.com/content/336/7657/1374>)

Thiruchelvam PT, Willicombe M, Hakim N, Taube D, Papalois V. Renal transplantation. *BMJ* 2011;343:d7300 (<http://dx.doi.org/10.1136/bmj.d7300>)

4 Terasaki PI, Cecka JM, Gjertson DW, Takemoto S. High survival rates of kidney transplants from spousal and living unrelated donors. *N Engl J Med* 1995;333:333-6. doi:10.1056/NEJM199508103330601 pmid:7609748.

5 Laupacis A, Keown P, Pus N, et al. A study of the quality of life and cost-utility of renal transplantation. *Kidney Int* 1996;50:235-42. doi:10.1038/ki.1996.307 pmid:8807593.

6 Cecka JM. The OPTN/UNOS Renal Transplant Registry. *Clin Transpl* 2005;1:1-16.pmid:17424721.

7 Roodnat JI, van Riemsdijk IC, Mulder PGH, et al. The superior results of living-donor renal transplantation are not completely caused by selection or short cold ischemia time: a single-center, multivariate analysis. *Transplantation* 2003;75:2014-8. doi:10.1097/01.TP.0000065176.06275.42 pmid:12829903.

8 Oniscu GC, Brown H, Forsythe JL. How great is the survival advantage of transplantation over dialysis in elderly patients? *Nephrol Dial Transplant* 2004;19:945-51. doi:10.1093/ndt/gfh022 pmid:15031354.

Resources for healthcare professionals

- Kidney Disease Improving Global Outcomes (KDIGO). *Clinical Practice Guideline on the Evaluation and Follow-up Care of Living Kidney Donors*, released for public consultation in November 2015 and expected to be formally published in 2016
- Joint Working Party of The British Transplantation Society and The Renal Association. *United Kingdom Guidelines for Living Donor Kidney Transplantation*. 3rd ed. 2011. <https://www.bts.org.uk/Documents/Guidelines/Active/UK%20Guidelines%20for%20Living%20Donor%20Kidney%20July%202011.pdf>
 - Details the UK guidelines for the evaluation of both living kidney donors and transplant recipients. Includes the legal framework and guidance on financial reimbursement
 - Plus “Addendum to Living Kidney Donor Guidelines.” 2015. http://www.bts.org.uk/BTS/Guidelines_Standards/Current/BTS/Guidelines_Standards/Current_Guidelines.aspx—Provides an update on the risks of living kidney donation based on recent research findings
- NHSBT Organ Donation and Transplantation. Living donor kidney transplantation <http://www.odt.nhs.uk/donation/living-donation/living-donor-kidney-transplantation>
 - UK’s national transplant website with information on all aspects of donation and transplantation, including guidelines and activity reports
- Kidney Disease Improving Global Outcomes (KDIGO). *Outcomes of Living Kidney Donation*. www.kdigo.org/clinical_practice_guidelines/LivingDonor/KDIGO%20Guideline%20Evidence%20Report.pdf
 - Reports findings from a systematic review of research into the outcomes of living kidney donation
- Thiruchelvam PT, Willcombe M, Hakim N, Taube D, Papalois V. Renal transplantation. *BMJ* 2011;343:d7300 (<http://dx.doi.org/10.1136/bmj.d7300>)
 - Provides further details on renal transplantation, including details of transplant candidate eligibility and details of deceased donor transplantation. However, the content regarding living donor transplantation has been updated for this review
- Reese PP, Boudville N, Garg AX. Living kidney donation: outcomes, ethics, and uncertainty. *Lancet* 2015;385:2003-13
 - Summarises data on long term donor outcomes and the ethical implications and challenges related to decision making for donors

Information for patients

- NHSBT Organ Donation. Living organ donation. <https://www.organdonation.nhs.uk/faq/living-organ-donation>
- Give a Kidney—One’s Enough. How to give a kidney. <http://www.giveakidney.org/2015/uncategorized/how-to-give-a-kidney>
- British Kidney Patient’s Association. Donating a kidney: what to expect. http://www.britishkidney-pa.co.uk/images/stories/downloads/patientinfopage/BKPA_leaflet_leaflet-donating.pdf
- healthtalk.org. People’s experiences: organ donation. <http://www.healthtalk.org/peoples-experiences/organ-donation-transplant/organ-donation/topics>—Video accounts of the experiences of living donors
- National Kidney Foundation. The big ask: the big give. <https://www.kidney.org/transplantation/livingdonors>—US website about living kidney donation

Ongoing research

Ongoing research aims to determine the long term outcomes for living kidney donors, especially for younger donors, those with comorbidities such as hypertension and obesity, and donors of all ethnic groups and socioeconomic positions. Determining the long term outcomes for donors is one of the “Top 10 kidney transplant research priorities” set by the UK James Lind Alliance and Centre for Evidence in Transplantation Priority Setting Partnership in February 2016 (<http://www.jla.nihr.ac.uk/priority-setting-partnerships/kidney-transplant/top-10-priorities/>)

Examples of active trials:

- Kidney Donor Outcomes Cohort (KDOC) Study – USA. ClinicalTrials.gov identifier: NCT01427452 (<https://clinicaltrials.gov/ct2/show/NCT01427452>)
- The Long Term Medical and Psychological Implications of Becoming a Living Kidney Donor: A Prospective Pilot Study – USA, Canada and Australia. ClinicalTrials.gov identifier: NCT00319579 (<https://clinicaltrials.gov/ct2/show/NCT00319579>)
- Long-term Effects of Becoming a Living Kidney Donor Study – Canada, Australia. ClinicalTrials.gov identifier: NCT00936078 (<https://clinicaltrials.gov/ct2/show/NCT00936078>)

Further research is also required to better understand barriers to living kidney donation, in order for socioeconomic and racial disparities in living donor kidney transplantation to be addressed and to facilitate living kidney donation when individuals are motivated to donate:

- LaPointe Rudow D, Hays R, Baliga P, et al. Consensus conference on best practices in live kidney donation: recommendations to optimize education, access, and care. *Am J Transplant* 2015;15: 914-22.

How patients were involved in the creation of this review

A living kidney donor was invited to provide an account of her experiences. Jacqueline Johnston had donated a kidney to her daughter, and her husband had also donated previously. Her account contrasts the experience her husband had of donor evaluation over several months with that of her own assessment, which took place in one day.

- Knoll GA. Kidney transplantation in the older adult. *Am J Kidney Dis* 2013;61:790-7. doi: 10.1053/j.ajkd.2012.08.049 pmid:23261121.
- NHS Blood and Transplant. Living donor kidney transplantation 2020: A UK strategy. 2014. www.odt.nhs.uk/pdf/ldkt_2020_strategy.pdf.
- Hariharan S, Johnson CP, Bresnahan BA, Taranto SE, McIntosh MJ, Stablein D. Improved graft survival after renal transplantation in the United States, 1988 to 1996. *N Engl J Med* 2000;342:605-12. doi:10.1056/NEJM200003023420901 pmid:10699159.
- Barnieh L, Manns BJ, Klarenbach S, McLaughlin K, Yilmaz S, Hemmelgarn BR. A description of the costs of living and standard criteria deceased donor kidney transplantation. *Am J Transplant* 2011;11:478-88. doi:10.1111/j.1600-6143.2010.03425.x pmid:21299831.
- Smith CR, Woodward RS, Cohen DS, et al. Cadaveric versus living donor kidney transplantation: a Medicare payment analysis. *Transplantation* 2000;69:311-4. doi:10.1097/00007890-200001270-00020 pmid:10670645.
- Meier-Kriesche HU, Port FK, Ojo AO, et al. Effect of waiting time on renal transplant outcome. *Kidney Int* 2000;58:1311-7. doi:10.1046/j.1523-1755.2000.00287.x pmid:10972695.
- Meier-Kriesche HU, Kaplan B. Waiting time on dialysis as the strongest modifiable risk factor for renal transplant outcomes: a paired donor kidney analysis. *Transplantation* 2002;74:1377-81. doi:10.1097/00007890-200211270-00005 pmid:12451234.
- NHS Blood and Transplant. Taking Organ Transplantation to 2020: A UK strategy. Economic case for organ transplantation. <http://www.nhsbt.nhs.uk/to2020/the-strategy/supporting-documents/economic-case-for-organ-transplantation.pdf>.

- 17 McKeigue S. Costs and savings to NHS from solid organ transplantation. Department of Health. 2015. http://www.odt.nhs.uk/pdf/Solid_Organ_Transplantation_Final.pdf.
- 18 Joint Working Party of the British Transplantation Society and the Renal Association. United Kingdom guidelines for living donor kidney transplantation. 2011. www.bts.org.uk/transplantation/standards-and-guidelines/.
- 19 Dor FJ, Massey EK, Frunza M, et al. New classification of ELPAT for living organ donation. *Transplantation* 2011;91:935-8. doi:10.1097/TP.0b013e3182129236 pmid:21423070.
- 20 Human Tissue Authority. Guidance to transplant teams and independent assessors. 2015. [https://www.hta.gov.uk/sites/default/files/Guidance_to_Transplant_Teams_and_Independent_Assessors%20\(1\)_0.pdf](https://www.hta.gov.uk/sites/default/files/Guidance_to_Transplant_Teams_and_Independent_Assessors%20(1)_0.pdf).
- 21 Frunza M, Van Assche K, Lennerling A, et al. Dealing With Public Solicitation of Organs From Living Donors--An ELPAT View. *Transplantation* 2015;99:2210-4. doi:10.1097/TP.0000000000000669 pmid:25769072.
- 22 British Transplantation Society. Guidelines for directed altruistic organ donation. 2014. [http://www.bts.org.uk/Documents/Guidelines/Active/Guidelines%20for%20Directed%20Altruistic%20Donation-%20Final%20Revised%20Website%20Version%2006-2014%20\(2\).pdf](http://www.bts.org.uk/Documents/Guidelines/Active/Guidelines%20for%20Directed%20Altruistic%20Donation-%20Final%20Revised%20Website%20Version%2006-2014%20(2).pdf).
- 23 Montgomery RA, Zachary AA, Racusen LC, et al. Plasmapheresis and intravenous immune globulin provides effective rescue therapy for refractory humoral rejection and allows kidneys to be successfully transplanted into cross-match-positive recipients. *Transplantation* 2000;70:887-95. doi:10.1097/00007890-200009270-00006 pmid:11014642.
- 24 Montgomery RA, Lonze BE, King KE, et al. Desensitization in HLA-incompatible kidney recipients and survival. *N Engl J Med* 2011;365:318-26. doi:10.1056/NEJMoa1012376 pmid:21793744.
- 25 British Transplantation Society. Guidelines for antibody incompatible transplantation. 2011 www.bts.org.uk/Documents/Guidelines/Active/AiT%20guidelines%20Jan%202011%20FINAL.pdf.
- 26 NHS Blood and Transplant. Organ donation and transplantation national living donor kidney sharing scheme. 2012. www.odt.nhs.uk/donation/living-donation/national-living-donor-kidney-sharing-scheme.
- 27 de Klerk M, Keizer KM, Claas FH, Witvliet M, Haase-Kromwijk BJ, Weimar W. The Dutch national living donor kidney exchange program. *Am J Transplant* 2005;5:2302-5. doi:10.1111/j.1600-6143.2005.01024.x pmid:16095513.
- 28 de Klerk M, Kal-van Gestel JA, Haase-Kromwijk BJ, Claas FH, Weimar W. Living Donor Kidney Exchange Program. Eight years of outcomes of the Dutch Living Donor Kidney Exchange Program. *Clin Transpl* 2011;287-90. doi:10.1111/j.1600-6143.2005.01024.x pmid:16095513.
- 29 Park K, Lee JH, Huh KH, Kim SI, Kim YS. Exchange living-donor kidney transplantation: diminution of donor organ shortage. *Transplant Proc* 2004;36:2949-51. doi:10.1016/j.transproceed.2004.12.013 pmid:15686667.
- 30 Delmonico FL, Morrissey PE, Lipkowitz GS, et al. Donor kidney exchanges. *Am J Transplant* 2004;4:1628-34. doi:10.1111/j.1600-6143.2004.00572.x pmid:15367217.
- 31 Tong A, Chapman JR, Wong G, Kanellis J, McCarthy G, Craig JC. The motivations and experiences of living kidney donors: a thematic synthesis. *Am J Kidney Dis* 2012;60:15-26. doi:10.1053/j.ajkd.2011.11.043 pmid:22305757.
- 32 Nolan MT, Walton-Moss B, Taylor L, Dane K. Living kidney donor decision making: state of the science and directions for future research. *Prog Transplant* 2004;14:201-9. doi:10.7182/prtr.14.3.440p7266x4524344 pmid:15495779.
- 33 Challenor J, Watts J. 'It seemed churlish not to': How living non-directed kidney donors construct their altruism. *Health (London)* 2014;18:388-405. doi:10.1177/1363459313501358 pmid:24026358.
- 34 Clarke A, Mitchell A, Abraham C. Understanding donation experiences of unspecified (altruistic) kidney donors. *Br J Health Psychol* 2014;19:393-408. doi:10.1111/bjhp.12048 pmid:23692296.
- 35 Khajehdehi P. Living non-related versus related renal transplantation--its relationship to the social status, age and gender of recipients and donors. *Nephrol Dial Transplant* 1999;14:2621-4. doi:10.1093/ndt/14.11.2621 pmid:10534501.
- 36 Malakoutian T, Hakemi MS, Nassiri AA, et al. Socioeconomic status of Iranian living unrelated kidney donors: a multicenter study. *Transplant Proc* 2007;39:824-5. doi:10.1016/j.transproceed.2007.04.018 pmid:17524823.
- 37 Moazzam F, Zaman RM, Jafarey AM. Conversations with kidney vendors in Pakistan: an ethnographic study. *Hastings Cent Rep* 2009;39:29-44. doi:10.1353/hcr.0.0136 pmid:19537621.
- 38 Awaya T, Siruno L, Toledano S, Aguilar F, Shimazono Y, De Castro L. Failure of informed consent in compensated non-related kidney donation in the Philippines. *Asian Bioethics Review*. 2009;1:138-43.
- 39 Ghods AJ, Savaj S. Iranian model of paid and regulated living-unrelated kidney donation. *Clin J Am Soc Nephrol* 2006;1:1136-45. doi:10.2215/CJN.00700206 pmid:17699338.
- 40 Segev DL, Muzaale AD, Caffo BS, et al. Perioperative mortality and long-term survival following live kidney donation. *JAMA* 2010;303:959-66. doi:10.1001/jama.2010.237 pmid:20215610.
- 41 Kortram K, Ijzermans JN, Dor FJ. Perioperative Events and Complications in Minimally Invasive Live Donor Nephrectomy: A Systematic Review and Meta-Analysis. *Transplantation* 2016. doi:10.1097/TP.0000000000001327 pmid:27428715.
- 42 Najarian JS, Chavers BM, McHugh LE, Matas AJ. 20 years or more of follow-up of living kidney donors. *Lancet* 1992;340:807-10. doi:10.1016/0140-6736(92)92683-7 pmid:1357243.
- 43 Rook M, Hofker HS, van Son WJ, Homan van der Heide JJ, Ploeg RJ, Navis GJ. Predictive capacity of pre-donation GFR and renal reserve capacity for donor renal function after living kidney donation. *Am J Transplant* 2006;6:1653-9. doi:10.1111/j.1600-6143.2006.01359.x pmid:16827867.
- 44 Ahmadi AR, Lafranica JA, Claessens LA, et al. Shifting paradigms in eligibility criteria for live kidney donation: a systematic review. *Kidney Int* 2015;87:31-45. doi:10.1038/ki.2014.118 pmid:24786706.
- 45 Lentine KL, Schnitzler MA, Xiao H, et al. Racial variation in medical outcomes among living kidney donors. *N Engl J Med* 2010;363:724-32. doi:10.1056/NEJMoa1000950 pmid:20818874.
- 46 Narkun-Burgess DM, Nolan CR, Norman JE, Page WF, Miller PL, Meyer TW. Forty-five year follow-up after uninephrectomy. *Kidney Int* 1993;43:1110-5. doi:10.1038/ki.1993.156 pmid:8510390.
- 47 Fehrman-Ekholm I, Elinder CG, Stenbeck M, Tydén G, Groth CG. Kidney donors live longer. *Transplantation* 1997;64:976-8. doi:10.1097/00007890-199710150-00007 pmid:9381544.
- 48 Okamoto M, Akioka K, Nobori S, et al. Short- and long-term donor outcomes after kidney donation: analysis of 601 cases over a 35-year period at Japanese single center. *Transplantation* 2009;87:419-23. doi:10.1097/TP.0b013e318192dc95 pmid:19202449.
- 49 Mjøs G, Hallan S, Hartmann A, et al. Long-term risks for kidney donors. *Kidney Int* 2014;86:162-7. doi:10.1038/ki.2013.460 pmid:24284516.
- 50 Muzaale AD, Massie AB, Wang MC, et al. Risk of end-stage renal disease following live kidney donation. *JAMA* 2014;311:579-86. doi:10.1001/jama.2013.285141 pmid:24519297.
- 51 Matas AJ, Woodle ES, Gaston RS, Forsythe JL, Wadström J, Stegall MD. Kidney donors at increased risk? Additional studies are needed. *Kidney Int* 2014;86:650. doi:10.1038/ki.2014.146 pmid:25168503.
- 52 Matas AJ, Wadström J, Ibrahim HN. Kidney donation and risk of ESRD. *JAMA* 2014;312:92-3. doi:10.1001/jama.2014.5515 pmid:25058223.
- 53 Niemi M, Mandelbrot DA. Kidney donation and risk of ESRD. *JAMA* 2014;312:92. doi:10.1001/jama.2014.5521 pmid:25058224.
- 54 Kiberd B, Tennankore K. Kidney donation and risk of ESRD. *JAMA* 2014;312:93. doi:10.1001/jama.2014.5518 pmid:25058225.
- 55 Grams ME, Sang Y, Levey AS, et al. Chronic Kidney Disease Prognosis Consortium. Kidney-failure risk projection for the living kidney-donor candidate. *N Engl J Med* 2016;374:411-21. doi:10.1056/NEJMoa1510491 pmid:26544982.
- 56 Boudville N, Prasad GV, Knoll G, et al. Donor Nephrectomy Outcomes Research (DONOR) Network. Meta-analysis: risk for hypertension in living kidney donors. *Ann Intern Med* 2006;145:185-96. doi:10.7326/0003-4819-145-3-200608010-00006 pmid:16880460.
- 57 Garg AX, Muirhead N, Knoll G, et al. Donor Nephrectomy Outcomes Research (DONOR) Network. Proteinuria and reduced kidney function in living kidney donors: A systematic review, meta-analysis, and meta-regression. *Kidney Int* 2006;70:1801-10. doi:10.1038/sj.ki.5001819 pmid:17003822.
- 58 Kidney Disease Improving Global Outcomes (KDIGO). Outcomes of living kidney donation: a systematic review for a clinical practice guideline by the Kidney Disease Improving Global Outcomes (KDIGO). www.kdigo.org/clinical_practice_guidelines/LivingDonor/KDIGO%20Guideline%20Evidence%20Report.pdf.
- 59 Garg AX, Meirambayeva A, Huang A, et al. Donor Nephrectomy Outcomes Research Network. Cardiovascular disease in kidney donors: matched cohort study. *BMJ* 2012;344:e1203. doi:10.1136/bmj.e1203 pmid:22381674.
- 60 Reese PP, Bloom RD, Feldman HI, et al. Mortality and cardiovascular disease among older live kidney donors. *Am J Transplant* 2014;14:1853-61. doi:10.1111/ajt.12822 pmid:25039276.
- 61 Ibrahim HN, Akkina SK, Leister E, et al. Pregnancy outcomes after kidney donation. *Am J Transplant* 2009;9:825-34. doi:10.1111/j.1600-6143.2009.02548.x pmid:19353771.
- 62 Reisaeter AV, Roislien J, Henriksen T, Irgens LM, Hartmann A. Pregnancy and birth after kidney donation: the Norwegian experience. *Am J Transplant* 2009;9:820-4. doi:10.1111/j.1600-6143.2008.02427.x pmid:18853953.
- 63 Garg AX, Nevis IF, McArthur E, et al. DONOR Network. Gestational hypertension and preeclampsia in living kidney donors. *N Engl J Med* 2015;372:124-33. doi:10.1056/NEJMoa1408932 pmid:25397608.
- 64 Timmerman L, Laging M, Westerhof GJ, et al. Mental health among living kidney donors: a prospective comparison with matched controls from the general population. *Am J Transplant* 2015;15:508-17. doi:10.1111/ajt.13046 pmid:25582231.
- 65 Johnson EM, Anderson JK, Jacobs C, et al. Long-term follow-up of living kidney donors: quality of life after donation. *Transplantation* 1999;67:717-21. doi:10.1097/00007890-199903150-00013 pmid:10096528.
- 66 Ku JH. Health-related quality of life of living kidney donors: review of the short form 36-health questionnaire survey. *Transpl Int* 2005;18:1309-17. doi:10.1111/j.1432-2277.2005.00231.x pmid:16297049.
- 67 Ibrahim HN, Foley R, Tan L, et al. Long-term consequences of kidney donation. *N Engl J Med* 2009;360:459-69. doi:10.1056/NEJMoa0804883 pmid:19179315.
- 68 Gross CR, Messersmith EE, Hong BA, et al. RELIVE Study Group. Health-related quality of life in kidney donors from the last five decades: results from the RELIVE study. *Am J Transplant* 2013;13:2924-34. doi:10.1111/ajt.12434 pmid:24011252.
- 69 Clemens KK, Thiessen-Philbrook H, Parikh CR, et al. Donor Nephrectomy Outcomes Research (DONOR) Network. Psychosocial health of living kidney donors: a systematic review. *Am J Transplant* 2006;6:2965-77. doi:10.1111/j.1600-6143.2006.01567.x pmid:17294524.
- 70 Lumsdaine JA, Wray A, Power MJ, et al. Higher quality of life in living donor kidney transplantation: prospective cohort study. *Transpl Int* 2005;18:975-80. doi:10.1111/j.1432-2277.2005.00175.x pmid:16008749.
- 71 Garcia MFFM, Andrade LGM, Carvalho MFC. Living kidney donors--a prospective study of quality of life before and after kidney donation. *Clin Transplant* 2013;27:9-14. doi:10.1111/j.1399-0012.2012.01687.x pmid:22831164.
- 72 Glotzer OS, Singh TP, Gallichio MH, Conti DJ, Siparsky NF. Long-term quality of life after living kidney donation. *Transplant Proc* 2013;45:3225-8. doi:10.1016/j.transproceed.2013.05.006 pmid:24182789.
- 73 Young A, Karpinski M, Treleaven D, et al. Differences in tolerance for health risk to the living donor among potential donors, recipients, and transplant professionals. *Kidney Int* 2008;73:1159-66. doi:10.1038/ki.2008.65 pmid:18354380.
- 74 Maple NH, Hadjianastassiou V, Jones R, Mamode N. Understanding risk in living donor nephrectomy. *J Med Ethics* 2010;36:142-7. doi:10.1136/jme.2009.031740 pmid:20211992.
- 75 Spital A, Spital M. Living kidney donation. Attitudes outside the transplant center. *Arch Intern Med* 1988;148:1077-80. doi:10.1001/archinte.1988.00380050081014 pmid:3284500.
- 76 Allen MB, Abt PL, Reese PP. What are the harms of refusing to allow living kidney donation? An expanded view of risks and benefits. *Am J Transplant* 2014;14:531-7. doi:10.1111/ajt.12599 pmid:24730047.
- 77 European Renal Best Practice Transplantation Guideline Development Group. ERBP guideline on the management and evaluation of the kidney donor and recipient. *Nephrol Dial Transplant* 2013;28(suppl 2):ii1-71. doi:10.1093/ndt/28(suppl_2)/ii1-71 pmid:24026881.
- 78 Ierino F, Boudville N, Kanellis J. CARL. The CARL guidelines. Donors at risk: hypertension. *Nephrology (Carlton)* 2010;15(Suppl S1):S114-20. doi:10.1111/j.1469-7610.2010.02591022.
- 79 US organ procurement and transplantation network policies - Policy 14: Living donation. 2016. p 175-89. https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf.
- 80 Kidney Disease: Improving Global Outcomes (KDIGO). *Summary of published living kidney donor guidelines*. www.kdigo.org/clinical_practice_guidelines/LivingDonor/Summary%20of%20Published%20Living%20Kidney%20Donor%20Guidelines.pdf.
- 81 Kidney Disease: Improving Global Outcomes (KDIGO). Clinical practice guideline on the evaluation and follow-up care of living kidney donors - draft. 2015. www.kdigo.org/home/guidelines/livingdonor.
- 82 Arunachalam C, Garrues M, Biggins F, Woywodt A, Ahmed A. Assessment of living kidney donors and adherence to national live donor guidelines in the UK. *Nephrol Dial Transplant* 2013;28:1952-60. doi:10.1093/ndt/gft096 pmid:23658249.

- 83 Rodrigue JR, Pavlakis M, Danovitch GM, et al. Evaluating living kidney donors: relationship types, psychosocial criteria, and consent processes at US transplant programs. *Am J Transplant* 2007;7:2326-32. doi:10.1111/j.1600-6143.2007.01921.x PMID:17845566.
- 84 Graham J, Courtney A. Oral Presentation: 5 years of 1-days: Outcomes of potential living kidney donors undergoing a 1-day assessment pathway. British Transplantation Society Congress; Scottish Exhibition and Conference Centre, Glasgow 2016.
- 85 Kher A, Mandelbrot DA. The living kidney donor evaluation: focus on renal issues. *Clin J Am Soc Nephrol* 2012;7:366-71. doi:10.2215/CJN.10561011 PMID:22223615.
- 86 Peters AM, Howard B, Neilly MD, et al. The reliability of glomerular filtration rate measured from plasma clearance: a multi-centre study of 1,878 healthy potential renal transplant donors. *Eur J Nucl Med Mol Imaging* 2012;39:715-22. doi:10.1007/s00259-011-2024-5 PMID:22223168.
- 87 MacDonald D, Kukla AK, Ake S, et al. Medical outcomes of adolescent live kidney donors. *Pediatr Transplant* 2014;18:336-41. doi:10.1111/petr.12238 PMID:24646177.
- 88 Wilson CH, Sanni A, Rix DA, Soomro NA. Laparoscopic versus open nephrectomy for live kidney donors. *Cochrane Database Syst Rev* 2011;11:CD006124. PMID:22071829.
- 89 Giacomoni A, Di Sandro S, Lauterio A, et al. Robotic nephrectomy for living donation: surgical technique and literature systematic review. *Am J Surg* 2016;211:1135-42. doi:10.1016/j.amjsurg.2015.08.019 PMID:26499052.
- 90 Gupta A, Ahmed K, Kynaston HG, Dasgupta P, Chlosta PL, Aboumarzouk OM. Laparoendoscopic single-site donor nephrectomy (LESS-DN) versus standard laparoscopic donor nephrectomy. *Cochrane Database Syst Rev* 2016;5:CD010850. PMID:27230690.
- 91 International Registry in Organ Donation and Transplantation IRODaT Newsletter - Final Numbers. 2014. December 2015 [cited 2016 June 3]. Available from: www.irodat.org/img/database/pdf/NEWSLETTER2015_December2.pdf.
- 92 Muller E, Dominguez-Gil B, Matesanz R, Delmonico F. Kidney transplantation across the globe: the good and bad. The Lancet Kidney Campaign. *Lancet*. <http://www.thelancet.com/campaigns/kidney/updates/kidney-transplantation-across-the-globe>.
- 93 Global Observatory on Donation and Transplantation (GODT) data, produced by the World Health Organization-Organización Nacional de Trasplantes collaboration. <http://www.transplant-observatory.org/Pages/home.aspx>.
- 94 Ambagtsheer F, Weimar W. Organ Trade: Knowledge, Awareness, and Nonlegislative Responses. *Transplantation* 2016;100:5-6. doi:10.1097/TP.0000000000001063 PMID:26674729.
- 95 Ambagtsheer F, Weimar W, eds. *Trafficking in human beings for the purpose of organ removal: results and recommendations. The HOTT Project*. Pabst Science Publishers, 2016.
- 96 Steering Committee of the Istanbul Summit. Organ trafficking and transplant tourism and commercialism: the Declaration of Istanbul. *Lancet* 2008;372:5-6. doi:10.1016/S0140-6736(08)60967-8 PMID:18603141.

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>

Figures

Fig 1 Paired and pooled living kidney donation. Donor-recipient pairs who are immunologically incompatible and between whom a direct transplant is not viable are registered in the national scheme to achieve a compatible transplant match with other donor-recipient pairs. When two pairs are involved it is termed paired donation; pooled donation comprises more than two pairs. Donor-recipient pairs who have poor compatibility or substantial age-disparity and would like to achieve a better match can also register in this scheme.

DMSA = dimercaptosuccinic acid; CT = computed tomography;
MRI = magnetic resonance imaging; MDT = multidisciplinary team.

Fig 2 Steps in the evaluation of potential living kidney donors. Adapted with permission from Graham JM, Courtney AE. Oral presentation: 5 years of 1-days: Outcomes of potential living kidney donors undergoing a 1-day assessment pathway. British Transplantation Society Congress; Scottish Exhibition and Conference Centre, Glasgow 2016⁸⁴

Fig 3 Kidney transplants from living donors 2014 (per million population). Adapted with permission from International Registry in Organ Donation and Transplantation IRODaT Newsletter - Final numbers 2014. December 2015⁹¹

Fig 4 Kidney transplants in 2014 (or last year with data available) for WHO member states (per million population). DCD=donation after circulatory death. DBD=donation after brain death. *Data from 2013. **Data from 2012. ***Data from 2011. ****Data from 2010. Data source: WHO-ONT Global Observatory on Organ Donation and Transplantation Adapted with permission from Elsevier and Muller E, et al. Kidney transplantation across the globe: the good and bad. The Lancet Kidney Campaign. *Lancet*⁶²