Benn, M., Tybjaerg-Hansen, A., Davey Smith, G., & Nordestgaard, B. G. (2016). High body mass index and cancer risk: a Mendelian randomisation study. *European Journal of Epidemiology*, 31(9), 879-892. <https://doi.org/10.1007/s10654-016-0147-5>

Peer reviewed version

Link to published version (if available):
[10.1007/s10654-016-0147-5](https://doi.org/10.1007/s10654-016-0147-5)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Springer at <http://link.springer.com/article/10.1007%2Fs10654-016-0147-5>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

High body mass index and cancer risk
- A Mendelian randomisation study

Marianne Benn^{1,2,6,7}, Anne Tybjærg-Hansen^{2,3,6,7}, George Davey Smith⁴, and Børge G. Nordestgaard^{2,5,6,7}

¹Department of Clinical Biochemistry, Gentofte Hospital, Copenhagen University Hospital, Denmark; ²The Copenhagen General Population Study, Herlev Hospital, Copenhagen University Hospital, Denmark; ³Department of Clinical Biochemistry, Rigshospitalet, Copenhagen University Hospital, Denmark; ⁴ MRC Integrative Epidemiology Unit (IEU) at the University of Bristol, School of Social and Community Medicine, University of Bristol, United Kingdom; ⁵Department of Clinical Biochemistry, Herlev Hospital, Copenhagen University Hospital, Denmark; ⁶The Copenhagen City Heart Study, Frederiksberg Hospital, Copenhagen University Hospital; and ⁷The Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.

Supplementary Table 1. Baseline characteristics of participants by gender and body mass index category.

	Body mass index, kg/m ²					
	<18.5	18.5-24.9	25-29.9	≥30	P trend	All
Women						
No. of individuals (%)	976 (1.6%)	31,244 (52.3%)	18,646 (31.2%)	8,906 (14.9%)		59,772 (100%)
Body mass index, kg/m ²	17.9 (17.3-18.3)	22.5 (21.2-23.7)	26.9 (25.9-28.2)	32.7 (31.1-35.4)		24.6 (22.3-27.8)
Age, years	54.0 (42.7-64.0)	53.4 (44.4-63.5)	58.9 (49.0-67.6)	59.2 (49.6-67.3)	<0.001	56.0 (46.3-65.5)
C-reactive protein, mg/L	1.1 (0.7-1.7)	1.3 (0.9-1.8)	1.6 (1.2-2.7)	2.6 (1.6-4.8)	<0.001	1.5 (1.0-2.5)
Current smokers	53%	73%	78%	79%	<0.001	75%
Smoking, pack-years (smokers only)	18 (10-30)	18 (9.6-30)	20 (10-32)	22 (11-35)	<0.001	19 (10-30)
Physical activity						
Low	58%	44%	47%	51%	<0.001	46%
Intermediate	36%	48%	46%	44%	<0.001	47%
High	7%	8%	6%	5%	<0.001	7%
Alcohol consumption, units per week	3 (0-8)	5 (2-10)	5 (2-10)	3 (1-8)	<0.001	5 (2-10)
Education						
Less than 10 years	23%	21%	31%	36%	<0.001	26%
10 to 13 years	45%	55%	51%	46%	<0.001	52%
13 years or more	16%	18%	13%	11%	<0.001	15%
Postmenopausal	62%	57%	70%	72%	<0.001	63%

Men

No. of individuals (%)	182 (0.4%)	17,143 (35.0%)	23,595 (48.1%)	8,120 (16.6%)		49,040 (100%)
Body mass index, kg/m ²	17.9 (17.4-18.3)	23.4 (22.2-24.2)	27.1 (26.0-28.3)	32.0 (30.8-33.9)		26.2 (24.1-28.7)
Age, years	54.0 (41.3-63.0)	54.0 (44.0-65.0)	57.8 (48.0-67.0)	58.9 (49.4-66.7)	<0.001	56.7 (46.9-66.2)
C-reactive protein, mg/L	1.3 (0.9-2.2)	1.2 (0.8-1.8)	1.5 (1.0-2.3)	2.0 (1.3-3.4)	<0.001	1.4 (1.0-2.3)
Current smokers	36%	66%	74%	76%	<0.001	71%
Smoking, pack-years (smokers only)	30 (20-41)	26 (15-40)	30 (16-44)	31 (18-48)	<0.001	29 (15-43)
Physical activity						
Low	70%	48%	45%	46%	<0.001	46%
Intermediate	23%	42%	44%	43%	<0.001	43%
High	8%	10%	11%	11%	<0.001	10%
Alcohol consumption, units per week	5.5 (0-16)	9 (4-18)	11 (4-20)	10 (3-20)	<0.001	10 (4-19)
Education						
Less than 10 years	19%	22%	29%	38%	<0.001	28%
10 to 13 years	32%	51%	49%	44%	<0.001	49%
13 years or more	20%	17%	15%	11%	<0.001	15%

Continuous values are median and interquartile range and categorical values are in percent. 1 unit alcohol=12g.

Supplementary Figure 1.

Distribution of body mass index in the general population (upper panel), the Copenhagen General Population Study and The Copenhagen City Heart Study, and the cumulative distribution function of body mass index (lower panel) showing the probability of a body mass index (y-axis) less than or equal to x as a function of number of body mass index elevating alleles. BMI=body mass index.

Supplementary Figure 2.

Associations of potential confounders with body mass index, allele score, and cancer. Potential confounders were dichotomized: age (≥ 55 years versus < 55 years), gender (men versus women), C-reactive protein concentrations (≥ 2 mg/L versus < 2 mg/L), smoking (smoking versus non-smoking), physical activity (intermediate and high versus low), alcohol consumption ($> 14/21$ versus $\leq 14/21$ units per week for women/men), education (≥ 10 years versus < 10 years education), and for women menopausal status (postmenopausal versus premenopausal). For each potential confounder, logistic regression analysis was used to calculate age and gender-adjusted odds ratios and p-values for, respectively, a 1 kg/m² increase in observational body mass index, a 1 unit increase in allele score, and for presence versus absence of cancer. BMI=body mass index, CRP=C-reactive protein.

Supplementary Figure 3A.

Supplementary Figure 3B.

Risk of cancer for a 10kg/m² higher, respectively, observational and causal genetically determined body mass index in all participants, and women and men separately. The hazard ratio for a 10 kg/m² higher observational body mass index was calculated using Cox regression, while the odds ratio for genetically elevated body mass index was derived from an instrumental variable analysis. P-values

are, respectively, significance of hazard ratios and odds ratios. Power denotes the odds ratio that can be excluded at a two-sided α of 0.05 and β of 80%.

CI=confidence interval, BMI=body mass index.

Supplementary Figure 4.

Risk of cancer for a 10kg/m² higher, respectively, observational and causal genetically determined body mass index in participants. The odds ratio for genetically elevated body mass index was derived from an instrumental variable analysis using a weighted sum of risk alleles across the five

genotypes applying a weight equal to the effect of each variant on body mass index reported by a previous genome wide association study[1]. The hazard ratio for a 10 kg/m² higher observational body mass index was calculated using Cox regression. P-values are, respectively, significance of hazard ratios and odds ratios. CI=confidence interval, BMI=body mass index.

Supplementary Figure 5.

Risk of cancer for a 10kg/m² higher, respectively, observational and causal genetically determined body mass index in participants. The odds ratio for genetically elevated body mass index was estimated using an allele score combining the *FTO*(rs9939609), *MC4R*(rs17782313),

TMEM18(rs6548238), and *GNPDA2*(rs10938397) genotypes and thus omitting the *BDNF*(rs10767664) genotype. This was done as a sensitivity analysis, since it has been suggested that this genotype may have a pleiotropic effect on smoking so that the allele associated with a high body mass index is associated with a higher propensity for smoking. The hazard ratio for a 10 kg/m² higher observational body mass index was calculated using Cox regression. P-values are, respectively, significance of hazard ratios and odds ratios. CI=confidence interval, BMI=body mass index.

References

1. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, Lango AH, Lindgren CM, Luan J, Magi R, Randall JC, Vedantam S, Winkler TW, Qi L, Workalemahu T, Heid IM, Steinthorsdottir V, Stringham HM, Weedon MN, Wheeler E, Wood AR, Ferreira T, Weyant RJ, Segre AV, Estrada K, Liang L, Nemesh J, Park JH, Gustafsson S, Kilpelainen TO, Yang J, Bouatia-Naji N, Esko T, Feitosa MF, Kutalik Z, Mangino M, Raychaudhuri S, Scherag A, Smith AV, Welch R, Zhao JH, Aben KK, Absher DM, Amin N, Dixon AL, Fisher E, Glazer NL, Goddard ME, Heard-Costa NL, Hoesel V, Hottenga JJ, Johansson A, Johnson T, Ketkar S, Lamina C, Li S, Moffatt MF, Myers RH, Narisu N, Perry JR, Peters MJ, Preuss M, Ripatti S, Rivadeneira F, Sandholt C, Scott LJ, Timpson NJ, Tyrer JP, van WS, Watanabe RM, White CC, Wiklund F, Barlassina C, Chasman DI, Cooper MN, Jansson JO, Lawrence RW, Pellikka N, Prokopenko I, Shi J, Thiering E, Alavere H, Alibrandi MT, Almgren P, Arnold AM, Aspelund T, Atwood LD, Balkau B, Balmforth AJ, Bennett AJ, Ben-Shlomo Y, Bergman RN, Bergmann S, Biebertmann H, Blakemore AI, Boes T, Bonnycastle LL, Bornstein SR, Brown MJ, Buchanan TA, Busonero F, Campbell H, Cappuccio FP, Cavalcanti-Proenca C, Chen YD, Chen CM, Chines PS, Clarke R, Coin L, Connell J, Day IN, den HM, Duan J, Ebrahim S, Elliott P, Elosua R, Eiriksdottir G, Erdos MR, Eriksson JG, Facheris MF, Felix SB, Fischer-Posovszky P, Folsom AR, Friedrich N, Freimer NB, Fu M, Gaget S, Gejman PV, Geus EJ, Gieger C, Gjesing AP, Goel A, Goyette P, Grallert H, Grassler J, Greenawalt DM, Groves CJ, Gudnason V, Guiducci C, Hartikainen AL, Hassanali N, Hall AS, Havulinna AS, Hayward C, Heath AC, Hengstenberg C, Hicks AA, Hinney A, Hofman A, Homuth G, Hui J, Igl W, Iribarren C, Isomaa B, Jacobs KB, Jarick I, Jewell E, John U, Jorgensen T, Jousilahti P, Jula A, Kaakinen M, Kajantie E, Kaplan LM, Kathiresan S, Kettunen J, Kinnunen L, Knowles JW, Kolcic I, Konig IR, Koskinen S, Kovacs P, Kuusisto J, Kraft P, Kvaloy K, Laitinen J, Lantieri O, Lanzani C, Launer LJ, Lecoeur C, Lehtimäki T, Lettre G, Liu J, Lokki ML, Lorentzon M, Luben RN, Ludwig B, Manunta P, Marek D, Marre

M, Martin NG, McArdle WL, McCarthy A, McKnight B, Meitinger T, Melander O, Meyre D, Midthjell K, Montgomery GW, Morken MA, Morris AP, Mulic R, Ngwa JS, Nelis M, Neville MJ, Nyholt DR, O'Donnell CJ, O'Rahilly S, Ong KK, Oostra B, Pare G, Parker AN, Perola M, Pichler I, Pietilainen KH, Platou CG, Polasek O, Pouta A, Rafelt S, Raitakari O, Rayner NW, Ridderstrale M, Rief W, Ruukonen A, Robertson NR, Rzehak P, Salomaa V, Sanders AR, Sandhu MS, Sanna S, Saramies J, Savolainen MJ, Scherag S, Schipf S, Schreiber S, Schunkert H, Silander K, Sinisalo J, Siscovick DS, Smit JH, Soranzo N, Sovio U, Stephens J, Surakka I, Swift AJ, Tammesoo ML, Tardif JC, Teder-Laving M, Teslovich TM, Thompson JR, Thomson B, Tonjes A, Tuomi T, van Meurs JB, van Ommen GJ. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 2010;**42**(11):937-948.