Chu, D., Yu, J., Tong, J., Benton, M. J., Song, H., Huang, Y., Song, T., & Tian, L. (2016). Biostratigraphic correlation and mass extinction during the Permian-Triassic transition in terrestrial-marine siliciclastic settings of South China. *Global and Planetary Change*, 146, 67–88. <https://doi.org/10.1016/j.gloplacha.2016.09.009>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.gloplacha.2016.09.009](https://doi.org/10.1016/j.gloplacha.2016.09.009)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <http://www.sciencedirect.com/science/article/pii/S0921818116300182>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Biostratigraphic correlation and mass extinction during the Permian-Triassic transition in terrestrial-marine siliciclastic settings of South China

Daoliang Chu^a, Jianxin Yu^{a,*}, Jinnan Tong^{a,*}, Michael J. Benton^b, Haijun Song^a, Yunfei Huang^c, Ting Song^a, Li Tian^a

^aState Key Laboratory of Biogeology and Environmental Geology, School of Earth Sciences, China University of Geosciences, Wuhan 430074, China

^bSchool of Earth Sciences, University of Bristol, Bristol, BS8 1RJ, UK

^cSchool of Geosciences, Yangtze University, Wuhan 430010, China

*Correspondence to: J.X. Yu and J.N. Tong, State Key Laboratory of Biogeology and Environmental Geology, School of Earth Sciences, China University of Geosciences, Wuhan, 430074, China.

E-mail addresses: yujianxin@cug.edu.cn (J.X. Yu); jntong@cug.edu.cn (J.N. Tong).

Abstract

The Permian-Triassic boundary marks the greatest mass extinction during the Phanerozoic, which was coupled with major global environmental changes, and is known especially from well-preserved marine fossil records and continuous carbonate deposits. However, the placement of the Permian-Triassic boundary in terrestrial sections and accurate correlation with the marine strata are difficult due to the absence of the key marine index fossils in terrestrial-marine siliciclastic settings. Here, we present detailed fossil data from four terrestrial sections, two paralic sections and one shallow marine section in South China. Our data show that the rapid mass disappearance of the *Gigantopteris* flora in various sections represents the end-Permian mass extinction and the base of the Permian-Triassic transitional beds in terrestrial-marine siliciclastic settings of South China. In

particular, we find a mixed marine and terrestrial biota from the coastal transitional sections of the Permian-Triassic transitional Kayitou Formation, which provides a unique intermediate link for biostratigraphic correlation between terrestrial and marine sequences. Accordingly, the *Euestheria gutta*-bearing conchostracan fauna and the *Pteria ussurica variabilis*-*Towapteria scythica*-*Eumorphotis venetiana* bivalve assemblage are proposed as markers of the Permian-Triassic transitional beds in terrestrial-marine siliciclastic settings of South China.

Keywords: Terrestrial facies; Permian-Triassic transitional beds; Biostratigraphic correlation; South China

1. Introduction

The Permian-Triassic (P-Tr) mass extinction event, the greatest such event in the Phanerozoic, eliminated more than 90% of marine species and up to 70% of terrestrial vertebrate species, entirely restructuring ecosystems (Erwin, 1993; Benton and Twitchett, 2003; Song et al., 2013; Benton, 2015). Further, major changes in plant communities have also been documented (Retallack, 1995; Looy et al., 1999, 2001; Hermann et al., 2011; Schneebeil-Hermann et al., 2015; Yu et al., 2015; Zhang et al., 2016). A large number of studies have proposed environmental stressors as potentially responsible for the unusual and extreme conditions during the P-Tr crisis and its aftermath, such as global warming, anoxia, ocean acidification, wildfire, enhanced terrestrial weathering, rapid sea level changes and others (Wignall and Twitchett, 1996; Isozaki, 1997; Benton and Twitchett, 2003; S.Z. Shen et al., 2006, 2011; Wu et al., 2006; Payne et al., 2007; Algeo and Twitchett, 2010; Algeo et al., 2011; Clapham and Payne, 2011; W.J. Shen et al., 2011; Hinojosa et al., 2012; Joachimski et al., 2012; Sun et al., 2012; Benton and Newell, 2014; Sedlacek et al., 2014; Tian et al., 2014; Yin et al.,

2014; Clarkson et al., 2015; Ikeda et al., 2015; Song et al., 2015; Rey et al., 2016). Biotic changes associated with the extinction have been documented in detail, including the Lilliput effect (Urbanek, 1993; Twitchett, 2007) in most marine organisms (Hayami, 1997; Fraiser and Bottjer, 2004; Chen et al., 2005; Payne, 2005; Twitchett et al., 2005, 2007; He et al., 2007, 2010, 2014; Luo et al., 2008; McGowan et al., 2009; Mutter and Neuman, 2009; Song et al., 2011; Forel et al., 2015; Romano et al., 2016) as well as in some terrestrial/freshwater organisms (Tverdokhlebov et al., 2002; Huttenlocker, 2014; Chu et al., 2015; Romano et al., 2016), cyanobacterial and microbialite blooms (Kershaw et al., 2002, 2012; Wang et al., 2005; Cao et al., 2009; Xie et al., 2010; Mata and Bottjer, 2012; Wu et al., 2014; Chu et al., 2015), and bursts of low-diversity, opportunistic, and cosmopolitan taxa (Schubert and Bottjer, 1992; Shen et al., 1995; Rodland and Bottjer, 2001; Peng et al., 2007; Huang and Tong, 2014). Recent reviews indicate that tetrapods on land suffered a massive loss during the P-Tr crisis, as severe as that for life in the oceans, but the extinction seems to have been less profound for plants and insects (Benton and Newell, 2014). Some works also suggested that the destruction of terrestrial ecosystems happened simultaneously with the decline in marine diversity during the P-Tr transition (Twitchett et al., 2001; S.Z. Shen et al., 2011; Metcalfe et al., 2015; Zhang et al., 2016). However, less is known about the specific changes in terrestrial ecosystems during the P-Tr transition. In fact, stratigraphic investigation of the terrestrial P-Tr boundary (PTB) succession lags behind that of marine environments because of more variable sedimentary facies and lower preservation of fossils (Peng and Shi, 2009; Benton and Newell, 2014).

Most work on the biostratigraphy of the terrestrial P-Tr transition has focused on vertebrates, palaeobotany and palynology, with key sections located in China, Russia, Central Europe, South Africa, Australia and Antarctica (Benton, 1985; Retallack, 1995, 2013; Retallack et al., 1996, 2003,

2011; Ward et al., 2000, 2005; Benton et al., 2004; Grauvogel-Stamm and Ash, 2005; Peng and Shi, 2009; Retallack, 2013; Benton and Newell, 2014; Benton, 2015; Chu et al., 2015; Yu et al., 2015; Zhang et al., 2016; Pereira et al., 2016). However, the placement of the PTB in terrestrial sections and accurate correlation with marine strata are difficult because typical marine index fossils, such as conodonts and ammonoids, are absent in terrestrial facies. So far, the methods of magnetostratigraphy, sequence stratigraphy, cyclostratigraphy, geochemistry and isotopic dating have been employed in the terrestrial PTB and Lower Triassic studies (Aigner and Bachmann, 1992; Szurlies et al., 2003, 2012; Bachmann and Kozur, 2004; Cao et al., 2008; Taylor et al., 2009; S.Z. Shen et al., 2011; Cui et al., 2015; Gastaldo et al., 2015; Metcalfe et al., 2015; Zhang et al., 2016). In addition, an eventostratigraphic correlation of the PTB successions has been adopted in terrestrial deposits using the mass extinction or environmental event index itself as a marker (Peng et al., 2001; Cao et al., 2008; Peng and Shi, 2009; Chu et al., 2015; Zhang et al., 2016), but such an approach is far from ideal and the reliability of such a correlation must be proved. High-precision age-dating associated with multidisciplinary data, especially biostratigraphy, would be the most effective method for the correlation of the PTB from the marine to terrestrial facies (S.Z. Shen et al., 2011; Gastaldo et al., 2015; Metcalfe et al., 2015).

Among all the terrestrial P-Tr transitional strata around the world, the joint area of western Guizhou and northeastern Yunnan in southwest China has been subject to much recent attention, being regarded as an ideal place to investigate the correlation between terrestrial and marine P-Tr sequences because it contains good outcrops of shallow marine, marginal or coastal marine, and terrestrial PTB sections, all in close geographic proximity (Wang and Yin, 2001; Peng et al., 2005; Yu et al., 2007, 2015; Peng and Shi, 2009; S.Z. Shen et al., 2011; Bercovici et al., 2015; Cui et al.,

2015; Zhang et al., 2016). Thus far, the Kayitou Formation has been regarded as comprising P-Tr transitional beds (PTTB), which record a typical P-Tr mixed biota and the process of terrestrial mass extinction, though the placement of the PTB within this succession is still controversial (Yu et al., 2007, 2015; S.Z. Shen et al., 2011; Zhang et al., 2016).

Here, we document new fossil data from four terrestrial sections, the Guanbachong section in northeastern Yunan, and the Chahe, Jiuchaichong and Xiaohebian sections in western Guizhou; two coastal transitional sections, the Jinzhong section in western Guizhou, and the Jijibang section in southern Sichuan; and one shallow marine section, the Wadu section in western Guizhou (Fig. 1). The PTB of these sections have been extensively studied to retrieve fossil data in order to establish a correlation among the various facies. A mixed marine and terrestrial biota from the Kayitou Formation of the paralic Jinzhong and Jijibang sections has been recognized, which provides a unique link from the marine to terrestrial facies. Moreover, seven P-Tr sections have been studied in order to understand the biofacies of transitional shallow marine to terrestrial settings

2. Materials and methods

We measured seven stratigraphic sections and collected over 1550 fossils, including bivalves, brachiopods, conchostracans and plants from the Xuanwei, Kayitou, Dongchuan and Feixianguan formations of the studied sections. Fossils were carefully hunted bed by bed throughout the stratigraphic sequences and all were recorded with their bed number and thickness level. All identified fossils were numbered and photographed. Detailed lithological logs of the seven sections are provided to mark the positions of the fossils recovered and support the correlation. Plant and bivalve fossils were photographed using a Canon EOS 7D digital camera, while the conchostracan

specimens were examined and photographed using a LEICA-DM-750P microscope equipped with an automatic camera stack-image system. Ultramicroscopic study of the conchostracan specimens was carried out under a Quanta-200 SEM. All of the fossil material in this study is stored in the palaeontological collection of the State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan (BGEG). The catalogue numbers of the studied specimens include the profile name of the section and specimen number, followed by the storage location. The Guanbachong, Chahe, Xiaohebian, Jiuchaichong, Jinzhong, Jinjibang and Wadu sections are abbreviated as GBC, CH, XHB, JCC, JZ, JJB and WD, respectively.

3. General lithostratigraphy

The Sichuan-Yunnan old landmass is recognized in southwestern China during the P-Tr interval (Fig. 1). Eastwards from this old landmass, an almost continuous series of P-Tr sections records a gradual change from terrestrial to marine facies via the terrestrial-marine siliciclastic facies in the adjoining area between western Guizhou and eastern Yunnan (WGEY). The terrestrial P-Tr succession in WGEY consists of three main formations: the Xuanwei, Kayitou and Dongchuan formations, in ascending order (Fig. 2). Of these, the Xuanwei Formation consists mainly of siliciclastic sandstone deposits with intercalations of organic-rich mudstones and coal beds or seams that lie unconformably on the Emeishan basalts and are of latest Permian Wuchiapingian and Changhsingian age (He, 2007). The key characteristic of the Xuanwei Formation in South China is the presence of a rich *Gigantopteris* flora and coal beds or seams (Yu et al., 2007). The Kayitou Formation is similar to the underlying Xuanwei Formation, mainly comprising a 10- to 70-m-thick succession of mudstone, shale and silty sandstone. The difference between the Kayitou and Xuanwei

formations is that the Kayitou Formation contains no coal beds or seams. The rocks of the Kayitou Formation, in ascending order, are yellowish green sandstone, siltstone, claystone and shale in the lower part, brownish-yellow siltstone, claystone and shale in the middle part, and purple-red siltstone increasing in proportion upwards in the upper part. The Dongchuan Formation is characterized by a thick succession (over 800 m thick) of purple-red siliciclastic sandstones, siltstones and mudstones, but contains some thin beds of marine muddy limestone at the top. In the eastern part of the area the lower lithostratigraphical unit is also the Xuanwei Formation, while the Kayitou and Dongchuan formations are laterally replaced by the Feixianguan Formation, comprising marine siliciclastic sandstones, mudstones and limestones.

4. Biostratigraphy of the studied sections

4.1. Guanbachong section

The Guanbachong section, located in Leju Country, Zhaotong City, Yunnan Province (Fig. 1), records a succession comprising terrestrial Late Permian to Early Triassic siliciclastic deposits of the Xuanwei, Kayitou and Dongchuan formations. For biostratigraphic investigation, well-preserved Late Permian or Palaeozoic-type plants with large leaves and entire margins were collected in the Xuanwei Formation and the lower part of the Kayitou Formation (Fig. 3), such as *Gigantopteris*, *Gigantonoclea*, *Lobatannularia*, *Annularia*, *Compsopteris*, *Paracalamites*, *Fascipteris*, and *Pecopteris*. Both the data of our palaeobotanical collection and the report by S.Z. Shen et al. (2011) indicate a dramatic reduction in the diversity and abundance of the *Gigantopteris* flora in the lower part of the Kayitou Formation of the Guanbachong section (Fig. 3). In addition, a negative organic carbon isotope anomaly occurs in the Kayitou Formation, only slightly younger than the

disappearance of the *Gigantopteris* flora (Zhang et al., 2016). The loss of the *Gigantopteris* flora and the organic carbon isotope negative shift represent the P-Tr mass extinction of plants as well as the collapse of the tropical rainforest ecosystem.

The PTB succession in the Kayitou Formation consists mainly of brownish-yellow siltstone, mudstone and purple-red siltstones, containing relatively few small conchostracans and plants. About 30 conchostracan specimens with small elliptical carapaces less than 3.5 mm were collected in the upper part of the Kayitou Formation, approximately 2 m below the top, and they are attributed to the genus *Euestheria* (Fig. 4). The carapace of *Euestheria* is characterized by a very small, strongly convex umbo and dense, narrow growth bands; in particular, well-preserved specimens show an indistinct, regular and very small reticular ($< 10 \mu\text{m}$) microsculpture (Fig. 4-15). In addition, plant fossils assigned to the genus *Peltaspermum* were found in association with the conchostracans (Fig. 3). The genus *Peltaspermum* was present in the Early Permian in Northwest China and has also been recorded in the Sunjiagou and Sunan formations of the Late Permian and in Early and Middle Triassic rocks in North China (Wang and Wang, 1990; Huang, 1996; Huang and Ding, 1998). Accordingly, the presence of the genus *Peltaspermum* in the Kayitou Formation indicates that this genus survived the P-Tr mass extinction (Yu et al., 2015). The immediately overlying deposits are largely barren of fossils, and only a few conchostracans and plant fragments were discovered.

4.2. Chahe section

The Chahe section, located near Chahe Town, Weining County, Guizhou Province (Fig. 1) is a classical terrestrial P-Tr section. It has received much recent attention and has been subjected to many high-resolution stratigraphic methods, including biostratigraphy, eventostratigraphy,

chronostratigraphy and chemostratigraphy, because of its continuous exposure and well-preserved
 fossils (Peng et al., 2005; Yu et al., 2007, 2015; Peng and Shi, 2009; S.Z. Shen et al., 2011; Cui et al.,
 2015; Bercovici et al., 2015; Zhang et al., 2016). The *Gigantopteris* flora consists of more than 30
 genera and 81 species in the Xuanwei Formation in the Chahe section, dominated by hygrophilous
 and thermophilous plant groups such as pectopterids, gigantopterids, Lycopsiales, and Equisetales
 (Fig. 5; Yu, 2008; Yu et al., 2008, 2015). In addition, further palaeobotanical study indicates a
 dramatic reduction in the diversity and abundance of the *Gigantopteris* flora in the topmost Bed 68
 (Fig. 6), a less than 3-cm-thick coal seam mixed with an ash bed, which is nearly consistent in age
 with Bed 25 in the Meishan section (252.30 ± 0.07 Ma, major mass extinction event horizon; S.Z.
 Shen et al., 2011). During our study of the Kayitou Formation in the Chahe section, we found 14
 genera and 20 species of plant fossils belonging to the *Gigantopteris* flora, including a certain
 number of large leaves with entire margins in Bed 69 and the base of Bed 70, such as *Gigantopteris*,
Gigantonoclea, *Pecopteris*, *Fascipteris*, *Compsopteris*, and *Lobatannularia*, which represent the last
 occurrence (LO) of the *Gigantopteris* flora at this site (Fig. 7).

In the overlying section, plant fossils surviving from the Late Permian disappear above Bed 71,
 whilst plants of the genus *Peltaspermum* occur in association with conchostracans in the
 middle-upper part of Bed 71, which is a conchostracan-rich horizon similar to that in the
 Guanbachong section (Fig. 6). A total of 45 conchostracan specimens were collected in an
 approximately 8-m-thick interval between Bed 71 and Bed 78, including the genera *Euestheria* and
Palaeolimnadia, similar to the assemblage from the Guanbachong section; *Palaeolimnadia* also have
 small-sized rounded carapaces but with low convexity and a smooth unsculptured surface (Chu et al.,
 2013). In the red-purple Dongchuan Formation, except in the basal part, neither conchostracans nor

any other fossils have been observed, but a few plant fragments.

4.3. Xiaohebian section

The Xiaohebian section is situated near Xiaohebian Village of Halahe Town, Guizhou Province (Fig. 1). It consists of a well-developed upper Permian coal-bearing exposure of the Xuanwei Formation, the overlying PTB succession of the brownish-yellow Kayitou Formation, and the red-purple Dongchuan Formation. In our study, an abundant *Gigantopteris* flora was recovered from the Xuanwei Formation below the last ash bed (Bed 8), consisting of *Gigantopteris*, *Gigantonoclea*, *Fascipteris*, *Pecopteris*, etc. (Fig. 8). A dramatic reduction in diversity and abundance of the *Gigantopteris* flora occurs at the top of Bed 8, although a few plant fragments belonging to the this flora are occasionally observed in the overlying strata (Fig. 8). Considering the similarity of the *Gigantopteris* flora with that in the Chahe section, the isotopic age of the youngest ash bed (Bed 8), above a 3-cm-thick coal seam, may be nearly consistent with Bed 68 of the Chahe section as well as Bed 25 of the Meishan section (S.Z. Shen et al., 2011), also representing the mass extinction horizon.

Plant fossils of the *Gigantopteris* flora disappear above Bed 9, while *Peltaspermum* and abundant conchostracans were collected in the interval between Bed 10 and the top of the Kayitou Formation (Fig. 8). More than 85 conchostracan specimens were collected from Beds 10 to 15, including *Euestheria gutta*, *Palaeolimnadia xuanweiensis* and *Euestheria* sp.; 65 specimens were identified as belonging to the genus *Euestheria* (Fig. 4).

4.4. Jiucaichong section

The Jiucaichong section is situated near Jiucaichong Village of Heishitou Town, Guizhou

Province (Fig. 1), and exposes a continuous outcrop of the upper part of the Xuanwei Formation, the Kayitou Formation and the lower part of the Dongchuan Formation. Similarly, several well-preserved plants of the *Gigantopteris* flora were collected in the Xuanwei Formation under the last appearance datum (LAD) of coal (Bed 23, Fig. 9). Above the LAD of coal, nine large-leaved plant species belonging to the *Gigantopteris* flora were found at the base of the Kayitou Formation, representing the LO of the *Gigantopteris* flora. In the overlying strata, only fragmentary *Gigantopteris* flora plant fossils in association with *Peltaspermum* were collected.

The dramatic mass extinction of the *Gigantopteris* flora is recorded at the base of the Kayitou Formation (Fig. 9). The strongly contrasting conchostracan fauna appears in the lower part of this formation; over 100 specimens were collected, and *Euestheria gutta* dominates (Fig. 4). Neither the plant nor conchostracan fossils were found in the overlying strata.

4.5. Jinzhong section

The Jinzhong section, located 15 km south of Jinzhong Town, Guizhou Province (Fig. 1), comprises a continuously exposed outcrop of the upper part of the Xuanwei Formation, the Kayitou Formation and the lower part of the Dongchuan Formation. The successions within the Xuanwei Formation were deposited chiefly in the alluvial-plain to fluvial and lacustrine facies. The *Gigantopteris* flora is commonly observed below the LAD of coal (Bed 7) in the Xuanwei Formation (Fig. 10). Above Bed 7, the diversity of the *Gigantopteris* flora decreases sharply, and only a few plant fragments were collected at the base of the Kayitou Formation in our study (Fig. 10).

The P-Tr transition is represented by the 65-m-thick Kayitou Formation, which corresponds to a shallow coastal lagoon evolving towards coastal transitional deposition, and has yielded a mixed

terrestrial-marine biota including plants, conchostracans, bivalves, brachiopods (lingulids), gastropods and insects (Fig. 11). From the base of Bed 9 to the top of the Kayitou Formation, we found the plant fossils *Annalepis* and *Peltaspermum* in abundance (Fig. 10). *Annalepis* is a Triassic lycopsid related to *Isoetes*, with similar growth habits to *Pleuromeia* (Grauvogel-Stamm and Düringer, 1983; Grauvogel-Stamm and Lugardon, 2001; Yu et al., 2010; Naugolnykh, 2013), and is regarded as an important marker fossil for identifying deposits of late Early and Middle Triassic age (Grauvogel-Stamm, 1999; Grauvogel-Stamm and Ash, 2005). Previous work in an adjacent area (Yu et al., 2010, 2015; Bercovici et al., 2015) suggested that the *Annalepis-Peltaspermum* assemblage represents the beginning of a new terrestrial flora after the disappearance of the Permian flora following the P-Tr mass extinction, as it contrasts significantly with the *Gigantopteris* flora recovered from the underlying Xuanwei Formation.

In addition, conchostracans are common at this locality and are very abundant on some bedding surfaces of the brownish-yellow mudstones in the Kayitou Formation. Over 200 collected conchostracan specimens are preserved as detailed impressions, commonly with a complete and lightly mineralized chitin carapace; these were identified as belonging to two genera, *Euestheria* and *Palaeolimnadia*, similar to the conchostracans collected from the aforementioned four sections (Fig. 4). Of these, *Euestheria gutta* was most abundant in all of the various levels collected.

Like the characteristically mixed terrestrial-marine biotic communities of the Kayitou Formation in this same facies, the marine fauna, including bivalves and brachiopods (lingulids), also occupies an important position. Several distinct marine bivalve species were recovered from the lower part of the Kayitou Formation (beds 9-13) and were identified as *Pteria ussurica variabilis*, *Neoschizodus orbicularis*, *Promyalina schamarae*, *Neoschizodus laevigatus*, *Eumorphotis venetiana* and

265 *Permophorus bregeri* (Fig. 11). These bivalves are well preserved as both internal and external
 266 moulds in weathered shale layers, and several specimens were collected in association with the
 267 lingulids. Most of the bivalve specimens are small to medium sized, ranging from 6–12 mm in height
 268 and 4–15 mm in length, commonly slightly longer than high. This bivalve assemblage represents an
 269 Early Triassic fauna. Lingulid brachiopods are very abundant at this site, and most are preserved as
 270 complete small-sized internal and external moulds of disassociated valves ranging from 4–8 mm in
 271 height and 3–5 mm in length (Fig. 11), assigned to the genus *Sinolingularia*, by the evidence of the
 272 shape of the shell and the value of the width/length ratio (Peng and Shi, 2008). This lingulid fauna is
 273 characterized by low taxonomic diversity, but high abundance of individuals, and some lingulid
 274 specimens are preserved associated with the conchostracans and plant fossils (Fig. 11). Early Triassic
 275 lingulids were adapted to a wide range of environmental conditions, including variable salinity,
 276 dissolved oxygen, and pH, from high-latitude to low-latitude and from the coastal to the relatively
 277 deep sea environments, and have been reported world-wide, having survived the catastrophic event
 278 as a result of being environmentally opportunistic (Rodland and Bottjer, 2001; Peng and Shi, 2007).
 279 Rare monospecific and similarly mm-sized and trochospiral gastropods were collected in association
 280 with the bivalves, and these might belong to a recorded Early Triassic micro-gastropod form (Fraiser
 281 and Bottjer, 2004).

282

283 4.6. Jinjibang section

284 The Jinjibang section is located near Jinji village, approximately 5 km southwest of Zhenzhou
 285 town, Junlian County, Sichuan Province (Fig. 1), and exposes a continuous outcrop of the upper part
 286 of the Xuanwei Formation and the lower part of the Feixianguan Formation. Latest Permian

deposition is represented by the Xuanwei Formation, which is composed of siliciclastic sandstones, mudstones, claystones, coal beds or seams associated with a diverse *Gigantopteris* flora, marine bivalves and brachiopods (lingulids). In previous studies, the coal-bearing Xuanwei Formation was locally divided into two parts in this area: in ascending order, the Junlian and Jinjibang formations. The overlying lower part of the Feixianguan Formation is composed of yellowish-green siltstone, claystone and shale, which contain marine bivalves, brachiopods, conchostracans and plants. The PTB succession is the lower part of the Feixianguan Formation, which corresponds to a shallow coastal lagoon evolving towards a coastal transitional deposition.

The latest Permian megaflora recovered from the Xuanwei Formation consists of 45 genera and 83 species, and these are dominated by pteridosperms, Equisetales, lycophytes and Filicales, such as *Gigantopteris*, *Gigantonoclea*, *Compsopteris*, *Lobatannularia*, *Paracalamites*, *Lepidodendron*, *Lepidostrobohyllum*, *Pecopteris*, and *Fasciapteris*, which represent a tropical rainforest community (Fig. 12; Li et al., 1982; Zhu et al., 1984). Similar to other sections, the diversity of the end-Permian rainforest-type vegetation is drastically reduced at the top of the Xuanwei Formation, associated with the disappearance of coal seams (Fig. 12). The PTB succession is represented by the yellowish-green siltstone, claystone and shale deposits of the lower part of the Feixianguan Formation, less than 10 m thick, yielding a mixed terrestrial-marine biota including plants, conchostracans, bivalves, brachiopods (lingulids), and gastropods. The abundant and monogeneric *Annalepis* flora replaces the *Gigantopteris* flora in this interval, associated with low-diversity marine bivalves including *Pteria ussurica variabilis* and *Neoschizodus orbicularis* (Fig. 12). In addition, a few conchostracans were collected 2 m above the LO of the *Gigantopteris* flora and were identified as *Euestheria*, dominated by the species *Euestheria gutta* (Fig. 11).

4.7. Wadu section

The Wadu section, located south of Wadu village, Panxian County, Guizhou Province (Fig. 1), records the well-developed Xuanwei and Feixianguan formations. The Xuanwei Formation is composed of sandstones, siltstones, claystones and coal seams, and approximately 160 m of outcrop was measured. In ascending order, the Feixianguan Formation consists mainly of yellowish green siltstone, claystone and shale in the bottom 10 m, brownish-yellow sandstone, siltstone and shale in the > 30 m of the middle part, and more than 500 m of purple-red sandstone in the upper part. The yellowish green deposits at the bottom of the Feixianguan Formation at this site laterally correlate with strata of the Kayitou Formation and represent the PTB succession.

Among the fossils recovered from the Xuanwei Formation, an abundant *Gigantopteris* flora with large leaves and entire margins was collected below the LAD of coal (Bed 17), including *Gigantopteris*, *Gigantonoclea*, *Lobatannularia*, *Stigmaria*, *Annularia*, *Paracalamites*, *Fasciapteris*, and *Pecopteris* (Fig. 13). Only a few fragmentary plant remains were collected from Bed 18 (15-cm-thick claystones), and the *Gigantopteris* flora disappears above this level (Fig. 13). The marine bivalve fauna begins to appear at the base of Bed 19, coupled with lingulid brachiopods, including *Pteria ussurica variabilis*, *Promyalina schamarae*, *P. putiatinensis*, *P. spathi*, *Neoschizodus orbicularis*, *Astartella* sp., *Unionites fassaensis*, *Claraia bioni*, *Eumorphotis venetiana*, *Permophorus bregeri*, *Neoschizodus laevigatus*, *Towapteria scythica*, *Leptochondria virgalensis*, and *Bakevellia goldfussi* (Fig. 14). Obviously, the mean size of these bivalves is larger than the value measured from the last two sections because of the appearance of larger forms. In particular, the lingulid brachiopods with articulated valves are vertically positioned in the sandstone of Bed 19,

which may indicate that the fossils were preserved *in situ*. In addition, bivalves and lingulid brachiopods are common in the upper deposits.

5. Biostratigraphic correlation and the end-Permian mass extinction

5.1. Biostratigraphic correlation among different siliciclastic settings in South China

In the four continental sections described here, the fossil records reveal the same pattern of ecosystem evolution during the Permian-Triassic transitional interval, namely that the diversity of the *Gigantopteris* flora is drastically reduced in the lower part of the Kayitou Formation, representing the terrestrial end-Permian mass extinction in South China. The plants of the *Gigantopteris* flora, with large leaves and entire margins, appear at intervals of 2.2 m, 1 m, 2 m and 0.7 m above the occurrence of the last coal bed or seam in the Kayitou Formation, in the Guanbachong, Chahe, Xiaohebian and Jiucaichong sections respectively, and this variability of thickness to the last coal bed or seam across space might reflect the changes in the local depositional environment. The fragmentary remains of the *Gigantopteris* flora commonly appear in overlying beds associated with abundant and low-diversity peltaspermalean foliage in the Kayitou Formation (Fig. 15). The *Euestheria gutta*-bearing conchostracan fauna appears in the Kayitou Formation at 1.8 m, 2 m and 5 m above the occurrence of the last coal bed or seam, associated with peltaspermalean foliage and *Gigantopteris* flora remains. These characteristics of fossil distribution appear to be common in the terrestrial Permian-Triassic transitional Kayitou Formation in South China.

The mixed marine-terrestrial biota from the Kayitou Formation of the Jinzhong and Jinjibang sections contains a range of fossil groups, including plants, conchostracans, bivalves, brachiopods (lingulids), gastropods and insects. The distribution of the mixed assemblage is represented by the

following characteristics: the diversity of the *Gigantopteris* flora decreases significantly at the base of the Kayitou Formation or the top of the Xuanwei Formation, corresponding to the horizon of the terrestrial end-Permian mass extinction in South China; the *Euestheria gutta*-bearing conchostracan fauna is recovered from the Kayitou Formation at the Jinzhong and Jinjibang sections and is related to the same assemblage from the terrestrial sections; the Triassic-type plant *Annalepis* associated with the peltaspermalean flora first appears at the base of the Kayitou Formation; the marine bivalve assemblage associated with the *Euestheria gutta*-bearing conchostracan fauna is recovered from the Kayitou Formation and is dominated by *Pteria ussurica variabilis*, *Neoschizodus orbicularis*, *Promyalina schamarae*, *Neoschizodus laevigatus*, *Eumorphotis venetiana* and *Permophorus bregeri*, assigned to the *Pteria ussurica variabilis*-*Eumorphotis venetiana*-*Promyalina schamarae* assemblage. The mixed terrestrial-marine biota records both the process of the mass extinction and the biostratigraphic evidence of the correlation of the Permian-Triassic transitional interval between marine and terrestrial successions.

In the latest Permian Xuanwei Formation, the Wudu section comprises terrestrial-marine sediments, which contain abundant plant fossils of the *Gigantopteris* flora, and also records the mass extinction of plants at the top of the Xuanwei Formation associated with the LAD of the coal bed. Typical Permian-Triassic transitional marine siliciclastic deposits contain abundant bivalves, brachiopods (lingulids), and ostracods, all recovered from the Wudu section in our study. At the base of the Feixianguan Formation, the marine bivalve fauna is represented by the *Pteria ussurica variabilis*-*Towapteria scythica*-*Eumorphotis venetiana* assemblage. This bivalve fauna is widespread in shallow-water facies PTB sections in South China (Sheng et al., 1984, 1987; Shen et al., 1995; Chen et al., 2009; Huang and Tong, 2014).

Therefore, we have provided evidence that the same fossil assemblage and variation trends appear in both the Kayitou Formation in the terrestrial and coastal sections, and in the lower part of the Feixianguan Formation in the shallow marine section. We infer that these intervals were deposited simultaneously and should be considered transitional from the latest Permian to earliest Triassic based on the P-Tr mixed character of the various fossils from these PTTB (Fig. 16).

5.2. The end-Permian mass extinction in the studied area

It is important to focus on plants during mass extinction intervals in terrestrial successions because plants are some of the most important elements and the primary producers in the terrestrial ecosystem, and plant abundance and diversity are key indices of ecosystem functioning in both geological and modern times. The terrestrial mass extinction during the P-Tr transitional interval is commonly associated with a significant decrease in the diversity of plant fossils (Retallack, 1995; Looy et al., 1999, 2001; S.Z. Shen et al., 2011; Benton and Newell, 2014; Yu et al., 2015). Furthermore, the current study has revealed that the PTB crisis had a much more profound effect on vegetation than the other mass extinctions that not only impacted individual organisms but also whole families, and so it can really be regarded as being a global mass extinction event that widely affected both the marine and terrestrial ecosystems (Cascales-Miñana and Cleal, 2014; Cascales-Miñana et al., 2015). Previous studies have suggested that the main terrestrial P-Tr crisis occurred at the base of the Kayitou Formation in the area of western Guizhou and eastern Yunnan, and this is based on biostratigraphic, isotopic and geochronological data (Yu et al., 2007, 2015; Peng and Shi, 2009; S.Z. Shen et al., 2011). Recently, a credible multidisciplinary study suggested that the Kayitou Formation records the process of the terrestrial Permian-Triassic mass extinction (Zhang et

al., 2016). In our studied sections, the rainforest-type *Gigantopteris* flora is commonly observed and mostly well-preserved, with large leaves and entire margins in the coal-bearing Xuanwei Formation (Fig. 16). Our detailed investigation of the plant fossils indicates a dramatic reduction in the diversity and abundance of this flora at the base of the Kayitou Formation, a few metres above the LAD of coal beds, in the Guanbachong, Chahe, Xiaohedian, and Jiuchaichong sections (Fig. 16). In the Jinzhong, Jinjibang and Kele sections, the *Gigantopteris* flora disappears at the top of the Xuanwei Formation or the base of the Feixianguan Formation, which corresponds to the level of the extinction event in the four terrestrial sections (Fig. 16). The absence of coal and plant fossils in the immediately overlying deposits, associated with the negative shift of $^{13}\text{C}_{\text{org}}$ in the Guanbachong and Chahe sections, indicates the terrestrial end-Permian mass extinction and the collapse of the tropical rainforest ecosystem (S.Z. Shen et al., 2011; Zhang et al., 2016).

From the plant fossil records in the series of PTB sections in WGEY, we find that the diversity and abundance of the rainforest-type *Gigantopteris* flora experiences a sharp decrease at the base of the PTTB and shows a similar pattern among the studied sections that record the terrestrial end-Permian biotic crisis (Fig. 16). Considering the high-precision isotopic age from the Chahe section and the correlation among the continuous sequences, the time horizon of the biotic crisis in the studied area is simultaneous with the end-Permian mass extinction or the first stage of the Permian-Triassic mass extinction (Song et al., 2013) in marine facies.

Most of the Permian-type plant taxa were wiped out in the PTB crisis, with only a few relicts persisting into the PTTB. These relict taxa of the *Gigantopteris* flora, however, are usually represented by only a few fragmentary specimens from the lower part of the Kayitou Formation (Fig. 15), suggesting long-distance transport or reworking. In the aftermath of the main biotic crisis, the

Gigantopteris flora is replaced by a low-diversity opportunistic flora dominated by the genera *Annalepis* and *Peltaspermum*, which demonstrates stressed environmental conditions on land. Meanwhile, conchostracans appear in abundance a few metres above the main extinction horizon in the Kayitou Formation of the four terrestrial sections and the two coastal Jinzhong and Jinjibang sections, but disappear immediately at the horizon of the lower part of the overlying Dongchuan Formation. The disappearance of conchostracans was probably caused by dramatically shrinking niches as the regional facies turned from a shallow lake or brackish lagoon into braided rivers within an alluvial plain, between the Kayitou and Dongchuan formations in western Guizhou and eastern Yunnan (Bercovici et al., 2015).

This sedimentary switch from low-energy meandering streams to high-energy braided streams has been noted before in the terrestrial red bed successions of Russia (Newell et al., 1999), South Africa (Ward et al., 2000), and Spain (Arche and López-Gómez, 2005) at or close to the PTB. The switch had been explained by changes in regional tectonics or increased rainfall, but the most likely explanation is that stripping of forests from the land, presumably by acid rain and global warming, released soils and led to massive erosion of uplands onto alluvial plains. The timing of the sedimentary regime switch is interesting (Fig. 16), whether it was precisely at the level of the end-Permian plant extinction (EPPE) or estimated-PTB (e-PTB), and these South China sections provide better age control than was possible elsewhere.

6. The importance of bivalve and conchostracan faunas for the correlation of the terrestrial-marine siliciclastic PTB

Bivalves are very common in the marine record, especially in shallow-water siliciclastic settings.

More or less complete latest Permian to earliest Triassic successions with abundant bivalve fossils enhance our understanding of the evolution of bivalves and benthic ecosystems during this period in various areas, such as South China, the Southern Alps, Pakistan and western North America (Broglia-Loriga et al., 1986; Schubert and Bottjer, 1995; Fraiser and Bottjer, 2007; Wasmer et al., 2012; Huang and Tong et al., 2014). Bivalve assemblage zones are generally valuable for Triassic biostratigraphy and regional or global stratigraphic correlation (Gao et al., 2009; McRoberts, 2010; Huang and Tong, 2014). For PTB sections that lack conodonts and ammonoids, biostratigraphic subdivisions and correlations are often based on bivalve assemblages. From the end of the Late Palaeozoic to the Late Mesozoic, conchostracans have the highest biostratigraphic significance of all terrestrial fossils, and in some intervals (Kozur and Weems, 2010, 2011; Scholze et al., 2015), they play the same role as ammonoids and conodonts in pelagic marine beds. They are distributed across a wide range of ecological dimensions and geographic regions, as well as in a variety of facies, such as freshwater deposits, brackish deposits, deltaic marginal marine beds, and some bedding planes or brackish intervals of very shallow marine deposits, which offer the greatest potential for correlation with marine facies (Kozur and Weems, 2010). Here, we discuss the biostratigraphic significance of the specific bivalve and conchostracan faunas that are widespread in the P-Tr transition interval from our studied area and propose that they are of great use for the definition and correlation of the PTB in South China, and even globally.

6.1 The *Pteria*-*Towapteria*-*Eumorphotis* bivalve fauna

We collected abundant bivalve fossils from the Jinzhong, Jinjibang, Kele and Wadu sections, including *Pteria ussurica variabilis*, *Promyalina schamarae*, *P. putiatinensis*, *P. spathi*,

Neoschizodus orbicularis, *Astartella* sp., *Unionites fassaensis*, *Claraia bioni*, *Eumorphotis venetiana*, *Permophorus bregeri*, *Neoschizodus laevigatus*, *Towapteria scythica*, *Leptochondria virgalensis*, and *Bakevellia goldfussi* (Figs, 11, 14). The *Pteria-Towapteria-Eumorphotis* bivalve fauna is widespread in shallow-water facies of conodont-free PTB sections in South China. The bivalve fauna of the transitional beds of the basal Triassic is represented by the *Pteria ussurica variabilis-Towapteria scythica-Eumorphotis venetiana* assemblage in South China (Yin, 1985; Shen et al., 1995; Chen et al., 2009; Huang and Tong, 2014). Other studies also recognized it as corresponding to the mixed fauna in Bed 26 at the Meishan section (Sheng et al., 1984, 1987; Zhang et al., 2016), which is clearly of latest Permian age. In addition, this assemblage is also regarded as the Permian-Triassic transitional bivalve fauna that flourished in the extinction interval, typified by opportunistic survivor taxa (Fang, 2004). In any case, the *Pteria ussurica variabilis-Towapteria scythica-Eumorphotis venetiana* bivalve assemblage can be considered as the marker of the P-Tr transitional interval in conodont-barren PTB sections and the end-Permian mass extinction horizon in shallow, siliciclastic facies PTB sections.

6.2 The *Euestheria gutta*-bearing conchostracan fauna

Chu et al. (2013) reported the occurrence of the *Euestheria gutta*-bearing conchostracan fauna from the Kayitou Formation in three terrestrial sections of South China. Meanwhile, four other species of Chu et al. (2013) were assigned to the genus *Euestheria*, and six species were assigned to the genus *Palaeolimnadia*. Considering deformation issues and intraspecific variation, diversity at the species level might have been overestimated in this previous study. In any case, *Euestheria gutta* dominated in this conchostracan fauna in association with a few *Palaeolimnadia xuanweiensis*

specimens. Here, we collected abundant and well-preserved conchostracan fossils from the Guanbachong, Chahe, Xiaohebian, Jiucaichong, Jinzhong and Jinjibang sections (Fig. 4). *Euestheria gutta* was the dominant conchostracan fossil in each section. Previous studies have shown that *Euestheria gutta* was widespread in Early Triassic deposits from different areas, including the Calvörde Formation and Bernburg Formation of the Germanic Basin (Kozur and Seidel, 1983), the Rybinskian horizon on the Russian platform (Lutkevitch, 1937), and the Vokhma Formation in the Moscow Syncline (Scholze et al., 2015). The *Euestheria gutta*-bearing conchostracan fauna has been assigned an Early Triassic stratigraphic range in Central Russia and Germany, though the real lowest stratigraphic extent of this species seems to be less certainly known, due to the poorly-preserved specimens that were found in the uppermost Zechstein Group (Fulda Formation; Late Permian) of the Germanic Basin (Kozur and Weems, 2010).

6.3. The definition of the PTB in siliciclastic settings of South China

The end-Permian mass extinction horizon at the base or lower part of the Kayitou Formation in this area corresponds to a significant decrease in the diversity of plant fossils, which is nearly consistent in age with Bed 25 in the Meishan section from isotopic dating and biostratigraphic data in South China. In the present study, we observed almost the same pattern of biological evolution in the various sections, i.e. the *Gigantopteris* flora became extinct in the similar interval that contained a specific ash bed (Shen et al., 2011), the subsequent occurrence of the low-diversity opportunistic flora, the conchostracan fauna and mixed terrestrial-marine biota in the Kayitou Formation. Accordingly, we propose that the collapse of end-Permian tropical marine and non-marine ecosystems is synchronous, at least, in this studied area. The synchronous collapse of end-Permian

tropical ecosystems in this area is an effective marker for the correlation between non-marine and marine facies (Zhang et al., 2016). In addition, the occurrence of Triassic-type uniform conchostracan and bivalve faunas in the Kayitou Formation or the base of the Feixinguan Formation in the studied sections is the key basis for the biostratigraphic correlation, which indicates that the Kayitou Formation is latest Permian to earliest Triassic in age, and the PTB can be placed at a horizon within the Kayitou Formation (Fig. 16).

In fact, comprehensive biostratigraphic and geochemical data have been documented from various terrestrial sections in South China, suggesting that the Kayitou Formation is the witness of the terrestrial end-Permian mass extinction, mostly or entirely of latest Changhsingian age, especially from the evidence of the last appearance of plant remains and charcoal fossils (Zhang et al., 2016). In addition, a marine transgression beginning at the top of the Xuanwei Formation or the base of the Kayitou Formation was recognized based on a number of terrestrial and transitional sections in South China, and it was also well documented to be latest Changhsingian in age in the marine carbonate sections (this study; Yin et al., 2014; Zhang et al., 2016), which demonstrates that the base of the Kayitou Formation corresponds to the latest Changhsingian rather than the earliest Triassic. However, clear criteria are still lacking for the definition of the PTB in terrestrial sections stemming from uncertainty in identifying and using index fossils.

We suggest that the occurrence of the *Euestheria gutta*-bearing conchostracan fauna marks the PTB in terrestrial sections in South China, and that the mixed biota of the *Pteria ussurica variabilis*-*Towapteria scythica*-*Eumorphotis venetiana* bivalve assemblage coupled with the *Euestheria gutta*-bearing conchostracan fauna marks the PTB in coastal transitional sections. The *Pteria ussurica variabilis*-*Towapteria scythica*-*Eumorphotis venetiana* bivalve assemblage can be

considered as a marker of the PTB in shallow water conodont-barren PTB sections. All of the above fossil assemblages appear at different levels in the Kayitou Formation or the lower part of the Feixianguan Formation in the different sections; for example, the *Euestheria gutta*-bearing conchostracan fauna appears in the upper part of the Kayitou Formation in the Guanbachong section, while it appears in the lower middle part of Kayitou Formation in the Chahe section, indicating time-transgressive formation boundaries (Fig. 16).

7. Conclusions

The collapse of terrestrial ecosystems probably happened simultaneously with the decline in marine diversity during the P-Tr mass extinction, from the latest Changhsingian to earliest Triassic, which corresponds to Beds 25 to 28 in the Meishan sections. The loss of the *Gigantopteris* flora and the organic carbon isotope negative shift represent the beginning of the mass extinction and the collapse of the tropical rainforest ecosystem in the lower part of the Kayitou Formation (this study; S.Z. Shen et al. 2011; Zhang et al., 2016). All previous publications agree with our detailed biostratigraphic studies in finding that the Kayitou Formation was the witness of the terrestrial Permian-Triassic mass extinction in South China, and that it documents the transition from the latest Permian to the earliest Triassic (S.Z. Shen et al. 2011; Yu et al., 2015; Zhang et al., 2016). In this study, we describe detailed biological records from seven studied PTB sections, including four terrestrial sections, three coastal transitional sections and one shallow marine section, and demonstrate biostratigraphic correlations in the PTTB among contrasting siliciclastic settings in South China. In particular, we conclude that the mixed marine and terrestrial biota from the coastal transitional PTB sections constitutes an intermediate link for biostratigraphic correlation between

terrestrial and marine deposits. Finally, we propose that the occurrence of the *Euestheria*
gutta-bearing conchostracan fauna and the *Pteria ussurica variabilis*-*Towapteria*
scythica-Eumorphotis venetiana bivalve assemblage can be treated as markers of the PTTB in
 terrestrial-marine siliciclastic settings of South China, and perhaps more widely.

Acknowledgments

We thank Xincheng Qiu, Dongdong Li, Sunquan Wang, Xiao Shi, Hui Li, and Hongfei Chi for
 help with sampling from the studied section. We also acknowledge undergraduate volunteers
 Wenchao Shu and Gan Liu for their fossil finding and company in the field. We thank Jean Broutin
 for plant identification. Daoliang Chu is thankful to Frank Scholze for valuable comments on earlier
 versions of this manuscript. The authors are grateful to Shuzhong Shen, the anonymous reviewer and
 editor Thomas Cronin for their critical comments and constructive suggestions, which have greatly
 improved the quality of the paper. This study was supported by the National Natural Science
 Foundation of China (40972002, 41272024, 41272372, 41530104), the 111 Project (B08030). It is a
 contribution to IGCP 630.

References

- Aigner, T., Bachmann, G.H., 1992. Sequence-stratigraphic framework of the German Triassic.
 Sedimentary Geology 80, 115–135.
- Algeo, T.J., Twitchett, R.J., 2010. Anomalous Early Triassic sediment fluxes due to elevated
 weathering rates and their biological consequences. Geology 38, 1023–1026.
- Algeo, T.J., Chen, Z. Q., Fraiser, M. L., Twitchett, R.J., 2011. Terrestrial-marine teleconnections in

- the collapse and rebuilding of Early Triassic marine ecosystems. *Palaeogeography, Palaeoclimatology, Palaeoecology* 308, 1–11.
- Arche, A., López-Gómez, J., 2005. Sudden changes in fluvial style across the Permian–Triassic boundary in the eastern Iberian Ranges, Spain: analysis of possible causes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 229, 104–126.
- Bachmann, G.H., Kozur, H.W., 2004. The Germanic Triassic: correlations with the international chronostratigraphic scale, numerical ages and Milankovitch cyclicity. *Hallesches Jahrbuch für Geowissenschaften B* 26, 17–62.
- Benton, M.J., 1985. Mass extinction among non-marine tetrapods. *Nature* 316, 811–814.
- Benton, M.J., 2015. *When Life Nearly Died: The Greatest Mass Extinction of All Time*, second edition. Thames & Hudson, London, 352p.
- Benton, M.J., Twitchett, R.J., 2003. How to kill (almost) all life: the end-Permian extinction event. *Trends in Ecology and Evolution* 18, 358–365.
- Benton, M.J., Tverdokhlebov, V.P., Surkov, M., 2004. Ecosystem remodelling among vertebrates at the Permian-Triassic boundary in Russia. *Nature* 432, 97–100.
- Benton, M.J., Newell, A.J., 2014. Impacts of global warming on Permo-Triassic terrestrial ecosystems. *Gondwana Research* 25, 1308–1337.
- Bercovici, A., Cui, Y., Forel, M.B., Yu, J.X., Vajda, V., 2015. Terrestrial paleoenvironment characterization across the Permian-Triassic boundary in South China. *Journal of Asian Earth Sciences* 98, 225–246.
- Broglia-Loriga, C., Neri, C., Posenato, R., 1986. The Lower Triassic of the Dolomites and Cadore. In: *Permian and Permian–Triassic Boundary in the South-Alpine Segment of Western Tethys*.

- 595 IGCP Project 203, Brescia, Excursion Guidebook, pp. 29–34.
- 596 Cao, C.Q., Wang, W., Liu, L.J., Shen, S.Z., Summons, R.E., 2008. Two episodes of ^{13}C -depletion in
 597 organic carbon in the latest Permian: Evidence from the terrestrial sequences in northern
 598 Xinjiang, China. *Earth and Planetary Science Letters* 270, 251–257.
- 599 Cascales-Miñana, B., Cleal, C.J., 2014. The plant fossil record reflects just two great extinction
 600 events. *Terra Nova* 26, 195–200.
- 601 Cascales-Miñana, B., Diez, J.B., Gerrienne, P., Cleal, C.J., 2015. A palaeobotanical perspective on
 602 the great end-Permian biotic crisis. *Historical Biology*, doi:10.1080/08912963.2015.1103237.
- 603 Chen, Z.Q., Tong, J., Zhang, K., Yang, H., Liao, Z., Song, H., Chen, J., 2009. Environmental and
 604 biotic turnover across Permian–Triassic boundary from shallow carbonate platform in western
 605 Zhejiang, South China. *Australian Journal of Earth Sciences* 56, 775–797.
- 606 Chen, Z.Q., Kaiho, K., George, A.D., 2005. Survival strategies of brachiopod faunas from the
 607 end-Permian mass extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 224,
 608 232–269.
- 609 Chu, D.L., Tong, J.N., Yu, J.X., Song, H.J., Tian, L., 2013. The conchostracan fauna from the
 610 Kayitou Formation of western Guizhou, China. *Acta Palaeontologica Sinica* 52, 265–276.
- 611 Chu, D.L., Tong, J.N., Song, H.J., Benton, M.J., Bottjer, D.J., Song, H.Y., Tian, L., 2015. Early
 612 Triassic wrinkle structures on land: stressed environments and oases for life. *Scientific Reports* 5,
 613 e10109.
- 614 Chu, D.L., Tong, J.N., Song, H.J., Benton, M.J., Song, H.Y., Yu, J.X., Qiu, X.C., Huang, Y.F., Tian,
 615 L., 2015. Lilliput effect in freshwater ostracods during the Permian-Triassic extinction.
 616 *Palaeogeography, Palaeoclimatology, Palaeoecology* 435, 38–52.

- 617 Clapham, M.E., Payne, J.L., 2011. Acidification, anoxia, and extinction: A multiple logistic
618 regression analysis of extinction selectivity during the Middle and Late Permian. *Geology* 39,
619 1059–1062.
- 620 Clarkson, M.O., Kasemann, S.A., Wood, R.A., Lenton, T.M., Daines, S.J., Richoz, S., Ohnemuehler,
621 F., Meixner, A., Poulton, A.W., Tipper, E.T., 2015. Ocean acidification and the Permo-Triassic
622 mass extinction. *Science* 348, 229–232.
- 623 Cui, Y., Bercovici, A., Yu, J.X., Kump, L., Freeman, K., Su, S.G., Vajda, V., 2015. Carbon cycle
624 perturbation expressed in terrestrial Permian–Triassic boundary sections in south China. *Global*
625 *and Planetary Change*, in press, doi: 10.1016/j.gloplacha.2015.10.018.
- 626 Erwin, D.H., 1993. The great Paleozoic crisis: life and death in the Permian. Columbia University
627 Press, New York, 327p.
- 628 Fang, Z.J., 2004. Approach to the extinction pattern of Permian Bivalvia of South China, in: Rong,
629 J.Y., Fang, Z.J. (Eds.), *Mass Extinction and Recovery Evidences from the Palaeozoic and*
630 *Triassic of South China*. University of Science and Technology of China Press, Heifei, pp. 571
631 –646.
- 632 Forel, M.B., Crasquin, S., Chitnarin, A., Angiolini, L., Gaetani, M., 2015. Precocious sexual
633 dimorphism and Lilliput effect of Neo-Tethyan Ostracoda (Crustacea) through the
634 Permian-Triassic boundary. *Palaeontology* 58, 409–454.
- 635 Fraiser, M.L., Bottjer, D.J., 2004. The non-actualistic Early Triassic gastropod fauna: a case study of
636 the Lower Triassic Sinbad Limestone member. *Palaios* 19, 259–275.
- 637 Fraiser, M.L., Bottjer, D.J., 2007. When bivalves took over the world. *Paleobiology* 33, 397–413.
- 638 Gao, Y.Q., Shi, G.R., Peng, Y.Q., 2009. A new bivalve fauna from the Permian-Triassic boundary

- 639 section of southwestern China. *Alcheringa* 33, 33–47.
- 640 Gastaldo, R.A., Kamo, S.L., Neveling, J., Geissman, J.M., Bamford, M., Looy, C.V., 2015. Is the
641 vertebrate-defined Permian-Triassic boundary in the Karoo Basin, South Africa, the terrestrial
642 expression of the end-Permian marine event? *Geology* 43, 939–942.
- 643 Grauvogel-Stamm, L., Düringer, P., 1983. *Annalepis zeilleri* Fliche 1910 emend., un organe
644 reproducteur de Lycophyte de la Lettenkohle de l'Est de la France. Morphologie, spores in situ
645 et paléoécologie. *Geologische Rundschau* 72, 23–51.
- 646 Grauvogel-Stamm, L., Lugardon, B., 2001. The Triassic lycopsids *Pleuromeia* and *Annalepis*:
647 relationships, evolution, and origin. *American Fern Journal* 91, 115–149.
- 648 Grauvogel-Stamm, L., 1999. *Pleuromeia sternbergii* (Münster) Corda, eine charakteristische Pflanze
649 des deutschen Buntsandsteins. In: Hauschke, N., Wilde, V. (Eds.), *Trias-eine ganz andere Welt*.
650 Europa im frühen Erdmittelalter. Verlag Dr. Friedrich Pfeil, München, pp. 271–281.
- 651 Grauvogel-Stamm, L., Ash, S.R., 2005. Recovery of the Triassic land flora from the end-Permian life
652 crisis. *Comptes Rendus Palevo* 14, 593–608.
- 653 Hayami, I., 1997. Size changes of bivalves and a hypothesis about the cause of mass extinction.
654 *Fossils* 62, 24–36 (in Japanese).
- 655 He, B., Xu, Y.G., Huang, X.L., Luo, Z.Y., Shi, Y.R., Yang, Q.J., Yu, S.Y., 2007. Age and duration of
656 the Emeishan flood volcanism, SW China: Geochemistry and SHRIMP zircon U–Pb dating of
657 silicic ignimbrites, post-volcanic Xuanwei Formation and clay tuff at the Chaotian section. *Earth*
658 *and Planetary Science Letters* 225, 306–323.
- 659 He, W.H., Shi, G.R., Feng, Q.L., Campi, M.J., Gu, S.Z., Bu, J.J., Peng, Y.Q., Meng, Y.Y., 2007,
660 Brachiopod miniaturization and its possible causes during the Permian-Triassic crisis in deep

- 661 water environments, South China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 252,
662 145–163.
- 663 Hermann, E., Hochuli, P.A., Bucher, H., Bruhwiler, T., Hautmann, M., Ware, D., Roohi, G., 2011.
664 Terrestrial ecosystems on North Gondwana following the end-Permian mass extinction.
665 *Gondwana Research* 20, 630–637.
- 666 Hinojosa, J.L., Brown, S.T., Chen, J., DePaolo, D.J., Paytan, A., Shen, S.Z., Payne, J.L., 2012.
667 Evidence for end-Permian ocean acidification from calcium isotopes in biogenic apatite.
668 *Geology* 40, 743–746.
- 669 Huang, B.H., 1996. The Angara flora from North China. *Palaeobotanist* 45, 309–314.
- 670 Huang, B.H., Ding, Q., 1998. The Angara flora from North China. *Acta Geoscientia Sinica*,
671 *Geologica Sciences* 19, 97–104.
- 672 Huang, Y.F., Tong, J.N., 2014. Advance in the study of the Permian-Triassic bivalves. *Advance in*
673 *Earth Sciences* 29, 922–933.
- 674 Huttenlocker, A.K., 2014. Body size reductions in nonmammalian eutheriodont therapsids
675 (Synapsida) during the end-Permian mass extinction. *PloS One* 9, e87553.
- 676 Ikeda, M., Hori, R.S., Okada, Y., Nakada, R., 2015. Volcanism and deep-ocean acidification across
677 the end-Triassic extinction event. *Palaeogeography, Palaeoclimatology, Palaeoecology* 440,
678 725–733.
- 679 Isozaki, Y., 1997. Permo-Triassic boundary superanoxia and stratified superocean: records from lost
680 deep sea. *Science* 276, 235–238.
- 681 Joachimski, M.M., Lai, X.L., Shen, S.Z., Jiang, H.S., Luo, G.M., Chen, B., Chen, J., Sun, Y.D., 2012.
682 Climate warming in the latest Permian and the Permian-Triassic mass extinction. *Geology* 40,

- 683 195–198.
- 684 Kershaw, S., Guo, L., Swift, A., Fan, J.S., 2002. Microbialites in the Permian-Triassic boundary
685 interval in Central China: Structure, age and distribution. *Facies* 47, 83–89.
- 686 Kershaw, S., Crasquin, S., Li, Y., Collin, P.Y., Forel, M.B., Mu, X.N., Baud, A., Wang, Y., Xie, S.C.,
687 Maurer, F., Guo, L., 2012. Microbialites and global environmental change across the
688 Permian–Triassic boundary: a synthesis. *Geobiology* 10, 25–47.
- 689 Kozur, H., Seidel, G., 1983. Revision der Conchostracen-Faunen des unteren und mittleren
690 Buntsandsteins. Teil I. *Zeitschrift für Geologische Wissenschaften* 11, 289–417.
- 691 Kozur, H.W., Weems, R.E., 2010. The biostratigraphic importance of conchostracans in the
692 continental Triassic of the northern hemisphere. *Geological Society, London, Special
693 Publications* 334, 315–417.
- 694 Kozur, H.W., Weems, R.E., 2011. Detailed correlation and age of continental late Changhsingian
695 and earliest Triassic beds: Implications for the role of the Siberian Trap in the Permian-Triassic
696 biotic crisis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 308, 22–40.
- 697 Li, Z.J., Zhu, J.N., Hu, Y.F., 1982. The new views on the correlation for the Late Permian
698 coal-bearing strata in Junlian area, South Sichuan. *Journal of Stratigraphy* 6, 174–182.
- 699 Looy, C.V., Brugman, W.A., Dilcher, D.L., Visscher, H., 1999. The delayed resurgence of equatorial
700 forests after the Permian-Triassic ecologic crisis. *Proceedings of the National Academy of
701 Sciences, U.S.A.* 96, 13857–13862.
- 702 Looy, C.V., Twitchett, R.J., Dilcher, D.L., Van Konijnenburg-Van Cittert, J.H., Visscher, H., 2001.
703 Life in the end-Permian dead zone. *Proceedings of the National Academy of Sciences, U.S.A.*
704 98, 7879–7883.

- 705 Luo, G., Lai, X., Shi, G.R., Jiang, H., Yin, H., Xie, S., Tong, J., Zhang, K., He, W., Wignall, P.B.,
 706 2008. Size variation of conodont elements of the *Hindeodus-Isarcicella* clade during the
 707 Permian–Triassic transition in South China and its implication for mass extinction.
 708 *Palaeogeography, Palaeoclimatology, Palaeoecology* 264, 176–187.
- 709 Lutkevitch, E.M., 1937. O nekotorych Phyllopoda SSSR (On some Phyllopoda of the USSR).
 710 *Ezhegodnik Vsesoyuznogo Paleontologicheskogo Obshchestva* 11, 60–70 (In Russian).
- 711 Mata, S.A., Bottjer, D.J., 2012. Microbes and mass extinctions: paleoenvironmental distribution of
 712 microbialites during times of biotic crisis. *Geobiology* 10, 3–24.
- 713 McGowan, A.J., Smith, A.B., Taylor, P.D., 2009. Faunal diversity, heterogeneity and body size in
 714 the Early Triassic: testing post-extinction paradigms in the Virgin Limestone of Utah, USA.
 715 *Australian Journal of Earth Sciences* 56, 859–872.
- 716 McRoberts, C.A., 2010. Biochronology of Triassic bivalves. Geological Society, London, Special
 717 Publications, 334, 201–219.
- 718 Metcalfe, I., Crowley, J.L., Nicoll, R.S., Schmitz, M., 2015. High-precision U-Pb CA-TIMS
 719 calibration of Middle Permian to Lower Triassic sequences, mass extinction and extreme
 720 climate-change in eastern Australian Gondwana. *Gondwana Research* 28, 61–81.
- 721 Mutter, R.J., Neuman, A.G., 2009. Recovery from the end-Permian extinction event: Evidence from
 722 “Lilliput *Listracanthus*”. *Palaeogeography, Palaeoclimatology, Palaeoecology* 284, 22–28.
- 723 Naugolnykh, S.V., 2013. The heterosporous lycopodiophyte *Pleuromeia rossica* Neuburg, 1960 from
 724 the Lower Triassic of the Volga River basin (Russia): organography and reconstruction
 725 according to the ‘whole plant’ concept. *Wulfenia* 20, 1–16.
- 726 Newell, A.J., Tverdokhlebov, V.P., Benton, M.J., 1999. Interplay of tectonics and climate on a

- 727 transverse fluvial system, Upper Permian, Southern Uralian Foreland Basin, Russia.
728 *Sedimentary Geology* 127, 11–29.
- 729 Payne, J.L., 2005. Evolutionary dynamics of gastropod size across the end-Permian extinction and
730 through the Triassic recovery interval. *Paleobiology* 31, 269–290.
- 731 Payne, J.L., Lehrmann, D.J., Follett, D., Seibel, M., Kump, L.R., Riccardi, A., Altiner, D., Sano, H.,
732 Wei, J., 2007. Erosional truncation of uppermost Permian shallow-marine carbonates and
733 implications for Permian-Triassic boundary events. *Geological Society of America Bulletin* 119,
734 771–784.
- 735 Peng, Y.Q., Tong, J.N., Shi, G.R., Hansen, H.J., 2001. The Permian-Triassic boundary stratigraphic
736 set: characteristics and correlation. *Newsletters on Stratigraphy*, 55–71.
- 737 Peng, Y.Q., Zhang, S.X., Yu, T.X., Yang, F.Q., Gao, Y.Q., Shi, G.R., 2005. High-resolution
738 terrestrial Permian-Triassic event stratigraphic boundary in western Guizhou and eastern
739 Yunnan, southwestern China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 215,
740 285–295.
- 741 Peng, Y.Q., Shi, G.R., Gao, Y.Q., He, W.H., Shen, S.Z., 2007. How and why did the Lingulidae
742 (Brachiopoda) not only survive the end-Permian mass extinction but also thrive in its aftermath?
743 *Palaeogeography, Palaeoclimatology, Palaeoecology* 252, 118–131.
- 744 Peng, Y.Q., Shi, G.R. 2008. New Early Triassic Lingulidae (Brachiopoda) genera and species from
745 South China. *Alcheringa* 32, 149–170.
- 746 Peng, Y.Q., Shi, G.R. 2009. Life crises on land across the Permian-Triassic boundary in South
747 China. *Global and Planetary Change* 65, 155–165.
- 748 Pereira, Z., Fernandes, P., Lopes, G., Marques, J., Vasconcelos, L., 2016. The Permian–Triassic

- 749 transition in the Moatize-Minjova Basin, Karoo Supergroup, Mozambique: A palynological
750 perspective. *Review of Palaeobotany and Palynology* 226, 1–19.
- 751 Rey, K., Amiot, R., Fourel, F., Rigaudier, T., Abdala, F., Day, M.O., Fernandez, V., Fluteau, F.,
752 France-Lanord, C., Rubidge, B.S., Smith, R.M., Viglietti, P.A., Zipfel, B., Lécuyer, C., 2016.
753 Global climate perturbations during the Permo-Triassic mass extinctions recorded by continental
754 tetrapods from South Africa. *Gondwana Research* 37, 384–396.
- 755 Retallack, G.J., 1995. Permian-Triassic life crisis on land. *Science* 267, 77–80.
- 756 Retallack, G.J., Veevers, J.J., Morante, R., 1996. Global coal gap between Permian-Triassic
757 extinction and Middle Triassic recovery of peat-forming plants. *Geological Society of America*
758 *Bulletin* 108, 195–207.
- 759 Retallack, G.J., Smith, R.M., Ward, P.D., 2003. Vertebrate extinction across Permian-Triassic
760 boundary in Karoo Basin, South Africa. *Geological Society of America Bulletin* 115,
761 1133–1152.
- 762 Retallack, G.J., Sheldon, N.D., Carr, P.F., Fanning, M., Thompson, C.A., Williams, M.L., Jones,
763 B.G., Hutton, A., 2011. Multiple Early Triassic greenhouse crises impeded recovery from Late
764 Permian mass extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 308, 233–251.
- 765 Retallack, G.J., 2013. Permian and Triassic greenhouse crises. *Gondwana Research* 24, 90–103.
- 766 Rodland, D.L., Bottjer, D.J., 2001. Biotic recovery from the end-Permian mass extinction: Behavior
767 of the inarticulate brachiopod *Lingula* as a disaster taxon. *Palaaios* 16, 95–101.
- 768 Romano, C., Koot, M.B., Kogan, I., Brayard, A., Minikh, A.V., Brinkmann, W., Bucher, H., Kriwet,
769 J., 2016. Permian-Triassic Osteichthyes (bony fishes): diversity dynamics and body size
770 evolution. *Biological Reviews* 91, 106–147.

- 771 Schneebeili-Hermann, E., Kürschner, W.M., Kerp, H., Bomfleur, B., Hochuli, P.A., Bucher, H., Ware,
 772 D., Roohi, G., 2015. Vegetation history across the Permian-Triassic boundary in Pakistan (Amb
 773 section, Salt Range). *Gondwana Research* 27, 911–924.
- 774 Schubert, J.K., Bottjer, D.J., 1992. Early Triassic stromatolites as post-mass extinction disaster forms.
 775 *Geology* 20, 883–886.
- 776 Schubert, J.K., Bottjer, D.J., 1995. Aftermath of the Permian-Triassic mass extinction event:
 777 Paleocology of Lower Triassic carbonates in the western USA. *Palaeogeography,*
 778 *Palaeoclimatology, Palaeoecology* 116, 1–39.
- 779 Scholze, F., Golubev, V.K., Niedzwiedzki, G., Sennikov, A.G., Schneider, J.W., Silantiev, V.V.,
 780 2015. Early Triassic conchostracans (Crustacea: Branchiopoda) from the terrestrial
 781 Permian-Triassic boundary sections in the Moscow syncline. *Palaeogeography,*
 782 *Palaeoclimatology Palaeoecology* 429, 22–40.
- 783 Sedlacek, A.R., Saltzman, M.R., Algeo, T.J., Horacek, M., Brandner, R., Foland, K., Denniston, R.F.,
 784 2014. $^{87}\text{Sr}/^{86}\text{Sr}$ stratigraphy from the Early Triassic of Zal, Iran: Linking temperature to
 785 weathering rates and the tempo of ecosystem recovery. *Geology* 42, 779–782.
- 786 Song, H.J., Tong, J.N., Chen, Z.Q., 2011. Evolutionary dynamics of the Permian-Triassic foraminifer
 787 size: Evidence for Lilliput effect in the end-Permian mass extinction and its aftermath.
 788 *Palaeogeography, Palaeoclimatology, Palaeoecology* 308, 98–110.
- 789 Song, H.J., Wignall, P.B., Tong, J.N., Yin, H.F., 2013. Two pulses of extinction during the
 790 Permian-Triassic crisis. *Nature Geoscience* 6, 52–56.
- 791 Song H.J., Wignall, P.B., Tong J.N., Song H.Y., Chen J., Chu D.L., Tian L., Luo M., Zong K.Q.,
 792 Chen Y.L., Lai X.L., Zhang K.X., Wang H.M., 2015. Integrated Sr isotope variations and global

- 793 environmental changes through the Late Permian to early Late Triassic. *Earth and Planetary*
 794 *Science Letters* 424, 140–147.
- 795 Shen, S.Z., He, Z.L., Shi, G.R., 1995. Biostratigraphy and correlation of several Permian-Triassic
 796 boundary sections in southwestern China. *Journal of Southeast Asian Earth Sciences* 12, 19–30.
- 797 Shen, S.Z., Cao, C.Q., Henderson, C.M., Li, W.Z., Zhang, H., 2006. Rapid deforestation across the
 798 Permian-Triassic boundary in South China. *Geological Society of America Abstracts with*
 799 *Programs*. Geological Society of America, Washington DC, p. 338.
- 800 Shen, S.Z., Crowley, J.L., Wang, Y., Bowring, S.A., Erwin, D.H., Sadler, P.M., Cao, C.Q., Rothman,
 801 D.H., Henderson, C.M., Ramezani, J., Zhang, H., Shen, Y., Wang, X.D., Wang, W., Mu, L., Li,
 802 W.Z., Tang, Y.G., Liu, X.L., Liu, L.J., Zeng, Y., Jiang, Y.F., Jin, Y.G., 2011. Calibrating the
 803 end-Permian mass extinction. *Science* 334, 1367–1372.
- 804 Shen, W.J., Sun, Y.G., Lin, Y.T., Liu, D.H., and Chai, P.X., 2011. Evidence for wildfire in the
 805 Meishan section and implications for Permian-Triassic events. *Geochimica et Cosmochimica*
 806 *Acta* 75, 1992–2006.
- 807 Sheng, J.Z., Chen, C.Z., Wang, Y.G., Rui, L., Liao, Z.T., Bando, Y., Ishii, K.I., Nakazawa, K.,
 808 Nakamura, K., 1984. Permian-Triassic boundary in middle and eastern Tethys. *Journal of the*
 809 *Faculty of Science, Hokkaido University, Series 4: Geology and Mineralogy* 21, 133–181.
- 810 Sheng, J.Z., Chen, C.Z., Wang, Y.G., Rui, L., Liao, Z.T., He, J.W., Jiang, N.Y., Wang, C.Y., 1987.
 811 New advances on the Permian and Triassic boundary of Jiangsu, Zhejiang and Anhui, in:
 812 *Palaeontology, N.I.o.G.a. (Ed.), Permian-Triassic boundary (1), stratigraphy and Palaeontology*
 813 *of systematic boundaries in China*. Nanjing University Press, Nanjing, pp. 1–21.
- 814 Sun, Y.D., Joachimski, M.M., Wignall, P.B., Yan, C.B., Chen, Y.L., Jiang, H.S., Wang, L.N., Lai,

- 815 X.L., 2012. Lethally hot temperatures during the early Triassic greenhouse. *Science* 338,
816 366–370.
- 817 Szurlies, M., Bachmann, G.H., Menning, M., Nowaczyk, N.R., Käding, K.C., 2003.
818 Magnetostratigraphy and high-resolution lithostratigraphy of the Permian-Triassic boundary
819 interval in Central Germany. *Earth and Planetary Science Letters* 212, 263–278.
- 820 Szurlies, M., Geluk, M.C., Krijgsman, W., and Kurschner, W.M., 2012. The continental
821 Permian–Triassic boundary in the Netherlands: Implications for the geomagnetic time scale:
822 *Earth and Planetary Science Letters* 317–318, 165–176.
- 823 Taylor, G.K., Tucker, C., Twitchett, R.J., Kearsley, T., Benton, M.J., Newell, A.J., Surkov, M.V., and
824 Tverdokhlebov, V.P., 2009. Magnetostratigraphy of Permian/Triassic boundary sequences in the
825 Cis-Urals, Russia: No evidence for a major temporal hiatus. *Earth & Planetary Science*
826 *Letters* 281, 36–47.
- 827 Tian, L., Tong, J. N., Algeo, T. J., Song, H. J., Song, H. Y., Chu, D. L., Shi, L., Bottjer, D. J., 2014.
828 Reconstruction of Early Triassic ocean redox conditions based on framboidal pyrite from the
829 Nanpanjiang Basin, South China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 412,
830 68–79.
- 831 Tverdokhlebov, V.P., Tverdokhlebova, G.I., Surkov, M.V., Benton, M.J., 2002. Tetrapod localities
832 from the Triassic of the SE of European Russia. *Earth-Science Reviews* 60, 1–66.
- 833 Twitchett, R.J., Looy, C.V., Morante, R., Visscher, H., Wignall, P.B., 2001. Rapid and synchronous
834 collapse of marine and terrestrial ecosystems during the end-Permian biotic crisis. *Geology* 29,
835 351–354.
- 836 Twitchett, R.J., Feinberg, J.M., O'Connor, D.D., Alvarez, W., McCollum, L.B., 2005. Early Triassic

- 837 ophiuroids: their paleoecology, taphonomy, and distribution. *Palaios* 20, 213–223.
- 838 Twitchett, R.J., 2007. The Lilliput effect in the aftermath of the end-Permian extinction event.
- 839 *Palaeogeography, Palaeoclimatology, Palaeoecology* 252, 132–144.
- 840 Urbanek, A., 1993. Biotic crises in the history of Upper Silurian graptoloids: a palaeobiological
- 841 model. *Historical Biology* 7, 29–50.
- 842 Wang, S.Y., Yin, H.F., 2001. Study on terrestrial Permian–Triassic boundary in eastern
- 843 Yunnan and western Guizhou. China University of Geosciences Press, Wuhan. 88p. (in
- 844 Chinese with English abstract).
- 845 Wang, Y.B., Tong, J.N., Wang, J.S., Zhou, X.G., 2005. Calcimicrobialite after end-Permian mass
- 846 extinction in South China and its palaeoenvironmental significance. *Chinese Science Bulletin* 50,
- 847 665–671.
- 848 Wang, Z.Q., Wang, L.X., 1990. Late Early Triassic fossil plants from upper part of the
- 849 Shiqianfeng Group in North China. *Shanxi Geology* 5, 97–154.
- 850 Ward, P.D., Montgomery, D.R., Smith, R., 2000. Altered river morphology in South Africa related to
- 851 the Permian-Triassic extinction. *Science* 289, 1740–1743.
- 852 Ward, P.D., Botha, J., Buick, R., De Kock, M.O., Erwin, D.H., Garrison, G.H., Kirschvink, J.L.,
- 853 Smith, R., 2005. Abrupt and gradual extinction among Late Permian land vertebrates in the
- 854 Karoo Basin, South Africa. *Science* 307, 709–714.
- 855 Wasmer, M., Hautmann, M., Hermann, E., Ware, D., Roohi, G., Ur-Rehman, K., Yaseen, A., Bucher,
- 856 H., 2012. Olenekian (Early Triassic) bivalves from the Salt Range and Surghar Range, Pakistan.
- 857 *Palaeontology* 55, 1043–1073.
- 858 Wignall, P.B., Twitchett, R.J., 1996. Oceanic anoxia and the end Permian mass extinction. *Science*

- 859 272, 1155–1158.
- 860 Wu, Y.S., Yu, G.L., Li, R.H., Song, L.R., Jiang, H.X., Riding, R., Liu, L.J., Liu, D.Y., Zhao, R.,
 861 2014. Cyanobacterial fossils from 252 Ma old microbialites and their environmental significance.
 862 Scientific Reports 4, e3820.
- 863 Xie, S.C., Pancost, R.D., Wang, Y.B., Yang, H., Wignall, P.B., Luo, G.M., Jia, C.L., Chen, L., 2010.
 864 Cyanobacterial blooms tied to volcanism during the 5 m.y. Permo-Triassic biotic crisis. Geology
 865 38, 447–450.
- 866 Yin, H.F., 1985. Bivalves near the Permian-Triassic boundary in South China. Journal of
 867 Paleontology 59, 572–600.
- 868 Yin, H.F., Jiang, H.S., Xia, W.C., Feng, Q.L., Zhang, N., Shen, J., 2014. The end-Permian regression
 869 in South China and its implication on mass extinction, Earth-Science Reviews 137, 19–33.
- 870 Yu, J.X., Peng, Y.Q., Zhang, S.X., Yang, F.Q., Zhao, Q.M., Huang, Q.S., 2007. Terrestrial events
 871 across the Permian-Triassic boundary along the Yunnan-Guizhou border, SW China. Global and
 872 Planetary Change 55, 193–208.
- 873 Yu, J., 2008. Floras and the evolutionary dynamics across the Permian-Triassic boundary nearby the
 874 border of Guizhou and Yunnan, south China, China University of Geosciences, Wuhan,
 875 Université Pierre et Marie Curie, Paris 6, 254 pp.
- 876 Yu, J., Li, H., Zhang, S., Yang, F. and Feng, Q., 2008. Timing of the terrestrial Permian–Triassic
 877 boundary biotic crisis: Implications from U-Pb dating of authigenic zircons. Science in China
 878 Series D: Earth Sciences 51, 1633–1645.
- 879 Yu, J.X., Broutin, J., Huang, Q.S., Grauvogel-Stamm, L., 2010. *Annalepis*, a pioneering lycopsid
 880 genus in the recovery of the Triassic land flora in South China. Comptes Rendus Palévol 9,

- 881 479–486.
- 882 Yu, J.X., Broutin, J., Chen, Z.Q., Shi, X., Li, H., Chu, D.L., Huang, Q.S., 2015. Vegetation
883 changeover across the Permian-Triassic Boundary in Southwest China. Extinction, survival,
884 recovery and palaeoclimate: A critical review. *Earth-Science Reviews*, in press, doi:
885 10.1016/j.earscirev.2015.04.005.
- 886 Zhang, H., Cao, C.Q., Liu, X.L., Mu, L., Zheng, Q.F., Liu, F., Xiang, L., Liu, L.J., Shen, S.Z., 2016.
887 The terrestrial end-Permian mass extinction in South China. *Palaeogeography,*
888 *Palaeoclimatology, Palaeoecology* 448, 108–124.
- 889 Zhu, J.N., Hu, Y.F., Li, Z.J., 1984. Upper Permian plants of Junlian area, South Sichuan.
890 *Professional Papers of Stratigraphy and Palaeontology* 11, 133–150.
- 891

Figure captions

Fig. 1. Investigated localities and simplified lithofacies of South China during the Permian-Triassic transition. A. Palaeogeographic map of South China during the Permian-Triassic transition (after Yin et al., 2014). The studied sections (marking by red dots) were located in different facies. B. The reconstruction of the depositional environment during the Permian-Triassic transition (modified from Bercovici et al., 2015), and position of the seven studied sections is indicated relative to the proximal/distal transect. Abbreviations: GL, Guangyuan-Liangping; XL, Xiakou-Lichuan; NMBY, north marginal basin of Yangtze Platform; HGG, Hunan-Guizhou-Guangxi; ZFG, Gejiang-Fujian-Guangdong.

Fig. 2. Permian-Triassic sections in terrestrial-marine siliciclastic settings of South China. 1. The Xuanwei Formation with abundant *Gigantopteris* flora at Chahe section; 2. The lithological boundary between the Xuanwei and Kayitou formations; the base of Kayitou Formation is marked by a thin coal seam and an ash bed which is dated as 252.30 ± 0.07 Ma (S.Z. Shen et al., 2011); 3. The lithological boundary between the Kayitou and Dongchuan formations in the Guanbachong section; 4, 5. Lithological boundaries between the Xuanwei and Kayitou formations in the Jiuchaichong and Jinzhong sections; 6. The Feixianguan Formation with abundant bivalves in the Wadu section.

Fig. 3. Lithostratigraphy and fossil distribution in the Guanbachong section. Solid lines indicate known stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards (Plant fossil data are from Zhang et al., 2016). Abbreviations: Fm, Formation; Th, Thickness. The list of the fossils is provided in the electronic supplementary material.

Fig. 4. *Euestheria gutta* Lutkevitch specimens with various deformation characteristics. 1-5, 8-10.

913 *Euestheria gutta* specimens collected from the middle part of the Kayitou Formation of the
 914 Jiucaichong section, JCC-26-1, JCC-26-1-1, JCC-26-7, JCC-26-13, JCC-26-22, JCC-27-4,
 915 JCC-27-10, JCC-27-17 ; 6, 7. *Euestheria gutta* specimens collected from the upper part of the
 916 Kayitou Formation of the Xiaohebian section, XHB-10-12a, XHB-10-12b; 11, 12. *Euestheria gutta*
 917 specimens collected from the lower part of the Kayitou Formation of the Chahe section, CH-71-7,
 918 CH-71-9; 13. *Euestheria gutta* specimens collected from the lower part of the Kayitou Formation of
 919 the Jinzhong section, JZ-9-11; 14-17. SEM images showing growth bands and reticular
 920 microsculptures. The scale bars in 1-13 represent 1 mm.

921 **Fig. 5.** Typical *Gigantopteris* flora from the Xuanwei Formation of the Chahe section. 1.
 922 *Lobatannularia cathaysiana* Yao, CH-18-29; 2. *Cladophlebis permica* Lee et Wang, CH-18-12a; 3.
 923 *Stigmara ficoides* (Sternb.) Brongnairt, CH-46-1; 4. *Fascipteris densata* Gu et Zhi, CH-18-42; 5.
 924 *Pecopteris orientalis* (Schenk) Potonie, CH-18-59; 6. *Compsopteris* sp., CH-58-5; 7. *Guizhoua*
 925 *gregalis* Zhao, CH-58-17b; 8. *Pecopteris marginata* Gu et Zhi, CH-58-14a; 9. *Compsopteris*
 926 *contracta* Gu et Zhi, CH-58-6; 10. *Gigantonoclea hallei* (Asama) Gu et Zhi, CH-29-1; 11. *Rajahia*
 927 *guizhouensis* Zhang, CH-67-13. The scale bars represent 1 cm.

928 **Fig. 6.** Lithostratigraphy and fossil distribution in the Chahe section. The age data based on S.Z.
 929 Shen et al. (2011). Solid lines indicate known stratigraphic ranges; dotted lines indicate uncertain
 930 ranges up from below or upwards. Abbreviations: Fm, Formation; Th, Thickness. The list of the
 931 fossils is provided in the electronic supplementary material.

932 **Fig. 7.** Typical *Gigantopteris* flora from the lower part of the Kayitou Formation of the Chahe and
 933 Jiucaichong sections. 1. *Gigantopteris dictyophylloides* Gu et Zhi, CH-69-13; 2. *Rajahia*

934 *guizhouensis* Zhang, CH-69-2; 3. *Stigmara ficoides* (Sternb.) Brongnairt, CH-69-36; 4.
 935 *Gigantopteris* sp., JCC-24-2; 5, 6. *Gigantonoclea plumosa* Mo, JCC-24-13; 7, 8. *Rajahia*
 936 *guizhouensis* Zhang, CH-69-7, JCC-24-6. The scale bars in 1-3, 7, 8 represent 1 cm.

937 **Fig. 8.** Lithostratigraphy and fossil distribution of the Xiaohebian section. Solid lines indicate known
 938 stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards. Abbreviations:
 939 Fm, Formation; Th, Thickness. The list of the fossils is provided in the
 940 electronic supplementary material.

941 **Fig. 9.** Lithostratigraphy and fossil distribution of the Jiucaichong section. Solid lines indicate known
 942 stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards. Abbreviations:
 943 Fm, Formation; Th, Thickness. The list of the fossils is provided in the
 944 electronic supplementary material.

945 **Fig. 10.** Lithostratigraphy and fossil distribution of the Jinzhong section. Solid lines indicate known
 946 stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards. Abbreviations:
 947 Fm, Formation; Th, Thickness. The list of the fossils is provided in the
 948 electronic supplementary material.

949 **Fig. 11.** Mixed marine and terrestrial biota from the coastal transitional Jinzhong and Jinjibang
 950 sections. 1. Conchostracans and marine bivalves preserved on the same bedding surface, collected
 951 from the base of the Feixianguan Formation in the Jinjibang section, JJB-158002; 2. *Euestheria gutta*
 952 Lutkevitch, close-up views of indicated portions of 1, JJB-158-1; 3. *Pteria ussurica variabilis* Chen,
 953 close-up views of indicated portions of 1, JJB-158-3; 4. Marine bivalves collected from the Kayitou

954 Formation in the Jinzhong section, JZ-9-13; 5. Conchostracans collected from the Kayitou Formation
 955 of the Jinzhong section, JZ-9-22; 6. *Peltaspermum* sp., collected from the Kayitou Formation of the
 956 Jinzhong section, JZ-10-2; 7. *Pecopteris* sp., collected from the Kayitou Formation of the Jinzhong
 957 section, JZ-9-14; 8. *Annalepis brevicystis* Meng, collected from the base of the Feixianguan
 958 Formation of the Jinjibang section, JJB-159-7; 9. *Annalepis zeilleri* Fliche, collected from the base of
 959 the Feixianguan Formation of the Jinjibang section, JJB-159-8; 10, 11. *Peltaspermum* cf. *martinsii*
 960 (Harris) Poort et Kerp, collected from the Kayitou Formation of the Jinzhong section, JZ-10-14,
 961 JZ-11-10; 12. lingulid brachiopod specimens, assigned to *Sinolingularia*, collected from the Kayitou
 962 Formation of the Jinzhong section, JZ-9-21.

963 **Fig. 12.** Lithostratigraphy and fossil distribution of the Jinjibang section. Solid lines indicate known
 964 stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards. Abbreviations:
 965 Fm, Formation; Th, Thickness. The list of the fossils is provided in the electronic supplementary
 966 material.

967 **Fig. 13.** Lithostratigraphy and fossil distribution of the Wadu section. Solid lines indicate known
 968 stratigraphic ranges; dotted lines indicate uncertain ranges up from below or upwards. Abbreviations:
 969 Fm, Formation; Th, Thickness. The list of the fossils is provided in the electronic supplementary
 970 material.

971 **Fig. 14.** Triassic-form bivalve fossils from the lower part of the Feixianguan Formation of the Wadu
 972 section. 1-4. *Claraia bioni* Nakazawa, WD-801190, WD-904059, WD-905002, WD-905006; 5-8.
 973 *Eumorphotis venetiana* Hauer, WD-902130a, WD-902130b, WD-801189, WD-801188; 9-11. *Pteria*
 974 *ussurica variabilis* Chen, WD-902120, WD-902124, WD-902126; 12. *Towapteria scythica* Wirth,

975 WD-801210; 13. *Pteria ussurica variabilis*-*Eumorphotis venetiana*-*Promyalina schamarae*
 976 assemblage, WD-190001; 14. *Neoschizodus laevigatus* Ziethen, WD-903108; 15. *Promyalina*
 977 *schamarae* Bittner, WD-903095; 16. *Promyalina putiatinensis* Kiparisova, WD-904061; 17.
 978 *Permophorus bregeri* Girty, WD-903090a/WD903090b. The scale bars represent 5 mm.

979 **Fig. 15.** *Gigantopteris* flora fragmented specimens from the lower part of the Kayitou Formation of
 980 the terrestrial sections. The scale bars represent 2 mm.

981 **Fig. 16.** The distribution of the *Gigantopteris* flora, conchostracans, bivalves, lingulid brachiopods,
 982 and LAD of the coal and ash bed, with ages showing the correlation among the studied sections in
 983 the Permian-Triassic transitional beds and the end-Permian mass extinction of the *Gigantopteris*
 984 flora. PTTB, Permian-Triassic transitional beds; e-PTB, estimated Permian-Triassic boundary; EPPE,
 985 end-Permian plant mass extinction; *Pte.-Towa.-Eumor.* assemblage, *Pteria-Towapteria-Eumorphotis*
 986 assemblage.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

Figure 14

Figure 15

Figure 16