



Richmond, R., Timpson, N., Felix, J. F., Palmer, T. M., Gaillard, R., McMahon, G., Davey Smith, G., Jaddoe, V. W. V., & Lawlor, D. (2017). Using genetic variation to explore the causal effect of maternal pregnancy adiposity on future offspring adiposity: a Mendelian randomization study. *PLoS Medicine*, 14(1), Article e1002221. <https://doi.org/10.1371/journal.pmed.1002221>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY

Link to published version (if available):  
[10.1371/journal.pmed.1002221](https://doi.org/10.1371/journal.pmed.1002221)

[Link to publication record on the Bristol Research Portal](#)  
PDF-document

## University of Bristol – Bristol Research Portal

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

# Using Genetic Variation to Explore the Causal Effect of Maternal Pregnancy Adiposity on Future Offspring Adiposity: A Mendelian Randomisation Study

Rebecca C. Richmond, Nicholas J. Timpson, Janine F. Felix, Tom Palmer, Romy Gaillard, George McMahon, George Davey Smith, Vincent W. Jaddoe, Debbie A. Lawlor

Published: January 24, 2017 • <http://dx.doi.org/10.1371/journal.pmed.1002221>

## Abstract

### Background

It has been suggested that greater maternal adiposity during pregnancy affects lifelong risk of offspring fatness via intrauterine mechanisms. Our aim was to use Mendelian randomisation (MR) to investigate the causal effect of intrauterine exposure to greater maternal body mass index (BMI) on offspring BMI and fat mass from childhood to early adulthood.

### Methods and Findings

We used maternal genetic variants as instrumental variables (IVs) to test the causal effect of maternal BMI in pregnancy on offspring fatness (BMI and dual-energy X-ray absorptiometry [DXA] determined fat mass index [FMI]) in a MR approach. This was investigated, with repeat measurements, from ages 7 to 18 in the Avon Longitudinal Study of Parents and Children (ALSPAC;  $n = 2,521$  to  $3,720$  for different ages). We then sought to replicate findings with results for BMI at age 6 in Generation R ( $n = 2,337$  for replication sample;  $n = 6,057$  for total pooled sample).

In confounder-adjusted multivariable regression in ALSPAC, a 1 standard deviation (SD, equivalent of  $3.7 \text{ kg/m}^2$ ) increase in maternal BMI was associated with a  $0.25 \text{ SD}$  (95% CI  $0.21\text{--}0.29$ ) increase in offspring BMI at age 7, with similar results at later ages and when FMI was used as the outcome.

A weighted genetic risk score was generated from 32 genetic variants robustly associated with BMI (minimum F-statistic = 45 in ALSPAC). The MR results using this genetic risk score as an IV in ALSPAC were close to the null at all ages (e.g.,  $0.04 \text{ SD}$  (95% CI  $-0.21\text{--}0.30$ ) at age 7 and  $0.03 \text{ SD}$  (95% CI  $-0.26\text{--}0.32$ ) at age 18 per SD increase in maternal BMI), which was similar when a 97 variant generic risk score was used in ALSPAC.

When findings from age 7 in ALSPAC were meta-analysed with those from age 6 in Generation R, the pooled confounder-adjusted multivariable regression association was  $0.22 \text{ SD}$  (95% CI  $0.19\text{--}0.25$ ) per SD increase in maternal BMI and the pooled MR effect (pooling the 97 variant score results from ALSPAC with the 32 variant score results from Generation R) was  $0.05 \text{ SD}$  (95% CI  $-0.11\text{--}0.21$ ) per SD increase in maternal BMI ( $p$ -value for difference between the two results =  $0.05$ ). A number of sensitivity analyses exploring violation of the MR results supported our main findings. However, power was limited for some of the sensitivity tests and further studies with relevant data on maternal, offspring, and paternal genotype are required to obtain more precise (and unbiased) causal estimates.

### Conclusions

Our findings provide little evidence to support a strong causal intrauterine effect of incrementally greater maternal BMI resulting in greater offspring adiposity.

## Author Summary

### Why Was This Study Done?

- › Being overweight or obese in pregnancy causes babies to be born larger.
- › Whether these babies go into childhood and adolescence more overweight as a result of their mothers being overweight in pregnancy has not been fully established.
- › We wanted to investigate whether genetically elevated body mass index (BMI) of mothers in pregnancy was causally associated with higher levels of fatness in their offspring in childhood and adolescence, over and above the association expected given genetic transmission of BMI-associated variants.

### What Did the Researchers Do and Find?

- › We used information on 6,057 mother–child pairs from two prospective birth cohort studies, the Avon Longitudinal Study of Parents and Children (ALSPAC) and the Generation R study to assess this.
- › Our findings provide little evidence to support the long-term impact of maternal BMI in pregnancy on the child's risk of fatness in childhood and adolescence.
- › Rather, most of the association between a mother's BMI in pregnancy and her child's fatness is explained by genetic transmission of BMI-associated variants.

### What Do These Findings Mean?

- These findings suggest that public health interventions directed at all family members and at different stages of the life course are likely to be important and are potentially more likely to halt the obesity epidemic than a focus on maternal overweight and obesity status in pregnancy.
- Although we found no large effects of being overweight or obese in pregnancy on offspring's risk of fatness in later life, further studies with relevant data are required to determine whether there might be smaller effects that our study was unable to detect.

**Citation:** Richmond RC, Timpson NJ, Felix JF, Palmer T, Gaillard R, McMahon G, et al. (2017) Using Genetic Variation to Explore the Causal Effect of Maternal Pregnancy Adiposity on Future Offspring Adiposity: A Mendelian Randomisation Study. *PLoS Med* 14(1): e1002221. doi:10.1371/journal.pmed.1002221

**Academic Editor:** Jenny E. Myers, University of Manchester, UNITED KINGDOM

**Received:** August 10, 2016; **Accepted:** December 14, 2016; **Published:** January 24, 2017

**Copyright:** © 2017 Richmond et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** Data used for this submission will be made available on request to the ALSPAC executive committee (alspac-exec@bristol.ac.uk). The ALSPAC data management plan (available here: <http://www.bristol.ac.uk/alspac/researchers/data-access/documents/alspac-data-management-plan.pdf>) describes in detail the policy regarding data sharing, which is through a system of managed open access. The Generation R study has an open policy in regard to collaboration with other research groups (<http://www.generationr.nl/researchers/collaboration.html>). Requests for data access and collaboration are discussed in the Generation R Study Management Team.

**Funding:** This work was supported by grants from the European Research Council (ObesityDevelop; 669545), the US National Institute of Health (R01 DK10324), and the Wellcome Trust (WT094529MA). RCR, NJT, GDS, and DAL work in a unit that receives funds from the University of Bristol and the UK Medical Research Council (MC\_UU\_12013/1, MC\_UU\_12013/3, and MC\_UU\_12013/5). RCR is supported by Cancer Research UK (C18281/A19169). DAL is a UK National Institute of Health Research Senior Investigator (NF-SI-0611-10196). The UK Medical Research Council and Wellcome Trust (102215/2/13/2) and the University of Bristol provide core support for Avon Longitudinal Study of Parents and Children (ALSPAC). A grant from the Wellcome Trust funded genotyping in the ALSPAC mothers (WT088806). ALSPAC offspring genome-wide association studies data were generated by Sample Logistics and Genotyping Facilities at the Wellcome Trust Sanger Institute and Laboratory Corporation of America (LabCorp) using support from 23andMe. The general design of Generation R Study is made possible by financial support from the Erasmus Medical Center, Rotterdam; the Erasmus University, Rotterdam; the Netherlands Organisation for Health Research and Development (ZonMw); the Netherlands Organisation for Scientific Research (NWO); the Ministry of Health, Welfare, and Sport; and the Ministry of Youth and Families. VWJ received an additional grant from the Netherlands Organization for Health Research and Development (VIDI 016.136.361) and a European Research Council Consolidator Grant (ERC-2014-CoG-648916). JFF has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 633595 (DynaHEALTH). The Generation R Study received funding from the European Union's Horizon 2020 research and innovation programme (733206, LIFECYCLE). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** I have read the journal's policy and the authors of this manuscript have the following competing interests: GDS is a member of the Editorial Board of *PLOS Medicine*. In addition to grant funding that is acknowledged in the paper and that was relevant to the conduct of the study reported in the paper, the University of Bristol has received funds from public, charity, and industry funders from grants on which DAL is the Principal applicant (UK Medical Research Council, UK Economic and Social Research Council, UK National Institute of Health Research, Wellcome Trust, British Heart Foundation, Roche Diagnostics, Ferring Pharmaceuticals, and Medtronic PLC.) These funders had no impact on any aspect of the work presented in this paper.


**Abbreviations:** ALSPAC, Avon Longitudinal Study of Parents and Children; BMI, body mass index; DXA, dual-energy X-ray absorptiometry; FMI, fat mass index; IV, instrumental variable; IVW, inverse-variance weighted; GWAS, genome-wide association studies; MR, Mendelian randomisation; SD, standard deviation

## Introduction

The developmental overnutrition hypothesis suggests mechanisms by which intrauterine conditions related to greater maternal adiposity might affect lifelong risk of offspring fatness [1]. Maternal body mass index (BMI) is positively associated with greater pregnancy-related increases in circulating glucose, lipids, and fatty acids [2,3], and in turn higher maternal gestational levels of these nutrients are associated with greater birth size [4–6]. In support of this hypothesis, strong evidence for a causal effect of greater maternal gestational adiposity and circulating fasting glucose, but not triglycerides, on birth weight and ponderal index at birth has recently been shown using a Mendelian randomisation (MR) approach [7]. As birth size is correlated with later size, it is possible that these effects will extend into later offspring postnatal life. In addition to this proposed tracking effect, it has been suggested that intrauterine exposure to higher levels of adiposity-related nutrients, such as glucose, results in permanent changes to offspring appetite control, neuroendocrine functioning, or energy metabolism, which subsequently result in greater adiposity in later life, irrespective of any effect on birth size [8–10]. Ascertaining whether greater maternal gestational adiposity results in greater offspring adiposity through intrauterine effects is important because if it does then that mechanism could result in acceleration of the obesity epidemic across generations [11,12] and would emphasise the importance of preconception or antenatal interventions in women of reproductive age to halt and reverse the obesity epidemic [13,14].

Several large cohort studies have shown that maternal pre- or early-pregnancy BMI is positively associated with offspring fatness, measured with BMI, waist circumference, or more direct assessments of fat mass, across the whole maternal BMI distribution [1,15–18]. However, due to the high heritability of adiposity and shared environmental and behavioural characteristics between mothers and their offspring, it is impossible to determine specific intrauterine effects from such studies [1,19,20]. An intergenerational MR design in which maternal genetic variants are used as instrumental variables (IVs) for environmentally modifiable intrauterine exposures, such as exposure to greater maternal adiposity, may be useful for providing insights into the causal effect of these exposures on later offspring outcomes (Fig 1) [21]. To our knowledge, this approach has only been used once to examine the causal effect of maternal BMI on offspring adiposity in childhood [22]. In that study, a variant in the *FTO* gene was

used as an IV for pre-pregnancy BMI. The results suggested no causal effect of maternal gestational BMI on offspring dual-energy X-ray absorptiometry (DXA)-determined fat mass at age 10 y once offspring genotype was taken into account. However, the CI for the IV estimate was very wide, and it could not be statistically distinguished from the positive multivariable association of maternal pregnancy BMI with offspring fat mass. That MR study and, indeed, most of the conventional multivariable association studies have examined associations only in infancy or early childhood rather than into adulthood [1]. Examining associations at older ages is important because the potential for this mechanism to accelerate the obesity epidemic relates to maternal gestational adiposity, influencing their daughters' adiposity during their reproductive years, such that they go into their pregnancies somewhat fatter and influence the next generation and so on through generations [1].



**Fig 1. Intergenerational MR analysis to investigate a causal intrauterine effect of maternal BMI on offspring adiposity.**  
<http://dx.doi.org/10.1371/journal.pmed.1002221.g001>

Our aim was to use MR to investigate the causal effect of intrauterine exposure to greater pre-/early-pregnancy maternal BMI on offspring BMI and fat mass using repeat measurements from childhood to early adulthood. The use of an allele score, generated from a large number of BMI genetic variants identified in genome-wide association studies (GWAS), can act as a stronger genetic instrument than a single variant [23,24]. Furthermore, we sought to replicate findings in an independent study to explore whether any results might be due to chance and, if results were replicated, to further increase study power by pooling results from the two studies.

## Methods

### Cohorts and Selection of Participants

We used data from mother–offspring pairs who participated in the Avon Longitudinal Study of Parents and Children (ALSPAC) [25,26] cohort in the main analysis and from the Generation R Study [27] in the replication analysis. In both studies, only singleton births were included because of the markedly different intrauterine growth patterns between singleton and multiple births. In ALSPAC, data were available on offspring BMI- and DXA-determined fat mass at multiple ages between 7 to 18 y ( $n = 2,521$  to 3,720 mother–offspring pairs for different ages), whereas in Generation R ( $n = 2,337$ ), offspring BMI at approximately 6 y of age was available.

### ALSPAC

The ALSPAC is a population-based prospective birth cohort study that enrolled 14,541 pregnant women residing in the former County of Avon, United Kingdom, with an expected delivery date between April 1, 1991 and December 31, 1992 [26,27]. The study website contains details of all available data through a fully searchable data dictionary (<http://www.bristol.ac.uk/alspac/researchers/data-access/data-dictionary/>). Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. Of the 13,678 live-born singleton offspring in ALSPAC, genotype data were available for 5,206 mother–offspring pairs. For the main analysis, self-reported pre-pregnancy BMI was available for 4,629 of the included mother–offspring pairs, and of these, 3,720 offspring BMI measures were obtained at a clinic when the offspring were a mean age of 7.5 y (referred to as 7-y assessment). In ALSPAC, adiposity measures were assessed at five further follow-up clinics. These clinics took place when the offspring were mean ages 9.8, 11.7, 13.8, 15.4, and 17.8 y (henceforth referred to as 10, 12, 14, 16, and 18 y). The number of eligible individuals at these later clinics for inclusion in this study varied from 3,496 at age 10 to 2,521 at age 18 due to loss to follow-up at the later time points.

### Generation R

The Generation R study is a multiethnic population-based prospective cohort study from early pregnancy onward based in Rotterdam, the Netherlands [27]. All pregnant women living in the study area with a delivery date between April 2002 and January 2006 were eligible for enrolment. The study protocol was approved by the Medical Ethical Committee of the Erasmus Medical Centre, Rotterdam. The cohort includes 9,778 mothers and their children (9,749 live-born children). Of these, 8,880 (91%) women were enrolled during pregnancy and 8,633 had singleton live births. Genotype data were available for 3,909 mother–offspring pairs. For the main analysis, self-reported pre-pregnancy BMI was available for 3,199 of the included mother–offspring pairs and of these, 2,337 offspring BMI measures at a 6-y clinic (offspring mean age 6.2 y) were available.

### Anthropometry

Self-reported pre-pregnancy weight and height were obtained for the mothers in both cohorts during pregnancy and used as the main exposure. In both studies, recorded weight at the first antenatal clinic visit correlated very highly with their maternal report of pre-pregnancy weight at recruitment (correlation coefficients = 0.96). For each study, age (in 1-y categories) z-scores for maternal BMI were derived using internal standardisation. In ALSPAC and Generation R, offspring height and weight were measured at research clinics and used to calculate offspring BMI. For each study, internally standardised sex and age (in month categories) z-scores for offspring BMI and (in ALSPAC only) DXA-determined fat mass index (FMI) were calculated.

### ALSPAC

After enrolment, the mother was asked to report her height and pre-pregnancy weight in a questionnaire administered at 12 wk gestation, from which pre-pregnancy BMI was calculated (as weight [in kilograms] divided by height [in meters] squared). The correlation of pre-pregnancy weight obtained by questionnaire and weight measured at the first antenatal visit (10–12 wk gestation) was 0.96. The women's partners (fathers of the children) reported their own heights and weights in questionnaires completed at the same time as the mothers; these were used to determine their BMI.

From age 7 onwards, offspring were invited to follow-up clinics where anthropometry was measured. Weight and height were measured with the child in light clothing and without shoes. Weight was measured to the nearest 0.1 kg using a Tanita Body Fat Analyser (Model TBF 305; Tanita UK Limited, Viewsley, UK) and height to the nearest 0.1 cm using a Harpenden Stadiometer (Holtain Limited, Dyfed, UK). BMI ( $\text{kg/m}^2$ ) was then calculated as weight (in kilograms) divided by height (in meters) squared. The age of the child when they attended the research clinic was recorded in months.

From the 9-y clinic onwards, total fat mass (kg) and total lean mass (kg) were assessed by whole body DXA with the use of a Lunar Prodigy DXA scanner (GE Medical Systems Lunar, Madison, WI). The scans were visually inspected and realigned where necessary. Once complete, the tester examined the scan to ensure its quality and, if necessary, repeated the scan. FMI was subsequently calculated as total fat mass (in kilograms) divided by height (in meters) squared.

#### Generation R

Information about maternal weight just before pregnancy was obtained by questionnaire, and height was measured without shoes and heavy clothing on enrolment, from which BMI ( $\text{kg/m}^2$ ) was calculated. The correlation of pre-pregnancy weight obtained by questionnaire and weight measured at the first antenatal visit was 0.96. Partner's height (cm) and weight (kg) were also measured at enrolment.

When the children were age 5 onwards, they were invited to participate in a hands-on assessment of anthropometry. Their height and weight without shoes and heavy clothing were measured and used to calculate BMI. Weight (kg) was measured using a mechanical personal scale (SECA, Almere, The Netherlands) and height (m) was determined in standing position using a Harpenden stadiometer (Holtain Limited, Dyfed, UK). The age of the child when they attended the research clinic was recorded in months.

#### Genotyping

The genotyping procedures and details of the BMI SNPs obtained from genotyping for the mothers and offspring in the two cohorts are outlined below.

#### ALSPAC

ALSPAC mothers were genotyped using the Illumina 660K quad SNP chip (Illumina Inc., San Diego, CA, US) at the Centre National de Genotypage, Paris. PLINK software [28] (v1.07) was used to carry out quality control (QC) measures. Individuals with incorrect sex assignments, unusual genome-wide or X-chromosome heterozygosity, disproportionate levels of individual missingness (>5%), evidence of cryptic relatedness (>12.5% identity-by-descent [IBD]), or those of non-European ancestry determined from multidimensional scaling analysis seeded with individuals from the International HapMap project [29] were excluded. The resulting dataset consisted of 8340 individuals [30].

SNPs with a minor allele frequency (MAF) of <1%, a call rate of <95%, or those not in Hardy–Weinberg equilibrium (HWE,  $p < 1 \times 10^{-6}$ ) were removed [30]. Imputation of the directly genotyped data has been conducted with MaCH (v1.0.16) Markov Chain Haplotyping software [31,32] using CEU individuals from HapMap phase 2 (release 22) as a reference set for autosomal imputation [29].

ALSPAC offspring were genotyped using the Illumina HumanHap550 quad genome-wide SNP genotyping platform (Illumina Inc., San Diego, CA, US) by the Wellcome Trust Sanger Institute (Cambridge, UK) and the Laboratory Corporation of America (Burlington, NC, US). A similar QC procedure to that performed in the ALSPAC mothers was carried out. Individuals with incorrect sex assignments, extreme heterozygosity (<0.320 and >0.345 for Wellcome Trust Sanger Institute data and <0.310 and >0.330 for LabCorp data), disproportionate levels of individual missingness (>3%), evidence of cryptic relatedness (>10% IBD), or those of non-European ancestry were excluded. The resulting dataset consisted of 8,365 individuals. [33,34]

SNPs with a MAF of <1%, a call rate of <95%, or those not in HWE ( $p < 5 \times 10^{-7}$ ) were removed [33,34]. Imputation of the directly genotyped data has been conducted in the same way as the mothers.

Established BMI variants [35,36] in the ALSPAC mothers and offspring were extracted from the data set imputed using the HapMap individuals as a reference set (in which all genotypes for BMI were present).

#### Generation R

Genome-wide data are currently not available for the mothers in Generation R, but DNA is available for genotyping in candidate gene or replication studies. Therefore, custom genotyping of 32 SNPs identified in a large-scale GWAS meta-analysis for BMI [35] was carried out by LGC Genomics (formerly Kbiosciences) using a Taqman allelic discrimination assay (Applied Biosystems, Foster City, CA) and Abgene QPCR ROX mix (Abgene, Hamburg, Germany) with a call rate of 99.3%, duplicate concordance of 99.8%, and all considered SNPs found to be in HWE, with the exception of rs4836133 (*ZNF608*;  $p = 3 \times 10^{-15}$ ). To confirm the accuracy of the genotyping results, 276 randomly selected samples were genotyped for a second time using the same method, with an error rate of <1%.

Genome-wide genotyping in the offspring was performed using either the Illumina HumanHap 610 or 660 Quad chips (Illumina Inc., San Diego, US) depending on time of DNA collection [37,38]. QC of the genotype and imputation process was performed in the study as previously described [38,39]. In brief, individuals with duplicate detection, low call rates (<97.5%), sex mismatches, and high heterozygosity (>4 standard deviations [SDs]) were excluded, and SNPs with low call rates (<98.0%), not in HWE ( $p < 1 \times 10^{-6}$ ), with a low MAF (< 0.1%) or those with differential missingness between the two chips ( $p < 1 \times 10^{-7}$ ) were excluded.

Ethnic composition of the sample was estimated by Identity-By-State analysis using principal components analysis (PCA) seeded with International HapMap Phase 2 release 22 individuals [29]. Participants were defined as being of non-European ancestry when they deviated more than 4 SDs from the CEU panel mean value in any of the first four principal components. Cryptic familial relationships were identified through IBD analysis.

MaCH imputation [32] of the offspring genetic data was done using both HapMap Phase 2 release 22 [29] and 1000 Genomes Phase 3 [40] reference panels, using all available haplotypes from the different populations in a "cosmopolitan" approach [38]. Established BMI variants [35] were extracted from the 1000 Genomes imputed data set due to missingness of three of the SNPs in the HapMap phase 2 data set.

From the maternal and offspring genotype data, weighted BMI allele scores were generated using 32 independent variants that have been shown to be reliably associated with BMI in both cohorts [41]. In ALSPAC, it was also possible to generate an additional allele score composed of 97 independent variants associated with BMI in a more recent GWAS [36], as ALSPAC has full genome-wide data on both mothers and offspring. Allele scores were derived using the dose of the effect (BMI-increasing) allele at each SNP, which was first weighted by the effect size of the variant in GWAS [36,41] and then summed:

$$\text{Weighted BMI score} = w_1 \times \text{SNP}_1 + w_2 \times \text{SNP}_2 + \dots + w_n \times \text{SNP}_n$$

where  $w$  is the weight (i.e., the beta-coefficient of association of the SNP with BMI from the published GWAS) and SNP is the dosage of BMI-raising alleles at that locus (i.e., 0, 1, or 2 BMI-raising alleles). The weights used are provided in S1 Table. The score was then rescaled to reflect the average number of BMI-increasing alleles carried by an individual using the formula described in Lin et al. [42]:

$$\text{Rescaled weighted BMI score} = \frac{\text{Weighted score} \times \text{Number of SNPs available}}{\text{Sum of weights of available SNPs}}$$

#### Other Variables

Parental social class and education, maternal smoking during pregnancy, parity, and paternal BMI were considered as potential confounding factors in the multivariable regression analyses. Ethnicity was also considered as a potential confounder in Generation R, a multiethnic cohort (in ALSPAC, 95% of participants were white European origin). Details of how each of these confounders were assessed in the two cohorts are outlined below.

#### ALSPAC

Parity, defined as the number of previous pregnancies resulting in a live or stillbirth, was recorded in a questionnaire completed at 18 wk gestation. In a questionnaire completed at 32 wk of gestation, mothers recorded the occupation and education of themselves and their partners. Highest occupation of the mother or their partner was used to allocate family social class groups (classes I [professional occupations], II [managerial and technical occupations], III non-manual [skilled non-manual occupations], III manual [skilled manual occupations], IV [partly skilled occupations], V [unskilled occupations]) using the 1991 British Office of Population Censuses and Surveys classification. Highest educational qualifications of the mother and father were treated as separate variables, and each was collapsed into one of four categories: education up to age 16 y with vocational training or certificate of secondary education, education up to age 16 y with general certification of education (ordinary level), education up to age 18 with general certificate of education (advanced level), and university degree. Information on mothers' smoking status during pregnancy was obtained in questionnaires administered at 18 and 32 wk of gestation. Data were used to generate a categorical variable: never smoked during pregnancy, smoked in early pregnancy only, and smoked throughout pregnancy. Maternal age at delivery was derived from the mother's date of birth, which was recorded at the time of recruitment, and the date of birth of her offspring. Offspring sex was recorded in the delivery room and abstracted from obstetric records and/or birth notifications.

#### Generation R

Information about the mother's parity, defined as the number of times that the woman had given birth to a foetus with a gestational age of 24 wk or more, was obtained by questionnaire at enrolment. Information about household income (euro/month) was obtained from a questionnaire administered during pregnancy. The highest completed education level (primary school, secondary school, higher education) for both mothers and fathers was obtained from questionnaires administered at enrolment. Information about maternal smoking in pregnancy was assessed by questionnaire in each trimester and was categorised into never smoked during pregnancy, smoked in early pregnancy only, and smoked throughout pregnancy. Maternal age was obtained in the questionnaire administered at enrolment. Using this and information about the gestational age of the foetus at enrolment and gestational age at birth, mother's age at delivery was derived. Offspring sex was obtained from midwife and hospital registries at birth.

#### Statistical Analysis

We examined multivariable regression associations and MR effects of maternal pre-pregnancy BMI on offspring BMI at ages 6–7 y in the two cohorts and both BMI and DXA fat mass at all ages with available data up to the age of 18 y in ALSPAC. We were unable to find any other study with relevant data on maternal and offspring genetic variants, maternal pre- or early-pregnancy BMI, and offspring BMI or fat mass in adolescence/early adulthood and so could not explore replication of ALSPAC findings at older ages. Results are presented in males and females combined as point estimates and looked very similar in both sexes, and there was no strong statistical evidence for an interaction by sex in either the multivariable or MR analyses (all  $p$ -values  $>0.07$ ).

We followed an analysis plan, which was written before analysing any data for the main analyses, while some sensitivity analyses were carried out post hoc (S1 Appendix). All analyses were undertaken using Stata (Stata Corp, TX, US), version 13.

#### Analysis of Effect of Maternal Pre-pregnancy BMI on Offspring Adiposity through Childhood to Young Adulthood in ALSPAC

Multivariable linear regression was performed to examine the association of maternal pre-pregnancy BMI with offspring BMI from ages 7 to 18 and FMI from ages 10 to 18. In the first model, maternal age, offspring age, and sex were controlled in the standardised exposure and outcome. In the second model, we additionally adjusted for potential confounding by parental social class, parental education, maternal smoking during pregnancy, parity, and paternal BMI.

MR analysis was first carried out using the maternal weighted BMI allele score composed of 32 SNPs as an IV for her pre-/early-pregnancy BMI to assess its causal effect on offspring BMI (from ages 7 to 18) and offspring FMI (from ages 10 to 18). Both multivariable and IV approaches examined the same relationship, i.e., the SD change in outcome per 1 SD increase in maternal pre-/early-pregnancy BMI.

We used two-stage least squares (TSLS) IV analysis for the MR approach. We assumed an additive genetic model as supported by the original GWAS [41]. The strength of the IV was assessed by examining the  $R^2$  and F-statistics from the first stage regression for each analysis [43]. Maternal age, offspring age, and sex were controlled in all MR analyses by the use of exposure and outcome measurements standardised for these characteristics. Despite evidence that the lifestyle and socioeconomic characteristics that commonly confound observational studies are randomly distributed with respect to genotype [44], we tested this assumption in our study by examining associations between the allele score and potential confounders of the observational association [45].

To obtain a causal estimate of the intrauterine influence of maternal BMI on offspring BMI, it is important to adjust for offspring genotype to exclude the possibility of another independent pathway between the genetic instrument and outcome, i.e., through genetic transmission from the mother to their offspring (Fig 1). This was achieved by adding the offspring weighted BMI allele score to the IV models. A z-test was used to test for a difference between this MR IV analysis and the confounder-adjusted multivariable regression analysis, with evidence for a difference between the two being indicative of the possibility of unobserved confounding in the multivariable analysis. The z-statistic was calculated by estimating the covariance between the multivariable regression and IV estimates using a bootstrapping procedure.

We generated an additional weighted allele score from 97 genetic variants, using SNP-specific weights taken from the recent GWAS meta-analysis in which they were all identified [36], in order to explore whether our main findings were consistent when using this score, with potentially greater power.

#### Replication in an Independent Cohort and Meta-Analysis

We undertook multivariable regression and MR IV analyses in the replication Generation R Study using the same methods as described above for ALSPAC, with the additional adjustment for ethnicity, by including the first 20 principal components obtained from the offspring genotype data as covariates in both analyses. We were only able to complete IV analyses using the 32-SNP weighted allele score in this cohort.

We used a fixed effects meta-analysis to combine the multivariable regression results from ALSPAC and Generation R and used Cochran's Q test and the  $I^2$  statistic to explore heterogeneity between the results from these two cohorts [46]. We took a similar approach to pooling the MR IV results. We did this by pooling the results from the two stages of the IV analyses separately: (i) the maternal allele score (with and without adjustment for offspring allele score) association with maternal BMI (first regression) and (ii) the maternal allele score (with and without adjustment for offspring allele score) association with offspring BMI at age 6 (second regression). We then combined these two pooled estimates to obtain the IV MR causal effect using the ratio estimate, i.e., the pooled results of (ii) ÷ the pooled results of (i) [47]. The standard errors of these estimates were calculated using a Taylor series approximation [48]. We pooled results using the 32-SNP scores generated in both studies and also results using the 97-SNP score in ALSPAC with the 32-SNP score in Generation R. We compared the pooled offspring allele score-adjusted MR IV results with the pooled confounder-adjusted multivariable estimates using a z-test and bootstrapping, as described above for ALSPAC.

To provide results that are more interpretable for clinical and public health use, we converted the results on the SD scale to BMI units by multiplying them by a representative value of the SD of pre-pregnancy BMI (3.7 kg/m<sup>2</sup>) and offspring BMI (2.0 kg/m<sup>2</sup>) taken from the Discovery and largest study (ALSPAC).

#### Sensitivity Analyses

In the MR analysis, it is necessary to adjust for offspring BMI allele score to separate the influence of genetic inheritance from the intrauterine effect of maternal adiposity during pregnancy. However, as was highlighted in a response to the previous paper where this method was used [49], adjustment may introduce a form of bias known as collider bias in estimating the exposure–outcome association. This is because by adjusting for offspring allele score in the MR IV analyses, we may induce an association via paternal genetic variants (which we do not have data on; S1 Fig). We explain this possibility in more detail in the appendix (S1 Appendix) and also describe how we undertook simulation studies to explore the likelihood of this biasing our main results with adjustment for offspring allele score. Furthermore, an alternative method for avoiding possible collider bias is to use only the maternal non-transmitted alleles [50]. We explored the use of this approach within ALSPAC and present results in the appendix (S1 Appendix) only given the relatively low statistical power associated with this method.

A high degree of heterogeneity between causal estimates of the individual SNPs comprising the allele scores could indicate violation of the MR assumption that there is no pleiotropy. We therefore performed inverse-variance weighted (IVW) meta-analysis of the individual SNP estimates in both ALSPAC and Generation R and calculated Cochran's Q and  $I^2$  statistic to estimate the degree of heterogeneity in the fixed effects meta-analysis [51]. We also investigated potential bias due to pleiotropy by performing MR Egger regression [52] for the model adjusted for offspring allele score. The intercept in this analysis provides a test for overall directional pleiotropy, and the coefficient provides a valid causal estimate in the presence of pleiotropy. Results were obtained for both ALSPAC (using both sets of 32 SNPs and 97 SNPs) and Generation R (using the set of 32 SNPs). The intercepts and slopes were meta-analysed and compared with those obtained using IVW of the individual SNPs. Analysis was performed using the mrobust Stata package [53].

Lastly, we investigated possible nonlinearity of the association in both the multivariable regression and MR analyses by overlaying a nonparametric loess smoother and a line of best fit on an augmented partial residual plot.

## Results

Table 1 shows the key characteristics of the Discovery and Replication cohort.

	ALSPAC (Discovery)	Generation R (Replication)	Generation R (European*)
n	5,720	2,207	1,200
Males (%)	48.8%	49.8%	49.8%
Offspring age in months (SD)	88.6 (1.5)	74.9 (0.1)	73.2 (0.6)
Offspring birth weight in kg (SD)	3.463 (0.11)	3.479 (0.07)	3.401 (0.06)
Offspring weight in kg (SD)	25.4 (4.4)	23.2 (4.1)	22.8 (3.4)
Offspring height in cm (SD)	102.7 (5.3)	118.8 (5.5)	118.0 (5.4)
Offspring BMI in kg/m <sup>2</sup> (SD)	16.2 (2.5)	18.2 (3.1)	15.9 (1.4)
Maternal BMI in kg/m <sup>2</sup> (SD)	22.9 (3.7)	23.9 (4.1)	23.2 (3.8)

\*Generation R is a multiethnic cohort; these are characteristics for those of European origin only (as defined by 4 SNPs from the HapMap CEU panel) mean values for all four principal components from the genetic data.

doi:10.1371/journal.pmed.1002221.t001

**Table 1. Characteristics of the offspring and their mothers in the Discovery and Replication cohorts.**

<http://dx.doi.org/10.1371/journal.pmed.1002221.t001>

#### Analyses from Childhood to Early Adulthood in ALSPAC

The sample size at each age from 7 to 18 decreased as a result of loss to follow-up, but the proportion of males and females and the distribution of birth weight (of those remaining in the cohort) were similar in each age group (S2 Table). Height, weight, BMI, fat mass, FMI, and lean mass increased with increasing age from 7 to 18 y as expected (S2 Table).

Maternal BMI was associated with characteristics that a priori we considered to be likely confounders (S3 Table). With one exception, the maternal BMI allele score was not associated with these confounders. There was a weak inverse association with paternal education (suggesting that on average each category increase of paternal education was associated with a -0.03 [95% CI -0.07–0.00] SD change in the weighted allele score [S3 Table]). The maternal BMI allele score was normally distributed and was robustly associated with maternal pre-pregnancy BMI, explaining 2.2% of the variation in BMI (S4 Table), and with first-stage F-statistics for each of the MR analyses all being >45 (S5 Table).

Table 2 shows the confounder-adjusted multivariable associations of maternal pre-pregnancy BMI with offspring BMI and FMI at each age and the equivalent MR results (with adjustment for offspring allele scores). S6 and S7 Tables show more detailed results, including the multivariable results unadjusted for offspring allele scores and MR results. In confounder-adjusted multivariable regression, a 1 SD (equivalent to 3.7 kg/m<sup>2</sup>) higher age-adjusted maternal BMI was associated with a 0.25 (0.21, 0.29) SD higher offspring BMI at age 7 and a 0.33 SD higher offspring BMI (0.28, 0.37) at age 18 (equivalent of 0.56 and 0.76 kg/m<sup>2</sup>, respectively; Table 2). Equivalent results for the genetic IV analyses with adjustment for offspring allele score were 0.04 (-0.21, 0.30) SD at age 7 and -0.03 (-0.32, 0.26) SD at age 18 (Table 2). Results for FMI from ages 10 to 18 were similar for both the multivariable regression and MR results to those seen with BMI (Table 2). We further evaluated, in a post hoc manner, the impact of additionally adjusting for maternal and offspring allele score in the multivariable regression models and found that these additional adjustments did not substantially change the point estimates for mean difference (S14 Table).

Offspring outcome	Confounder-adjusted multivariable regression results		MR (genetic IV maternal allele score adjusted for offspring allele score) results		P <sup>a</sup> difference <sup>b</sup>
	n	Difference in mean offspring outcome (SD) per 1SD increase maternal BMI (95% CI)	n	Difference in mean offspring outcome (SD) per 1SD increase maternal BMI (95% CI)	
BMI age 7	2,585	0.25 (0.21–0.29)	2,570	0.04 (-0.21–0.30)	0.13
BMI age 10	2,587	0.31 (0.27–0.36)	2,587	0.03 (-0.23–0.29)	0.05
BMI age 12	2,611	0.32 (0.29–0.36)	2,600	0.01 (-0.26–0.26)	0.02
BMI age 14	2,604	0.30 (0.28–0.32)	2,627	-0.07 (-0.34–0.20)	0.01
BMI age 16	1,979	0.34 (0.30–0.38)	2,006	-0.10 (-0.41–0.20)	0.003
BMI age 18	1,798	0.33 (0.29–0.37)	2,071	-0.03 (-0.30–0.24)	0.01
FMI age 10	2,613	0.30 (0.28–0.33)	2,606	0.13 (-0.13–0.39)	0.071
FMI age 12	2,675	0.31 (0.27–0.36)	2,644	0.04 (-0.22–0.30)	0.062
FMI age 14	2,233	0.30 (0.28–0.34)	2,192	0.03 (-0.23–0.29)	0.043
FMI age 16	1,927	0.33 (0.29–0.38)	2,715	-0.10 (-0.40–0.21)	0.001
FMI age 18	1,730	0.33 (0.29–0.37)	2,400	0.03 (-0.27–0.30)	0.008

In all analyses, offspring BMI has been standardised on their sex and age and maternal BMI has been standardised on her age.  
<sup>a</sup>The following were adjusted for in multivariable regression analyses: parental social class, parental education, maternal smoking during pregnancy, parity, and paternal BMI.  
<sup>b</sup>Testing the null hypothesis that the multivariable regression analysis results do not differ from the MR results.  
Abbreviations: BMI, body mass index; FMI, fat mass index.  
doi:10.1371/journal.pmed.1002221.t002

**Table 2. Confounder-adjusted multivariable and genetic IV (MR) associations of maternal pregnancy BMI with offspring BMI and FMI from ages 7 to 18 in ALSPAC (Discovery sample).**  
<http://dx.doi.org/10.1371/journal.pmed.1002221.t002>


There was strong statistical evidence that the MR IV analysis results differed from the multivariable regression analysis results, with the exception of offspring BMI at age 7 and FMI at age 10, where the IV associations had point estimates that were smaller than the multivariable regression analysis results but were statistically consistent with those results (Table 2). Results were similar when the 97-SNP BMI allele score was used as an IV (S8 and S9 Tables).

**Replication and Meta-Analysis with an Additional Independent Cohort**

Maternal BMI was associated with potential confounders in Generation R (S10 Table). The maternal BMI allele score was normally distributed with similar mean and SD as in ALSPAC and was robustly associated with maternal pre-pregnancy BMI (S4 Table). Similar to ALSPAC, maternal BMI allele score was weakly inversely related to paternal education in Generation R but not to other observed confounders (S10 Table).

Age- and sex-adjusted results were similar to those in ALSPAC, although there was evidence of heterogeneity in the confounder-adjusted multivariable regression results (I<sup>2</sup> = 82%; S2 Fig). Pooling results from both studies showed that a 1 SD (equivalent of 3.7 kg/m<sup>2</sup>) increase in age-adjusted maternal BMI was associated with a 0.22 (0.19, 0.25) SD (equivalent of 0.44 [0.38, 0.50] kg/m<sup>2</sup>) increase in offspring BMI in the confounder-adjusted model. The maternal BMI allele score was similarly positively associated with offspring BMI in both studies but with adjustment for offspring allele score the associations attenuated to the null, with no evidence for heterogeneity between the estimates (S3 Fig).

There were similar associations of the maternal allele score with maternal BMI and with offspring BMI between the two cohorts (ALSPAC and Generation R; S3 Fig). Although the MR effect when pooling the offspring-adjusted results based on the 32-SNP allele score in both cohorts appeared weaker than our confounder-adjusted multivariable estimate, 0.10 (-0.11, 0.31) SD (equivalent of 0.20 [-0.22, 0.62] kg/m<sup>2</sup>) versus 0.22 (0.19, 0.25) SD (equivalent of 0.44 [0.38, 0.50] kg/m<sup>2</sup>) per 1 SD greater maternal pre-pregnancy BMI, there was no strong statistical evidence that these two estimates differed from each other (p(diff) = 0.34; Fig 2). When we pooled the MR results using the 97-SNP allele score from ALSPAC with those from Generation R using the 32-SNP allele score, the point estimates were closer to the null and were more precisely estimated (0.05 [-0.11, 0.21] SD per 1 SD greater maternal pre-pregnancy BMI), and there was increasing statistical evidence for a difference between these MR results and the confounder-adjusted multivariable estimate (p(diff) = 0.05; Fig 2).



**Fig 2. Relationship of maternal pregnancy BMI with offspring BMI at age 7: pooled ALSPAC and Generation-R multivariable and MR analyses.**  
<http://dx.doi.org/10.1371/journal.pmed.1002221.g002>

All results are difference in mean offspring BMI at age 7 (SD units) per increase in 1 SD maternal pregnancy BMI pooling the results from ALSPAC (Discovery cohort) with those from Generation R Study (Replication cohort). In all analyses, offspring age and sex are controlled for through standardisation of the offspring BMI, and maternal age is controlled for by standardisation of BMI on age. The multivariable regression analyses results are controlled for parental social class, parental education, maternal smoking



during pregnancy, parity, and paternal BMI and in Generation R only for ethnicity (adjusting for top 20 principal components of offspring genome-wide array data). The MR results are adjusted for offspring allele score and in Generation R only for ethnicity (adjusting for top 20 principal components of offspring genome-wide data).

Given the weak association of the maternal allele score with paternal education in both cohorts, we re-ran all of the IV analyses with additional adjustment for paternal education. Additional adjustment for paternal education did not alter the null MR estimate following adjustment for offspring allele score (S4 Fig). In addition, analyses restricted to those of European ancestry in Generation R were similar to those including all Generation R participants (S5 Fig).


#### Sensitivity Analyses

Our investigation of the possibility of introducing bias via paternal genotype suggested that the offspring allele score-adjusted MR estimate was the least biased result of the causal effect of maternal pre-pregnancy BMI on offspring BMI. Although the inability to also adjust for paternal genetic variants meant this was somewhat biased towards the null, it was unlikely to have markedly changed our findings (S1 Appendix and S11 Table). Furthermore, we explored the use of the non-transmitted (to offspring) haplotype approach [50] in ALSPAC and both the 32-SNP and 97-SNP haplotype scores revealed that the non-transmitted maternal haplotype score did not show strong evidence of association with offspring while the transmitted maternal haplotype score was strongly associated, indicating the expected offspring genetic influence on their own BMI but providing little support for a maternal BMI intrauterine effect on offspring BMI (S1 Appendix and S12 Table).

Results of the IVW MR approach showed estimates approximately equal to the TSLS allele score approach. Furthermore, this approach showed no clear evidence for heterogeneity of effect estimates of the individual SNPs comprising the score in ALSPAC or Generation R ( $I^2 = 0\%$ ; S6 Fig, S13 Table). In addition, the MR Egger method gave no indication of directional pleiotropy influencing the results of the MR analysis (intercept = 0.005 [-0.003, 0.013],  $p = 0.13$ ) for the meta-analysis of 97 SNPs in ALSPAC and 32 SNPs in Generation R) and provided evidence for a lack of consistent causal effect of maternal BMI on offspring BMI in the models adjusted for offspring genotype (coefficient = 0.08 SD [-0.12, 0.28],  $p = 0.07$ ) in the meta-analysis of estimates from ALSPAC and Generation R, although there was some degree of heterogeneity between the two studies ( $I^2$  for slope = 70%; S6 Fig, S13 Table). There was no strong evidence for departure from linearity in the relation of maternal BMI with offspring BMI in either multivariable or MR analyses (S7 Fig and S8 Fig).

## Discussion

In ALSPAC, we found positive associations of maternal pre-pregnancy BMI with offspring BMI and FMI at all ages from childhood to early adulthood in confounder-adjusted multivariable regression analyses. However, we found no evidence for these associations being causal in MR IV analyses using a weighted allele score of 32 (or 97) genetic variants known to be robustly associated with BMI. There was statistical evidence that the confounder-adjusted multivariable analyses differed from the offspring genetic variant-adjusted MR analyses, and results were virtually identical when we used FMI measured by DXA scan. ALSPAC findings were replicated in the independent Generation R Study with BMI assessed at age 6 y. Taken together, these results do not support an important causal intrauterine effect of greater maternal BMI on later offspring adiposity. This is in contrast to evidence of a causal effect of greater maternal adiposity on birth weight and ponderal index at birth identified using MR in a previous study that included both of the cohorts used here [7], potentially indicating a diminishing effect of this intrauterine exposure over the life course (Fig 3).



**Fig 3. Diminishing causal effect of developmental overnutrition across the life course.**

Multivariable and IV effect estimates from ALSPAC at ages 7–18 (Table 2) compared with those obtained when investigating the effect of maternal BMI on offspring ponderal index ( $\text{kg}/\text{m}^3$ ) at birth in the same cohort.

<http://dx.doi.org/10.1371/journal.pmed.1002221.g003>

Findings of a null causal effect from MR analysis in both ALSPAC (at all ages from childhood to early adulthood) and Generation R (in childhood) are consistent with most of the negative control studies, which show a similar magnitude of association between maternal pre-pregnancy BMI and offspring BMI as that between paternal BMI and offspring BMI [54–59], although none of those studies assessed outcomes in early adulthood. They are also consistent with a large within-sibling analysis, which suggested that shared familial characteristics confounded the positive maternal early-pregnancy BMI–offspring BMI association at age 18, as this disappeared when analyses were conducted within siblings with close matching of shared family characteristics [60]. However, that study was performed in male offspring only. Lastly, the results are consistent with the previous MR study in ALSPAC conducted using the *FTO* variant only as an IV for maternal BMI and examining offspring BMI- and DXA-determined fat mass at age 10 [22]. By combining a large number of genetic variants in an allele score, in this study, we have increased statistical power and shown a difference between the multivariable and IV analysis in ALSPAC at multiple ages (from childhood to early adulthood), which was not the case with the previous ALSPAC publication. We have also shown independent replication of our findings in Generation R, investigated the use of multiple instruments, and tested for potential pleiotropic effects.

For the MR analysis, we thoroughly investigated any violation of the main assumptions of this approach, which are that the IV is robustly associated with the exposure of interest, not related to confounding factors of the exposure–outcome association, and not related to the outcome independent of its effect on the exposure [45].

The allele scores derived from established variants for BMI were strongly associated with the exposure of interest (maternal BMI) in the full cohorts, and large F-statistics indicated sufficient strength as IVs, enabling a more precise assessment of the effect of maternal BMI than was previously possible with the use of a single genetic variant [22].

The allele scores were not consistently associated with a range of socioeconomic factors, and this illustrates a key strength of the MR approach. In addition, by adjusting for offspring genotype, intergenerational MR is able to distinguish between causal maternal effects on offspring adiposity from genetic transmission of adiposity variants.

The existence of pleiotropy, in which a genetic instrument has an effect on an outcome (offspring BMI) independent of its effect on the exposure (maternal BMI), would have implications for the assumptions made in the MR analysis. Similarly, if a genetic variant in the score was in linkage disequilibrium with another genetic variant that influences the outcome through a pathway that is unrelated to the exposure, this could bias the causal estimate. The fact that some BMI variants relate to glucose and other metabolites [61,62] is an example of type 2 or spurious pleiotropy and is unlikely to bias the IV effects [63]. We have attempted to control for the effect of offspring genotype, which is related to maternal genotype, on offspring BMI by adjusting for offspring genotype. It is hard to determine a real pleiotropic effect that would produce the null effects seen following this adjustment. In addition, the consistency of IV estimates obtained using two allele scores in this study suggests that pleiotropy is unlikely. Furthermore, there was no strong evidence for heterogeneity of individual SNP estimates comprising the allele scores, and the MR Egger method gave no indication of directional pleiotropy in ALSPAC or Generation R.

Although maternal genetic variants are being used as an unconfounded proxy for an intrauterine exposure, these variants will also impact on maternal BMI after pregnancy and so any causal effect, had it been identified, might also incorporate postnatal effects. For example, mothers who are genetically predisposed to have a higher BMI might influence their offspring's adiposity through postnatal characteristics such as maternal feeding behaviours [64]. The heterogeneity observed in the confounder-adjusted multivariable models could represent variation in prevailing postnatal environmental contexts, for example cultural influences on parents' child-feeding behaviour [65]. However, the MR effects were null in both ALSPAC and Generation R, suggesting that the causal effect of postnatal maternal BMI on offspring adiposity is at most very small. The weak inverse association of maternal BMI allele scores with their partners (but not their own) education that we saw in both cohorts is intriguing and worth further exploration, but adjusting for this did not alter our MR results.

It is important to adjust for offspring genotype in the MR analyses, but in doing so we might introduce a path between maternal genotype and offspring BMI via paternal genotype. Our simulations suggested that this would result in a weak bias towards the null, but this bias would not fully explain the null effects we observe as the results in the simulation study with adjustment for offspring allele score were close to the true simulated result (S11 Table). Furthermore, our findings were corroborated with results from the non-transmitted haplotype analysis [50], applied in ALSPAC, which revealed that the non-transmitted maternal haplotype score did not show strong evidence of association with offspring BMI.

In the MR meta-analysis using the 97-SNP allele score in ALSPAC and the 32-SNP allele score in Generation R, with adjustment for offspring genotype ( $n = 6,057$ ), we had 82% power to detect a causal effect the size of that seen in the multivariable analysis controlled for age and sex (0.28 SD) with a two-sided  $\alpha = 0.05$  in our MR analysis. Therefore, we were adequately powered to detect an effect the size of that seen observationally, and the heterogeneity we see between the multivariable and MR results is unlikely to be due to chance. However, we would need a much larger number of participants to completely rule out evidence for a weaker causal effect.

Furthermore, power was limited for some of the sensitivity tests such as the evaluation of SNP heterogeneity, the MR Egger analysis, and the evaluation of heterogeneity in causal estimates between ALSPAC and Generation R. Further studies with relevant data on maternal, offspring, and paternal genotype are required to obtain more precise (and unbiased) causal estimates.

Further limitations to the MR analysis include the possibility of population stratification and canalisation [66]. Although in ALSPAC population stratification is unlikely because the participants are unrelated individuals of European ancestry, the Generation R Study is a multiethnic cohort. However, attempts made to adjust for population stratification by including principal components and analyses restricted to those of European ancestry were similar on the whole (S3 Fig). Developmental canalisation, in which systems develop differently to counterbalance the effects of a particular genotype, may pose a problem for conventional MR. However, this is less of an issue in intergenerational MR analysis. This is because when maternal genotype is used as an indicator of the intrauterine environment, then this will only influence the developmental environment of the offspring through the exposure of interest and not by competing mechanisms [67].

Maternal pre-pregnancy BMI was self-reported in both cohorts. Although these measures of self-report have been shown to correlate strongly with measured BMI in early pregnancy [68–70], the possibility of systematic under-reporting, for example, if those who are heavier systematically under-report their weight, might bias findings. However, previous analysis in ALSPAC has shown that misreporting is similar for the majority of participants and is not markedly influenced by mean weight [71]. Furthermore, the magnitude of the association of the weighted allele score with maternal BMI in this study was similar to that seen for its relationship to BMI based on measured weight and height from other studies [7].

A general limitation of the longitudinal measures in ALSPAC is loss to follow-up of the sample over time, from 3,720 participants at age 7 to 2,521 at age 17. Nonetheless, the distribution of birth weight was similar in groups at each age. In addition, loss to follow-up bias is unlikely to influence the MR estimates, as genetic associations are unlikely to be biased by missing data [72]. This is supported by the fact that the allele scores used in this study were not strongly associated with a range of risk factors associated with loss to follow-up.

In this study we have looked at outcomes at each time point separately in order to explore whether magnitudes are similar at each age. It might also be valuable to examine whether maternal exposures relate to the rate of change in offspring adiposity across childhood and into adulthood using multilevel models. Methods of applying genome-wide data to such models have recently been developed [73]. However, we are not aware of them being used in an MR IV framework, and we feel the approach we have adopted is relevant to our aim and produces results that are easy to interpret.

Our study examined the effect of linear (incremental) increases in maternal BMI with offspring outcomes. Although methods for assessing nonlinear effects using MR have been recently developed [74], these require very large sample sizes and we are not able to apply these to our data. Thus, we cannot rule out the possibility of, for example, a nonlinear threshold effect of extreme maternal obesity having a causal intrauterine effect on offspring adiposity. However, we did not find any clear departures from

linearity for either the multivariable regression or MR analyses (S7 Fig and S8 Fig). Lastly, although the pattern of positive multivariable regression and null MR results was consistent across ages in ALSPAC, we acknowledge that it would be valuable to show further replication of our results in large independent cohorts, particularly at older ages.

Given our results, together with those from sibling comparison [60] and negative control studies [54–59,65], it seems unlikely that subtle incremental differences in maternal pre- or early-pregnancy BMI play a key role in initiating or perpetuating the obesity epidemic [11,12]. Although some negative control [22,75–77] and sibling comparisons [78,79] suggest weak positive effects of maternal pre- or early-pregnancy BMI on offspring childhood BMI, in general, those studies are smaller than the ones finding null effects and have not explored associations into adulthood [1]. Our results showed no effect up to the age of 18 in females and males in the ALSPAC cohort. This finding is important given that a lack of effect in offspring entering their reproductive life course suggests that incrementally greater maternal adiposity across the population in pregnancy is unlikely to fuel the obesity epidemic across generations, although we cannot rule out an effect of more extreme phenotypes, such as extreme obesity or gestational diabetes [1]. In addition, we have not been able to investigate the causal effect of gestational weight gain during pregnancy on offspring adiposity using MR, given the current absence of robust genetic instruments for this exposure [80].

These findings suggest that overreliance on interventions in pregnancy to reduce population obesity may not be warranted and that consensus statements [14], which direct public health interventions to all family members [81,82] and at different stages of the life course, and not just intrauterine or early life [13], are likely to be important. For example, interventions aimed at the whole population (i.e., all family members at all life course stages), such as proposals for excess tax on obesogenic foods [83], are potentially more likely to halt the obesity epidemic than a focus on maternal pre-pregnancy BMI.

## Supporting Information

### **S1 Appendix. Supplementary methods for investigating potential collider bias and the analysis plan.**

doi:10.1371/journal.pmed.1002221.s001  
(DOCX)

### **S1 Fig. Schematic diagram showing relationships for the MR analysis and how “collider bias” might occur when adjusting for offspring genotype.**

doi:10.1371/journal.pmed.1002221.s002  
(DOCX)

### **S2 Fig. Forest plot of association between maternal BMI and offspring BMI using multivariable linear regression in ALSPAC and Generation R.**

doi:10.1371/journal.pmed.1002221.s003  
(DOCX)

### **S3 Fig. Forest plot of association between maternal BMI allele score and offspring BMI using multivariable linear regression in ALSPAC and Generation R.**

doi:10.1371/journal.pmed.1002221.s004  
(DOCX)

### **S4 Fig. Multivariable and instrument variable meta-analysis to assess the association between maternal BMI and offspring BMI in ALSPAC and Generation R, additionally adjusted for paternal education in the IV analysis.**

doi:10.1371/journal.pmed.1002221.s005  
(DOCX)

### **S5 Fig. Multivariable and instrument variable meta-analysis to assess the association between maternal BMI and offspring BMI in ALSPAC and Generation R, restricted to European ancestry.**

doi:10.1371/journal.pmed.1002221.s006  
(DOCX)

### **S6 Fig. Scatterplot of maternal genetic associations with offspring BMI (adjusted for offspring genetic variants) against maternal genetic associations with maternal BMI, with causal estimates of maternal BMI on offspring BMI estimated by MR-Egger (solid line) and IVW (dashed line) methods.**

doi:10.1371/journal.pmed.1002221.s007  
(DOCX)

### **S7 Fig. Augmented partial residual plots to evaluate departures from linearity in observational and IV models from the ALSPAC main analysis.**

doi:10.1371/journal.pmed.1002221.s008  
(DOCX)

### **S8 Fig. Augmented partial residual plots to evaluate departures from linearity in observational and IV models from the Generation R main analysis.**

doi:10.1371/journal.pmed.1002221.s009  
(DOCX)

### **S1 Table. Details of the BMI SNPs obtained from genotyping and imputation.**

doi:10.1371/journal.pmed.1002221.s010  
(DOCX)

### **S2 Table. Descriptive characteristics of included participants in ALSPAC at each age.**

doi:10.1371/journal.pmed.1002221.s011  
(DOCX)

**S3 Table. Associations between maternal BMI and maternal BMI allele score and possible confounding factors in ALSPAC.**

doi:10.1371/journal.pmed.1002221.s012  
(DOCX)

**S4 Table. Characteristics of maternal BMI allele score and association with maternal BMI.**

doi:10.1371/journal.pmed.1002221.s013  
(DOCX)

**S5 Table. F-statistics for first-stage regression of IV analyses in ALSPAC and Generation R.**

doi:10.1371/journal.pmed.1002221.s014  
(DOCX)

**S6 Table. Associations between maternal BMI and offspring BMI from ages 7 to 18 using multivariable and IV methods with a 32-SNP allele score in ALSPAC (Discovery) cohort.**

doi:10.1371/journal.pmed.1002221.s015  
(DOCX)

**S7 Table. Associations between maternal BMI and offspring FMI from ages 10 to 18 using multivariable and IV methods with a 32-SNP allele score in the ALSPAC (Discovery) cohort.**

doi:10.1371/journal.pmed.1002221.s016  
(DOCX)

**S8 Table. Association between maternal BMI and offspring BMI from ages 7 to 18 using multivariable and IV methods with a 97-SNP allele score.**

doi:10.1371/journal.pmed.1002221.s017  
(DOCX)

**S9 Table. Association between maternal BMI and offspring FMI from ages 7 to 18 using multivariable and IV methods with a 97-SNP allele score.**

doi:10.1371/journal.pmed.1002221.s018  
(DOCX)

**S10 Table. Associations between maternal BMI and maternal BMI allele score and possible confounding factors in Generation R.**

doi:10.1371/journal.pmed.1002221.s019  
(DOCX)

**S11 Table. Simulation to investigate the direction and magnitude of bias in observational multivariable and IV regression models.**

doi:10.1371/journal.pmed.1002221.s020  
(DOCX)

**S12 Table. Association between transmitted and non-transmitted maternal haplotype scores and offspring BMI at age 7 in ALSPAC (Total n = 3,720).**

doi:10.1371/journal.pmed.1002221.s021  
(DOCX)

**S13 Table. Regression estimates for the application of MR-Egger and IVW methods for MR in ALSPAC and Generation R.**

doi:10.1371/journal.pmed.1002221.s022  
(DOCX)

**S14 Table. Confounder-adjusted multivariable associations of maternal pregnancy BMI with offspring BMI and FMI from ages 7 to 18 in ALSPAC (Discovery sample).**

doi:10.1371/journal.pmed.1002221.s023  
(DOCX)

## Acknowledgments

ALSPAC: We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

Generation R: The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, Rotterdam, the Rotterdam Homecare Foundation, Rotterdam, and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond (STAR-MDC), Rotterdam. We gratefully acknowledge the contribution of children and parents, general practitioners, hospitals, midwives, and pharmacies in Rotterdam. The generation and management of GWAS genotype data for the Generation R Study were done at the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, the Netherlands. We would like to thank Karol Estrada, Dr. Tobias A. Knoch, Anis Abuseiris, Luc V. de Zeeuw, and Rob de Graaf for their help in creating GRIMP, BigGRID, MediGRID, and Services@MediGRID/D-Grid for access to their grid computing resources. We thank Mila Jhamai, Manoushka Ganesh, Pascal Arp, Marijn Verkerk, Lizbeth Herrera, and Marjolein Peters for their help in creating, managing, and QC of the GWAS database. Also, we thank Karol Estrada for his support in creation and analysis of imputed data.

We would like to thank Thorkild Sørensen and Ellen Aagaard Nøhr for their helpful comments on an earlier draft of this paper, and Rachel Freathy, Jessica Tyrrell, Camilla Morgen, Maribel Casas, Mariona Bustamante, and others from the Early Growth Genetics (EGG) consortium for their insights.

## Author Contributions

**Conceptualization:** RCR NJT GDS VWJ DAL.  
**Formal analysis:** RCR TP GM.  
**Funding acquisition:** DAL GDS VWJ JFF NJT.  
**Methodology:** RCR JFF TP RG GM.  
**Software:** TP GM.  
**Supervision:** DAL.  
**Validation:** RCR JFF RG VWJ.  
**Visualization:** RCR.  
**Writing – original draft:** RCR DAL.  
**Writing – review & editing:** NJT JFF RG GDS VWJ GM TP.

## References

1. Lawlor DA. The Society for Social Medicine John Pemberton Lecture 2011. Developmental overnutrition—an old hypothesis with new importance? *International journal of epidemiology*. 2013;42(1):7–29. Epub 2013/03/20. doi: 10.1093/ije/dys209. pmid:23508404  
View Article • PubMed/NCBI • Google Scholar
2. Shin D, Song WO. Prepregnancy body mass index is an independent risk factor for gestational hypertension, gestational diabetes, preterm labor, and small- and large-for-gestational-age infants. *Journal of Maternal-Fetal & Neonatal Medicine* 2014:1–8. Epub 2014/09/12.  
View Article • PubMed/NCBI • Google Scholar
3. Harmon KA, Gerard L, Jensen DR, Kealey EH, Hernandez TL, Reece MS, et al. Continuous glucose profiles in obese and normal-weight pregnant women on a controlled diet: metabolic determinants of fetal growth. *Diabetes care*. 2011;34(10):2198–204. Epub 2011/07/22. PubMed Central PMCID: PMC3177740. doi: 10.2337/dc11-0723. pmid:21775754  
View Article • PubMed/NCBI • Google Scholar
4. Pederson J. Weight and length at birth of infants of diabetic mothers. *Acta Endocrinologica*. 1954;16:330–42. pmid:13206643  
View Article • PubMed/NCBI • Google Scholar
5. Jackson WP. The prediabetic syndrome; large babies and the (pre) diabetic father. *Journal of Clinical Endocrinology and Metabolism*. 1954;14(2):177–83. Epub 1954/02/01. doi: 10.1210/jcem-14-2-177. pmid:13130664  
View Article • PubMed/NCBI • Google Scholar
6. Freinkel N. Banting Lecture 1980. Of pregnancy and progeny. *Diabetes*. 1980;29(12):1023–35. Epub 1980/12/01. pmid:7002669  
View Article • PubMed/NCBI • Google Scholar
7. Tyrrell J, Richmond RC, Palmer TM, Feenstra B, Rangarajan J, Metrustry S, et al. Genetic evidence for causal relationships between maternal obesity-related traits and birth weight. *JAMA*. 2015;(In Press)  
View Article • PubMed/NCBI • Google Scholar
8. Gillman MW. A life course approach to obesity. In: Kuh D, Ben-Shlomo Y, editors. *A Lifecourse Approach to Chronic Disease Epidemiology*. 2nd edn. Oxford: Oxford University Press; 2004. p. 189–217.
9. Taylor PD, Poston L. Developmental programming of obesity in mammals. *Experimental Physiology*. 2007;92(2):287–98. Epub 2006/12/16. doi: 10.1113/expphysiol.2005.032854. pmid:17170060  
View Article • PubMed/NCBI • Google Scholar
10. Remmers F, Delemarre-van de Waal HA. Developmental programming of energy balance and its hypothalamic regulation. *Endocrine Reviews*. 2011;32(2):272–311. Epub 2010/11/06. doi: 10.1210/er.2009-0028. pmid:21051592  
View Article • PubMed/NCBI • Google Scholar
11. Pettitt DJ, Knowler WC. Diabetes and obesity in the Pima Indians: a cross-generational vicious cycle *Journal of Obesity and Weight Regulation*. 1988; (7):61–75.  
View Article • PubMed/NCBI • Google Scholar
12. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002;360(9331):473–82. Epub 2002/09/21. doi: 10.1016/S0140-6736(02)09678-2. pmid:12241736  
View Article • PubMed/NCBI • Google Scholar
13. Lawlor DA, Chaturvedi N. Treatment and prevention of obesity—are there critical periods for intervention? *International journal of epidemiology*. 2006;35(1):3–9. doi: 10.1093/ije/dyi309. pmid:16396899  
View Article • PubMed/NCBI • Google Scholar

14. Poston L, Harthoorn LF, Van Der Beek EM, Workshop CttIE. Obesity in pregnancy: implications for the mother and lifelong health of the child. A consensus statement. *Pediatric Research*. 2011;69(2):175–80. Epub 2010/11/16. doi: 10.1203/PDR.0b013e3182055ede. pmid:21076366  
View Article • PubMed/NCBI • Google Scholar
15. Nelson SM, Matthews P, Poston L. Maternal metabolism and obesity: modifiable determinants of pregnancy outcome. *Human Reproduction Update*. 2010;16(3):255–75. Epub 2009/12/08. PubMed Central PMCID: PMC2849703. doi: 10.1093/humupd/dmp050. pmid:19966268  
View Article • PubMed/NCBI • Google Scholar
16. Drake AJ, Reynolds RM. Impact of maternal obesity on offspring obesity and cardiometabolic disease risk. *Reproduction*. 2010;140(3):387–98. Epub 2010/06/22. doi: 10.1530/REP-10-0077. pmid:20562299  
View Article • PubMed/NCBI • Google Scholar
17. Lawlor DA, Relton C, Sattar N, Nelson SM. Maternal adiposity—a determinant of perinatal and offspring outcomes? *Nat Rev Endocrinol*. 2012;8(11):679–88. doi: 10.1038/nrendo.2012.176. pmid:23007319  
View Article • PubMed/NCBI • Google Scholar
18. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis. *PLoS ONE*. 2013;8(4):e61627. Epub 2013/04/25. PubMed Central PMCID: PMC3628788. doi: 10.1371/journal.pone.0061627. pmid:23613888  
View Article • PubMed/NCBI • Google Scholar
19. Davey Smith G. Assessing intrauterine influences on offspring health outcomes: can epidemiological studies yield robust findings? *Basic and Clinical Pharmacology & Toxicology*. 2008;102(2):245–56. Epub 2008/01/30.  
View Article • PubMed/NCBI • Google Scholar
20. Gaillard R, Felix JF, Duijts L, Jaddoe VW. Childhood consequences of maternal obesity and excessive weight gain during pregnancy. *Acta Obstetrica et Gynecologica Scandinavica*. 2014;93(11):1085–9. Epub 2014/09/19. doi: 10.1111/aogs.12506. pmid:25231923  
View Article • PubMed/NCBI • Google Scholar
21. Davey Smith G, Ebrahim S. "Mendelian randomisation": can genetic epidemiology contribute to understanding environmental determinants of disease? *International journal of epidemiology*. 2003;32:1–22. pmid:12689998  
View Article • PubMed/NCBI • Google Scholar
22. Lawlor DA, Timpson NJ, Harbord RM, Leary S, Ness A, McCarthy MI, et al. Exploring the developmental overnutrition hypothesis using parental-offspring associations and FTO as an IV. *PLoS Med*. 2008;5(3):484–93.  
View Article • PubMed/NCBI • Google Scholar
23. Burgess S, Thompson SG. Use of allele scores as instrumental variables for Mendelian randomization. *International journal of epidemiology*. 2013;42(4):1134–44. doi: 10.1093/ije/dyt093. pmid:24062299  
View Article • PubMed/NCBI • Google Scholar
24. Evans DM, Brion MJA, Paternoster L, Kemp JP, McMahon G, Munafò M, et al. Mining the Human Phenome Using Allelic Scores That Index Biological Intermediates. *PLoS Genet*. 2013;9(10).  
View Article • PubMed/NCBI • Google Scholar
25. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2013;42(1):111–27. Epub 2012/04/18. PubMed Central PMCID: PMC3600618. doi: 10.1093/ije/dys064. pmid:22507743  
View Article • PubMed/NCBI • Google Scholar
26. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al. Cohort Profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol*. 2013;42(1):97–110. Epub 2012/04/18. PubMed Central PMCID: PMC3600619. doi: 10.1093/ije/dys066. pmid:22507742  
View Article • PubMed/NCBI • Google Scholar
27. Jaddoe VW, van Duijn CM, Franco OH, van der Heijden AJ, van Ijzendoorn MH, de Jongste JC, et al. The Generation R Study: design and cohort update 2012. *European journal of epidemiology*. 2012;27(9):739–56. Epub 2012/10/23. doi: 10.1007/s10654-012-9735-1. pmid:23086283  
View Article • PubMed/NCBI • Google Scholar
28. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics*. 2007;81(3):559–75. Epub 2007/08/19. PubMed Central PMCID: PMC1950838. doi: 10.1086/519795. pmid:17701901  
View Article • PubMed/NCBI • Google Scholar
29. International HapMap Consortium, Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, et al. A second generation human haplotype map of over 3.1 million SNPs. *Nature*. 2007;449(7164):851–61. Epub 2007/10/19. PubMed Central PMCID: PMC2689609. doi: 10.1038/nature06258. pmid:17943122  
View Article • PubMed/NCBI • Google Scholar
30. Evans DM, Zhu G, Dy V, Heath AC, Madden PAF, Kemp JP, et al. Genome-wide association study identifies loci affecting blood copper, selenium and zinc. *Hum Mol Genet*. 2013;22(19):3998–4006. doi: 10.1093/hmg/ddt239. pmid:23720494  
View Article • PubMed/NCBI • Google Scholar

31. Li Y, Willer C, Sanna S, Abecasis G. Genotype Imputation. *Annu Rev Genom Hum G.* 2009;10:387–406.  
View Article • PubMed/NCBI • Google Scholar
32. Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR. MaCH: Using Sequence and Genotype Data to Estimate Haplotypes and Unobserved Genotypes. *Genet Epidemiol.* 2010;34(8):816–34. doi: 10.1002/gepi.20533. pmid:21058334  
View Article • PubMed/NCBI • Google Scholar
33. Paternoster L, Zhurov AI, Toma AM, Kemp JP, St Pourcain B, Timpson NJ, et al. Genome-wide association study of three-dimensional facial morphology identifies a variant in PAX3 associated with nasion position. *American journal of human genetics.* 2012;90(3):478–85. Epub 2012/02/22. PubMed Central PMCID: PMC3309180. doi: 10.1016/j.ajhg.2011.12.021. pmid:22341974  
View Article • PubMed/NCBI • Google Scholar
34. Kemp JP, Medina-Gomez C, Estrada K, St Pourcain B, Heppe DH, Warrington NM, et al. Phenotypic dissection of bone mineral density reveals skeletal site specificity and facilitates the identification of novel loci in the genetic regulation of bone mass attainment. *PLoS Genet.* 2014;10(6):e1004423. Epub 2014/06/20. PubMed Central PMCID: PMC4063697. doi: 10.1371/journal.pgen.1004423. pmid:24945404  
View Article • PubMed/NCBI • Google Scholar
35. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature genetics.* 2010;42(11):937–48. Epub 2010/10/12. PubMed Central PMCID: PMC3014648. doi: 10.1038/ng.686. pmid:20935630  
View Article • PubMed/NCBI • Google Scholar
36. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature.* 2015;518(7538):197–206. Epub 2015/02/13. PubMed Central PMCID: PMC4382211. doi: 10.1038/nature14177. pmid:25673413  
View Article • PubMed/NCBI • Google Scholar
37. Kruihof CJ, Kooijman MN, van Duijn CM, Franco OH, de Jongste JC, Klaver CC, et al. The Generation R Study: Biobank update 2015. *European journal of epidemiology.* 2014;29(12):911–27. Epub 2014/12/21. doi: 10.1007/s10654-014-9980-6. pmid:25527369  
View Article • PubMed/NCBI • Google Scholar
38. Medina-Gomez C, Felix JF, Estrada K, Peters MJ, Herrera L, Kruihof CJ, et al. Challenges in conducting genome-wide association studies in highly admixed multi-ethnic populations: the Generation R Study. *European journal of epidemiology.* 2015;30(4):317–30. Epub 2015/03/13. PubMed Central PMCID: PMC4385148. doi: 10.1007/s10654-015-9998-4. pmid:25762173  
View Article • PubMed/NCBI • Google Scholar
39. Jaddoe VW, van Duijn CM, van der Heijden AJ, Mackenbach JP, Moll HA, Steegers EA, et al. The Generation R Study: design and cohort update 2010. *European journal of epidemiology.* 2010;25(11):823–41. Epub 2010/10/23. PubMed Central PMCID: PMC2991548. doi: 10.1007/s10654-010-9516-7. pmid:20967563  
View Article • PubMed/NCBI • Google Scholar
40. 1000 Genomes Project Consortium, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature.* 2012;491(7422):56–65. Epub 2012/11/07. PubMed Central PMCID: PMC3498066. doi: 10.1038/nature11632. pmid:23128226  
View Article • PubMed/NCBI • Google Scholar
41. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature Genetics.* 2010;42(11):937–48. Epub 2010/10/12. PubMed Central PMCID: PMC3014648. doi: 10.1038/ng.686. pmid:20935630  
View Article • PubMed/NCBI • Google Scholar
42. Lin X, Song K, Lim N, Yuan X, Johnson T, Abderrahmani A, et al. Risk prediction of prevalent diabetes in a Swiss population using a weighted genetic score—the CoLaus Study. *Diabetologia.* 2009;52(4):600–8. doi: 10.1007/s00125-008-1254-y. pmid:19139842  
View Article • PubMed/NCBI • Google Scholar
43. Staiger D, Stock JH. Instrumental variables regression with weak instruments. *Econometrica.* 1997;65(3):557–86.  
View Article • PubMed/NCBI • Google Scholar
44. Davey Smith G, Lawlor DA, Harbord R, Timpson N, Day I, Ebrahim S. Clustered environments and randomized genes: a fundamental distinction between conventional and genetic epidemiology. *PLoS Med.* 2007;4(12):e352. Epub 2007/12/14. PubMed Central PMCID: PMC2121108. doi: 10.1371/journal.pmed.0040352. pmid:18076282  
View Article • PubMed/NCBI • Google Scholar
45. Didelez V, Meng S, Sheehan NA. Assumptions of IV Methods for Observational Epidemiology. *Stat Sci.* 2010;25(1):22–40.  
View Article • PubMed/NCBI • Google Scholar
46. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Brit Med J.* 2003;327(7414):557–60. doi: 10.1136/bmj.327.7414.557. pmid:12958120  
View Article • PubMed/NCBI • Google Scholar
47. Palmer TM, Lawlor DA, Harbord RM, Sheehan NA, Tobias JH, Timpson NJ, et al. Using multiple genetic variants as instrumental variables for modifiable risk factors. *Statistics in medicine.* 2012;21(3):223–242. doi: 10.1177/0962280210394459. pmid:21216802  
View Article • PubMed/NCBI • Google Scholar

48. Thomas DC, Lawlor DA, Thompson JR. Re: Estimation of bias in nongenetic observational studies using "Mendelian triangulation" by Bautista et al. *Annals of epidemiology*. 2007;17(7):511–3. doi: 10.1016/j.annepidem.2006.12.005. pmid:17466535  
View Article • PubMed/NCBI • Google Scholar
49. Ding EL, Hu FB. Determining origins and causes of childhood obesity via Mendelian randomization analysis. *PLoS Med*. 2008;5(3):e65. PubMed Central PMCID: PMC2265761. doi: 10.1371/journal.pmed.0050065. pmid:18336066  
View Article • PubMed/NCBI • Google Scholar
50. Zhang G, Bacelis J, Lengyel C, Teramo K, Hallman M, Helgeland O, et al. Assessing the Causal Relationship of Maternal Height on Birth Size and Gestational Age at Birth: A Mendelian Randomization Analysis. *PLoS Med*. 2015;12(8).  
View Article • PubMed/NCBI • Google Scholar
51. Del Greco MF, Minelli C, Sheehan NA, Thompson JR. Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Stat Med*. 2015;34(21):2926–40. doi: 10.1002/sim.6522. pmid:25950993  
View Article • PubMed/NCBI • Google Scholar
52. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *International journal of epidemiology*. 2015;44(2):512–25. doi: 10.1093/ije/dyv080. pmid:26050253  
View Article • PubMed/NCBI • Google Scholar
53. Palmer T. mrrobust package: Stata commands for MR-Egger, IVW, and weighted median estimators. <https://github.com/remlapmot/mrrobust2016>.
54. Davey Smith G, Steer C, Leary S, Ness A. Is there an intrauterine influence on obesity? Evidence from parent-child associations in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Arch Dis Child*. 2007;92(10):876–80. doi: 10.1136/adc.2006.104869. pmid:17595200  
View Article • PubMed/NCBI • Google Scholar
55. Patel R, Martin RM, Kramer MS, Oken E, Bogdanovich N, Matush L, et al. Familial associations of adiposity: findings from a cross-sectional study of 12,181 parental-offspring trios from Belarus. *PLoS ONE*. 2011;6(1):e14607. Epub 2011/02/08. PubMed Central PMCID: PMC3029263. doi: 10.1371/journal.pone.0014607. pmid:21298034  
View Article • PubMed/NCBI • Google Scholar
56. Subramanian SV, Ackerson LK, Davey Smith G. Parental BMI and childhood undernutrition in India: an assessment of intrauterine influence. *Pediatrics*. 2010;126(3):e663–71. Epub 2010/08/18. doi: 10.1542/peds.2010-0222. pmid:20713473  
View Article • PubMed/NCBI • Google Scholar
57. Fleten C, Nystad W, Stigum H, Skjaerven R, Lawlor DA, Davey Smith G, et al. Parent-offspring body mass index associations in the Norwegian Mother and Child Cohort Study: a family-based approach to studying the role of the intrauterine environment in childhood adiposity. *Am J Epidemiol*. 2012;176(2):83–92. Epub 2012/07/10. PubMed Central PMCID: PMC3493198. doi: 10.1093/aje/kws134. pmid:22771730  
View Article • PubMed/NCBI • Google Scholar
58. Vik KL, Romundstad P, Carslake D, Davey Smith G, Nilsen TI. Comparison of father-offspring and mother-offspring associations of cardiovascular risk factors: family linkage within the population-based HUNT Study, Norway. *International journal of epidemiology*. 2014;43(3):760–71. Epub 2013/12/25. doi: 10.1093/ije/dyt250. pmid:24366488  
View Article • PubMed/NCBI • Google Scholar
59. Corsi DJ, Subramanian SV, Ackerson LK, Davey Smith G. Is there a greater maternal than paternal influence on offspring adiposity in India? *Arch Dis Child*. 2015.  
View Article • PubMed/NCBI • Google Scholar
60. Lawlor DA, Lichtenstein P, Langstrom N. Association of maternal diabetes mellitus in pregnancy with offspring adiposity into early adulthood: sibling study in a prospective cohort of 280,866 men from 248,293 families. *Circulation*. 2011;123(3):258–65. Epub 2011/01/12. doi: 10.1161/CIRCULATIONAHA.110.980169. pmid:21220735  
View Article • PubMed/NCBI • Google Scholar
61. Freathy RM, Timpson NJ, Lawlor DA, Pouta A, Ben-Shlomo Y, Ruukonen A, et al. Common variation in the FTO gene alters diabetes-related metabolic traits to the extent expected given its effect on BMI. *Diabetes*. 2008;57(5):1419–26. doi: 10.2337/db07-1466. pmid:18346983  
View Article • PubMed/NCBI • Google Scholar
62. Wurtz P, Wang Q, Kangas AJ, Richmond RC, Skarp J, Tiainen M, et al. Metabolic Signatures of Adiposity in Young Adults: Mendelian Randomization Analysis and Effects of Weight Change. *PLoS Med*. 2014;11(12).  
View Article • PubMed/NCBI • Google Scholar
63. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet*. 2014;23(R1):R89–R98. Epub 2014/07/30. doi: 10.1093/hmg/ddu328. pmid:25064373  
View Article • PubMed/NCBI • Google Scholar
64. Baker JL, Michaelsen KF, Rasmussen KM, Sorensen TIA. Maternal prepregnant body mass index, duration of breastfeeding, and timing of complementary food introduction are associated with infant weight gain. *Am J Clin Nutr*. 2004;80(6):1579–88. pmid:15585772  
View Article • PubMed/NCBI • Google Scholar



65. Patro B, Liber A, Zalewski B, Poston L, Szajewska H, Koletzko B. Maternal and paternal body mass index and offspring obesity: a systematic review. *Annals of Nutrition & Metabolism*. 2013;63(1–2):32–41. Epub 2013/07/28.  
View Article • PubMed/NCBI • Google Scholar
66. Lawlor DA, Harbord RM, Sterne JAC, Timpson NJ, Davey Smith G. Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Statistic in Medicine*. 2008;27:1133–63.  
View Article • PubMed/NCBI • Google Scholar
67. Davey Smith G. Mendelian Randomization for Strengthening Causal Inference in Observational Studies: Application to Gene x Environment Interactions. *Perspect Psychol Sci*. 2010;5(5):527–45. doi: 10.1177/1745691610383505. pmid:26162196  
View Article • PubMed/NCBI • Google Scholar
68. Brion MJ, Zeegers M, Jaddoe V, Verhulst F, Tiemeier H, Lawlor DA, et al. Intrauterine effects of maternal prepregnancy overweight on child cognition and behavior in 2 cohorts. *Pediatrics*. 2011;127(1):e202–11. Epub 2010/12/29. PubMed Central PMCID: PMC3605781. doi: 10.1542/peds.2010-0651. pmid:21187310  
View Article • PubMed/NCBI • Google Scholar
69. Gaillard R, Durmus B, Hofman A, Mackenbach JP, Steegers EA, Jaddoe VW. Risk factors and outcomes of maternal obesity and excessive weight gain during pregnancy. *Obesity*. 2013;21(5):1046–55. Epub 2013/06/21. doi: 10.1002/oby.20088. pmid:23784909  
View Article • PubMed/NCBI • Google Scholar
70. Nohr EA, Timpson NJ, Andersen CS, Davey Smith G, Olsen J, Sorensen TI. Severe obesity in young women and reproductive health: the Danish National Birth Cohort. *PLoS ONE*. 2009;4(12):e8444. Epub 2009/12/31. PubMed Central PMCID: PMC2793537. doi: 10.1371/journal.pone.0008444. pmid:20041193  
View Article • PubMed/NCBI • Google Scholar
71. Sharp GC, Lawlor DA, Richmond RC, Fraser A, Simpkin A, Suderman M, et al. Maternal pre-pregnancy BMI and gestational weight gain, offspring DNA methylation and later offspring adiposity: findings from the Avon Longitudinal Study of Parents and Children. *International journal of epidemiology*. 2015;10.1093/ije/dyv042. Epub 2015/04/10.  
View Article • PubMed/NCBI • Google Scholar
72. Bhatti P, Sigurdson AJ, Wang SS, Chen J, Rothman N, Hartge P, et al. Genetic variation and willingness to participate in epidemiologic research: data from three studies. *Cancer Epidemiology, Biomarkers & Prevention*. 2005;14(10):2449–53.  
View Article • PubMed/NCBI • Google Scholar
73. Warrington NM, Howe LD, Paternoster L, Kaakinen M, Herrala S, Huikari V, et al. A genome-wide association study of body mass index across early life and childhood. *Int J Epidemiol*. 2015;44(2):700–12. Epub 2015/05/09. PubMed Central PMCID: PMC4469798. doi: 10.1093/ije/dyv077. pmid:25953783  
View Article • PubMed/NCBI • Google Scholar
74. Silverwood RJ, Holmes MV, Dale CE, Lawlor DA, Whittaker JC, Davey Smith G, et al. Testing for non-linear causal effects using a binary genotype in a Mendelian randomization study: application to alcohol and cardiovascular traits. *International journal of epidemiology*. 2014;43(6):1781–90. doi: 10.1093/ije/dyu187. pmid:25192829  
View Article • PubMed/NCBI • Google Scholar
75. Catalano PM, Farrell K, Thomas A, Huston-Presley L, Mencin P, de Mouzon SH, et al. Perinatal risk factors for childhood obesity and metabolic dysregulation. *Am J Clin Nutr*. 2009;90(5):1303–13. Epub 2009/09/18. PubMed Central PMCID: PMC2762159. doi: 10.3945/ajcn.2008.27416. pmid:19759171  
View Article • PubMed/NCBI • Google Scholar
76. Lawlor DA, Davey Smith G, O'Callaghan M, Alati R, Mamun AA, Williams GM, et al. Epidemiologic evidence for the fetal overnutrition hypothesis: findings from the mater-university study of pregnancy and its outcomes. *Am J Epidemiol*. 2007;165(4):418–24. Epub 2006/12/13. doi: 10.1093/aje/kwk030. pmid:17158475  
View Article • PubMed/NCBI • Google Scholar
77. Gaillard R, Steegers EA, Duijts L, Felix JF, Hofman A, Franco OH, et al. Childhood cardiometabolic outcomes of maternal obesity during pregnancy: the Generation R Study. *Hypertension*. 2014;63(4):683–91. Epub 2014/01/01. doi: 10.1161/HYPERTENSIONAHA.113.02671. pmid:24379180  
View Article • PubMed/NCBI • Google Scholar
78. Kral JG, Biron S, Simard S, Hould FS, Lebel S, Marceau S, et al. Large maternal weight loss from obesity surgery prevents transmission of obesity to children who were followed for 2 to 18 years. *Pediatrics*. 2006;118(6):e1644–9. Epub 2006/12/05. doi: 10.1542/peds.2006-1379. pmid:17142494  
View Article • PubMed/NCBI • Google Scholar
79. Smith J, Cianflone K, Biron S, Hould FS, Lebel S, Marceau S, et al. Effects of maternal surgical weight loss in mothers on intergenerational transmission of obesity. *Journal of Clinical Endocrinology and Metabolism*. 2009;94(11):4275–83. Epub 2009/10/13. doi: 10.1210/jc.2009-0709. pmid:19820018  
View Article • PubMed/NCBI • Google Scholar
80. Warrington NN, Evans DM, Lawlor DA, Consortium EGG. Does genetic variation contribute to the architecture of gestational weight gain?. *Journal of Developmental Origins of Health and Disease*. 2015;6(S2):S6.  
View Article • PubMed/NCBI • Google Scholar
81. Richardson SS, Daniels CR, Gillman MW, Golden J, Kukla R, Kuzawa C, et al. Don't blame the mothers. *Nature*. 2014;512(7513):131–2. doi: 10.1038/512131a. pmid:25119222

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

82. Warin M, Zivkovic T, Moore V, Davies M. Mothers as smoking guns: Fetal overnutrition and the reproduction of obesity. *Fem Psychol.* 2012;22(3):360–75.

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

83. Tedstone A, Targett V, Allen R, England PH. *Sugar Reduction: The evidence for action* London: Public Health England; 2015.