



Candler, T., Mahmoud, O., Edge, J., & Hamilton-Shield, J. (2017). Hypercholesterolaemia screening in Type 1 Diabetes: a difference of opinion. *Diabetic Medicine*, 34(7), 983-986.  
<https://doi.org/10.1111/dme.13322>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY-NC

Link to published version (if available):  
[10.1111/dme.13322](https://doi.org/10.1111/dme.13322)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at <http://onlinelibrary.wiley.com/doi/10.1111/dme.13322/abstract>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Table 1. Question's from Survey

Question 1	Do you measure cholesterol in your patients with Type 1 diabetes?	Yes=1 No=0
Question 1a	If yes, do you do measure?	1=Fasting sample routinely, 2 = non-fasting sample routinely 3= fasting if routine high
Question 1b	If yes, in which groups of children and young people?	1=all every year 2= all at diagnosis 3= age >12 every year 4=> 12 at diagnosis 5 = other
Question 1c	If yes, do you also measure/calculate	1= Triglyceride 2 = HDL 3= LDL
Question 2	Do you use any guidelines to decide treatment or further investigation?	1=yes 2= No
Question 2a	If yes, which guidelines do you use	Open response
Question 2b	If no, at what level of total or LDL cholesterol would you take further action?	Open response
Question 3	What would the further action be if above threshold for treatment?	1=refer to lipidologist 2= start statin 3= referral to dietician
Question 4	Have any children or young people in your clinic with T1DM been started on Statin therapy for high lipids that were not due to Familial Hypercholesterolemia?	Open response
Question 4a	If yes, how many in the	

	last 5 years?	
Question 4b	Excluding those with Familial Hypercholesterolemia, approximately how many patients have you identified in your clinic in last 5 years with a high LDL or total cholesterol that you felt required referral or treatment based on guidelines you use?	