



Pritchard, M. F., Powell, L. C., Khan, S., Griffiths, P. C., Mansour, O. T., Schweins, R., Beck, K., Buurma, N. J., Dempsey, C. E., Wright, C. J., Rye, P. D., Hill, K. E., Thomas, D. W., & Ferguson, E. L. (2017). The antimicrobial effects of the alginate oligomer OligoG CF-5/20 are independent of direct bacterial cell membrane disruption. *Scientific Reports*, 7, [44731]. <https://doi.org/10.1038/srep44731>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1038/srep44731](https://doi.org/10.1038/srep44731)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Nature at <https://www.nature.com/articles/srep44731>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>


The antimicrobial effects of the alginate oligomer OligoG CF-5/20 are independent of direct bacterial cell membrane disruption

Manon F. Pritchard*¹; Lydia C. Powell¹; Saira Khan¹; Peter C. Griffiths²; Omar T. Mansour²; Ralf Schweins³; Konrad Beck¹; Niklaas J. Buurma⁴; Christopher E. Dempsey⁵; Chris J. Wright⁶; Philip D. Rye⁷; Katja E. Hill¹; David W. Thomas^{†1}; Elaine L. Ferguson^{†1}


¹Advanced Therapies Group, Oral and Biomedical Sciences, School of Dentistry, College of Biomedical and Life Sciences, Cardiff University, Heath Park, Cardiff, UK. ²Department of Pharmaceutical, Chemical and Environmental Sciences, Faculty of Engineering and Science, University of Greenwich, Medway Campus, Central Avenue, Chatham Maritime, UK.

³Institut Laue-Langevin, DS/LSS group, 6 rue Jules Horowitz, 38042 Grenoble Cedex 9, France. ⁴Physical Organic Chemistry Centre, School of Chemistry, Cardiff University, Cardiff, UK. ⁵School of Biochemistry, Biomedical Sciences Building, University Walk, Clifton, BS8 1TD, UK. ⁶Centre for NanoHealth, Systems and Process Engineering Centre, College of Engineering, Swansea University, Swansea, UK. ⁷AlgiPharma AS, Sandvika, Norway.


*Corresponding author (email: PritchardMF@cardiff.ac.uk). †These authors jointly supervised this work.



Supplementary Fig. S1. Structural analysis of LPS. Small-angle neutron scattering from LPS (10 mg/ml) in D_2O containing (a) 0.001 M NaCl, (b) 0.01 M NaCl and (c) 0.1 M NaCl at pH 5-9.



Supplementary Fig. S2. Self-aggregation of OligoG CF-5/20. Enthalpogram for the dilution of 20 mg/ml OligoG CF-5/20 at 37°C in buffer (a) 20 mM phosphate pH 7, 100 mM NaCl, 1 mM EDTA, or (b) in 20 mM phosphate pH 7, 100 mM NaCl, 1 mM CaCl₂ (c) or in 20 mM phosphate pH 7, 100 mM NaCl, 1 mM EDTA, 2 mM CaCl₂.



Supplementary Fig. S3. Effect of OligoG CF-5/20 on *S. mutans* cell surface charge at pH 5 and pH 7 (2.6 mM NaCl \pm 60 mg/ml OligoG CF-5/20). Mean zeta potential measurements (mV) of *S. mutans*, 60 mg/ml OligoG CF-5/20 and both combined following hydrodynamic shear.