



Bedard, A., Northstone, K., Henderson, J., & Shaheen, SO. (2017). Maternal intake of sugar during pregnancy and childhood respiratory and atopic outcomes. *European Respiratory Journal*, 50(1), [1700073]. <https://doi.org/10.1183/13993003.00073-2017>

Peer reviewed version

Link to published version (if available):  
[10.1183/13993003.00073-2017](https://doi.org/10.1183/13993003.00073-2017)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via ERS Publications at <http://erj.ersjournals.com/content/50/1/1700073>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

## **Maternal intake of sugar during pregnancy and childhood respiratory and atopic outcomes**

Annabelle Bédard<sup>1</sup>, PhD, Kate Northstone<sup>2,3</sup>, PhD, A John Henderson<sup>3\*</sup>, MD, Seif O Shaheen<sup>1\*</sup>, PhD

<sup>1</sup>Centre for Primary Care and Public Health, Barts and The London School of Medicine and Dentistry, London, Queen Mary University of London, UK

<sup>2</sup>NIHR CLAHRC West, Bristol, UK

<sup>3</sup>School of Social and Community Medicine, University of Bristol, UK

\* Joint senior authors

### **Corresponding author:**

Dr Annabelle Bédard, Centre for Primary Care and Public Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, 58 Turner Street, London E1 2AB.

Email: [a.bedard@qmul.ac.uk](mailto:a.bedard@qmul.ac.uk)

Tel: +44 (0)20 7882 3863; Fax: +44 (0)20 7882 2252

**Take-home message:** A higher maternal intake of sugar in pregnancy may increase the risk of allergy and allergic asthma in the offspring.

**Word count:** 3,463

**ABSTRACT**

The possible role of maternal consumption of free sugar during pregnancy in the inception of respiratory and atopic diseases has not been studied. We aimed to study the relationship between maternal intake of free sugar during pregnancy and respiratory and atopic outcomes in the offspring in the population-based birth-cohort, the Avon Longitudinal Study of Parents and Children (ALSPAC). We analysed associations between maternal intake of free sugar in pregnancy (estimated by food frequency questionnaire), and current doctor-diagnosed asthma, wheezing, hayfever, eczema, atopy, serum total IgE, and lung function in children at 7-9 years (n=8,956 with information on maternal diet in pregnancy and at least one outcome of interest). After controlling for potential confounders, maternal intake of free sugar was positively associated with atopy (odds ratio for highest vs lowest quintile of sugar intake (95% CI) 1.38 (1.06-1.78), *P*-trend 0.006) and atopic asthma (OR 2.04 (1.24-3.35), *P*-trend 0.003). These associations were not confounded by intake of sugar in early childhood, which was unrelated to these outcomes. Our results suggest that a higher maternal intake of free sugar during pregnancy is associated with an increased risk of atopy and atopic asthma in the offspring, independently of sugar intake in early childhood.

**Word count:** 199 words

**Keywords:** asthma; atopy; early-life determinants; birth-cohort; prenatal; ALSPAC

## INTRODUCTION

There has been considerable interest in the role of maternal diet in pregnancy in the aetiology of childhood asthma and atopy(1). Studies have focused particularly on the potentially beneficial effects of antioxidants, following the hypothesis that a declining intake of antioxidants in Westernized countries has led to a reduction in pulmonary antioxidant defences, and hence to an increase in prevalence of asthma and atopy in recent decades(2). An alternative hypothesis, which has received less attention, is that the epidemic of asthma and atopy in the West could partly be explained by an increasing dietary intake of foods and constituents which may be harmful. Between 1970 and 2000, there has been a 25% increase in the per capita consumption of all refined sugars in the United States, matching a worldwide trend(3). Current international dietary guidelines advise people to reduce their consumption of sugar, and more particularly free sugars, which comprise sugars (monosaccharides and disaccharides) added to foods or drinks by the manufacturer, cook or consumer, and sugars naturally present in honey, syrups, and unsweetened fruit juices(4). Whilst in children a high consumption of sugar-sweetened beverages(5–7) and fruit juice(7, 8) has been linked to asthma, and particularly atopic asthma(7), the relation between total maternal consumption of free sugar during pregnancy and respiratory and atopic outcomes in the offspring has not been studied. One ecological study reported a correlation between perinatal consumption of sugar and severe childhood asthma symptoms(9), but could not specifically address maternal sugar intake in pregnancy. A recent Danish birth cohort study investigated the relation between soft drink consumption, but not total free sugar intake, during pregnancy and childhood asthma and allergic rhinitis(10).

We have investigated whether a high intake of free sugar in pregnancy is associated with adverse respiratory and atopic outcomes in the offspring in a large population-based UK birth cohort.

## **METHODS**

### **Participants**

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a population-based birth cohort that recruited 14,541 predominantly white pregnant women resident in Avon, UK with expected dates of delivery 1<sup>st</sup> April 1991 to 31<sup>st</sup> December 1992. These pregnancies resulted in 13,972 singleton or twin children who were alive at one year of age. The cohort has been followed since birth with annual questionnaires and, since age 7 years, with objective measures in annual research clinics. The study protocol has been described previously(11, 12) and further information can be found at: <http://www.alspac.bris.ac.uk>, which contains details of all the data that are available: <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>. Ethics approval was obtained from the ALSPAC Ethics and Law Committee (IRB 00003312) and the Local NHS Research Ethics Committees.

### **Exposure assessment**

Data on maternal diet in pregnancy were collected by a food frequency questionnaire (FFQ) at 32 weeks gestation, covering all the main foods consumed in Britain(13). The questionnaire asked about their current weekly frequency of consumption of 43 food groups and food items. More detailed questions were asked about daily consumption of a further eight basic foods (including sugar, coffee and tea). The FFQ was used to estimate total energy intake and daily nutrient intake, by multiplying the daily frequency of consumption of a food by the nutrient content(14) of a standard portion(15) of that food, and summing this for all the foods consumed. In this way free sugar intake was estimated. Free sugar does not include lactose when naturally present in milk and milk products, or the sugars contained within the cellular structure of fruits and vegetables). Information on child's free sugar

consumption at age 3 years, as well as maternal and partner's sugar consumption at 4 years post-partum, was collected by a similar FFQ.

Information from a questionnaire at recruitment and from the obstetric records was used to classify women into four mutually exclusive categories: no evidence of glycosuria or diabetes, existing diabetes mellitus before the pregnancy, gestational diabetes and persistent glycosuria during pregnancy. For the purposes of analysis we combined the last three categories to create a binary maternal 'diabetes' variable (see online supplement for further details).

## **Outcome assessment**

### *Binary variables*

Current doctor-diagnosed asthma was defined in children at 7.5 years (primary outcome) if mothers responded positively to the question 'Has a doctor *ever* actually *said* that your study child has asthma?' and to one or both of the questions 'Has your child had any of the following in the past 12 months: wheezing with whistling; asthma?.'

Current wheezing, eczema and hay fever in children at 7.5 years were defined by a positive answer to the question: 'Has your child had any of the following in the past 12 months: wheezing with whistling; eczema; hay fever?.'

Atopy at 7 years was defined as a positive reaction (maximum diameter of any detectable weal) to *D.pteronysinus*, cat or grass (after subtracting positive saline reactions from histamine and allergen weals, and excluding children unreactive to 1% histamine).

### *Categorical variables*

Children were further classified, post hoc, according to their asthmatic/atopic status, thus defining a 4-category variable (no atopy or asthma, atopy only, non-atopic asthma, and atopic

asthma) and according to their number of positive reactions to cat, grass and dust mite allergens (0, 1,  $\geq 2$ ).

Data on child's asthma status at 7 and at 14 years were used to derive asthma status phenotypes between 7-14 years (none, remitting, incident and persisting) (16).

### *Continuous variables*

Serum total IgE (kU/l) was measured by fluoroimmunoassay using the Pharmacia UNICAP system (Pharmacia and Upjohn Diagnostics AB, Uppsala, Sweden).

Lung function was measured by spirometry (Vitalograph 2120) at age 8½ years after withholding short-acting bronchodilators for at least 6 hours and long-acting bronchodilators and theophyllines for at least 24 hours. The best of three reproducible flow-volume curves was used to measure forced expiratory volume in 1 second (FEV<sub>1</sub>), forced vital capacity (FVC) and maximal mid-expiratory flow (FEF<sub>25-75</sub>), which were further transformed to age, height and gender adjusted standard deviation units(17). The tests adhered to American Thoracic Society (ATS) criteria for standardisation and reproducibility of flow-volume measurement(18), with the exception of ATS recommendations for duration of expiration(19); as many children did not fulfil forced expiratory time >6 seconds end of test criteria, a minimal volume change over the final 1 second was used.

### **Potential confounders**

We selected potential confounding factors which are known (from existing literature) to be associated with one or more of the outcomes of interest(20). These included maternal age at delivery, sex of child, multiple pregnancy, season of birth, maternal history of atopic diseases (hay fever, asthma, eczema, allergies, or attacks of wheezing with whistling on the chest or attacks of breathlessness in the past two years), parity, highest educational qualification,

housing tenure, financial difficulties, ethnicity, breastfeeding duration, and maternal factors during pregnancy (smoking status, anxiety score [Crown-Crisp Experiential Index], paracetamol use, antibiotic use, infections [urinary infection, influenza, rubella, thrush, genital herpes, other], supplement use and total energy intake [kJ/day]). Smoking status was categorised as the maximum exposure during pregnancy (never, passive smoking only, 1-9 cigarettes per day, 10-19 cigarettes per day,  $\geq 20$  cigarettes per day).

### **Statistical analyses**

We compared the distributions of child and maternal variables across maternal free sugar intake quintiles using F-statistics for differences in continuous variables and chi-squared tests for differences in categorical variables. Logistic regression, multinomial logistic regression, and linear regression were used to analyse relations between maternal free sugar intake in pregnancy and binary, categorical and continuous outcomes, respectively. After log transforming total IgE, linear regression was used to estimate geometric mean ratios (GMR) for IgE; confidence limits were calculated using Huber variances. We analysed free sugar intake in quintiles, first as a categorical variable using the lowest quintile as reference to allow for a non-linear pattern of association, and second as a continuous variable to test for linear trend (ie per quintile effect). For all regression analyses, two stages of adjustment were used. In Model 1 we adjusted for total energy intake only. In Model 2 we adjusted additionally for all potential confounders listed above.

When evidence for associations persisted, we considered other factors which can be considered either as potential confounders or potential mediators of associations between maternal free sugar intake in pregnancy and childhood outcomes, namely, prematurity(21, 22), impaired fetal growth(21, 23), maternal obesity and weight gain(24–26) and offspring obesity(27, 28). We therefore adjusted additionally for maternal pre-pregnancy body mass



index (BMI) (self-reported), gestational age at delivery, birth weight, maternal weight gain during pregnancy (all abstracted from obstetric records) and child's BMI at 7 (based on measured height and weight at clinic) (See Figure E1 in the online supplement showing a directed acyclic graph). In order to assess confounding arising from other dimensions of diet, we additionally adjusted separately for maternal intake of vitamin E, zinc, selenium, n-3 polyunsaturated fatty acids (PUFAs), n-6 PUFAs, and total fruit and total vegetables in pregnancy(1, 29–31).

To investigate confounding by postnatal sugar intake we adjusted additionally for child's sugar intake at age 3 years. In order to investigate potential unmeasured confounding by genetic or shared environmental or lifestyle factors, we used a parental comparison approach, whereby effect estimates for maternal sugar intake in pregnancy were compared with effect estimates for maternal and paternal sugar intake after pregnancy. If there is a causal intra-uterine effect, one would expect a stronger association with maternal intake in pregnancy than with maternal postnatal intake or paternal intake (the latter two exposures cannot have a direct biological effect on offspring asthma risk) (32, 33) (see further details online).

As sensitivity analyses, we repeated analyses after exclusion of mothers with implausible energy intakes ( $<2,500$  or  $>25,000$  kJ/day(34)), and after exclusion of mothers with diabetes (whose offspring will have experienced high fetal exposure to glucose). To correct for potential loss to follow-up bias, we used inverse probability weighting and assigned to each woman a weight that is the inverse of the probability of her selection for given values of covariates (see further details online) (35). All statistical analyses were carried out using Stata version 12.1 (StataCorp LP, USA).

## RESULTS

Of the 13,972 singleton or twin children alive at one year of age, information on maternal diet was available for 12,078, of whom there was information on at least one of the outcomes of interest for 8,956 (Figure E2). Characteristics of the 8,956 mother-child pairs who were included in the analyses, and those of the 3,122 mother-child pairs with information on maternal diet who were excluded because of incomplete outcome data, are compared in supplementary Table E1. Among children with available information, 12.2% had current doctor diagnosed asthma, 10.7% had current wheezing with whistling, 8.8% had current hayfever, 16.2% had current eczema, 21.5% had atopy and 61.8% did not have any of these five outcomes.

Maternal characteristics which differed across quintiles of free sugar intake during pregnancy included age, parity, pregnancy size, season of birth, breastfeeding duration, educational level, ethnicity, housing tenure, financial difficulties, anxiety level, tobacco exposure, and infection during pregnancy. Women in the highest quintile of total sugar intake during pregnancy had a lower pre-pregnancy BMI, higher total energy intake, and gained more weight during pregnancy than women in the lowest quintile. Their offspring were more likely to have weighed less at birth and to have had a lower BMI at 7 (Table 1). After adjustment for potential confounders, there was weak evidence for positive associations between maternal free sugar intake in pregnancy and childhood doctor-diagnosed asthma, and childhood wheeze (odds ratio comparing highest versus lowest quintile 1.31 (95% CI: 0.98-1.75), *P*-trend 0.09 and 1.42 (1.05-1.92), *P*-trend 0.08, respectively), and stronger evidence for a positive association with atopy at 7 years (OR 1.38 (1.06-1.78), *P*-trend 0.006) (Table 2). There was no association with eczema, hay fever, total IgE, FEV<sub>1</sub>, FVC or FEF<sub>25-75</sub> (Tables 2 and E2). *Post hoc* analysis showed a positive association between maternal intake

of free sugar and atopic asthma (OR 2.01 (1.23, 3.29), *P*-trend 0.004) (Table 3). The main positive findings of our study are summarized in Figure 1.

*Further investigation of potential confounding/mediation of main findings, and sensitivity analyses*

Additional separate adjustment for maternal pre-pregnancy BMI, gestational age at delivery, birth weight, maternal weight gain during pregnancy and child's BMI at 7 did not substantially alter the main findings and therefore no further formal mediation analysis was conducted (Table E3). Additional separate adjustment for maternal intake of vitamin E, zinc, selenium, n-3 PUFAs, n-6 PUFAs, and total fruits and vegetables in pregnancy did not substantially alter the main findings (data not shown), nor did additional separate adjustment for child's free sugar intake at age 3 (Table E3). The latter exposure was not associated with any outcome (data not shown).

In subsets of the cohort with complete data for paternal (respectively maternal) free sugar intake after pregnancy, no association was found between paternal (respectively maternal) free sugar intake after pregnancy and childhood atopy or atopic asthma. The significant associations of maternal free sugar intake during pregnancy with childhood atopy and atopic asthma remained, unattenuated, on mutual adjustment for paternal (respectively maternal) postnatal exposure (Table 4 and Table E4, respectively).

When we analysed the association between maternal free sugar intake and the number of positive reactions to cat, grass and dust mite allergens, we observed a stronger association for children with  $\geq 2$  positive reactions (Table 5). We studied associations between maternal intake of free sugar intake in pregnancy and childhood asthma status phenotypes and did not observe any association (Table E5).

The exclusion of 17 women with implausible energy intake estimates did not alter the main results, nor did exclusion of mothers with diabetes. Maternal 'diabetes' was not associated with any respiratory or atopic outcome, but was associated with higher birth weight. The inverse probability weighting analysis also produced similar results (data not shown).

## DISCUSSION

In this population-based birth cohort study, we have found that a higher maternal intake of free sugar during pregnancy was associated with an increased risk of atopy and atopic asthma in the offspring, independently of sugar intake in early childhood.

To our knowledge these are novel findings. Whilst a previous ecological study reported a positive correlation between perinatal consumption of sugar and severe childhood asthma symptoms(9), the limitations of ecological studies for inferring causality are well known (not least because of the high likelihood of confounding)(36). Furthermore, that study was unable to specifically investigate the potential role of maternal intake of sugar during pregnancy, nor the specific role of free sugar. Our longitudinal findings linking maternal free sugar intake in pregnancy to childhood atopy and atopic asthma extend the ecological results, and allow stronger causal inference. Interestingly, the findings for atopy became stronger when we examined the association with multiple sensitisation. Although previous cross-sectional studies have reported a positive association between childhood consumption of sugar-containing drinks, including fruit juice, and asthma(5–8), intake of free sugar in early childhood in our study was not associated with any respiratory or atopic outcome.

### *Mechanisms*

We speculate that high maternal fructose consumption may underlie the positive associations between maternal intake of free sugar and childhood atopy and atopic asthma. Fructose, which is a major component of added sugars, and is present naturally in fruit juice and in sweetened drinks as added sucrose (ratio of fructose/glucose: 50/50%) or isolated fructose, has been mooted as driving previous cross-sectional findings linking sugar-containing beverage consumption to asthma in children(7, 8). Fructose consumption, in the form of high

fructose corn syrup (ratio of fructose/glucose: 60/40%), has increased from near zero to near 30% of per capita consumption of refined sugars in the US between the 1970s and 2000, whereas the consumption of sucrose and glucose has declined or remained constant(3).

A prospective randomized controlled trial in adults showed that dietary sugar, and especially fructose, increased levels of C-reactive protein(37). Fructose also causes generation of uric acid(38), and experimental evidence in mice suggests that uric acid may be an essential initiator and amplifier of Th2 immunity and allergic inflammation, through activation of inflammatory dendritic cells(39). Alternatively, fructose might influence atopic immune responses by conditioning the gut microbiome(40, 41). The potential of maternal diet in pregnancy to influence inception of offspring allergic airways disease through this mechanism was recently confirmed in a mouse model(42). We therefore propose that one explanation for our main findings is that high fetal exposure to fructose may cause persistence of Th2 immune responses postnatally, and allergic inflammation in the developing lung.

In contrast to a previous study which reported a link between gestational diabetes and risk of atopic eczema and atopy in early childhood(43), we found no association between maternal diabetes during pregnancy and any outcome in the offspring although, as expected(44), maternal diabetes was associated with higher birth weight. The lack of a relation with maternal diabetes would suggest that higher fetal exposure to glucose is unlikely to explain our main findings. Whilst high fructose consumption has been proposed as a risk factor for obesity(45), we found no evidence to suggest that the associations between maternal free sugar intake and atopy and atopic asthma in the offspring were mediated by maternal BMI, gestational weight gain or child's BMI, nor by prematurity or low birth weight, assuming key assumptions necessary for mediation analyses(46, 47) were met.

### *Strengths and limitations*

Strengths of the ALSPAC birth cohort include its size and population-based prospective design, rich information on numerous potential lifestyle and dietary confounders (including information on childhood free sugar intake and parental sugar intake outside of pregnancy), and detailed phenotypic outcome measurements.

Although the FFQ that we used had not been formally calibrated against other instruments such as diet diaries, it was based on the one used by Yarnell et al which has been validated against weighed dietary records(48), and modified in the light of a more recent weighed dietary survey(13). The FFQ lacked quantitative information on soft drink consumption, and this will have led to underestimation of maternal free sugar intake during pregnancy. However, as misclassification of maternal free sugar intake in pregnancy is likely to have been random with respect to childhood outcomes, the strength of associations may have been underestimated. We were unable to assess associations with maternal sugar intake in early pregnancy; however, intakes in early and late pregnancy are likely to be highly correlated. We were unable to assess whether associations between maternal intake of free sugar in pregnancy and childhood atopy and atopic asthma persist beyond the age of 7, as no data on atopy (only data on asthma status) have been collected in ALSPAC children after the age of 7.

We think that confounding of the main findings by lifestyle or other aspects of maternal diet in pregnancy is unlikely, as we controlled for numerous potential confounders in the analyses, including nutrients and foods that have been previously linked to childhood asthma and atopy. Importantly, the main findings were not confounded by the offspring's free sugar intake in early childhood. Whilst the possibility of residual confounding cannot be ruled out, the null findings for maternal and paternal free sugar intakes after pregnancy make

confounding by unmeasured familial behaviours linked to sugar intake and asthma risk a less likely explanation.

As with any longitudinal study, we cannot rule out the possibility that exclusion of mother-child pairs without complete information might have biased our findings. However, it could be argued that, for our results to be totally spurious in those included in our analysis (and for the associations to be truly null in the population as a whole), associations in the excluded mother-child pairs would have to be in the opposite direction and much stronger, compared to the positive associations we reported in the included mother-child pairs, which seems extremely unlikely. Furthermore, loss to follow-up bias has been shown to only slightly modify associations in longitudinal studies, including in ALSPAC (49), and the results of our inverse probability weighting analysis(35) confirmed that loss to follow up is unlikely to have biased our results. In view of the multiple analyses carried out and the *post hoc* nature of the findings for atopic asthma, we cannot exclude the possibility that the main findings occurred by chance; hence they should be interpreted with caution. Given the *a priori* nature of the hypothesis being tested, and the fact that some outcomes of interest are highly correlated, it did not seem appropriate to correct for multiple testing. However, we plan to re-examine this hypothesis in another birth cohort to see if we can replicate the main findings.

### *Conclusions and public health implications*

We conclude that a higher maternal intake of free sugar during pregnancy may increase the risk of atopy and atopic asthma in the offspring. If these findings are replicated we would design an appropriate intervention study in pregnancy to establish or refute causality. Given the very high levels of sugar consumption currently in the West, where childhood allergy and



asthma are so prevalent, confirmation of a causal link would raise exciting prospects for the primary prevention of these disorders.

**Acknowledgements**

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The authors would like to thank especially Raquel Granell for preliminary analysis and data collection. This paper is the work of the authors, and John Henderson and Seif Shaheen will serve as guarantors for its contents. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

**Sources of support:**

The UK Medical Research Council, the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol currently provide core support for ALSPAC. AB is funded by a European Respiratory Society Long-Term Research Fellowship (Fellowship ID LTRF 2015 – 5838). KN is supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care West (CLAHRC West) at University Hospitals Bristol NHS Foundation Trust.

**Author's contributions:**

AB and SS conceived the study and drafted the manuscript. All authors were involved in the analysis strategy, KN gave advice on the dietary data, and AB performed the statistical analyses. AJH was responsible for all clinical respiratory and allergy data collection. All authors participated in the interpretation of the findings, reviewed the manuscript and revised it critically before submission. All authors have seen and approved the final version of the manuscript.

## REFERENCES

1. Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol* 2011;127:724-33-30.
2. Seaton A, Godden DJ, Brown K. Increase in asthma: a more toxic environment or a more susceptible population? *Thorax* 1994;49:171-174.
3. Cordain L, Eaton SB, Sebastian A, Mann N, Lindeberg S, Watkins BA, O'Keefe JH, Brand-Miller J. Origins and evolution of the Western diet: health implications for the 21st century. *Am J Clin Nutr* 2005;81:341-54.
4. Scientific Advisory Committee on Nutrition. The Scientific Advisory Committee on Nutrition's recommendations on sugars. *July 2015* at <[https://www.nutrition.org.uk/attachments/article/872/sugars\\_factsheet.pdf](https://www.nutrition.org.uk/attachments/article/872/sugars_factsheet.pdf)>.
5. Stensson M, Wendt L-K, Koch G, Nilsson M, Oldaeus G, Birkhed D. Oral health in pre-school children with asthma--followed from 3 to 6 years. *Int J Paediatr Dent* 2010;20:165-72.
6. Park S, Blanck HM, Sherry B, Jones SE, Pan L. Regular-Soda Intake Independent of Weight Status Is Associated with Asthma among US High School Students. *J Acad Nutr Diet* 2013;113:106-111.
7. Berentzen NE, van Stokkom VL, Gehring U, Koppelman GH, Schaap L, Smit H, Wijga A. Associations of sugar-containing beverages with asthma prevalence in 11-year-old children: the PIAMA birth cohort. *Eur J Clin Nutr* 2014;69:303-308.
8. DeChristopher LR, Uribarri J, Tucker KL. Intakes of apple juice, fruit drinks and soda are associated with prevalent asthma in US children aged 2-9 years. *Public Health Nutr* 2015;19:1-8.
9. Thornley S, Stewart A, Marshall R, Jackson R. Per capita sugar consumption is associated with severe childhood asthma: An ecological study of 53 countries. *Prim Care Respir J* 2011;20:75-78.
10. Maslova E, Strøm M, Olsen SF, Halldorsson TI. Consumption of Artificially-Sweetened Soft Drinks in Pregnancy and Risk of Child Asthma and Allergic Rhinitis. *PLoS One* 2013;8:e57261.
11. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, Molloy L, Ness A, Ring S, Smith GD. Cohort profile: The "Children of the 90s"-The index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013;42:111-127.
12. Fraser A, Macdonald-wallis C, Tilling K, Boyd A, Golding J, Davey smith G, Henderson J, Macleod J, Molloy L, Ness A, Ring S, Nelson SM, Lawlor DA. Cohort profile: The avon longitudinal study of parents and children: ALSPAC mothers cohort. *Int J Epidemiol* 2013;42:97-110.
13. Rogers I, Emmett P. Diet during pregnancy in a population of pregnant women in South West England. ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. *Eur J Clin Nutr* 1998;52:246-250.
14. McCance and Widdowson. *The Composition of Foods, 5th Edition*. Royal Society of Chemistry/Ministry of Agriculture Fisheries and Food; 1991.
15. Ministry of Agriculture and Food. *Food Portion Sizes*. London:HMSO: 1991.
16. Granell R, Henderson AJ, Sterne JA. Associations of wheezing phenotypes with late asthma outcomes in the Avon Longitudinal Study of Parents and Children: A population-based birth cohort. *J Allergy Clin Immunol* 2016;138:1060-1070.
17. Chinn S, Rona RJ. Height and age adjustment for cross sectional studies of lung function in children aged 6-11 years. *Thorax* 1992;47:707-714.

18. Standardization of Spirometry, 1994 Update. American Thoracic Society. *Am J Respir Crit Care Med* 1995;152:1107–1136.
19. Arets HGM, Brackel HJL, Van Der Ent CK. Forced expiratory manoeuvres in children: Do they meet ATS and ERS criteria for spirometry? *Eur Respir J* 2001;18:655–660.
20. Nurmatov U, Nwaru BI, Devereux G, Sheikh a. Confounding and effect modification in studies of diet and childhood asthma and allergies. *Allergy* 2012;67:1041–59.
21. Lenders CM, Hediger ML, Scholl TO, Khoo CS, Slap GB, Stallings VA. Gestational age and infant size at birth are associated with dietary sugar intake among pregnant adolescents. *J Nutr* 1997;127:1113–1117.
22. Jaakkola JJK, Ahmed P, Ieromnimon A, Goepfert P, Laiou E, Quansah R, Jaakkola MS. Preterm delivery and asthma: A systematic review and meta-analysis. *J Allergy Clin Immunol* 2006;118:823–830.
23. Brooks AM, Byrd RS, Weitzman M, Auinger P, McBride JT. Impact of low birth weight on early childhood asthma in the United States. *Arch Pediatr Adolesc Med* 2001;155:401–6.
24. Tielemans MJ, Erler NS, Leermakers ETM, van den Broek M, Jaddoe VW V, Steegers EAP, Kiefte-de Jong JC, Franco OH. A Priori and a Posteriori dietary patterns during pregnancy and gestational weight gain: The generation R study. *Nutrients* 2015;7:9383–9399.
25. Forno E, Young OM, Kumar R, Simhan H, Celedon JC. Maternal Obesity in Pregnancy, Gestational Weight Gain, and Risk of Childhood Asthma. *Pediatrics* 2014;134:e535–e546.
26. Bédard A, Dumas O, Kauffmann F, Garcia-Aymerich J, Siroux V, Varraso R. Potential confounders in the asthma-diet association: how causal approach could help? *Allergy* 2012;67:1461-2-3.
27. Phelan S, Hart C, Phipps M, Abrams B, Schaffner A, Adams A, Wing R. Maternal behaviors during pregnancy impact offspring obesity risk. *Exp Diabetes Res* 2011;2011:985139.
28. Litonjua A a, Gold DR. Asthma and obesity: common early-life influences in the inception of disease. *J Allergy Clin Immunol* 2008;121:1075-84–6.
29. Varraso R. Nutrition and asthma. *Curr Allergy Asthma Rep* 2012;12:201–10.
30. Allan K, Devereux G. Diet and asthma: nutrition implications from prevention to treatment. *J Am Diet Assoc* 2011;111:258–68.
31. Beckhaus AA, Garcia-Marcos L, Forno E, Pacheco-Gonzalez RM, Celedón JC, Castro-Rodriguez JA. Maternal nutrition during pregnancy and risk of asthma, wheeze, and atopic diseases during childhood: a systematic review and meta-analysis. *Allergy* 2015;70:1588–604.
32. Smith GD. Assessing intrauterine influences on offspring health outcomes: Can epidemiological studies yield robust findings? *Basic Clin Pharmacol Toxicol* 2008;102:245–256.
33. Shaheen SO, Newson RB, Smith GD, Henderson AJ. Prenatal paracetamol exposure and asthma: Further evidence against confounding. *Int J Epidemiol* 2010;39:790–794.
34. Maslova E, Rytter D, Bech BH, Henriksen TB, Rasmussen MA, Olsen SF, Halldorsson TI. Maternal protein intake during pregnancy and offspring overweight 20 y later. *Am J Clin Nutr* 2014;100:1139–1148.
35. Hernán MA, Hernandez-Diaz S, Robins JM. A structural approach to selection bias. *Epidemiology* 2004;15:615–625.
36. Piantadosi S, Byar DP, Green SB. The ecological fallacy. *Am J Epidemiol* 1988;127:893–904.

37. Aeberli I, Gerber PA, Hochuli M, Kohler S, Haile SR, Gouni-berthold I, Berthold HK, Spinas GA, Berneis K. Low to moderate sugar-sweetened beverage consumption impairs glucose and lipid metabolism and promotes inflammation in healthy young men: a randomized controlled trial. *Am Soc Nutr* 2011;94:479–485.
38. Johnson RJ, Nakagawa T, Sanchez-Lozada LG, Shafiu M, Sundaram S, Le M, Ishimoto T, Sautin YY, Lanaspa MA. Sugar, uric acid, and the etiology of diabetes and obesity. *Diabetes* 2013;62:3307–3315.
39. Kool M, Willart MAM, van Nimwegen M, Bergen I, Pouliot P, Virchow JC, Rogers N, Osorio F, Reis e Sousa C, Hammad H, Lambrecht BN. An Unexpected Role for Uric Acid as an Inducer of T Helper 2 Cell Immunity to Inhaled Antigens and Inflammatory Mediator of Allergic Asthma. *Immunity* 2011;34:527–540.
40. Payne AN, Chassard C, Lacroix C. Gut microbial adaptation to dietary consumption of fructose, artificial sweeteners and sugar alcohols: implications for host-microbe interactions contributing to obesity. *Obes Rev* 2012;13:799–809.
41. McLoughlin RM, Mills KHG. Influence of gastrointestinal commensal bacteria on the immune responses that mediate allergy and asthma. *J Allergy Clin Immunol* 2011;127:1097–107.
42. Thorburn AN, McKenzie CI, Shen S, Stanley D, Macia L, Mason LJ, Roberts LK, Wong CHY, Shim R, Robert R, Chevalier N, Tan JK, Mariño E, Moore RJ, Wong L, McConville MJ, Tull DL, Wood LG, Murphy VE, Mattes J, Gibson PG, Mackay CR. Evidence that asthma is a developmental origin disease influenced by maternal diet and bacterial metabolites. *Nat Commun* 2015;6:7320.
43. Kumar R, Ouyang F, Story RE, Pongratic JA, Hong X, Wang G, Pearson C, Ortiz K, Bauchner H, Wang X. Gestational diabetes, atopic dermatitis, and allergen sensitization in early childhood. *J Allergy Clin Immunol* 2009;124:1031–1038.
44. Silverman BL, Rizzo T, Green OC, Cho NH, Winter RJ, Ogata ES, Richards GE, Metzger BE. Long-term prospective evaluation of offspring of diabetic mothers. *Diabetes* 1991;40 Suppl 2:121–5.
45. Lakhan SE, Kirchgessner A. The emerging role of dietary fructose in obesity and cognitive decline. *Nutr J* 2013;12:114.
46. Robins JM, Greenland S. Identifiability and exchangeability for direct and indirect effects. *Epidemiology* 1992;3:143–55.
47. Cole SR, Hernán MA. Fallibility in estimating direct effects. *Int J Epidemiol* 2002;31:163–165.
48. Yarnell JW, Fehily AM, Milbank JE, Sweetnam PM, Walker CL. A short dietary questionnaire for use in an epidemiological survey: comparison with weighed dietary records. *Hum Nutr Appl Nutr* 1983;37:103–12.
49. Howe LD, Tilling K, Galobardes B, Lawlor DA. Loss to Follow-up in Cohort Studies. *Epidemiology* 2013;24:1–9.

**Figure legends**

**Figure 1.** Summary of the main findings for the associations between maternal free sugar intake in pregnancy and childhood outcomes (atopy and atopic asthma)

\* Controlling for energy intake, smoking, infections, supplements, antibiotics and paracetamol use during pregnancy; maternal educational level, housing tenure, financial difficulties, ethnicity, age, parity, history of atopic diseases, anxiety; sex of child, season of birth, multiple pregnancy, breastfeeding duration

**Table 1.** Characteristics of mothers and offspring who had information on at least one of the outcomes of interest (wheeze, asthma, atopy, eczema, hayfever, total IgE, lung function) by maternal free sugar intake in pregnancy (n=8,956)

Mother and offspring characteristics	Free sugar intake (g/day)					P
	Quintile 1 [1.6-34.0]	Quintile 2 [34.0-46.6]	Quintile 3 [46.6-60.8]	Quintile 4 [60.8-82.4]	Quintile 5 [82.4-345.1]	
<b>Mother's age (years), m (sd)</b>	29.4 (4.7)	29.1 (4.5)	29.2 (4.6)	28.9 (4.5)	27.9 (4.7)	< .001
<b>Parity, %</b>						
0	43.2	46.1	48.6	44.9	44.0	0.001
1	35.4	37.2	35.1	36.3	35.9	
≥2	21.4	16.7	16.3	18.8	20.0	
<b>Sex of child, %</b>						
Male	51.2	52.3	50.5	52.3	49.5	0.39
Female	48.8	47.7	49.5	47.7	50.5	
<b>Multiple pregnancy, %</b>						
Singleton	97.3	96.7	98.0	98.3	97.3	0.01
Twin	2.7	3.3	2.0	1.7	2.7	
<b>Season of birth, %</b>						
Winter	15.0	17.6	16.6	15.7	16.0	< .001
Spring	23.6	25.3	26.8	29.1	30.7	
Summer	33.1	29.3	31.8	28.9	27.1	
Autumn	28.3	27.8	24.9	26.4	26.2	
<b>Breastfeeding duration, %</b>						
Never	21.3	18.9	17.5	22.1	27.1	< .001
<3 months	34.0	30.0	30.1	29.2	34.2	
3-6 months	13.1	14.9	14.6	13.3	12.8	
≥6 months	31.5	36.3	37.4	35.4	25.9	
<b>Mother's educational level, %</b>						
Certificate of Secondary Education	16.7	12.9	13.4	14.5	20.4	< .001
Vocational	8.6	9.2	8.1	8.5	10.8	

Ordinary level	36.2	35.0	32.8	35.1	38.8	
Advanced level	24.6	26.6	26.3	26.0	21.2	
Degree	14.0	16.2	19.4	15.9	8.9	
<b>Maternal ethnicity, %</b>						
White	97.8	97.2	98.5	98.4	98.8	0.003
Non-white	2.2	2.8	1.5	1.6	1.2	
<b>Housing tenure, %</b>						
Owned/mortgaged	82.6	86.5	86.5	85.1	76.8	< .001
Council rented	9.9	7.2	6.6	8.8	15.8	
Non-council rented	7.5	6.3	7.0	6.1	7.5	
<b>Financial difficulties, %</b>						
No	82.6	82.8	84.3	85.3	78.8	< .001
Yes	17.4	17.2	15.7	14.7	21.2	
<b>Maternal history of atopic diseases, %</b>						
No	32.2	32.0	33.7	31.4	28.9	0.05
Yes	67.8	68.0	66.3	68.6	71.1	
<b>Maternal anxiety score in pregnancy, %</b>						
0-9	24.0	23.4	23.1	20.8	14.3	< .001
10-14	25.1	27.5	25.7	26.1	24.3	
15-20	24.4	25.1	27.3	27.1	25.4	
≥20	26.5	24.1	24.0	26.1	36.1	
<b>Maximum maternal tobacco exposure, %</b>						
None	27.8	27.6	28.9	28.2	19.2	< .001
Passive only	45.4	48.5	47.3	46.0	42.3	
1-9 cig/day	7.6	7.7	8.1	8.0	8.1	
10-19 cig/day	12.5	10.1	9.4	10.1	15.2	
20+ cig/day	6.7	6.2	6.2	7.7	15.2	
<b>Maternal paracetamol use during pregnancy, %</b>						
No	38.0	38.0	37.7	38.0	36.4	0.86
Yes	62.0	62.0	62.3	62.0	63.6	
<b>Maternal antibiotic use during pregnancy, %</b>						
No	83.7	85.4	84.0	83.7	82.9	0.34



Yes	16.3	14.6	16.0	16.3	17.1	
<b>Maternal supplement use during pregnancy, %</b>						
No	44.5	43.1	44.4	41.9	41.6	0.26
Yes	55.5	56.9	55.6	58.1	58.4	
<b>Maternal infections in pregnancy, %</b>						
No	57.2	55.1	55.8	53.3	49.3	< .001
Yes	42.8	44.9	44.2	46.7	50.7	
<b>Total energy intake (kJ/day), m (sd)</b>	5567 (1376)	6499 (1326)	7166 (1360)	7931 (1488)	9317 (1994)	< .001
<b>Maternal pre-pregnancy BMI, %</b>						
<18.50 kg/m <sup>2</sup>	2.7	2.8	4.4	4.0	8.1	< .001
18.50-24.99 kg/m <sup>2</sup>	70.8	74.3	77.7	76.7	77.4	
25.00-29.99 kg/m <sup>2</sup>	17.7	17.1	13.5	14.8	11.6	
≥30.00 kg/m <sup>2</sup>	8.8	5.8	4.4	4.5	2.9	
<b>Birth weight, %</b>						
<2500 g	4.2	4.0	3.6	4.3	5.6	0.02
2500-2999 g	14.2	13.2	13.9	12.8	15.2	
3000-3499 g	34.8	34.9	35.1	36.7	35.6	
3500-3999 g	32.2	33.6	33.5	33.6	33.0	
≥4000 g	14.7	14.4	13.8	12.6	11.0	
<b>Gestational age (weeks), m (sd)</b>	39.4 (1.8)	39.5 (1.7)	39.5 (1.7)	39.5 (1.8)	39.4 (1.9)	0.28
<b>Child's BMI at 7, %</b>						
<15.00 kg/m <sup>2</sup>	25.8	26.3	30.3	27.7	30.7	
15.00-17.49 kg/m <sup>2</sup>	51.9	52.3	51.2	54.4	52.6	< .001
17.50-20.49 kg/m <sup>2</sup>	17.2	16.6	14.2	14.3	13.6	
≥20.50 kg/m <sup>2</sup>	5.2	4.8	4.3	3.5	3.1	
<b>Maternal weight gain during pregnancy, %</b>						
Quartile 1	30.3	23.7	24.5	23.5	24.5	< .001

---

Quartile 2	24.3	25.0	25.1	24.7	25.0
Quartile 3	24.2	27.0	26.3	26.4	23.4
Quartile 4	21.3	24.3	24.1	25.4	27.1

---

**Table 2.** Associations between maternal free sugar intake during pregnancy and asthma, wheeze, eczema, hay fever and atopy in the offspring

	Maternal free sugar intake during pregnancy					Per quintile	<i>P</i> -trend
	1 <sup>st</sup> quintile	2 <sup>nd</sup> quintile	3 <sup>rd</sup> quintile	4 <sup>th</sup> quintile	5 <sup>th</sup> quintile		
<b>Asthma</b> (n=7,677)							
OR <sup>a</sup> (95% CI)	1.00	1.14 (0.91-1.41)	0.97 (0.77-1.23)	1.22 (0.96-1.56)	1.37 (1.04-1.81)	1.07 (1.00-1.14)	0.04
OR <sup>b</sup> (95% CI)	1.00	1.18 (0.94-1.47)	1.02 (0.80-1.29)	1.24 (0.97-1.58)	1.31 (0.98-1.75)	1.06 (0.99-1.13)	0.09
<b>Wheeze</b> (n=7,762)							
OR <sup>a</sup> (95% CI)	1.00	1.21 (0.96-1.52)	1.05 (0.82-1.35)	1.19 (0.92-1.54)	1.38 (1.03-1.85)	1.06 (0.99-1.13)	0.09
OR <sup>b</sup> (95% CI)	1.00	1.25 (0.99-1.58)	1.11 (0.86-1.42)	1.22 (0.94-1.59)	1.42 (1.05-1.92)	1.06 (0.99-1.14)	0.08
<b>Eczema</b> (n=7,748)							
OR <sup>a</sup> (95% CI)	1.00	1.19 (0.98, 1.44)	1.13 (0.93, 1.38)	1.05 (0.84, 1.30)	0.91 (0.70, 1.17)	0.97 (0.92, 1.03)	0.33
OR <sup>b</sup> (95% CI)	1.00	1.17 (0.96, 1.42)	1.15 (0.94, 1.41)	1.07 (0.86, 1.33)	0.97 (0.74, 1.26)	0.99 (0.93, 1.05)	0.70
<b>Hay fever</b> (n=7,728)							
OR <sup>a</sup> (95% CI)	1.00	1.07 (0.82, 1.38)	1.25 (0.96, 1.63)	1.23 (0.93, 1.63)	1.22 (0.88, 1.69)	1.06 (0.98, 1.14)	0.13
OR <sup>b</sup> (95% CI)	1.00	1.03 (0.79, 1.33)	1.24 (0.95, 1.62)	1.21 (0.91, 1.60)	1.25 (0.89, 1.75)	1.07 (0.99, 1.15)	0.10
<b>Atopy</b> (n=6,117)							
OR <sup>a</sup> (95% CI)	1.00	0.99 (0.81-1.21)	1.09 (0.89-1.34)	1.19 (0.96-1.47)	1.24 (0.97-1.60)	1.06 (1.01-1.13)	0.03
OR <sup>b</sup> (95% CI)	1.00	0.98 (0.80-1.20)	1.10 (0.89-1.35)	1.20 (0.96-1.49)	1.38 (1.06-1.78)	1.09 (1.02-1.15)	0.006

OR: Odds ratio

<sup>a</sup> Controlling for energy intake<sup>b</sup> Controlling for energy intake, smoking, infections, supplements, antibiotics and paracetamol use during pregnancy; maternal educational level, housing tenure, financial difficulties, ethnicity, age, parity, history of atopic diseases, anxiety; sex of child, season of birth, multiple pregnancy, breastfeeding duration

**Table 3.** Associations between maternal free sugar intake during pregnancy and atopy without asthma, non-atopic and atopic asthma\* in the offspring

	Maternal free sugar intake during pregnancy					Per quintile	P-trend	P-trend
	1 <sup>st</sup> quintile	2 <sup>nd</sup> quintile	3 <sup>rd</sup> quintile	4 <sup>th</sup> quintile	5 <sup>th</sup> quintile			
<b>Atopy without asthma</b> (n=3063)								
Maternal free sugar intake in pregnancy	1.00	0.78 (0.61, 1.01)	1.05 (0.82, 1.34)	0.96 (0.73, 1.25)	1.09 (0.80, 1.50)	1.04 (0.97, 1.12)		0.30
OR <sup>a</sup> (95% CI)	1.00	0.78 (0.61, 1.01)	1.05 (0.82, 1.34)	0.96 (0.73, 1.25)	1.09 (0.80, 1.50)	1.04 (0.97, 1.12)		0.30
OR <sup>b</sup> (95% CI)	1.00	1.04 (0.78, 1.38)	1.16 (0.87, 1.55)	1.33 (0.98, 1.82)	1.64 (1.14, 2.37)	1.13 (1.04, 1.23)	0.004	0.19
<b>Non-atopic asthma</b> (n=301)	1.00	1.02 (0.77-1.36)	1.15 (0.86-1.53)	1.31 (0.96-1.79)	1.61 (1.11-2.33)	1.13 (1.04-1.22)	0.005	
OR <sup>a</sup> (95% CI)	1.00	0.82 (0.57, 1.16)	0.67 (0.46, 0.99)	0.77 (0.52, 1.16)	0.83 (0.52, 1.34)	0.95 (0.85, 1.06)		0.34
OR <sup>b</sup> (95% CI)	1.00	0.87 (0.61, 1.25)	0.73 (0.49, 1.08)	0.78 (0.52, 1.18)	0.71 (0.43, 1.15)	0.92 (0.82, 1.03)		0.14
<b>Atopic asthma</b> (n=337)								
OR <sup>a</sup> (95% CI)	1.00	1.66 (1.14, 2.41)	1.17 (0.78, 1.77)	2.09 (1.39, 3.14)	1.79 (1.11, 2.90)	1.14 (1.03, 1.27)		0.01
OR <sup>b</sup> (95% CI)	1.00	1.75 (1.20, 2.56)	1.27 (0.84, 1.93)	2.18 (1.45, 3.30)	2.01 (1.23, 3.29)	1.17 (1.05, 1.30)		0.004

OR: Odds ratio

\* No atopy or asthma was considered as baseline category (n=3,796)

<sup>a</sup> Controlling for energy intake<sup>b</sup> Controlling for energy intake, smoking, infections, supplements, antibiotics and paracetamol use during pregnancy; maternal educational level, housing tenure, financial difficulties, ethnicity, age, parity, history of atopic diseases, anxiety; sex of child, season of birth, multiple pregnancy, breastfeeding duration**Table 4.** Comparison of associations of childhood atopy and atopic asthma with maternal free sugar intake during pregnancy vs paternal

Paternal free sugar intake after pregnancy								
OR <sup>a</sup> (95% CI)	1.00	1.12 (0.84-1.50)	1.28 (0.95-1.73)	1.36 (0.99-1.86)	1.16 (0.81-1.68)	1.05 (0.97-1.14)	0.22	
OR <sup>b</sup> (95% CI)	1.00	1.10 (0.82-1.47)	1.25 (0.93-1.69)	1.30 (0.95-1.78)	1.10 (0.76-1.59)	1.04 (0.96-1.13)	0.34	
<b>Atopic asthma</b> (n=2,830)								
Maternal free sugar intake in pregnancy								
OR <sup>a</sup> (95% CI)	1.00	2.11 (1.26-3.52)	1.18 (0.66-2.11)	2.82 (1.60-4.96)	2.01 (1.01-4.00)	1.17 (1.02-1.36)	0.03	
OR <sup>b</sup> (95% CI)	1.00	2.13 (1.27-3.57)	1.17 (0.65-2.11)	2.80 (1.58-4.94)	1.96 (0.98-3.93)	1.17 (1.01-1.35)	0.04	
Paternal free sugar intake after pregnancy								
OR <sup>a</sup> (95% CI)	1.00	0.99 (0.60-1.62)	1.26 (0.75-2.11)	1.47 (0.87-2.48)	0.92 (0.48-1.77)	1.04 (0.90-1.21)	0.55	
OR <sup>b</sup> (95% CI)	1.00	0.96 (0.59-1.59)	1.23 (0.73-2.06)	1.39 (0.82-2.35)	0.87 (0.45-1.69)	1.03 (0.89-1.19)	0.72	

intake after pregnancy

OR: Odds ratio

<sup>a</sup> Controlling only for previously mentioned potential confounders

<sup>b</sup> Mutually adjusting for maternal free sugar intake in pregnancy and paternal free sugar intake after pregnancy, in addition to previously mentioned potential confounders

**Table 5.** Associations between maternal free sugar intake in pregnancy and number of positive skin-prick tests\*

	<b>1 positive SPT</b> (n=783)		<b>≥2 positive SPT</b> (n=531)	
	OR <sup>a</sup> (95% CI)	<i>P</i> trend	OR <sup>a</sup> (95% CI)	<i>P</i> trend
<b>Free sugar</b>				
1 <sup>st</sup> quintile	1.00		1.00	
2 <sup>nd</sup> quintile	0.87 (0.68, 1.12)		1.17 (0.87, 1.57)	
3 <sup>rd</sup> quintile	1.07 (0.83, 1.37)		1.14 (0.84, 1.56)	
4 <sup>th</sup> quintile	1.09 (0.83, 1.43)		1.37 (0.99, 1.90)	
5 <sup>th</sup> quintile	1.19 (0.86, 1.63)		1.73 (1.18, 2.52)	
Per quintile	1.06 (0.98, 1.14)	0.12	1.13 (1.03, 1.23)	0.006

OR: Odds ratio

\*No positive skin prick-test was considered as baseline category (n=4,797)

<sup>a</sup>Controlling for energy intake, smoking, infections, supplements, antibiotics and paracetamol use during pregnancy; maternal educational level, housing tenure, financial difficulties, ethnicity, age, parity, history of atopic diseases, anxiety; sex of child, season of birth, multiple pregnancy, breastfeeding duration