



Carswell, A., Hill, P. W., Jones, D. L., Blackwell, M. S. A., Johnes, P., Dixon, E., & Chadwick, D. (2018). Impact of microbial activity on the leaching of soluble N forms in soil. *Biological Fertility of Soils*, 54(1), 21-25. <https://doi.org/10.1007/s00374-017-1250-9>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1007/s00374-017-1250-9](https://doi.org/10.1007/s00374-017-1250-9)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Springer at <https://link.springer.com/article/10.1007%2Fs00374-017-1250-9> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Impact of microbial activity on the leaching of soluble N forms in soil: Online resource

A.M. Carswell^{a*}, P.W. Hill^b, D.L. Jones^b, M.S.A. Blackwell^a, P.J. Johnes^c, E.R. Dixon^a, D.R. Chadwick^b

^a *Rothamsted Research, Sustainable Agricultural Systems, Okehampton EX20 2SB, Devon, UK*

^b *School of Environment, Natural Resources & Geography, Bangor University, Gwynedd, LL57 2UW, UK*

^c *School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8 1SS, UK*

Corresponding author: Tel +44 (0) 1837 883 500

e-mail address: alison.carswell@rothamsted.ac.uk (A.M. Carswell)

Table S1. Initial soil properties

Parameter	
Soil type (FAO classification*)	Dystric Cambisol*
Soil texture	Loam
Soil pH in water	5.24 ± 0.05
Soil total C (g kg ⁻¹ DM)	18.6 ± 0.02
Soil total N (g kg ⁻¹ DM)	3.04 ± 0.02
Soil ¹³ C (Atom% of total C)	1.08 ± 0.00
Soil ¹⁵ N (Atom% of total N)	0.37 ± 0.00
Soil respiration (µg C g ⁻¹ DM h ⁻¹)	0.50 ± 0.07
Soil water (g g ⁻¹)	0.20 ± 0.01
Soil solution free amino acids (µM N)	13.6 ± 1.63
Soil solution urea (µM N)	6.22 ± 0.20
Soil solution NO ₃ -N (mM N)	0.20 ± 0.15
Soil solution NH ₄ -N (µM N)	13.1 ± 1.31
Soil solution DOC (mM C)	2.63 ± 0.31
Soil solution DON (mM N)	0.14 ± 0.02

*as classified by Harrod and Hogan (2008), values are mean ± SEM (*n* = 3)

Laboratory analyses

Soil ¹⁵N and ¹³C, and total N and total C were determined on subsamples of the <2 mm fraction using a Carlo Erba NA 2000 linked to a Sercon 20/22 isotope ratio mass spectrometer (Sercon, Crewe, UK; Carlo Erba, CE Instruments, Wigan, UK). Soil respiration and gravimetric soil water content were measured on field-moist <2 mm sieved soil. Soil respiration was determined using an SR-1 automated multi-channel soil respirometer (PP Systems Ltd, Hitchin, UK). Soil solution N analyses included NO₃-N (Mulvaney 1996), NH₄-N (Miranda et al. 2001), free amino acids (FAA; Jones et al. 2002), dissolved organic carbon (DOC; as non-purgeable organic C) and total N (Multi N/C analyser, Analytic Jena Multi N/C 2100s, Jena, Germany). Dissolved organic N was determined via subtraction of the mineral N values from the total dissolved N values. Soil sterility after autoclaving was tested in triplicate by inoculating sterilised soils onto 90 mm petri dishes containing potato dextrose agar (PDA; Sigma) or Luria Bertani agar (LB; Lab M). Triplicate plates containing living soil were set up as a control. The plates were sealed and incubated at 21°C under a 16 h fluorescent light regime for 60

h, before examining for culturable organisms. No living organisms were observed in the sterile soil cultures. Nonetheless this method does not account for the presence of viable but non-culturable organisms (Kell et al. 1998).

References

Harrod TR, Hogan DV (2008) The soils of North Wyke and Rowden: Revised edition of Harrod TR (1981) Soils in Devon V: Sheet SS61 (Chumleigh) Soil Survey Record no. 70 ed.

Jones DL, Owen AG, Farrar JF (2002) Simple method to enable the high resolution determination of total free amino acids in soil solutions and soil extracts. *Soil Biol Biochem* 34:1893-1902.

Kell DB, Kaprelyants AS, Weichart DH, Harwood CR, Barer MR (1998) Viability and activity in readily culturable bacteria: a review and discussion of the practical issues. *Antonie Van Leeuwenhoek* 73:169-187.

Miranda KM, Espey MG, Wink DA (2001) A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5:62-71.

Mulvaney RL (1996) Nitrogen - inorganic forms. In: Sparks DL, Page AL, Helmke PA, Loeppert RH (Eds) *Methods of Soil Analysis, part 3: Chemical Methods*. Madison, WI, pp 1123-1184