



Matsubara, V. H., Wang, Y., Bandara, H. M. H. N., Mayer, M. P. A., & Samaranyake, L. P. (2016). Probiotic lactobacilli inhibit early stages of *Candida albicans* biofilm development by reducing their growth, cell adhesion, and filamentation. *Applied Microbiology and Biotechnology*, 100(14), 6415-6426. <https://doi.org/10.1007/s00253-016-7527-3>

Peer reviewed version

Link to published version (if available):  
[10.1007/s00253-016-7527-3](https://doi.org/10.1007/s00253-016-7527-3)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Probiotic strain	Biofilm viable cells reduction (%)							
	<i>C. albicans</i> ATCC SC5314				<i>C. albicans</i> 75			
	Initial Colonization		Maturation		Initial Colonization		Maturation	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>L. rhamnosus</i> LR32	32.0	6.4	61.8*	8.3	31.8	12.2	39.8	6.2
<i>L. casei</i> L324m	59.0	6.9	54.7	4.6	27.7	5.0	36.6	3.3
<i>L. acidophilus</i> NCFM	56.3	4.2	34.2*	7.0	27.4	5.5	25.5	9.1

\* Significant difference between initial colonization and maturation (p<0.05).

Table 1. Percentage of reduction of viable *C. albicans* cells on biofilms after treatment with probiotic bacteria suspension ( $1 \times 10^7$  probiotic cells/ml) during initial colonization (24 h) and maturation (48 h) phases.