



Jefferys, A., Lenguerrand, E., Cornthwaite, K., Johnson, A., Lynch, M., Hyde, J., Draycott, T., & Siassakos, D. (2017). Deflation versus maintained inflation of gastric band in pregnancy: A national cohort study. *Obstetric Medicine*, 10(2), 67-73.
<https://doi.org/10.1177/1753495X16681200>

Peer reviewed version

Link to published version (if available):
[10.1177/1753495X16681200](https://doi.org/10.1177/1753495X16681200)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Sage at <http://journals.sagepub.com/doi/10.1177/1753495X16681200> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/user-guides/explore-bristol-research/ebr-terms/>

Table 1. Demographic characteristics of women at booking

| | Total (n=107) | Deflated* (n=42) | Inflated† (n=54) | Unknown (n=11) |
|----------------------------------|-------------------------|----------------------------|----------------------------|--------------------------|
| Age | | | | |
| Mean (y) +/- SD | 31.8 +/- 4.9 | 32.5 +/- 4.5 | 31.3 +/- 4.8 | 31.2 +/- 6.5 |
| ≥35y, n= (%) | 33 (30.8) | 14 (33.3) | 15 (27.8) | 4 (36.4) |
| ≥40y, n= (%) | 8 (7.5) | 3 (7.1) | 4 (7.4) | 1 (9.1) |
| Missing, n= | 0 | 0 | 0 | 0 |
| Parity | | | | |
| Nulliparity, n= (%) | 50 (46.7) | 19 (45.2) | 27 (50.0) | 4 (36.4) |
| Missing, n= | 0 | 0 | 0 | 0 |
| Ethnicity | | | | |
| White, n= (%) | 100 (93.5) | 40 (95.2) | 50 (92.6) | 10 (90.9) |
| Asian, n= (%) | 4 (3.7) | 2 (4.8) | 2 (3.7) | 0 (0.0) |
| Black African, n= (%) | 1 (0.9) | 0 (0.0) | 0 (0.0) | 1 (9.1) |
| Mixed, other, n= (%) | 2 (1.9) | 0 (0.0) | 2 (1.9) | 0 (0.0) |
| Missing, n= | 0 | 0 | 0 | 0 |
| Weight at booking | | | | |
| Mean (kg) +/-SD | 101.7 +/- 20.9 | 101.7 +/- 21.0 | 101.2 +/- 20.7 | 103.6 +/- 23.6 |
| Missing, n= | 1 | 1 | 0 | 0 |
| Booking BMI | | | | |
| Mean (kg/m ²) +/- SD | 36.3 +/- 7.2 | 36.3 +/- 7.4 | 36.1 +/- 7.1 | 37.7 +/- 7.6 |
| Missing, n= | 1 | 1 | 0 | 0 |
| Employment status | | | | |
| Unemployed, n= (%) | 26 (25.0) | 10 (24.4) | 16 (30.2) | 0 (0.0) |
| Missing, n= | 3 | 1 | 1 | 1 |
| Smoking status | | | | |

| | | | | |
|--|-----------|---------|-----------|----------|
| Yes, n= (%) | 19 (17.8) | 4 (9.5) | 10 (18.5) | 5 (45.5) |
| Missing, n= | 0 | 0 | 0 | 0 |
| Diabetic status[‡] | | | | |
| Diabetes, n= (%) | 7 (6.5) | 2 (4.8) | 4 (7.4) | 1 (0.9) |
| Missing, n= | 0 | 0 | 0 | 0 |
| Hypertensive status[§] | | | | |
| Hypertensive, n= (%) | 11 (10.3) | 3 (7.1) | 8 (14.8) | 0 (0.0) |
| Missing, n= | 0 | 0 | 0 | 0 |

*Pregnant women who underwent band deflation pre- or during pregnancy. †Pregnant women who had band inflation maintained throughout pregnancy. ‡Pre-existing diabetes. §Pre-existing hypertensive disease