



Katz, B., Collin, S., Murphy, G., Moss-Morris, R., Bruun Wyller, V., Wensaas, K-A., Hautvast, J., Bleeker-Rovers, C., Vollmer-Conna, U., Buchwald, D., Taylor, R., Little, P., Crawley, E., White, P., & Lloyd, A. (2018). The international collaborative on fatigue following infection (COFFI). *Fatigue: Biomedicine, Health and Behavior*.
<https://doi.org/10.1080/21641846.2018.1426086>

Peer reviewed version

Link to published version (if available):
[10.1080/21641846.2018.1426086](https://doi.org/10.1080/21641846.2018.1426086)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Taylor & Francis at <http://www.tandfonline.com/doi/full/10.1080/21641846.2018.1426086> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/user-guides/explore-bristol-research/ebr-terms/>

Table 1: COFFI Research Questions

Q1	Do different infections trigger different post-infection syndromes?
Q2	What are the predictors of long-term symptoms following infection?
Q3	Are there qualitative differences in fatigue following infection and, if so, what factors predict any such differences?
Q4	Are there predisposing risk factors for post-infection fatigue before or during the febrile phase?
Q5	Are there early or late perpetuating risk factors for post-infection fatigue?
Q6	How do the risk factors and/or perpetuating factors interact and change over time?
Q7	What are therapeutic options for post-infection fatigue?