



Lewis, C., Raisanen, L., Bisson, J. I., Jones, I., & Zammit, S. (2018). Trauma exposure and undetected posttraumatic stress disorder among adults with a mental disorder. *Depression and Anxiety*, 35(2), 178-184. <https://doi.org/10.1002/da.22707>

Peer reviewed version

Link to published version (if available):
[10.1002/da.22707](https://doi.org/10.1002/da.22707)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/abs/10.1002/da.22707> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Number of figures: 0

Number of tables: 3

Trauma Exposure and Undetected Post-Traumatic Stress Disorder (PTSD) among Adults with a Mental Disorder

Catrin Lewis PhD¹; Lawrence Raisanen PhD¹; Jonathan
I. Bisson DM¹; Ian Jones MD¹; Stan Zammit PhD^{1,2}

¹ Institute of Psychological Medicine and Clinical Neuroscience, Cardiff University, Hadyn
Ellis Building, Maindy Road, Cathays, Cardiff, CF11 9QB

² Centre for Academic Mental Health, University of Bristol, School of Social and Community
Medicine, Oakfield House, Oakfield Grove, Bristol, BS8 2BN

Work conducted at the Institute of Psychological Medicine and Clinical Neurosciences, Cardiff

University School of Medicine, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ

Correspondence to: Dr. Catrin Lewis, Hadyn Ellis Building, Maindy Road, Cathays, Cardiff, CF24 4HQ.

+44 7878211716 – LewisCE7@Cardiff.ac.uk

Short title: Trauma and undetected PTSD among adults with a mental disorder

Keywords: Comorbidity; epidemiology; stress disorders, post-traumatic

Disclosures and acknowledgements: SZ is supported by the Bristol NIHR Biomedical Research Centre

Abstract

Background: Trauma exposure and post-traumatic stress disorder (PTSD) are common among individuals with a mental disorder, but symptoms often go undetected and untreated.

Methods: The aim of this study was to determine the prevalence of PTSD among a large sample of adults with psychiatric diagnoses and to establish factors associated with symptoms going undetected. Participants were 1946 adults recruited by the National Centre for Mental Health (NCMH). Structured interviews and validated self-report questionnaires were used to ascertain clinical and demographic information for analysis.

Results: The prevalence of participants screening positive for PTSD that had not been detected by clinical services was 13.9% [12.4%-15.5%, 95% CI]. Factors associated with undetected PTSD were female gender; younger age of first contact with psychiatric services; and lower household income. Especially poor rates of detection were observed after traumatic events such as child abuse and sexual assault.

Conclusions: Our findings demonstrate the need for routine assessment of trauma histories and symptoms of PTSD among individuals with any mental disorder.

Introduction

Epidemiological research suggests that 60% of men and 50% of women experience at least one serious traumatic event through the course of their lives (Kessler, Sonnega et al. 1995). The lifetime prevalence of trauma exposure is higher still among individuals with a mental disorder (Mueser, Goodman et al. 1998, Switzer, Dew et al. 1999, McFarlane, Bookless et al. 2001, Howgego, Owen et al. 2005, Cusack, Grubaugh et al. 2006, Mc Farlane, Schrader et al. 2006, Ford and Fournier 2007). This may be indicative of an aetiological role of trauma in the development of disorders such as schizophrenia and bipolar disorder (Read, Perry et al. 2001, Etain, Henry et al. 2008). It is also a likely consequence of individuals with mental disorders being more at risk of traumatic events such as interpersonal violence and victimisation (Mueser, Rosenberg et al. 2002). Previous studies have reported rates of trauma exposure among individuals with serious mental illnesses that range from 49 to 100% within study samples, including a high prevalence of physical and sexual assault (Grubaugh, Zinzow et al. 2011). The elevated risk of trauma exposure, combined with other vulnerability factors shared between PTSD and other psychiatric disorders results in an increased risk of PTSD for those with other primary mental health diagnoses (Mueser, Rosenberg et al. 2002, Grubaugh, Zinzow et al. 2011). Despite these findings, PTSD often goes undetected and untreated in this population due to symptom overlap and a tendency for clinicians not to fully examine traumatic stress symptoms during clinical assessment (Mueser, Goodman et al. 1998). This represents a significant clinical concern, given that undetected PTSD is associated with higher rates of mental and physical health problems; greater use of general medical and psychiatric services; higher levels of alcohol and substance misuse; and a poorer long-term prognosis (Mueser, Rosenberg et al. 2002, Howgego, Owen et al. 2005, Ford and Fournier 2007, Lommen and Restifo 2009, Grubaugh, Zinzow et al. 2011). Previous studies have reported elevated rates of undetected PTSD among individuals with primary diagnoses such as schizophrenia and bipolar disorder (Mueser, Rosenberg et al. 2002, Howgego, Owen et al. 2005, Ford and Fournier 2007, Lommen and Restifo 2009, Grubaugh, Zinzow et al. 2011), but sample sizes have generally been small, and few studies have sought to establish

factors associated with the disorder going undetected. We used data collected by the National Centre for Mental Health (NCMH) to examine the prevalence of undetected PTSD in a large sample of adults with a history of mental disorder and to establish associated factors that may help predict the likelihood of undetected PTSD.

Method

Data source

Data was obtained from the NCMH, a Welsh Government-funded Research Centre that investigates neurodevelopmental and psychiatric disorders across the lifespan. The Centre is operated by Cardiff, Swansea and Bangor Universities, in partnership with NHS Health Boards across Wales. The cohort of participant volunteers primarily includes individuals who have experienced a mental disorder, but some individuals without such a history have also been recruited into the cohort.

Participants were recruited using a variety of systematic approaches in primary and secondary health care services, including (a) the identification of potential participants by clinical care teams; and (b) screening of clinical notes. Non-systematic recruitment approaches included advertising in local/national media and engaging third-sector organisations to support and promote the research.

All adult participants with sufficient mental capacity provided written informed consent to participate. Trained researchers then administered a standardised interview assessment to ascertain details related to the participant's personal and family history of mental illness. Participants were given a pack of validated self-report questionnaires to complete and return to the research team after the initial assessment.

Sample

The sample for analysis consisted of all adults over the age of 18, who had taken part in a standardised face-to-face interview assessment and completed the Trauma Screening Questionnaire (TSQ) (Brewin, Rose et al. 2002) as part of a questionnaire pack that was posted back to the research team, usually within two weeks of the assessment. The questionnaire pack was given to 3524 out of 4024 adult participants who had sufficient capacity to consent for themselves and complete self-report questionnaires. The questionnaire pack was returned by 2001 (57%) participants, of which 1946 had a diagnosis (current or past) of a mental disorder.

Measures

Demographic and clinical variables

Information was collected by means of a standardised face-to-face interview assessment. This included information on gender; ethnicity; qualifications; employment; household income; alcohol and illegal drug use; diagnoses of mental disorders; age of first contact with mental health services; family history of mental disorder; and the number of admissions to psychiatric hospitals. Participants were asked to self-report if they had been given any psychiatric diagnoses by a health professional. Where uncertainty surrounded a diagnosis, this was verified with the individual's clinical care team if appropriate consent had been given to do so.

Trauma exposure and PTSD

Trauma exposure and PTSD symptoms were assessed using a self-report questionnaire provided at the time of the interview assessment and returned by post. Participants were asked whether they had ever experienced a 'major' traumatic event and, if so, were asked to provide more information. To take a conservative approach to the assessment of PTSD symptoms and to anchor reported traumatic stress symptoms to a single traumatic event, exposure to traumatic events was assessed by means of the free-text question: 'What is the worst traumatic event you have experienced in

your life?’ The descriptions of traumatic events were coded against a list of traumas compiled from the Life Events Checklists for DSM-5 (LEC) (Weathers 2013), with the addition of some additional traumatic events reported by the cohort that were not covered by the LEC (for example, suicide attempts). This resulted in 23 traumatic events (see supplemental table 1) that were categorised into 6 composite categories (assaultive violence; life-threatening illness or injury; witnessing death or life-threatening illness/injury; death of a child/grandchild; other injury or traumatic event; learned-about traumas). DSM-5 non-qualifying events were also coded and these included: non-life-threatening events; death of a partner; death of a parent; death other than parent/partner/child/grandchild; divorce/separation/child custody. When participants reported multiple events, the first listed trauma was taken as the ‘worst’ event. Two researchers independently coded each of the descriptions. If coders disagreed on any of the classifications, this was discussed with a third researcher and a consensus was reached.

All participants who reported a traumatic event completed the TSQ (Brewin, Rose et al. 2002), a valid and reliable tool used to screen for PTSD (Brewin, Rose et al. 2002). The TSQ was used to assess traumatic stress symptoms that had occurred at least twice in the past week. A cut-point off score of 6 was used, i.e. participants were required to self-report the occurrence of at least 6 symptoms occurring at least twice in the past week. Participants were considered to suffer from probable PTSD if they described a DSM-5 criterion-A fulfilling stressor for PTSD and a positive screening score according to the TSQ was obtained. Four groups were created to explore traumatic events and PTSD within the cohort: (1) Those with a lifetime diagnosis of PTSD who screened positive for the disorder (current detected PTSD); (2) those with a lifetime diagnosis of PTSD who did not screen positive for the disorder (past PTSD); (3) those without a lifetime diagnosis of PTSD who screened positive for the disorder (undetected PTSD) and (4) those without a lifetime diagnosis of PTSD who did not screen positive for the disorder (no PTSD).

Statistical procedures

After ascertaining the prevalence of probable undetected PTSD (according to the TSQ) within the cohort, rates were compared according to demographic and clinical characteristics, type of traumatic event, and primary self-reported psychiatric diagnosis. Differences in demographic characteristics were analysed with chi-square tests or one-way ANOVAs, as appropriate, to evaluate whether any differences were associated with undetected PTSD. To explore whether specific types of trauma exposure were more strongly associated with undetected PTSD than others, a chi-square test was conducted to examine the type of 'worst trauma' categorised into groups for those with undetected PTSD compared to rates in the other groups. Finally, to explore whether frequency of undetected-PTSD was higher in certain disorders than others, percentages by group were calculated and differences compared using chi-square.

Results

Trauma exposure

796 (40.9% [38.7% – 43.1%, 95% CI]) participants reported a DSM-5 PTSD qualifying traumatic event. The most commonly reported 'worst' traumas were childhood abuse (7.5%), transportation accidents (4.3%) and sexual assault (4.0%).

PTSD prevalence

438 (23%) participants screened positive for PTSD on the TSQ. 247 (13%) reported a diagnosis of PTSD given to them by a clinician. 169 (9%) both screened positive and reported a PTSD diagnosis. 78 (4%) reported a PTSD diagnosis but did not screen positive for PTSD. 271 individuals screened positive for PTSD but did not report a clinician PTSD diagnosis, i.e. the prevalence of undetected PTSD within the cohort, based on the TSQ, was 13.9% [12.4%-15.5%, 95% CI]. A subgroup analysis explored whether the prevalence was higher among participants who were recruited into the study systematically via the NHS (n = 920) versus those who were recruited non-systematically from the

general population (n=1026). The percentage with undetected PTSD according to the TSQ in the group recruited from the NHS was 15% (n=134) and in the non-systematically recruited group 13% (n=137) (p=0.441).

Demographic and clinical characteristics

The distribution of demographic and clinical characteristics in relation to undetected PTSD compared to detected PTSD (both current and past) is summarised in Table 1. There was strong evidence that undetected PTSD was more prevalent in females than males ($p = 0.003$). Undetected PTSD was not associated with age ($p = 0.210$), but there was strong evidence that those with a younger age of first contact with psychiatric services were more likely to have undetected PTSD ($p = 0.004$). There was no evidence of a difference in the prevalence of undetected PTSD according to lifetime occupational attainment ($p = 0.311$), but undetected PTSD was more common in individuals with a lower family income ($p = 0.018$) and in those with a lower level of educational attainment ($p = 0.007$). We were unable to examine whether prevalence of undetected PTSD varied by ethnicity since over 99% of the sample were categorised as being White Caucasian.

There was no evidence of a difference between detected and undetected PTSD in terms of history of alcohol use disorder ($p=0.844$), substance use disorders ($p=0.141$), number of involuntary admissions to psychiatric hospital ($p = 0.3309$) or family history of PTSD ($p=0.338$).

Insert Table 1 here

Psychiatric diagnoses

There was evidence that the prevalence of undetected PTSD varied across diagnostic groups ($p=0.003$) (Table 2). The prevalence of undetected PTSD was highest among those with a personality disorder (27.8% (19.2-37.9)). The prevalence of both detected and undetected PTSD were high among this diagnostic group. The prevalence of undetected PTSD was also high among those with a primary diagnosis of an anxiety disorder (16.5% (13.8-19.6)), bipolar disorder (14.5% (11.4-18.1)), a

psychotic disorder (16.7% (12.7-21.3)) or a depressive disorder (16.0% (13.8-18.3)). The greatest differences between the prevalence of detected versus undetected PTSD were found for bipolar disorder, psychotic disorders and personality disorders.

Insert Table 2 here

Trauma types

There was a strong relationship between PTSD and trauma grouping ($p < 0.001$) (see table 3). The most common trauma type for both the detected and undetected PTSD groups was assaultive violence, though this represented a higher proportion of the trauma types for the undetected PTSD (45%) compared to the detected PTSD group (36%). 23.25% of the cases of undetected PTSD were accounted for by child abuse (see supplementary table 1). The other specific traumas most strongly associated with undetected PTSD were sexual assault (8.12% of undetected PTSD cases) and domestic abuse (7.75% of cases) (both categorised as assaultive violence) and transportation accidents (7.38% of cases) (categorised as other injury or traumatic event). The other trauma types that were substantially more common in those with undetected PTSD compared to those with detected PTSD included death of a child or grandchild (5.17% vs 1.68), and witnessing an illness or death in someone close (5.90% vs 0.42).

Inset Table 3 here

Discussion

Main findings

Exposure to a PTSD qualifying traumatic event was reported by 40% of participants, and the prevalence of participants screening positive for current PTSD on the TSQ was 22%. These rates are lower than those reported by many previous studies of individuals with mental disorders, and is also slightly lower than many of the population-based studies conducted in the United States (1, 22). This may be reflective of the sample being drawn from the United Kingdom (1, 22). It may also be a consequence of the stringent approach taken to the determination of trauma exposure. Participants were asked whether they had experienced what they perceived to be a 'major' traumatic event and were only required to elaborate if a positive response was given. This may have resulted in underreporting of events that met theoretical criteria for PTSD on the basis that participants did not perceive the event to be 'majorly traumatic'.

Further analysis revealed that 13% of the cohort had probable undetected PTSD (i.e. participants screened positive for PTSD on the TSQ, but did not report a diagnosis of the disorder). This suggests that PTSD is commonly missed by healthcare services. This prevalence is lower than reported by many of the smaller studies to date, which largely focused on complex populations, such as inpatients, veterans and individuals with substance use disorders (Dansky, Roitzsch et al. 1997, Calhoun, Stechuchak et al. 2007, Bonn-Miller, Bucossi et al. 2012). These findings are more in keeping with the smaller number of large-scale studies that included a more representative sample of individuals with a mental disorder (Tagay, Herpertz et al. 2005, de Bont, van den Berg et al. 2015). The estimate of undetected PTSD within the cohort is likely to indicate a more accurate and realistic estimate of the rates of undetected PTSD among a broader population of individuals with a mental disorder. Consistent with a recent systematic review, our findings suggest that PTSD is common among individuals with a mental disorder and that PTSD is often overlooked.

This work went further than previous studies to indicate factors associated with undetected PTSD. It was found that female gender and younger age of first contact with psychiatric services were associated with undetected PTSD.

In terms of the gender difference, males and females within the cohort had a similar prevalence of PTSD, which contrasts with findings from the general population (Kessler, Sonnega et al. 1995), but is consistent with findings from studies of persons with a mental disorder (Grubaugh, Zinzow et al. 2011). Despite comparable rates of PTSD, the disorder was more likely to go undetected in females. This may be due, at least in part, to females reporting a greater proportion of 'worst' traumas such as childhood abuse and sexual assault, which were associated with poor rates of detection. Lower age of first contact with psychiatric services was associated with higher rates of PTSD and lower rates of detection, but this may be indicative of a more severe or complex course of mental disorder, which may in turn increase the likelihood of undetected PTSD.

Also in keeping with findings from the general population, undetected PTSD was associated with poorer socio-economic status, as indicated by lower household income and lower levels of educational attainment. Rates of PTSD and its detection were not associated with the number of involuntary admissions to psychiatric hospital. In terms of trauma type, undetected PTSD was commonly associated with traumas such as childhood abuse and sexual assault. This may be indicative of events that individuals have a reluctance to divulge and that clinicians are wary to ask about. Undetected PTSD was uncommon after combat exposure, perhaps due to more routine screening for PTSD in this population.

Strengths and limitations

To our knowledge, this is the largest study to date to explore factors associated with undetected PTSD within a cohort of individuals with a broad range of mental disorders. Since previous studies have largely been conducted in the US, this research highlights that undetected PTSD is equally a problem in the UK, and hence likely so in other countries. We included a broad spectrum of participants with a range of diagnoses, varying in terms of complexity and severity. This gives a more general estimate of the likely extent of undetected PTSD in psychiatric settings than studies focused

on small samples of participants with specific diagnoses such as substance use disorder. Unlike other studies, we anchored the screening tool to specific traumatic events, which were judged by two researchers independently to determine whether or not they met DSM-5 criteria for a PTSD qualifying event.

There were a number of limitations to our study. Firstly, we did not administer a standardised interview such as the Clinician Administered PTSD Scale (CAPS), although the TSQ has been shown to be a valid and reliable way to screen for the disorder, demonstrating high levels of sensitivity and specificity when using a cut-off score of 6 (Brewin, Rose et al. 2002). However, as the TSQ was anchored to only one traumatic event the impact of multiple traumas was not explored and therefore the extent of symptoms may have been under-reported. It is also possible that some of the symptoms endorsed by participants were features of their primary mental illness as opposed to comorbid PTSD. Secondly, the wide-ranging recruitment strategies used to build the cohort make it difficult to estimate the non-response rate of the study. Thirdly, participants were asked to self-report diagnoses of PTSD by a health professional. It is possible that the true frequency of PTSD diagnoses documented in medical records was underreported. Equally, the study only screened for current PTSD and the rates of undetected past-PTSD are therefore unknown. Studies using more robust methods of ascertaining diagnoses from clinical teams and the addition of a measure of screening for lifetime PTSD would be desirable.

Clinical implications

The high prevalence of participants who screened positive for PTSD on the TSQ but did not report a clinical diagnosis, suggest that traumatic stress symptoms are often overlooked by mental health

services. This is an issue that has been documented and highlighted as clinically problematic for decades, yet rates of PTSD detection remain far from adequate within mental health services. Prevalence of undetected PTSD varied across primary diagnoses and was highest for personality disorders (27.8%). It is surprising that trauma histories are not routinely explored and that PTSD is not screened for, given that individuals with comorbid mental disorders and PTSD are likely to exhibit a poorer prognosis and increased use of general medical and mental health services (Grubaugh, Zinzow et al. 2011). Trauma exposure and post-traumatic stress symptoms may precipitate, maintain or exacerbate whatever mental health issue an individual presents with, and are important to include in biopsychosocial formulations and to inform appropriate management plans.

There are many possible reasons for the high frequency of undetected PTSD. A contributing factor may be that individuals with a mental disorder attribute symptoms to their primary diagnosis and do not seek specific treatment or support for PTSD. It may also be the case that clinicians misdiagnose or find difficulty in disentangling traumatic stress symptoms from the more salient symptoms of the primary mental disorder. For example distinguishing complex PTSD from borderline personality disorder can be difficult as problems with emotion regulation, affective instability, and disturbed self-concept are characteristic of both, whilst frequent flashbacks and re-living during dissociative episodes that can occur in PTSD can be classed as psychotic phenomena if the clinician is unaware of the trauma context in which these experiences arise.

Under-detection may also stem from the previously advocated notion of a hierarchical diagnostic system, whereby diagnoses lower in the hierarchy are not made if higher level diagnoses are indicated (Foulds and Bedford 1975). Additionally, previous studies have reported that clinicians are often reluctant to discuss traumatic events for fear of complicating or exacerbating symptoms (Frueh, Cusack et al. 2006). This is compounded by documented evidence that PTSD diagnosis in

psychiatric settings is positively related to clinician confidence in the ability to effectively treat traumatic stress symptoms (Salyers, Evans et al. 2004).

Although there are effective treatments for PTSD and clear guidelines for its management, the majority of the studies that inform current guidance exclude individuals with comorbidities such as severe depression, psychosis and substance abuse (Bisson, Roberts et al. 2013). However, novel treatments for complex presentations of PTSD have been developed and there is evidence to support the efficacy of these interventions (Cloitre, Courtois et al. 2011). Increasing clinician confidence in their ability to ask about trauma history and offer appropriate evidence-based treatment may reduce the issue of undetected PTSD. This work indicates the continued and pressing need to identify individuals with trauma histories and PTSD and improve patient outcomes. Routine screening seems warranted given that PTSD is a treatable but commonly missed disorder. Validated diagnostic tools exist that enable detection of the disorder easily and quickly.

Research Implications

Future work should seek to disentangle the complex relationship between PTSD and other mental disorders, in order to determine how the effect is mediated and whether it differs cross-diagnostically. Since trauma exposure and undetected PTSD among individuals with a mental disorder are common and remain understudied, it is necessary to determine the extent of the problem amongst different diagnostic groups and across different clinical settings to establish indicators that may help predict those at greatest risk, where screening might be particularly important. Further research using longitudinal designs is needed to unravel the intricate issues of cause and effect; to determine factors underlying comorbidity; and to explore the long-term implications of undetected PTSD. While trauma focused psychological interventions show promise for the treatment of comorbid traumatic stress symptoms, we know less regarding the direct and indirect effects of these interventions on symptoms of depression and psychosis (Grubaugh, Zinzow

et al. 2011). Further work is needed to evaluate, develop and disseminate appropriate treatments for comorbid PTSD. Suboptimal PTSD detection rates within mental health services currently pose a significant barrier to effective intervention and unnecessarily add to the burden of mental disorder.

References

- Bisson, J., N. Roberts, M. Andrew, R. Cooper and C. Lewis (2013). "Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults (Review)." Cochrane Database of Systematic Reviews **12**(CD003388).
- Bonn-Miller, M. O., M. M. Bucossi and J. A. Trafton (2012). "The underdiagnosis of cannabis use disorders and other Axis-I disorders among military veterans within VHA." Military medicine **177**(7): 786-788.
- Breslau, N., G. C. Davis, P. Andreski and E. Peterson (1991). "Traumatic events and posttraumatic stress disorder in an urban population of young adults." Archives of General Psychiatry **48**(3): 216-222.
- Brewin, C. R., S. Rose, B. Andrews, J. Green, P. Tata, C. McEVEDY, S. Turner and E. B. Foa (2002). "Brief screening instrument for post-traumatic stress disorder." The British Journal of Psychiatry **181**(2): 158-162.
- Calhoun, P. S., K. M. Stechuchak, J. L. Strauss, H. B. Bosworth, C. E. Marx and M. I. Butterfield (2007). "Interpersonal trauma, war zone exposure, and posttraumatic stress disorder among veterans with schizophrenia." Schizophrenia Research **91**(1/3): 210-216.
- Cloitre, M., C. A. Courtois, A. Charuvastra, R. Carapezza, B. C. Stolbach and B. L. Green (2011). "Treatment of complex PTSD: Results of the ISTSS expert clinician survey on best practices." Journal of Traumatic Stress **24**(6): 615-627.
- Courtois, C. A. and J. D. Ford (2009). Treating complex traumatic stress disorders: An evidence-based guide, Cambridge Univ Press.
- Cusack, K. J., A. L. Grubaugh, R. G. Knapp and B. C. Frueh (2006). "Unrecognized trauma and PTSD among public mental health consumers with chronic and severe mental illness." Community mental health journal **42**(5): 487-500.
- Dansky, B. S., J. C. Roitzsch, K. T. Brady and M. E. Saladin (1997). "Posttraumatic stress disorder and substance abuse: Use of research in a clinical setting." Journal of Traumatic Stress **10**(1): 141-148.
- de Bont, P. A., D. P. van den Berg, B. M. van der Vleugel, C. de Roos, A. de Jongh, M. van der Gaag and A. van Minnen (2015). "Predictive validity of the Trauma Screening Questionnaire in detecting post-traumatic stress disorder in patients with psychotic disorders." The British Journal of Psychiatry **206**(5): 408-416.
- Etain, B., C. Henry, F. Bellivier, F. Mathieu and M. Leboyer (2008). "Beyond genetics: childhood affective trauma in bipolar disorder." Bipolar disorders **10**(8): 867-876.
- Ford, J. D. and D. Fournier (2007). "Psychological trauma and post-traumatic stress disorder among women in community mental health aftercare following psychiatric intensive care." Journal of Psychiatric Intensive Care **3**(01): 27-34.
- Foulds, G. and A. Bedford (1975). "Hierarchy of classes of personal illness." Psychological Medicine **5**(02): 181-192.
- Frueh, B. C., K. J. Cusack, A. L. Grubaugh, J. A. Sauvageot and C. Wells (2006). "Clinicians' perspectives on cognitive-behavioral treatment for PTSD among persons with severe mental illness." Psychiatric Services **57**(7): 1027-1031.
- Grubaugh, A. L., H. M. Zinzow, L. Paul, L. E. Egede and B. C. Frueh (2011). "Trauma exposure and posttraumatic stress disorder in adults with severe mental illness: A critical review." Clinical psychology review **31**(6): 883-899.
- Howgego, I. M., C. Owen, L. Meldrum, P. Yellowlees, F. Dark and R. Parslow (2005). "Posttraumatic stress disorder: An exploratory study examining rates of trauma and PTSD and its effect on client outcomes in community mental health." BMC psychiatry **5**(1): 21.
- Kessler, R. C., A. Sonnega, E. Bromet, M. Hughes and C. B. Nelson (1995). "Posttraumatic stress disorder in the National Comorbidity Survey." Archives of General Psychiatry **52**(12): 1048-1060.
- Lommen, M. J. and K. Restifo (2009). "Trauma and posttraumatic stress disorder (PTSD) in patients with schizophrenia or schizoaffective disorder." Community mental health journal **45**(6): 485-496.

McFarlane, A., G. Schrader, C. Bookless and D. Browne (2006). "Prevalence of victimization, posttraumatic stress disorder and violent behaviour in the seriously mentally ill." Australian and New Zealand journal of psychiatry **40**(11-12): 1010-1015.

McFarlane, A. C., C. Bookless and T. Air (2001). "Posttraumatic stress disorder in a general psychiatric inpatient population." Journal of traumatic stress **14**(4): 633-645.

McManus, S., H. Meltzer, T. Brugha, P. Bebbington and R. Jenkins (2009). "Adult psychiatric morbidity in England, 2007: results of a household survey."

Mueser, K. T., L. B. Goodman, S. L. Trumbetta, S. D. Rosenberg, F. C. Osher, R. Vidaver, P. Auciello and D. W. Foy (1998). "Trauma and Posttraumatic Stress Disorder in Severe Mental Illness." Journal of Consulting and Clinical Psychology **66**(3): 493.

Mueser, K. T., S. D. Rosenberg, L. A. Goodman and S. L. Trumbetta (2002). "Trauma, PTSD, and the course of severe mental illness: an interactive model." Schizophrenia research **53**(1): 123-143.

Read, J., B. D. Perry, A. Moskowitz and J. Connolly (2001). "The contribution of early traumatic events to schizophrenia in some patients: a traumagenic neurodevelopmental model." Psychiatry **64**(4): 319-345.

Salyers, M. P., L. J. Evans, G. R. Bond and P. S. Meyer (2004). "Barriers to assessment and treatment of posttraumatic stress disorder and other trauma-related problems in people with severe mental illness: Clinician perspectives." Community Mental Health Journal **40**(1): 17-31.

Switzer, G. E., M. A. Dew, K. Thompson, J. M. Goycoolea, T. Derricott and S. D. Mullins (1999). "Posttraumatic stress disorder and service utilization among urban mental health center clients." Journal of traumatic stress **12**(1): 25-39.

Tagay, S., S. Herpertz, M. Langkafel and W. Senf (2005). "Posttraumatic stress disorder in a psychosomatic outpatient clinic: Gender effects, psychosocial functioning, sense of coherence, and service utilization." Journal of Psychosomatic Research **58**(5): 439-446.

Weathers, F. W., Blake, D.D., Schnurr, P.P., Kaloupek, D.G., Marx, B.P., & Keane, T.M. (2013). "The Clinician-Administered PTSD Scale for DSM-5 (CAPS-5). Interview available from the National Center for PTSD at www.ptsd.va.gov."

Weathers, F. W., Blake, D.D., Schnurr, P.P., Kaloupek, D.G., Marx, B.P., & Keane, T.M. . (2013). " The Life Events Checklist for DSM-5 (LEC-5)."

Tables

Table 1: Frequency (percentage) or mean (range) of demographic and clinical characteristics in individuals with undetected PTSD compared to those with self-reported clinician diagnosis (past or current) PTSD

Variable	Detected PTSD (current and past) (N = 247)		Undetected PTSD (N = 271)		P value
	N (%) or mean (range)		N	%	
Gender					
Male	112	(45.3%)	90	32.1%	0.002
Female	135	54.7%	190	67.9%	
Age	48.1 [46.5-49.7]		49.7 [48.0-51.4]		0.023
Age of first contact with psychiatric services	32.85 [31.0-34.7]		28.96 [27.1-30.8]		0.004
Highest level of qualification					
None/less than equivalent to GCSE	38	15.8%	82	30.9%	0.007
GCSE or equivalent	85	35.3%	72	27.2%	
A level or equivalent	62	25.7%	66	24.9%	
Degree or above	56	23.2%	45	17.0%	
Professional occupation (lifetime)					
Yes	63	25.9%	59	22.1%	0.311
No	180	74.1%	208	77.9%	
Approximate gross household income					
Up to £10,000	69	32.2%	93	42.5%	0.018
£10,000 - £20,000	63	29.4%	69	31.5%	
£20,000 - £30,000	34	15.9%	20	9.1%	
Over £30,000	48	22.4%	37	16.9%	
Family history of PTSD					
Yes	24	10.5%	21	8.0%	0.338
No	205	89.5%	242	92.0%	
History of alcohol use disorder					
Yes	34	13.9%	36	13.3%	0.844
No	211	86.1%	235	86.7%	
History of drug use disorder					
Yes	12	4.9%	22	8.1%	0.141

No	233	95.1%	249	91.9%	
Number of involuntary admissions to psychiatric hospital	1.37 (0 - 48)	1.81 (0 - 72)			0.331

Table 2: Frequency and percentage (95% confidence intervals) of detected or undetected PTSD within each diagnostic category

Primary Diagnosis	Detected PTSD (current and past)		Undetected PTSD		Total	P value
	N	%	N	%	N	
Anxiety Disorder	71	10.8% (8.5–13.4)	109	16.5%(13.8-19.6)	180	0.003
Autistic Spectrum Disorders and Learning Disabilities	3	5.3% (1.1-14.6)	5	8.8% (2.9-19.3)	8	
Bipolar Disorder	22	4.9% (3.1-7.3)	65	14.5% (11.4-18.1)	87	
Schizophrenia and other Psychotic Disorders	15	4.8% (2.7-7.8)	52	16.7% (12.7-21.3)	67	
Depressive Disorder	134	12.6% (10.7-14.8)	170	16.0% (13.8-18.3)	304	
Personality Disorder	15	15.5% (8.9-24.2)	27	27.8% (19.2-37.9)	42	
Other	8	6.8% (3.0-12.9)	19	16.1% (10.0-24.0)	27	

Table 3: Frequency and percentage of types of traumatic event within detected or undetected PTSD groups

	Detected PTSD		Undetected PTSD		Significance
Traumatic event	Freq	%	Freq	%	
Assaultive violence	89	36.0%	122	45.0%	<i>p</i> < 0.001
Other injury or traumatic event	71	28.7%	53	19.6%	
Witnessing death or illness	13	5.3%	28	10.3%	
Death or child/ grandchild	7	2.8%	23	8.5%	
Life threatening illness or injury	9	3.6%	14	5.2%	
Learned-about traumas	8	3.2%	31	11.4%	
DSM-5 non-qualifying events	22	8.9%	0	0	
Not divulged/missing	28	11.2%	0	0	

Supplemental material

Supplementary table 1: Frequency and percentage of types of traumatic event within detected or undetected PTSD groups

Traumatic event	Detected PTSD		Undetected PTSD	
	Freq	%	Freq	%
Assaultive violence				
Physical assault/threatened physical assault	16	6.72	16	5.90
Sexual assault or other unwanted sexual contact	25	10.50	22	8.12
Child abuse	39	16.39	63	23.25
Domestic abuse	9	3.78	21	7.75
Other injury or traumatic event				
Natural or manmade disaster/terrorism	2	0.84	4	1.48
Other serious accident ^a	8	3.36	5	1.85
Transportation accident/incident	23	9.66	20	7.38
Traumatic childbirth	2	0.84	3	1.11
Causing death or injury to another	1	0.42	0	0.00
Combat exposure/exposure to a war zone	32	13.45	3	1.11
Other life threatening traumatic events ^b	2	0.84	3	1.11
Suicide attempt	1	0.42	15	5.54
Witnessing death or illness				
Witnessed violent or unexpected death	7	2.94	4	1.48
Witnessed illness and death of someone close	1	0.42	16	5.90
Finding a loved one dead	2	0.84	3	1.11
Witnessing life threatening injury/accident/medical	2	0.84	3	1.11
Witnessing violence/attack	1	0.42	1	0.37
Death of child/ grandchild				
Death of child or grandchild	4	1.68	14	5.17
Miscarriage/ still born/ death of new born	3	1.26	9	3.32
Life threatening illness or injury				
Life threatening illness or injury	9	3.78	14	5.17
Learned about traumas				
Learned about: sexual assault	1	0.42	4	1.48
Learned about: transportation accident/other accident	1	0.42	9	3.32
Learned about: violent or accidental death of someone close	0	0	0	0
DSM V non-qualifying events				
Mental illness/breakdown/inpatient experiences/psychotic	2	0.84	0	0.00
Learned about: life threatening illness or injury/suicide attempt	2	0.84	0	0.00
Other non-life threatening traumatic events	7	2.94	0	0.00
Death of parent	2	0.84	0	0.00
Death other than parent/partner/child/grandchild or an	7	2.94	0	0.00
Divorce/ separation/ custody	2	0.84	0	0.00