



Skreden, M., Hillesund, E. R., Wills, A. K., Brantsæter, A. L., Bere, E., & Øverby, N. C. (2018). Adherence to the New Nordic Diet during pregnancy and subsequent maternal weight development: a study conducted in the Norwegian Mother and Child Cohort Study (MoBa). *British Journal of Nutrition*, 119(11), 1286-1294.  
<https://doi.org/10.1017/S0007114518000776>

Peer reviewed version

Link to published version (if available):  
[10.1017/S0007114518000776](https://doi.org/10.1017/S0007114518000776)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge University Press at <https://doi.org/10.1017/S0007114518000776> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>




Figure 2: Predicted marginal mean BMI trajectory for each category of NND adherence score. The plot on the left is from the unadjusted model; the middle plot on the confounder adjusted model (a)\*; and the right plot the confounder plus mediator model (b)\*. For clarity, only the 95% CI for NND tertile 1 (lowest) is shown (grey).

\*Model a: adjusted for maternal age, parity, education, pre-pregnant smoking, physical activity, energy intake and offspring birthweight  
 Model b: model a + exclusive breast feeding duration.